

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
http://ageconsearch.umn.edu
aesearch@umn.edu

Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.

Economic and Environmental Possibilities of Sugar Beet In Spain: Towards Bio-ethanol Production?

Authors

Pedro Pablo Perez Hernandez. Faculty of Management and Economics Sciences, Loyola University of Andalusia (Spain). Escritor Castilla Aguayo, 4, Cordoba, 14004. Spain

Phone Number: 0034-957-222110 Fax Number: 0034-957-222101 Email: ppperez@etea.com

Melania Salazar Ordoñez. General Economy. Faculty of Management and Economics Sciences, Loyola University of Andalusia (Spain). Escritor Castilla Aguayo, 4. Cordoba,

14004, Spain

Phone Number: 0034-957-222100

Email: msalazar@etea.com

José Manuel Martin Lozano. Management and Quantitative Methods. Faculty of Management and Economics Sciences, Loyola University of Andalusia (Spain). Escritor

Castilla Aguayo, 4. Cordoba, Spain **Phone Number:** 0034-957-222100

Email: jmmartin@etea.com

Selected Poster prepared for presentation at the International Association of Agricultural Economists (IAAE) Triennial Conference, Foz do Iguaçu, Brazil, 18-24 August, 2012.

Copyright 2012 by Pérez, P.P, Salazar, M. & Martin, J.M. All rights reserved. Readers may make verbatim copies of this document for non-commercial purposes by any means, provided that this copyright notice appears on all such copies.

Economic and Environmental Possibilities of Sugar Beet In Spain: Towards Bio-ethanol Production?

Pedro Pablo Pérez Hernández[†], Melania Salazar Ordoñez², José Manuel Martín Lozano³

[†]Dpt. Applied Economics Universidad Loyola Andalucía, Cordoba (Spain)

E-mail: ppperez@etea.com¹, msalazar@etea.com² jmmartin@etea.com³

100 | 40.43 | 40.43 | 10.72 | 8.41 | 100.0 | 0.0

ntroduction

LOYOLA

SAMPLE

Since 2006 sugar beet in Europe has suffered a deep material for sugar production, being a familiar farming. In that context, the possible survival of great part of this farming is strategy. We have made an empirical approach to learn about future viability of the sector with environmental efficiency analysis (Greenhouse Gases emissions) caused by sugar beet farms with the aim of studying environmental non-market goods derived from bio-ethanol production. Goods that take part of the MDB= Greenhouse gases reduction by bio-ethanol use in CO₂ t beet equivalents; environmental function of this economic activity as a positive MDP= nitrous oxides emissions (N2O) by fertilizers use in CO2 t N/kg equivalents externality.

Sugar beet farming in Spain: the survey

THE QUESTIONNAIRE: three basic parts: a) owner identification and farms general characteristics; b) sugar beet farms expenses and c) income linked to sugar beet farming. SAMPLE DESIGN: was based on five basic principles: a) Sugar beet farming surface and territorial distribution, b) Farms size, c) Production economic value, d) Minimum number of individual observations, Geographical distribution. We for using opted 1998). They were proportional sample (Festinger, analyzed two productive regions with a level of significance and representativeness: Andalusia and Castilla-León.

CHARACTERIZATION:

farming/farm; dedicating an average of 27.5% of its surface to sugar beet farming; 77% of farms sugar beet farming is less than 40% of the total agricultural area and only 13.25% of them dedicate the whole area to this farming. Total expense per ha in Castilla-León is significantly less than in Andalusia (Andalusia salary expenses per ha are higher). North farm dimensions are smaller and familiar than South farm. In Castilla-León productivity level is higher (27%) in the north than in the south because varieties planted in the north have a higher average productivity per ha. So sugar beet farming in the north present a better economic situation. EXPERTS' OPINION: three Focus Groups. concluded that sugar beet future is conditioned and, especially from an energetic point of view, only suitable varieties that could reduce 10% of expenses and

achieve a significant increase of yield (to 150 t/ha).

Environmental efficiency analysis: DEA

Data Envelopment Analysis (DEA) is a mathematical programming measure technique which assesses the relative efficiency of comparable decision-making units -producers- or DMUs. A transformation. In Spain, sugar beet has been the main raw DMU is efficient if and only if the performances of other DMUs do not improve the input allocation or output production without worsening some others inputs or outputs (Cooper et a 2004). In this research, DEA is used to model the environmental efficiency from GG emissions they have zero gaps (s^{-*}=0 y s^{+*} =0). Efficiency media index has been 90.55% derived to bio-ethanol elaboration in according to the new EU of using sugar beet to bio-ethanol production in Spain, corrected by means of the nitrogen fertilizer emissions in cultivation (Piot-lepetit et al., 1997, and Zhang, 2008). Three bad inputs are exclusively considered.

INPUTS AND OUTPUTS: The optional input variables are the following: SUP= Hectares surface; SYP= Seeds and seedlings annual costs; FER=Fertilizers annual costs; FIT= Phytosanitary annual costs. And the outputs were the following:

| Woods y Bauen, 2003 | 45-80

different

Studies on GG emissions by substituting petrol for bio-ethanol										
Research	Crop (t/ha)	· Ineriarmance Kia-einanai (I/I raw		Emissions saving (CO ₂ t/ha equivalent) Mixtures E10	% Km saving Mixtures s E10					
Lovy 1002	66	5.28	101.3	6,600	3.54	35%				
Levy, 1993	78	6.24	101.3	7,800	7.22	56%				
European Commission, 1994	87	3.75	54.1		2.12	50%				
GM et al., 2002					6.50	41%				
Rajagopal y Zilberman 2007	100		110	7,370						

Conclusions

4,800-7,800

0-70% (2)


As a consequence of CAP application, certain beet producers have been expelled from the market, being practically limited to two zones in Spain: Andalusia and Castilla-León. In this context, farmers, professional associations and Spanish research centres for beet cultivation have made a great effort which has led to an increase on crop yield per ha obtained from the I+D+i investment with important varietal and technological improvements. Three are the main expenditures: seeds, fertilizers and phytosanitary. Average expenditure per hectare in the North zone, Castilla-León, is higher than in the South zone, Andalusia.


On the other side, energetic farming support can be granted whenever it is destined not to food production. Besides, this support can be joined to environmental goods provision because sugar beet production destined to ethanol production helps greenhouse gases reduction. From GG emissions point of view and considering variable scale yields, the 22% of farms have been efficient and those which are inefficient need to reduce fertilizers and phytosanitary use to improve nitrous oxide emissions to atmosphere.


Larger farms could seem to be more efficient in that aspect. However, small and medium size farms are the most efficient ones. Andalusia farms present a higher GG emission saving. Sugar beet consumption is situated over European production and surface destined to Bioenergetics farming has been considerably reduced since 2008. Future trend seems to be directed to import any type of crop to be used as raw material suitable for bio-ethanol production. Therefore, there is the possibility of disappearance of sugar beet production in the European agrarian model.


RESULTS

In the estimated model, there are 66 efficient farms from the GG emissions point of view which represents the 22% of samples. So, these DMU represent an efficiency of 100% (θ *=1) and they do not have to make improvements as and the dispersion, measured through the typical deviation is 7.6, situating most farms index over 85% while 14% of the farms have efficiencies lower than 80% We should emphasize that only two are situated under 60%. Distribution of efficiency values is shown in Figure 1









			Example						
	DMUs	Ha.	Eficiency		Inpu	ts weight		Outputs	s wei
			(%)	W _{Area}	W _{Seed}	W _{Phytosanit}	W _{Fertilizer}	$\mathbf{w}_{\mathbf{GG}}$	\mathbf{W}_{I}
	46	3.0	100	74.31	18.22	0.0	7.46	58.61	41
	93	2.0	100	94.52	5.47	0.0	0.0	15.44	84
	253	2.5	100	32.24	67.75	0.0	0.0	29.10	70
	254	12.0	100	79.84	1.43	12.94	5.77	67.54	32
	92	20.0	100	96.42	3.57	0.0	0.0	72.51	27
	7	40.0	100	97.4	0.0	0.0	2.57	89.74	10

Examples of inefficient farms

DMUs	DMUs Ha. Efficiency (%) 201 25.0 48.33		$\mathbf{w_{Area}}$	$\mathbf{w}_{\mathbf{Seed}}$	W _{Phytosanit}	W _{Fertilizer}	$\mathbf{w_{GG}}$	$\mathbf{w_{N2O}}$	
201			75.4	16.9	0.0	76.0	100.0	0.0	
DMUs Reference				S _{Area} (%)	S _{Seed} (%)	S _{Phytosanit} (%)	S _{Fertilizer} (%)	S _{GG} (%)	S _{N2O} (%)
195	195 207 239 243		0.0	0.0	-49.8	0.0	106.9	184.3	

DIVIUS	11a.	Efficiency (70)	** Area	** Seed	Phytosanit	** Fertilizer	w GG	** N2O
204	30.0 50.99		0.0	97.5	2.4	0.0	100	0.0
DM	IUs Ref	ference	S _{Area} (%)	S _{Seed} (%)	SPhytosanit (%)	S _{Fertilizer} (%)	S _{GG} (%)	S _{N2O} (%)
195	207	239	-3.1	0.0	0.0	-40.2	96.2	104.4
DMUs	На.	Efficiency (%)	W _{Area}	$\mathbf{w}_{\mathbf{Seed}}$	W _{Phytosanit}	W _{Fertilizer}	$\mathbf{w}_{\mathbf{GG}}$	$\mathbf{w_{N2O}}$

						Alta ('')	"Beeu (' ')	- I ny tosamt ()	Terunzer (**)	1 00 (11)	1120 (11)
	66	73	3 1	11	268	0.0	-14.7	0.0	-42.1	65.3	64.3
e	DMUs		MUs Ha.		acy (%)	$\mathbf{W_{Area}}$	W _{Seed}	WPhytosanit	W _{Fertilizer}	$\mathbf{w}_{\mathbf{GG}}$	W _{N2O}
	187 14.0		14.0	64.8		0.0	97.5	2.4	0.0	100	0.0
		DMUs Reference		S _{Area} (%)	s _{Seed} (%)	S _{Phytosanit} (%)	S _{Fertilizer} (%)	S _{GG} (%)	S _{N2O} (%)		
	55	185	195	207	246	0.0	0.0	0.0	-32.6	56.0	54.1

SArea (%) | Seed (%) | Sephytosopit (%) | Septilizer (%)

References

DMUs Reference

ADEME (2010). Life Cycle Analyses Applied to First Generation Bio-fuels Used in France. Major insights and learnings. Agence de l'environnement, Maîtrise de l'Energie, the Ministère de l'Ecologie, Energie, du Développement Durable et de la Mer, the Ministère de l'Alimentation, de l'Agriculture et de la Pêche, y FranceAgriMer (FAM). Paris.

Cooper, W. W.; Seidford, L. M.; Zhu, J. (2004). Handbook on Data Envelopment Analysis. Nueva York: Springer.

Festinger L.; Katz, L. (1998). Los métodos de investigación en las ciencias sociales. Barcelona: Hurape.

IPCC (2006). IPCC Guidelines for National Greenhouse Gas Inventories. Volume 4 Agriculture, Forestry and Other Land Use. Intergovernmental Panel on Climate Change. Ginebra.

Levy, R. H. (1993). Les biocarburants. Commission Consultative pour la production des Carburants de Substitution. París.

Piot-lepeti, I., Vermersch, D. and Weaver, R. D. (1997). Agriculture's environmental externalities: DEA evidence for French agriculture, Applied Economics, 29, 331–338

Rajapogal, D.; Zilberman, D. (2007). Review of environmental, economic and policy aspects of bio-fuels. Policy Research Working Paper nº 4341. Sustainable Rural and Urban Development Division, Development Economics Research Group Department, Banco Mundial. Nueva York.

Woods, H.; Bauen, A. (2003). Technology status and carbon abatement potential of renewable transport fuel in U. K. B/U2/00785REP URN 03/982. Imperial College of London and Centre for Energy Policy and Technology. Londres.

Zhang, T. (2008): Environmental performance in China's agricultural sector: a case study in corn production, Applied Economics Letters, 15:8, 641-645