AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

Modeling Perennial Crop Supply: An Illustration from the Pecan Industry

Abdelmoneim H. Elnagheeb and Wojciech J. Florkowski*

Abstract

Two methodological approaches were applied to estimating the number of non-bearing trees in the absence of such data using data for the Southern USA pecan industry. The first approach distinguished between bearing and non-bearing phases of a tree life and directly estimated the number of non-bearing trees. The second focused on indirect estimating of the non-bearing tree number from changes in production. This approach relaxed the assumption of maintaining maximum yields for infinite period as used in earlier studies. Empirical applications used two data sets from the pecan industry. The comparison of empirical results suggested that the first method was more accurate than the alternative approach in predicting the number of newly planted trees over an extended period of time. Additional data collection will allow for further application of available methodology to the pecan industry.

Key words: non-bearing trees, pecans, tree crop, tree yields, volume produced.

Introduction

The literature on perennial crop supply has developed over the last two decades. The first attempt to model perennial crop supply was French's study of the Michigan and U.S.A. apple production. Later, French and Bressler developed a supply response model for lemons specifying planting and removal relationships. French and Matthews developed a structural multi-equation model for the supply response of asparagus. The equations were subsequently combined to give a reduced-form, single equation for output. Wickens and Greenfield modeled the investment and harvest decisions separately for the Brazilian coffee. However, the separate equations were combined to give a reduced-form, single-output equation which used a lagged-variable technique for estimating the model.

Some researchers advocate the structural approach for estimating separate equations for planting, removal, and harvesting (Akiyama and Trivedi; Bellman and Hartley; French, King, and Minami; and Hartley, Nerlove, and Peters). However, a common difficulty in estimating a structural model for perennial crops is data availability. For many perennial crops, data on the number (or area) of new plantings, tree age-distribution, and removals are not available. Because of data limitations the reduced-form, single-equation approach has dominated perennial crop-supply studies.

This paper utilizes and compares two alternative approaches that mitigate the paucity of data on new plantings. The two models, which are described below, were estimated using data from Georgia's pecan industry. Pecans are an important

*A. H. Elnagheeb is a postdoctoral associate and W. J. Florkowski is an associate professor. Both are at the department of agricultural and applied economics, The University of Georgia College of Agricultural and Environmental Sciences, Griffin, GA.

source of income for many farmers in the Southeast. Georgia produces about 40 percent of the annual total pecan output in the U.S. Hence, it is important to understand how pecan supply responds to different policies. Previous pecan studies focused on pecan quality at the retail level (Williams, et al.), predicting pecan prices (Epperson and Allison), estimating pecan price flexibilities (Wells, et al.), evaluating the impact of pecan crop forecasts on pecan prices and value (Shafer) and differences in pecan prices by variety (Okunade and Cochran). This study expands the studies of the pecan industry by providing estimates of the pecan supply response.

The Model

Previous studies on perennial crop supply faced the problem of lack of data on new plantings and tree age-distribution. To overcome this problem, researchers modeled the unobserved variables as functions of observable variables (e.g. expected prices, costs, and output). These equations were substituted into other structural equations to obtain a single-equation, reduced-form model (Bateman; Behrman; French and Matthews; and Wickens and Greenfield).

The current study faces the similar data problem for new pecan plantings and the age distribution of trees. Two models were constructed. Estimates were compared and the performance of each model evaluated. In both models, the number of newly planted trees at time t , I_t , is assumed to be a linear function of lagged prices, P_{t-1} , and costs, C_t :

$$(1) \quad I_t = \alpha_0 + \alpha_1 P_{t-1} + \alpha_2 C_t$$

The first model uses a study by French and Matthews as its basis. The French and Matthews' (FM) model starts with the identity:

$$(2) \quad B_t = B_{t-1} + aN_{t-k} + R_{t-1}$$

where B_t is the total bearing acreage, N_t is new plantings in acres, R_{t-1} is bearing acreage removed at $t-1$, a is a constant with a value slightly less than

one to account for plantings removed before reaching bearing age, and k is the number of years necessary for a tree to reach a bearing age.

The FM model formulates desired new plantings as a function of expected long-run profits and a disturbance term. Such formulation reduces the problem of limited data availability on new plantings and removals. A function accounting for removals included a short-run profit expectations measure, a measure of acreage exceeding the age at which productivity typically begins a significant decline, and a disturbance term. New plantings and removal functions replaced explicit measures in identity (2). Because nonlinear forms for the new planting and removal functions will yield an equation difficult to estimate empirically, the FM model assumes linear functional forms. French and Willett expressed all variables, including B_t and B_{t-1} , in the log form as an approximation for an unknown nonlinear relationship.

The approach proposed in this study is similar to the FM's approach. Pecan tree life encompasses bearing and non-bearing phases. A study by Gorman et al. shows pecan trees as bearing at the age of six years. Thus, the following relationship was formulated:

$$(3) \quad NB_t = \sum_{i=0}^5 I_{t-i} + e_t$$

where NB_t is the number of non-bearing trees at time t , I_t is the number of trees planted at time t , and e_t is an independently identically distributed error term with zero mean and a constant variance σ^2 . The error term, e_t , can be construed as the error in measurement. Equation (3) states that at time t , the group of non-bearing trees consists of trees from zero to five years old. It was assumed that no removal occurs in the non-bearing group. The removal could be incorporated into equation (3) by multiplying the I_{t-i} by a , where a is slightly less than one, as used in the FM's model. However, a will be unidentifiable when the model derived below is estimated. In the case of pecans, the removal of trees occurs seldom and therefore a should be very close to one.

Substituting equation (1) into equation (3) and rewriting, we obtain:

$$(4) \quad NB_t = 6\alpha_0 + \alpha_1 P_t^* + \alpha_2 C_t^* + e_t$$

where,

$$(5) \quad \begin{aligned} P_t^* &= \sum_{i=0}^5 P_{t-1-i} \\ C_t^* &= \sum_{i=0}^5 C_{t-i} \end{aligned}$$

Equation (4) states that the number of non-bearing trees is influenced by past pecan prices and costs. Equation (4) can be estimated by ordinary least squares or generalized least squares, if the error terms are not homoscedastic, yielding consistent and unbiased estimates of the unknown parameters.

The second approach benefits from the models developed by Bateman; Behrman; Baritelle and Price; and Wickens and Greenfield. In those studies, authors measured output as the summation of the product of different bearing age tree groups (or acres) and the yield per bearing tree (acre)

$$(6) \quad Q_t^* = \sum_{i=0}^n y_{it} I_{t-i}$$

where Q_t^* is the total potential output, y_{it} is the yield per tree (or per acre) for trees in age group i , and I_{t-i} is the number of trees (or acres) in age group i at time t . Baritelle and Price used I_{t-i} as the number of trees, while Bateman used I_{t-i} to denote the number of bearing acres. More specifically, Bateman's model (p. 389) assumes that n equals infinity and yield is represented by a two-step process as described by a yield function:

$$(7) \quad Q_t^* = b_1 \left(\sum_{i=k}^{s-1} I_{t-i} \right) + b_2 \left(\sum_{i=s}^{\infty} I_{t-i} \right)$$

where Q_t^* is as defined above, b_1 the output per acre attained after the first increase in yield, b_2 the output per acre after reaching the plateau, k is the age at which trees begin to bear, and s is the year in which the second distinct increase in yield occurs. The assumption made by Bateman is that plants maintain peak yields for an infinite length of time.

Because the quantity harvested may differ from that produced, Bateman allowed for the effects of economic and climatic factors, by incorporating producer's price (P_t) and climatic variables, W_t (e.g., rainfall and humidity) into equation (7):

$$(8) \quad Q_t = b_1 \left(\sum_{i=k}^{s-1} I_{t-i} \right) + b_2 \left(\sum_{i=s}^{\infty} I_{t-i} \right) + c'W_t + dP_t$$

where Q_t is the amount harvested at time t . A problem with equation (8) is the infinite horizon which presents estimation difficulties. To overcome this problem, Bateman took the first difference of equation (8) to obtain:

$$(9) \quad \Delta Q_t = b_1 I_{t-k} + (b_2 - b_1) I_{t-s} + c' \Delta W_t + d \Delta P_t$$

where Δ indicates a change in the respective variable from one period to the next. Since data on newly planted acreage were not available, Bateman expressed I_t as a function of prices and costs and substituted that equation into equation (9). In this study, I_t represents the number of trees rather than the number of acres.

In this paper, equation (8) is modified to:

$$(10) \quad Q_t = \gamma_0 + \gamma_1 Q_t^* + \sum_{i=1}^m \beta_i P_{t-i} + \beta_{m+1} C_t + \mu'W_t + \phi Q_{t-1} + u_t$$

where Q_t , P_t , C_t , and W_t are as defined before; Q_t^* is the potential total output as given by equation (6); u_t is independently identically distributed error term with zero mean and constant variance, σ_u^2 . The differences between equations (8) and (10) include the addition of the term ϕQ_{t-1} into equation (10) and the scaling of Q_t^* term with the parameter γ_1 which was set equal to one in equation (8). The lagged dependent variable in equation (10) was added to capture the alternate bearing effect which characterizes many perennial crops.

Taking the first difference of equation (10), yields:

$$(11) \quad \Delta Q_t = \gamma_1 \Delta Q_t^* + \sum_{i=1}^m \beta_i \Delta P_{t-i} + \beta_{m+1} \Delta C_t + \mu' \Delta W_t + \phi \Delta Q_{t-1} + \Delta u_t$$

where

$$(12) \quad \Delta Q_t^* = y_k I_{t-k} + \sum_{i=k+1}^n (y_i - y_{i-1}) I_{t-i}$$

k is the age when a pecan tree starts bearing, and n is the age of the oldest tree (or tree group) at time t . Equation (12) showed that Bateman's assumption about maintaining yield for an infinite period of time (as in equation 7) was unnecessary.

Pecan trees start bearing significant amount at about six years of age and thereafter yields increase with age. However, the increase in yield, $(\mu_i = y_i - y_{i-1})$, becomes insignificant after about 13 years (see Gorman et al.). Hence, the summation in equation (12) covers the period from 7 to 13 years of age.

Substituting equation (1) into (12) and back into equation (11), we obtain:

$$(13) \quad \Delta Q_t = \alpha_0^* + \alpha_1^* \hat{P}_t + \alpha_2^* \hat{C}_t + \sum_{i=1}^m \beta_i \Delta P_{t-i} + \beta_{m+1} \Delta C_t + \mu' \Delta W_t + \phi \Delta Q_{t-1} + \Delta u_t$$

where

$$\begin{aligned} \alpha_0^* &= \gamma_1 \alpha_0 (y_6 + \sum_{i=7}^{13} \mu_i) \\ \alpha_i^* &= \gamma_1 \alpha_i, \quad i=1,2 \\ (14) \quad \hat{P}_t &= y_6 P_{t-7} + \sum_{i=7}^{13} \mu_i P_{t-1-i} \\ \hat{C}_t &= y_6 C_{t-6} + \sum_{i=7}^{13} \mu_i C_{t-i} \end{aligned}$$

Data on yield, y_t , were obtained from Gorman, et al. and were used to calculate the transformations in equations (14).

Estimation Procedures

Equations (4) and (13) represent the reduced forms of the two models to be estimated. For the model in equation (4), data on non-bearing pecan trees are available every five years. That is, the index on NB in equation (5) is $t+i$ where $i = 0, 5, 10, 15, \dots$. Because the error term, e_t , was assumed to be independently and identically distributed (iid), equation (4) can be estimated with ordinary least squares (OLS). However, initial results indicated the presence of serial correlation and the model was re-estimated using Prais-Winsten's method to correct for serial correlation.

For ease of exposition, the model in equation (13) is rewritten as:

$$(15) \quad M_t = \alpha_0^* + X_t' \pi + \phi M_{t-1} + \varepsilon_t$$

where $M_t = Q_t - Q_{t-1}$, $\varepsilon_t = u_t - u_{t-1}$, and X_t contains all other explanatory variables in equation (13) and π is a conformably defined vector of parameters. Applying OLS to equation (15) encounters two problems. First, the explanatory variables contain a lagged dependent variable. The OLS estimates are consistent but may suffer from small sample bias (Johnston). The second problem is the serial correlation of the error terms which will result in inconsistent OLS estimates (Johnston). Although the u_t 's were assumed iid, the error terms, ε_t 's, are not:

$$(16) \quad E[\varepsilon_t \varepsilon_{t+s}] = \begin{matrix} 2\sigma_u^2 & s = 0 \\ -\sigma_u^2 & s = \pm 1 \\ 0 & |s| \geq 2 \end{matrix}$$

For estimation purposes, the model in equation (15) is rewritten as:

$$(17) \quad W_t = \alpha_0^* + X_t' \pi + \phi M_{t-1} + W_{t-1}$$

where $W_t = Q_t - u_t$. Repeated substitution for W_{t-1} into equation (17) allows us to write equation (17) as:

$$(18) \quad W_t = \alpha_0^* \sum_{i=1}^t 1 + \sum_{i=1}^t X_i' \pi + \phi \sum_{i=0}^{t-1} M_i + W_0$$

Substituting for $W_t = Q_t - u_t$ and $\sum_{i=0}^{t-1} M_i = \sum_{i=0}^{t-1} \{Q_i - Q_{i-1}\} = Q_{t-1} - Q_{-1}$, leads to:

$$(19) \quad Q_t = \alpha_0^* t + \sum_{i=1}^t X_i' \pi + \phi Q_{t-1} + W_0 + \phi Q_{-1} + u_t$$

Define:

$$(20) \quad \begin{aligned} Z_t &= \sum_{i=1}^t X_i \\ \lambda &= W_0 + \phi Q_{-1} = Q_0 - u_0 + \phi Q_{-1} \end{aligned}$$

Now, the model in equation (19) can be written as:

$$(21) \quad Q_t = \lambda + \alpha_0^* t + Z_t' \pi + \phi Q_{t-1} + u_t$$

In equation (21), $\lambda = Q_0 - u_0 + \phi Q_{-1}$ is assumed to be a constant. OLS applied to equation (21) will now result in consistent estimates of parameters (Johnston).

Data and Empirical Results

The empirical specification of the model in equation (4) is given by:

$$(22) \quad NB_t = 6\alpha_0 + \alpha_1 P_t^* + \alpha_2 C_t^* + e_t$$

where, NB_t is the number of non-bearing pecan trees in 1,000 and P_t^* and C_t^* are as defined in equations (5).

Using equations (19) - (21), the empirical model of equation (13) is given by:

$$(23) \quad \begin{aligned} Q_t &= \lambda + \alpha_0^* t + \alpha_1^* \sum_{i=1}^t \hat{P}_i + \\ &\alpha_2^* \sum_{i=1}^t \hat{C}_i + \sum_{j=1}^m \beta_j \sum_{i=1}^t \Delta P_{i-j} + \\ &\beta_{m+1} \sum_{i=1}^t \Delta C_i + \phi Q_{t-1} + u_t \end{aligned}$$

where Q_t is harvested pecans in 1,000 pounds, P_t is the average annual price of pecan in cents/pound; C_t is the index of input prices paid by farmers; α_0^* , α_1^* , α_2^* , $\hat{P}_t$ and $\hat{C}_t$ are as defined in equations (14). Weather variables (annual rainfall and temperature) were included in the initial model but dropped from the final specification because their coefficients were statistically insignificant.

A series of Georgia Agricultural Facts provided data on pecan prices and quantities and the number of non-bearing trees. Agricultural Statistics supplied the series regarding the index numbers of input prices. National Agricultural Census provided data on bearing and non-bearing pecan trees which were reported every five years. The data

for the first model (equation 22) cover the period 1924 to 1987 with a five year gap, while data for the second model (equation 23) cover the period 1962-1987.

The model in equation (22) was estimated by the Prais-Winsten's method because initial results suggested first order autocorrelation as indicated by the Durbin-Watson's statistic. The results are presented in table 1. The coefficients on the price and cost variables were both significantly different from zero as indicated by the t -statistics. The signs were as expected. The number of non-bearing trees increased with an increase in last year's pecan prices, while it decreased with an increase in the index of input prices paid by farmers.

Table 1. Parameter Estimates of the First Model (equation 22)

Variable	Coefficient	Coefficient estimate	t -ratio
Constant X 6	$6\alpha_0$	150.940	0.670
P_t^*	α_1	2.9655 ^c	3.203
C_t^*	α_2	-0.17032 ^c	-2.162
Rho ^a	ρ	0.82154 ^c	4.992
R^2		0.44	
Durbin-Watson ^b		2.375	

^a Rho is the autocorrelation coefficient. Prais-Winsten's method was used to estimate the parameters.

^b D-W statistic is for the transformed residuals.

^c denotes significance at the 5 percent level.

The second model (equation 23) was estimated by OLS and results were corrected for heteroscedasticity (White). The value of the Durbin's h -statistic indicated no serial correlation could be detected (table 2). The value of m (lag length - see equation 23) was empirically determined. The value for m that gave the best statistical results was 3. The high R^2 indicated a good explanatory power of the model.

The α coefficients, together, reflect the indirect impact of past plantings on the quantity supplied. This impact was significant as indicated by the corresponding t statistics. The

signs of the α coefficients were as expected. An increase in the lagged pecan price positively influenced current plantings and hence, had an indirect positive influence on the supply of pecan. On the other hand, an increase in the index of input prices had a negative effect on new plantings and consequently, an indirect negative effect on pecan supply. The β coefficients represent the direct effects of pecan prices and the index of input prices on pecan supply. The interpretation of these coefficients should be according to the original model of equation (10). Increases in lagged pecan prices had positive effects on pecan supply. However, the greatest effect of current price on supply occurred after two years (β_2 is the

Table 2. Parameter Estimates of The Second Model (equation 23)

Variable	Coefficient	Coefficient estimate	t -ratio
Constant	λ	106020.000 ^c	8.869
t	α_0^*	-7346.200 ^b	-1.873
P_{t^*}	α_1^*	32.995 ^c	2.015
C_{t^*}	α_2^*	-2.775 ^b	-1.803
P_{t-1}	β_1	338.34	1.165
P_{t-2}	β_2	1624.600 ^c	5.720
P_{t-3}	β_3	319.830	0.954
C_t	β_4	-9.448	-0.231
Q_{t-1}	ϕ	-0.680 ^c	-6.578

$R^2 = 0.86$

Durbin's h -statistic = 0.261

Note: Results are corrected for heteroscedasticity (White).

^a For variable definitions see text and equations (14) and (23).

^b denotes significance at 10% level.

^c denotes significance at 5% level.

statistically significant coefficient). The direct effect of the index of input prices on pecan supply was negative as expected. The coefficient of the lagged dependent variable was negative and significantly different from zero. The result suggests the presence of the alternate bearing phenomenon in pecan production. An increase in the supply of pecans last year lowered supplies in the current year.

New Plantings Prediction

In order to predict new plantings, the estimates of the structural parameters of equation (1), the α parameters, must be recovered. Plugging these estimates back into equation (1) and using values of the observed price and cost variables, allows the prediction of new plantings.

The first model (equation 22) gives direct estimates of the α parameters (α_0 is obtained by dividing the intercept estimate by 6). However, the second model (equation 23) gives estimates of the transformations of the α parameters as described in equations (14). From equations (14) one could calculate the estimates of the α parameters if the value of γ_1 is known. In the following analyses γ_1 was assumed to be equal to one. That is, it was assumed that all potential output would be harvested, an assumption which seems to be reasonable in case of pecans.

The only report on new pecan plantings was for the year 1929 (Jones et al.). A survey of over two million trees of improved varieties reported that about 109,000 (or 5.4 percent of the total number of pecan trees) were planted in 1929. However, the estimate of the total number of trees reported by Jones et al. was 49 percent higher than the National Agricultural Census estimate for the same year.

Using the α estimates from the first model (equation 22), the predicted new planting was 77,816 trees in 1929. This estimate is equal to about 71 percent of the estimate reported in Jones et al. The second model (equation 23) predicted planting of 280,230 trees or about 257 percent of the estimate reported by Jones et al. However, if Jones et al. overestimated new plantings by the

same percentage as for the total number of trees, then the adjusted estimate of new trees is 73,154. After this adjustment, the first model predicted about 106 percent and the second model about 383 percent of Jones et al.'s estimate of new plantings in 1929. Moreover, while the Jones et al. estimate was for the improved pecan varieties only, the estimates in this study are for both improved varieties and seedlings. This may explain why both models predicted larger new plantings than reported by Jones et al. after adjustment particularly because at that time the common knowledge suggested planting of several cultivars including seedlings for improved pollination.

Because non-negativity constraints were not placed on either model, negative prediction of new plantings were possible due to the linear functional form of the planting equation (equation 1). However, negative predictions of new plantings were only among the results obtained using the second model. All predictions (for the period 1924 to 1987) from the first model were positive.

The above results suggest that the first model performed better in terms of predicting the number of new plantings. The first model is simpler and more direct than the second model. Hence, results indicate that simple approaches to model pecan new planting may outperform the complex ones, especially if detailed data are not available.

Concluding Comments

This study focused on improving methods of estimating the number of non-bearing trees, modifying and adopting the models to specific data limitations, and tree crop characteristic. The modeling of the tree crop planting response such as tree nuts is difficult due to gaps in data on the number of trees, new additions to existing orchards and removal of trees. The first of the proposed methods accounted directly for new plantings and the non-bearing and bearing growing stage. The second approach considered the dynamic yield-tree age relationship allowing for yield increases as a tree matures.

The application of both approaches to the pecan industry provided empirical illustration. The majority of the parameter estimates were statistically significant. The first approach more accurately estimated the pecan planting response in case of 1929 data than the model based on the second approach.

From practical standpoint, the first approach contains the basic information readily available to a grower as it includes past prices and a measure of production costs. This approach, less complex, predicted a positive number of new plantings confirmed by the actual observation of the developments in the pecan industry over time. Therefore, the first approach may be used in generating information for making investment decisions. The second approach incorporates more information by combining the harvesting and

investment decisions and by incorporating the dynamic yield-tree age relationship. However, predictions of new plantings obtained using the second approach were occasionally negative.

The models offer a methodological framework for estimating the planting response of the pecan industry. The empirical applications of the models could be expanded as more data become available. Progress in data reporting for pecan industry has been slow but steady as illustrated by initiating pecan storage reports in the past and including Arizona and California as individual areas in pecan price and production reports. The importance of data needs and progress in understanding pecan tree physiological mechanism will balance efforts in developing methodological framework enhancing the ability to provide increasingly accurate information needed for the practical decision-making.

References

Agricultural Statistics, USDA, various issues.

Akiyama, T. and P. K. Trivedi. "Vintage Production Approach to Perennial Crop Supply: An Application to Tea in Major Producing Countries." *J. Econometrics*, 36(1987):133-161.

Baritelle, J. L. and D. W. Price. "Supply Response and Marketing Strategies for Deciduous Crops." *Amer. J. Agr. Econ.*, 56(1974):245-53.

Bateman, M. J. "Aggregate and Regional Supply Functions for Ghanaian Cocoa, 1946-1962." *J. Farm Econ.*, 47(1965):384-401.

Behrman, J. R. "Monopolistic Cocoa Pricing." *Amer. J. Agr. Econ.* 50(1968):702-719.

Bellman, R. E. and M. J. Hartley. "The Tree-Crop Problem." Report No. DRD134, The World Bank, 1985.

Epperson, J. E. and J. R. Allison. "Price Futures for the United States Pecan Industry." *HortScience*, 15(4):475-478, 1980.

French, B. C. "The Long-Term Price and Production Outlook for Apples in the United States and Michigan." Michigan Agric. Exp. Stat. Tech. Bul. 255, 1956.

French, B. C. and R. G. Bressler "The Lemon Cycle." *J. Farm Econ.* 44(1962): 1021-1036.

French, B. C. and J. L. Matthews. "A Supply Response Model for Perennial Crops." *Amer. J. Agr. Econ.*, 53(1971):478-90.

- French, B. C. and L. S. Willet. "An Econometric Model of the U.S. Asparagus Industry." Giannini Res. Report No. 340, Univ. of California, Davis, September 1989.
- French, B. C., G. A. King, and D. D. Minami. "Planting and Removal of Perennial Crops." *Amer. J. Agr. Econ.*, 67(1985):215-223.
- Georgia Agricultural Facts*. Georgia Agricultural Extension Service, University System of Georgia, various issues.
- Gorman, W. D., D. F. Landrum, and S. D. Hicks. "Pecan Orchard Costs and Returns for 40 and 320 Acres, Southern Rio Grande Valley, New Mexico, 1979." Agr. Exp. Sta., Res. Report No. 413, New Mexico State University, May 1980.
- Hartley, M. J., M. Nerlove, and R. K. Peters. "An Analysis of Rubber Supply in Sri Lanka." *Amer. J. Agr. Econ.*, 69(1987):755-761.
- Johnston, J. *Econometric Methods*. New York: McGraw-Hill Book Co., 1972.
- Jones, S. A., V. C. Childs, R. S. Washburn, B. H. Thibodeaux, J. W. Park, and H. E. Rutland. "An Economic Study of the Pecan Industry." USDA Technical Bul. No. 324, 1932.
- Nerlove, M. "Distributed Lags and Estimation of Long-Run supply and Demand Elasticities: Theoretical Considerations." *J. Farm Econ.* 40:301-310.
- Okunade, A. A., M. J. Cochran. "Functional Forms and Form-level Demand for Pecans by Variety." *S. J. Agr. Econ.* 23(2):95-102, 1991.
- Shafer, C. E. "Price and Value Effects of Pecan Crop Forecasts, 1971-1987". *S. J. Agr. Econ.*, 21(1):97-103, 1989.
- Stern, R. M. "The Determinants of Cocoa supply in West Africa." R. E. Stewart, H. W. Ord, eds., *African Primary Products and International Trade*, Edinburgh University Press, 1965.
- Wells, G. J., S. E. Miller, and C. S. Thompson. "Farm Level Demand for Pecans Reconsidered." *S. J. Agr. Econ.*, 16(1):157-160, 1984.
- White, H. "A Heteroscedasticity Consistent Covariance Matrix and a Direct Test for Heteroscedasticity." *Econometrica*, 46(1978):817-838.
- Wickens, M. R., J. N. Greenfield. "The Econometrics of Agricultural Supply: An Application to the World Coffee Market." *Rev. Econ. Stat.* 55(19):533-440.
- Williams, F. W., M. G. LaPlante, and E. K. Heaton. "The Consumer Market for Pecans and Competing Nuts." *S. J. Agr. Econ.*, 4(1):101-106.