



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

Chalkiness Characters and Scanning Electron Microscope Observation of Rice Grain Endosperm of *Japonica* Varieties in Southern Henan

Bo PENG^{1*}, Yanfang SUN¹, Dongyan KONG¹, Ruihua PANG¹, Xiaohua SONG², Huilong LI², Jintao LI¹, Qiyong ZHOU¹, Qiu HAN¹, Shizhi SONG², Hongyu YUAN^{1#}

1. College of Life Sciences, Institute for Conservation and Utilization of Agro-bioresources in Dabie Mountains, Xinyang Normal University, Xinyang 464000, China; 2. Xinyang Academy of Agricultural Science, Xinyang 464000, China

Abstract Chalkiness characters affect not only the grain appearance, milling, eating and cooking qualities but also the grain nutritional quality in rice, thus it is one of the most important traits in rice. It is very important for us to investigate the relation of the chalkiness formation and the development of endosperm structure and starch granule of different rice varieties. Here, we have investigated the chalkiness characters such as chalkiness rate, chalkiness degree and chalkiness area in 15 *japonica* rice varieties from southern Henan. Furthermore, the endosperm structure and starch granules of rice grain were also observed with scanning electron microscope. The results showed that the 15 *japonica* varieties have a significantly linear relationship between the chalkiness rate and chalkiness degree. Among the varieties, the biggest difference is the chalkiness rate, the second is the chalkiness area, and the last is the chalkiness degree. Moreover, there is a certain correlation between the distribution of starch granules, the arrangement of endosperm cells and the occurrence of grain chalkiness in the different rice varieties. For the same variety, the starch granules of chalky and non-chalky grains have obvious difference, while the starch granules from the transparent part of chalky rice and non-chalky rice do not have significant difference. The results would provide useful references for the improvement of grain quality in rice.

Key words Chalkiness, Scanning electron microscope, *Japonica* rice, Starch granule, Endosperm

1 Introduction

Oryza sativa L. is one of the most important grain crops in the world, and more than half of the population in the world and 2/3 of the population in China take rice as staple food^[1–2]. With rapid development of national economy and continuous improvement of living level, people's demand on high-quality grain is also continuously increasing^[3–6]. However, rice quality is an extremely complex trait, including appearance, milling, nutrient, eating and cooking qualities^[2,7], while appearance quality is mainly decided by grain size, transparency and chalkiness^[8–9]. Chalkiness characters not only affect the grain appearance, milling, nutrient, eating and cooking qualities^[10–13], but also cause rice yield decline^[14–15]. Therefore, chalkiness characters directly decide grain quality and market price of rice to a large extent. Chalkiness is one of main problems existed in rice production and consumption, especially in South China, rice always has the problem of high chalkiness rate and chalkiness degree^[16]. Previous results showed

that grain chalkiness characters of rice are controlled by multiple genes^[3,17–18], have additive effect and interaction between additive effect and environment, and are affected by external environmental factors^[10–11,19–24]. For example, high temperature induction^[5,25], daily average temperature, diurnal temperature difference, sunshine and cultivation measures at grain filling stage all affect chalkiness characters of rice to different degree^[26–29]. At present, a batch of genes affecting chalkiness characters of rice have been isolated and cloned by using natural population or mutant, such as *GW2*^[30], *Chalk5*^[19], *OsPPDKB*^[31], *SSIIIa*^[32–33], *GIF1*^[34], *OsRab5a*^[35] and *flo2*^[36]. Chalkiness is white opaque part which is formed by aeration because of loose arrangement of starch granules when the development and enrichment of endosperm starch granules and protein of rice at grain filling stage are affected^[29,37–38]. Different varieties and qualities of rice have the fixed starch granule morphology, and grain quality could be identified initially by scanning electron microscope (SEM)^[16,39–41]. Based on prior researches, chalkiness characters of grain endosperm of the 15 *japonica* rice varieties were investigated in this paper, and SEM observation of grain endosperm cell and starch granule in rice was conducted to study the relationship between chalkiness characters formation of different *japonica* rice varieties and morphology and development of endosperm cell and starch granule. The results have paved the way for further studying chalkiness formation mechanism of *japonica* rice and cultivating high-quality new rice varieties.

Received: January 4, 2017 Accepted: February 16, 2017

Supported by Key Project of Science and Technology in Henan Province (152102110100, 152102110036); National Natural Science Foundation of China (U1604110, U1404319, 31270727, 31600992); Nanhu Scholars Program for Young Scholars of XYNU (2016056); Major Science and Technology Project in Henan Province (121100110200); Students Scientific Research Fund of Xinyang Normal University (2015-DXS-158); Fund of Institute for Conservation and Utilization of Agro-bioresources in Dabie Mountains (2016020).

Corresponding author. E-mail: *pengbo@xynu.edu.cn, #hongyu92@163.com

2 Materials and methods

2.1 Materials There were 15 *japonica* rice varieties for test, including 5 *japonica* rice varieties from Henan Province (Xiangnuo 1862, Xianggeng 33, Zhengdao 18, Xindao 18, Aiganxiangdaowan), 3 varieties of Liaoning Province (Yanfeng 47, Liaoxing 1 and Longyou 1875), 2 varieties of Jiangsu Province (Wuyungeng 23 and Nangeng 9108), 2 *japonica* rice varieties of Heilongjiang Province (Longgeng 31 and Longgeng 21), 2 varieties of Zhejiang Province (Zhegeng 88 and Xiushui 134) and Xinfeng 3 of Anhui Province. According to different growth periods of these varieties, they were sown in the same test field of Xinyang Academy of Agricultural Sciences in different batches of 2015, thereby guaranteeing that the 15 *japonica* rice varieties bloomed in last of August, and the temperature was relatively consistent at grain filling stage. Each variety was sown for two rows, and each row had 12 plants, and plant spacing was 16.5 cm × 26.4 cm. Conventional cultivation management of field was implemented from sowing to mature period of grain. After maturity, rice grain was tested after natural drying and conservation for three months at room temperature.

2.2 Methods

2.2.1 Determination of chalky traits. Chalkiness rate, chalkiness area and chalkiness degree of different *japonica* rice varieties were determined by the national standard GB/T17891-1999. 100 grains of full and complete polished rice from each variety were taken randomly, and the rice grain with chalkiness was picked out, which was repeated for three times, and the mean was the chalkiness rate of each variety. White belly rate and white core rate (the grain with white back was not observed) were determined by the method of chalkiness rate. 100 grains of full and complete polished rice were taken randomly, and the rice grain with white belly and white core was picked out, which was repeated for three times, and the mean was white belly rate and white core rate of each variety. 10 grains of chalky rice of each *japonica* rice variety were taken randomly, and the percentage of chalky part to whole rice grain area was estimated, which was repeated for three times, and the mean was chalkiness area. Chalkiness degree = Chalkiness rate × Chalkiness area. SPSS software was used for variance analysis of the measured data.

2.2.2 SEM observation. The chalky and non-chalky rice was taken from each *japonica* rice variety randomly. The blade back was used to knock middle part of the rice lightly, making it naturally split, and then the split site was cut off and made into 2–3 mm thickness of sample. A part was observed by ordinary optical microscope, and the other part was adhered to copper sample table by conductive adhesive. HUS-5GB type of vacuum coating apparatus was used to plate the section, and the sample was set under scanning electron microscope (Hitachi S-4800) to observe different parts of rice section and take a photo.

3 Results and analysis

3.1 Difference of chalkiness traits from different *japonica* rice varieties The measurement results of chalkiness traits from

15 *japonica* rice varieties were shown as Fig. 1. It was clear that chalkiness rate had the maximum difference among varieties, followed by chalkiness area, and the last was chalkiness degree. The chalkiness rate of Aiganxiangdaowan reached 93.1%, which was the highest. The chalkiness traits could not be observed from Xinfeng 3, and chalkiness rates of the 15 *japonica* rice varieties changed between 0 and 93.1%. Variance analysis showed that chalkiness rates of Yanfeng 47, Longyou 1875 and Nangeng 9108 had insignificant difference; chalkiness rates of Xindao 18, Longgeng 31 and Xianggeng 33 had insignificant difference; the difference of chalkiness rates of Longgeng 21 and Liaoxing 1 was insignificant; the difference of chalkiness rates of Xiushui 134 and Zhengdao 18 was insignificant. Chalkiness degree changed between 0 and 52.0%, and change trend among different varieties was very similar with that of chalkiness rate. Grain chalkiness of Zhegeng 88, Longgeng 21, Wuyungeng 23, Nangeng 9108, Liaoxing 1 and Longyou 1875 mainly occurred in belly, while chalkiness of other varieties mainly occurred in center part of rice endosperm. Chalkiness area changed between 0 and 63.0%, and the change trend was different from that of chalkiness rate and chalkiness degree. Chalkiness rates of Xianggeng 33 and Zhengdao 18 were not very high, but their chalkiness area surpassed 15.0%. It is noteworthy that chalkiness traits were not observed in Xinfeng 3. Chalkiness rate of Aiganxiangdaowan reached 93.1%, and its chalkiness area was 63.0%, while the chalkiness degree was 52.0%. Further analysis displayed that chalkiness rate showed obvious linear relationship with chalkiness degree, and chalkiness degree of the variety with high chalkiness rate was also higher.




Fig. 1 Chalkiness characters of the 15 *japonica* rice varieties

3.2 Clustering of chalkiness traits of different *japonica* rice varieties

The system clustering analysis of chalkiness rate (white belly rate and white core rate), chalkiness degree and chalkiness area of the 15 *japonica* rice varieties was conducted, and the results found that the tested *japonica* rice varieties could be divided into 4 classes (Fig. 2). Xindao 18, Longgeng 31, Xinfeng 3, Xiushui 134, Xianggeng 33 and Zhengdao 18 were clustered into Class I, and their chalkiness rates changed between 0 and 7.4%. Chalkiness area changed between 0 and 19.2%, and chalkiness degree changed between 0 and 10.2%. Zhegeng 88, Wuyungeng 23 and Xiangnuo 1862 were clustered into Class III, and their chalkiness rates changed between 47.7% and

49.5%. Chalkiness area changed between 25.3% and 36.7%, and chalkiness degree changed between 28.7% and 34.6%. Aiganxiangdaowan belonged to Class IV, and its chalkiness rate was 93.1%. Chalkiness area was 63.0%, and chalkiness degree was 52.0%. Longgeng 21, Liaoxing 1, etc. were clustered into Class II, and their chalkiness rates changed between 14.4% and 19.6%. Chalkiness area changed between 10.1% and 22.2%, and chalkiness degree changed between 11.5% and 19.0%. Therefore, for 6 varieties of Class I, their chalkiness rates were lower, and chalkiness degree and chalkiness area were smaller, while Aiganxiangdaowan of Class IV showed contrary chalkiness traits.





Fig. 2 Clustering of the chalkiness characters of the tested *japonica* rice varieties

3.3 Endosperm morphology of different *japonica* rice varieties

The representative milled grains were selected, and ordinary optical microscope and SEM observation of endosperm morphology of 15 different *japonica* rice varieties was conducted from arrangement manner, morphology, layer number of rice endosperm cell and starch granule distribution of its cross section, and the results were shown as Fig. 3. Arrangement manner of endosperm cell at cross section mainly contained four classes. Class I showed obvious radiation shape from middle cross section of rice endosperm to all around, and the radiation was longer. The variety dominated by white core belonged to the situation, such as Xianggeng 33, Xiangnuo 1862 and Aiganxiangdaowan (Fig. 3A, E, I). Class II also showed obvious radiation shape from middle cross section to all around, but endosperm cell arranged radially was not straight, and some bent together. The variety dominated by white belly belonged to the situation, such as Nangeng 9108, Longgeng 21, Liaoxing 1, Wuyungeng 23 (Fig. 3B, F, J). Although Class III showed radiation shape from middle cross section of rice endosperm to all around, the radiation was not obvious. The varieties with white core and white belly belonged to the situation, such as Yanfeng 47, Longyou 1875, Zhegeng 88 (Fig. 3C, G, K). Radia-

tion shape could not be observed from middle part of cross section of Class IV rice endosperm, and the varieties with lower chalkiness rate or without chalkiness traits belonged to the situation, such as Xinfeng 3 and Xindao 18 (Fig. 3D, H, L). Under scanning electron microscope, it could see that the varieties dominated by white core had multilayer polygonal columnar cell, and there were many oval cell all around, such as Xianggeng 33, Xiangnuo 1862, Aiganxiangdaowan (Fig. 3A). The varieties dominated by white belly had irregular cell group, and multilayer polygonal columnar cells were at peripheral layer, such as Nangeng 9108, Longgeng 21, Liaoxing 1, Wuyungeng 23 (Fig. 3B). The varieties with white core and white belly had oval cell group, and polygonal columnar cells were at peripheral layer, such as Yanfeng 47, Longyou 1875 and Zhegeng 88 (Fig. 3C). Polygonal columnar cells, oval cells and irregular cells of the varieties with lower chalkiness rate or without chalkiness traits were not obvious, such as Xinfeng 3 and Xindao 18 (Fig. 3D).

Electron microscope was used to observe distribution situation of starch granules at cross section of rice grains. It could be seen that the distribution of starch granules of chalky *japonica* rice were obviously uneven, and starch granules mainly concentrated in center part and dorsal abdominal diameter. The density of starch granule from different chalkiness types of *japonica* rice varieties was also different. Starch granule of *japonica* rice variety dominated by white core was mainly distributed in center part, while starch granule of *japonica* rice variety dominated by white belly was distributed in belly position of rice grain, and starch granule of *japonica* rice variety with white core and white belly was distributed in grain belly and center part, which was uneven. Starch granule of *japonica* rice variety without chalkiness or with very low chalkiness rate was obviously distributed in whole cross section, and the density of starch granule was larger (Fig. 3). The above results showed that arrangement manner of rice endosperm cells and distribution of starch granules had certain correlation with occurrence of chalkiness traits.




Note: A, E, I were Xianggeng 33; B, F, J were Longgeng 21; C, G, K were Longyou 1875; D, H, L were Xindao 18; Bar = 50 μ m.

Fig. 3 Low magnification view of the transversely fractured mid-region of the tested rice

3.4 Starch granule morphology of chalky and non-chalky rice

SEM observation of chalky and non-chalky rice from 15 *japonica* rice varieties was conducted. The results showed that starch granules at chalky part were spherical or ellipsoidal, with unobvious edge. The size of starch granule was different, with smaller diameter, and they accumulated with irregular shape. The space among starch granules became larger, with loose arrangement, and they were all single starch granule at free state (Fig. 4A, D). By contrasting transparent part of chalky rice grain with starch granule of non-chalky rice, it was found that there was not obvious difference, and most of starch granules had consistent size, and they were regular diamond or equal polygon, and had polyhedral edges. Moreover, the space among starch granules was very small, and the arrangement was very close (Fig. 4B, C, E, F). The above results showed that chalkiness formation was closely related to the shape and arrangement of endosperm starch granules. At the part with low chalkiness degree, its starch granule developed well, on the contrary, the development of starch granule was not good.




Note: A, D showed the starch grains in the chalky rice; B, E showed the starch grains in the transparent parts of chalky rice; C, F showed the starch grains in the non-chalky rice; Bar = 30 μm .

Fig.4 Scanning electron microscopic observation on starch granules in endosperm cells of the chalky and non-chalky rice

3.5 Morphology of starch granule at different parts of chalky rice grain

According to occurrence part of chalkiness in rice endosperm, chalkiness could be divided into three types: white belly, white core and white back. SEM observation on belly, core and back of chalky rice from 15 *japonica* rice varieties was conducted. It was found that the space among starch granules at center part of endosperm from Xianggeng 33, Xiangnuo 1862 and Aiganxiangdaowan was larger, with loose arrangement, and the development of starch granule was bad, while starch granule at endosperm belly and back developed better (Fig. 51A, 1B, 1C). Starch granule at endosperm belly of Nangeng 9108, Longgeng 21, Liaoxing 1, Wuyungeng 23 developed bad, while starch granule at endosperm core and back developed well (Fig. 52A, 2B, 2C). Starch granule at endosperm core and back of Yanfeng 47, Longyou 1875, Zhegeng 88 arranged loosely and developed bad, while

the development of starch granule at back was better (Fig. 53A, 3B, 3C). Starch granule at endosperm core, belly and back of Xinfeng 3 and Xindao 18 developed well (Fig. 54A, 4B, 4C).



Note: 1–4 were Xiangnuo 1862, Wuyungeng 23, Yanfeng 47 and Xinfeng 3 respectively; A showed starch grains in the center of the endosperm; B showed starch grains in the belly of the endosperm; C showed starch grains on the back of the endosperm; Bar = 10 μm .

Fig.5 Scanning electron microscopic observation on starch granules in different parts of the tested *japonica* rice varieties

4 Conclusions and discussions

Chalkiness trait is an extremely important quality. Chalkiness affects not only appearance quality of rice but also milling, nutrient, eating and cooking qualities^[10–13]. Morphology of starch granules in rice could be observed by scanning electron microscope, and rice quality could be identified preliminarily^[16,39–41]. Seen from scanning electron microscope results of endosperm cells from 15 *japonica* rice varieties, starch granule distribution at cross section of the variety with higher chalkiness rate and chalkiness degree (Xiangnuo 1862, Aiganxiangdaowan and Zhegeng 88) was obviously uneven, and it was more obvious at center part of endosperm. For the variety with lower chalkiness rate and chalkiness degree, starch granule distribution at center part, belly and back of endosperm was even. By comprehensively analyzing chalkiness traits of endosperm from 15 *japonica* rice varieties and cell morphology at cross section, it was found that if starch granule distribution at cross section was even, chalkiness traits were not easy to occur in the variety, and chalkiness rate and chalkiness degree were lower. If starch granule distribution at cross section was uneven, the space among starch granules was larger, with loose ar-

rangement, and the development of starch granule was bad, so chalkiness traits were very easy to occur in the variety, and the chalkiness rate and chalkiness degree at corresponding position were higher. Therefore, distribution situation of starch granules at cross section of rice endosperm had the relationship with the generation of chalkiness traits.

By observing chalkiness traits of endosperm from 15 *japonica* rice varieties, it was found that morphology and arrangement of starch granule at different endosperm parts of the same variety were also different. Compared with transparent part or starch granule of non-chalky rice grain, starch granules at chalky part of endosperm mainly existed freely, and the space among starch granules was larger, so it could not further develop into starch granule complex with certain shape. If single starch granule could realize ordered arrangement and become complex starch granule with certain geometrical shape, chalkiness traits could not appear at the part. Ellipsoidal or spherical starch grains could not arrange orderly at chalky part, and the accumulation showed as disordered shape. Appearance embodiment of the ordered and disordered accumulation of starch grain was if rice endosperm had chalkiness and showed as transparent state. Previous studies found that the development of rice endosperm at the back was better than that of center part and belly, so chalkiness traits of back were rarely to occur^[16,39,42-45]. In endosperm of the tested 15 *japonica* rice varieties, white back did not appear, and starch granule of back was regular diamond or equal polygon, with polyhedral edge, and the space among starch granules was very small, with tight arrangement, showing that the development of back starch granule was good. Maybe it was because that nutrients were transported firstly to the back of rice grain, and then center part and belly of endosperm, so the development of endosperm back was better than center part and belly. Therefore, chalkiness traits of rice endosperm were closely related to the development of starch granules.

Previous studies showed that chalkiness trait of rice grain was a complex number character, was commonly controlled by multiple genes, and was affected by environmental factors, especially temperature at grain filling stage significantly affected chalkiness traits^[19-21,46-47]. Chalkiness traits had obvious difference among the tested 15 *japonica* rice varieties. By combining our previous results^[7,48], it was found that chalkiness traits of endosperm were mainly controlled by inheritance. The tested 15 *japonica* rice varieties had better representation of ecological type. To decrease the influence of external environmental factors on chalkiness traits of the 15 *japonica* rice varieties as much as possible, the 15 *japonica* rice varieties were planted in the same tested field. Via prior pre-test and the manner of batch sowing, it was ensured that they simultaneously bloomed in last dekad of August, 2015, and temperature was relatively consistent at grain filling stage. From first dekad of August, 2015 to first dekad of September, 2015, temperature was relatively higher in Xinyang City, which might cause that chalkiness traits of some *japonica* rice varieties could be more serious than that in origin production site. But external environmental

conditions of the 15 *japonica* rice varieties at grain filling stage were basically same. The test results displayed that they showed similar some common chalkiness characters. However, the occurrence of chalkiness traits was a complex process, and the chalkiness formation may involve the relationship of "source", "storehouse" and "streams" in rice^[47]. Their any change or uncoordinated relationship may cause or increase the formation of chalkiness traits in rice endosperm. By combining our previous results, it was thought that rice endosperm cell and the structure of starch granule at cross section decided appearance performance of chalkiness traits of endosperm, while endosperm structure was decided by the development of rice endosperm cell. All of those were controlled by inherited factors^[48]. Therefore, the occurrence of chalkiness traits of rice endosperm was closely related to the distribution and development of starch granules, and the occurrence, development process, latter deposition mode and distribution situation of starch granules in rice endosperm may be very important. However, how to regulate prior occurrence, development process and latter deposition of starch granule in rice endosperm? How many genes join in the important process? These may be key problems further clarifying occurrence mechanism of chalkiness traits, which needs further research and exploration.

References

- [1] KIM J, KIM B, LEE J, *et al.* Protein content and composition of *waxy* rice grains[J]. Pakistan Journal of Botany, 2013, 45:151-156.
- [2] TIAN ZX, QIAN Q, LIU QQ, *et al.* Allelic diversities in rice starch biosynthesis lead to a diverse array of rice eating and cooking qualities[J]. Proceedings of the National Academy of Sciences of the United States of America, 2009, 106:21760-21765.
- [3] TAO YJ, XU MB, WANG F, *et al.* QTL mapping of chalkiness using chromosome single segment substituted lines in rice[J]. North China Journal of Agriculture, 2015, 30 (1):1-8.
- [4] PENG B, KONG HL, LI YB, *et al.* *OsAAP6* functions as an important regulator of grain protein content and nutritional quality in rice[J]. Nature Communications, 2014, 5:4847.
- [5] FITZGERALD MA, MCCOUCH SR, HALL RD. Not just a grain of rice: the quest for quality[J]. Trends in Plant Science, 2009, 14:133-139.
- [6] CHENG SH, ZHUANG JY, FAN YY, *et al.* Progress in research and development on hybrid rice: A super-domesticated in China[J]. Annals of Botany, 2007, 100:959-966.
- [7] PENG B, WANG LQ, FAN CC, *et al.* Comparative mapping of chalkiness components in rice using five populations across two environments [J]. BMC Genetics, 2014, 15:49.
- [8] ZHANG QF. Strategies for developing green super rice[J]. Proceedings of the National Academy of Sciences of the United States of America, 2007, 104:16402-16409.
- [9] WAN XY, WENG JF, ZHAI HQ, *et al.* Quantitative trait loci (QTL) analysis for rice grain width and fine mapping of an identified QTL allele *gw-5* in a recombination hotspot region on Chromosome 5[J]. Genetics, 2008, 179:2239-2252.
- [10] ZHENG LN, ZHANG WW, LIU SJ, *et al.* Genetic relationship between grain chalkiness, protein content, and paste viscosity properties in a backcross inbred population of rice [J]. Journal of Cereal Science, 2012, 56:153-160.
- [11] LIU XL, WAN XY, MA XD, *et al.* Dissecting the genetic basis for the effect of rice chalkiness, amylose content, protein content, and rapid viscosity analyzer profile characteristics on the eating quality of cooked

- rice using the chromosome segment substitution line population across eight environments[J]. *Genome*, 2011, 54:64–80.
- [12] YOSHIOKA Y, IWATA H, TABATA M, *et al.* Chalkiness in rice: Potential for evaluation with image analysis[J]. *Crop Science*, 2007, 47: 2113–2120.
 - [13] SHEN XP, SHEN XY, GU L, *et al.* Effect of seeding time on chalkiness of liangyoupeijiu in Jiangsu rice growing areas at different latitudes [J]. *Chinese Journal of Rice Science*, 2007, 21:677–680.
 - [14] ZHAO XQ, DAYGON VD, MCNALLY KL, *et al.* Identification of stable QTLs causing chalk in rice grains in nine environments[J]. *Theoretical and Applied Genetics*, 2016, 129:141–153.
 - [15] ZHAO XQ, FITZGERALD MA. Climate change: Implications for the yield of edible rice[J]. *PLoS One*, 2013, 8:e66218.
 - [16] KANG HQ, CHANG HY. Study on chalkiness characters and endosperm structures of the main parents kernel of hybrid rice[J]. *Chinese Agricultural Science Bulletin*, 2007, 23(4):180–185.
 - [17] QIU XJ, YUAN ZH, CHEN K, *et al.* Genetic dissection of grain chalkiness in *indica* mini-core germplasm using genome-wide association method[J]. *Acta Agronomica Sinica*, 2015, 41(7):1007–1016.
 - [18] QIU XJ, YUAN ZH, HE WJ, *et al.* Progress in genetics and breeding research on rice chalkiness[J]. *Journal of Plant Genetic Resources*, 2014, 15(5):992–998.
 - [19] LI YB, FAN CC, XING YZ, *et al.* *Chalk5* encodes a vacuolar H⁺ – translocating pyrophosphatase influencing grain chalkiness in rice[J]. *Nature Genetics*, 2014, 46:389–404.
 - [20] TSUKAGUCHI T, LIDA Y. Effects of assimilate supply and high temperature during grain-filling period on the occurrence of various types of chalky kernel in rice plants (*Oryza sativa* L.) [J]. *Plant Production Science*, 2008, 11:203–210.
 - [21] PENG S, HUANG J, SHEEHY JE, *et al.* Rice yields decline with higher night temperature from global warming[J]. *Proceedings of the National Academy of Sciences of the United States of America*, 2004, 101:9971–9975.
 - [22] MEI DY, ZHU YJ, YU YH, *et al.* Quantitative trait loci for grain chalkiness and endosperm transparency detected in three recombinant inbred line populations of *indica* rice[J]. *Journal of Integrative Agriculture*, 2013, 12:1–11.
 - [23] YAMAKAWA H, EBITANI T, TERAOKA T. Comparison between locations of QTLs for grain chalkiness and genes responsive to high temperature during grain filling on the rice chromosome map[J]. *Breeding Science*, 2008, 58:337–343.
 - [24] WAN XY, WAN JM, WENG J, *et al.* Stability of QTLs for rice grain dimension and endosperm chalkiness characteristics across eight environments[J]. *Theoretical and Applied Genetics*, 2005, 110:1334–1346.
 - [25] WANG ZM, LI HX, LIU XF, *et al.* Reduction of pyruvate orthophosphate dikinase activity is associated with high temperature-induced chalkiness in rice grains[J]. *Plant Physiology and Biochemistry*, 2015, 89:76–84.
 - [26] HASHIMOTO Z, MORI N, KAWAMURA M, *et al.* Genetic diversity and phylogeny of Japanese sake-brewing rice as revealed by AFLP and nuclear and chloroplast SSR markers[J]. *Theoretical and Applied Genetics*, 2004, 109:1586–1596.
 - [27] TAMAKI M, KURITA S, TOYOMARU M, *et al.* Difference in the physical properties of white-core and non-white-core kernels of the rice varieties for sake brewing is unrelated to starch properties [J]. *Plant Production Science*, 2006, 9:78–82.
 - [28] KAMARA J, HOSHINO M, SATOH Y, *et al.* Japanese sake-brewing rice cultivars show high levels of globulin-like protein and a chloroplast stromal HSP70[J]. *Crop Science*, 2009, 49:2198–2206.
 - [29] ZHOU L, LIANG S, PONCE K, *et al.* Factors affecting head rice yield and chalkiness in *indica* rice[J]. *Field Crops Research*, 2015, 172:1–10.
 - [30] SONG XJ, HUANG W, SHI M, *et al.* A QTL for rice grain width and weight encodes a previously unknown RING-type E3 ubiquitin ligase [J]. *Nature Genetics*, 2007, 39:623–630.
 - [31] KANG HG, PARK S, MATSUOKA M, *et al.* White-core endosperm floury endosperm-4 in rice is generated by knockout mutations in the C-type pyruvate orthophosphate dikinase gene (*OsPPDKB*) [J]. *Plant Journal*, 2005, 42:901–911.
 - [32] FUJITA N, YOSHIDA M, KONDO T, *et al.* Characterization of *SSIIIa*-deficient mutants of rice: The function of *SSIIIa* and pleiotropic effects by *SSIIIa* deficiency in the rice endosperm[J]. *Plant Physiology*, 2007, 144:2009–2023.
 - [33] RYOO N, YU C, PARK CS, *et al.* Knockout of a starch synthase gene *OsSSIIIa/Flo5* causes white-core floury endosperm in rice (*Oryza sativa* L.) [J]. *Plant Cell Reports*, 2007, 26:1083–1095.
 - [34] WANG ET, XU X, ZHANG L, *et al.* Duplication and independent selection of cell-wall invertase genes *GIF1* and *OsCIN1* during rice evolution and domestication[J]. *BMC Evolutionary Biology*, 2010, 10:108.
 - [35] WANG YH, REN YL, LIU X, *et al.* *OsRab5a* regulates endomembrane organization and storage protein trafficking in rice endosperm cells[J]. *Plant Journal*, 2010, 64:812–824.
 - [36] SHE KC, KUSANO H, KOIZUMI K, *et al.* A novel factor *FLOURY ENDOSPERM2* is involved in regulation of rice grain size and starch quality[J]. *Plant Cell*, 2010, 22:3280–3294.
 - [37] SHI CH, WU JG, LOU XJ, *et al.* Genetic analysis of transparency and chalkiness area at different filling stages of rice (*Oryza sativa* L.) [J]. *Field Crops Research*, 2002, 76:1–9.
 - [38] PATINDOL J, WANG YJ. Fine structures and physicochemical properties of starches from chalky and translucent rice kernels[J]. *Journal of Agricultural and Food Chemistry*, 2003, 51:2777–2784.
 - [39] CHANG HY, KANG HQ, FU HL, *et al.* Different rice varieties (lines) chalky endosperm difference and cytological study[J]. *Journal of Sichuan University (Natural Science Edition)*, 2006, 43(4):927–932.
 - [40] ZHAI B, XU YQ, FU LX. Identification to quality of rice grain with scanning electron microscope[J]. *Journal of Huazhong Agricultural University*, 1991, 10(40):404–408.
 - [41] LIANG JK, WANG XH, LU ND. Scanning electron microscope observation on starch grains of hybrid rice and their parents[J]. *Chinese Journal of Rice Science*, 1996, 10(2):79–84.
 - [42] LISLE AJ, MATIN M, FITZGERALD MA. Chalky and translucent rice grains differ in starch composition an structure and cooking properties [J]. *Cereal Chemistry*, 2000, 67:458–463.
 - [43] SING HN, SODHI NS, KAUR M, *et al.* Physico-chemical, morphological, thermal, cooking and textural properties of chalky and translucent rice kernels[J]. *Food Chemistry*, 2003, 82:433–439.
 - [44] TANG SX, JIANG YZ, LI SS. Observation on the amyloplasts in endosperm of early *indica* rice with scanning electron microscope[J]. *Acta Agronomica Sinica*, 1999, 25(2):269–271.
 - [45] KIM KS, KANG HJ, HWANG IK, *et al.* Comparative structure of ilpumbyeo, a high-quality *japonica* rice, and its mutant suweon 464: scanning and transmission electron microscopy studies[J]. *Journal of Agricultural and Food Chemistry*, 2004, 52:3876–3883.
 - [46] CHEN N, LI TG, LUO YK. Effect of the changes of temperature on the formation of chalkiness of early *indica* rice during endosperm development[J]. *Acta Agriculturae Zhejiangensis*, 2001, 13(2):103–106.
 - [47] LIN WH, XIAO LT, HUANG JL, *et al.* Relation between kernel chalkiness formation and source-sink characteristics in early *indica* rice[J]. *Acta Agriculturae Nucleatae Sinica*, 2003, 17:462–465.
 - [48] PENG B, SUN YF, LI QR, *et al.* Progress in genetic research on rice chalkiness[J]. *Journal of Xinyang Normal University (Natural Science Edition)*, 2016, 29:304–312.