



AgEcon SEARCH

RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

AGRO PRODUCTIVIDAD

Bioremediation

alternatives for total petroleum hydrocarbon removal in agricultural soil

pág. 213

Año 17 • Volumen 17 • Número 5 • mayo, 2024

Response of improved common bean (<i>Phaseolus vulgaris</i> L.) varieties to intermittent drought	3
Determination of geometric properties of cocoa beans (<i>Theobroma cacao</i> L.)	11
Interspecific grafting of <i>Pinus patula</i>	23
Impact of <i>Lippia palmeri</i> S. Watson during kid suckling and growth	31
Root density and accumulation of Myrobalan plum tree grafted with Methley Japanese plum	39
Response of chickpea genotypes (<i>Cicer arietinum</i> L.) to the fungi complex that causes wilt	49

y más artículos de interés...



Colegio de Postgraduados

Biofortification of forages through the application of selenium nanoparticles

Escalona-Tenorio K.¹; Medina-Pérez G.^{1*}; Morales-Rodríguez I.¹; Afanador-Barajas L. N.²; López-Palestina C. U.¹; Pérez-Ríos S.¹

¹ Autonomous University of the State of Hidalgo. Institute of Agricultural Sciences, Tulancingo Hidalgo Mexico, 43600.

² Universidad Central, Engineering and Basic Sciences Faculty, Bogotá 110311, Colombia.

* Correspondence: gabriela_medina@uah.edu.mx

ABSTRACT

Objective: This review paper examines the application of nanoparticles in agriculture, through biofortification with Se NPs in plants.

Methodology: A comprehensive review of the literature on the use of Se NPs in plant biofortification was made.

Results: According to documented studies, the foliar application of NPs significantly improves the morphological and physicochemical characteristics of plants. They promote root development, plant growth and increase the content of raw protein and ether extract in forages. An increase in the production of amino acids and essential fatty acids is also observed.

Limitations: The findings are promising, but there is still slight research on how NPs affect the environment and the safety of their use in the food chain. Further studies are needed to fully understand the long-term effects.

Conclusions: The application of Se NPs for biofortification can improve the nutritional quality of forages, contributing to more efficient and sustainable livestock production. However, to ensure the safety of its implementation it is necessary to continue investigating.

Keywords: sustainable agriculture, animal nutrition, nanotechnology.

Citation: Escalona-Tenorio, K., Medina-Pérez, G., Morales-Rodríguez, I., Afanador-Barajas, L.N., López-Palestina, C. U., & Pérez-Ríos, S. (2024). Biofortification of forages through the application of selenium nanoparticles.

Agro Productividad. <https://doi.org/10.32854/agrop.v17i5.2752>

Academic Editor: Jorge Cadena Iñiguez

Guest Editor: Daniel Alejandro Cadena Zamudio

Received: November 23, 2023.

Accepted: MArch 16, 2024.

Published on-line: June 13, 2024.

Agro Productividad, 17(5). May. 2024. pp: 173-181.

This work is licensed under a Creative Commons Attribution-Non-Commercial 4.0 International license.



INTRODUCTION

Livestock farming faces the challenge of increasing its production in the face of expectations of world population growth due to the products and by-products that come from these animal sources and are required to feed the population, with forecasts of reaching close to 10 billion people in 2050 [1]. The expectations to meet this challenge are high, but this requires more efficient livestock production with fewer polluting emissions, both greenhouse gases and substances that contaminate soils and waters.

Forages are the primary support of sustainable agriculture and constitute one of the leading resources for the success of the animal production system [2]. Selenium (Se) is an essential micronutrient for animals; soil typically contains low levels of Se. Plants can benefit their growth and development by increasing their tolerance to different types of stress and synthesizing phytochemicals with antioxidant properties [3, 4]. Therefore, Se

may be present, but in chemical forms, it is not available to organisms; as a result, Se content is deficient in plants, animals, and other organisms in the food chain [5, 6]. Biofortification increases the nutritional value of food crops by increasing the mineral and vitamin content in the plants [7]. Biofortification has been carried out in agricultural products (wheat, sorghum, peas, rice, and vegetables) and some forage varieties, such as alfalfa, due to the lack of Se in the soil [8]. Several biofortification methods are used for plants, but the foliar application method is more effective than soil fertilization [9]. In recent years, the benefits that Se provides for the health of farm animals have been demonstrated, such as improving reproductive performance and immunological and homeostatic function [10]. Worldwide, several studies relate the absence of consumption of this element to decreased development, growth, and reproductive and metabolic problems, especially in ruminants [11, 12].

Nanotechnology and agriculture

Nanotechnology is a multidisciplinary scientific research area that works on the design, characterization, manufacturing, and application of structures created by controlled size and shape manipulation at the nanometric scale (less than 100 nm) that produce complex systems in which they modify some specific characteristics [13]. Due to the above, many applications have been developed, including medicine, genetics, pharmaceuticals, and agriculture. In 2013, the FAO published information on the state of the art of nanotechnology in food and agriculture. Its report documented ten-year research focused on sustainability and solutions to environmental problems. These investigations focus on releasing active ingredients (disease management and crop protection), minimizing fertilization loss and increasing yield, and producing bionano compounds from traditional crops [14]. Reports of such research on possible applications in agriculture have been both positive and negative. However, there are reports in which favorable results have been found, for example, in the case of fertilization, germination, growth promoters, sensors of contaminants and other substances, and recovery and treatment of soil and water [15, 16]a. Our knowledge about the interactions of nanoparticles for agricultural use with the environment still needs to be improved. Due to its complexity, there is still a long way to go to understand it completely. The UN 2013 raised the need to increase food production since by 2050, there is an estimated population of 9.1 billion people. There is currently a trend towards producing energy crops derived from efforts to mitigate climate change and due to the imminent reduction of hydrocarbon reserves. At the same time, in the expert forum organized in 2009 by FAO, "How to feed the world in 2050," it was proposed that it will be necessary to increase cereal production by 70% by 2050 [17]. Nanotechnology could provide tools for modern agriculture and even solve future problems related to food production and energy demand, focusing on sustainability. However, some NPs have been shown to affect the development and yields of crops.

Nanotechnology and caring for the environment

The current food system worldwide is not sustainable and, for decades, has caused environmental damage, including greenhouse gas emissions, the irrational use of drinking

water, the contamination of soils and aquifers by nitrogen and phosphorus, the use of pesticides, etc. However, recent research on environmental nanotechnology shows it can be a valuable tool for reducing environmental pollution. [18] developed nano encapsulates and nanocomposites for food and feed additives, biocides, pesticides, and contact materials. [19] applied nanomaterials as catalysts in phytoremediation processes and used stabilizers to improve their performance.

Applications of nanoparticles in agriculture

Applied science in agricultural systems presents a series of adversities and a wide range of challenges, such as a decrease in crop yields, low efficiency in the use of macro and micronutrients, a decrease in soil organic matter, deficiencies in multi-nutritional foods, climate change, the reduction in the availability of arable land, as well as the scarcity of water and labor for the countryside [20]. Various experiments have been carried out regarding the agricultural sector to know the optimal size, shape, and concentration of NPs to be applied to plants trying to improve their penetration and vascular translocation through the vascular bundles of the xylem and phloem.

The literature mentions that concentrations or doses less than 5 ppm of NPs can promote more significant plant growth. The size of the NPs is a relevant factor that significantly intervenes in the action of their penetration and translocation inside the plant tissues; therefore, the larger the size of the NPs, the less penetration will be into the vascular system in the plants when foliar spray techniques are used [21]. When applied to plant leaves, NPs can penetrate plant tissues as an aerosol or spray.

Evidence shows that 14.7% of the nanomaterial applied to plants is lost when using the spray or aerosol technique, compared to 32.5% or more with conventional agricultural products applied traditionally. An alternative is the application of foliar spray for the manipulation of nanoscale materials. When fertilizers are applied in a foliar way, they can easily penetrate the interior of the plants through the opening of the stomata, which have micrometric sizes fluctuating from 10 μm to 60 μm [22, 23].

Pastures and forages

Mexico has the eighth-largest cattle population in the world, with approximately 31 million heads, representing 2.31% of the world herd. However, Mexico has a greater diversity of ruminants, such as sheep and goats [24].

In the tropics of Mexico, the exploitation of livestock, specifically cattle, is carried out mainly by grazing native grasses and introducing grasses for livestock feeding. Pastures are grass and legume plants that grow in the pasture and serve to feed livestock. At the same time, forage is defined as any grass or legume harvested to be supplied as food to animals, whether green, dry, or processed. They can have different nutritional characteristics (hay, silage, stubble, saccharin, ammonification) [25, 26].

Selenium and Selenium nanoparticles

Selenium (Se) is a trace element that is scarce in the Earth's crust and is distributed in all parts of the Earth, being an element of volcanic origin. It accompanies sulfur and

progress, yield and nutritional value of crops, (3) biofortifying crops using Se to increase their content, (4) alleviating abiotic stress and (5) increase the nutraceutical quality of consumable foods [32].

Uptake of Se NPs in plants

Concerning the phyto-uptake of SeNPs and their translocation, determining the safety and toxicity of NPs provokes a deep understanding of their uptake in plants (Figure 2). There are different ways for the plant to capture this trace element in the form of NP. An absorption mechanism of SeNPs occurs through the cell wall and penetration of the plasma membrane. Only NPs with sizes smaller than 100 nm can enter through the different pores (stomata) and pass through the cell wall successfully. The plant cell walls act as a barrier, preventing the entry of external substances or materials, including NPs, into the plant cell walls [33-35]. It has been shown that NPs can adhere to surfaces such as plant roots and could influence plant chemical and physical absorption [33]. One of the explanations that demonstrates the most outstanding scientific solidity is the translocation of nanomaterials, which can move intracellularly and extracellularly between plant tissues until they reach the xylem [36]. Once the NPs are within the plant's vascular system, the designed NPs could be transported to the aerial parts along with plant water transpiration and nutritional flux in nutrient transmission.[37] applied selenium nanoparticles in grass plants, *Festuca arundinacea*, reporting significant differences in biomass parameters, root length, and leaf length ($p < 0.05$) between the treatments (1.5, 3.0, and 4.5 ppm NPs_{Se}) and the control. The increase in root length, leaf length, and biomass production of se-treated plants could be attributed to the application of selenium nanoparticles; Se stimulates the growth and development of plants because it improves the synthesis of photosynthetic pigments and organogenesis [3, 38]. Some authors [39] observed that foliar application of different amounts of SeNPs affected morphological characteristics such as shoot length, root size, and fresh weight of the bean plant. In other study [40] reported that by applying SeNPs foliar, plant height, fresh weight of the shoots, and root size were increased when concentrations of 10 mg SeNPs were used in wheat plants, attributing that the SeNPs stimulated the organogenesis and growth of the plant roots.




Figure 2. Uptake of Selenium NPs in plants, foliar, and soil route of adsorption.

Proximal composition of fortified forages

This research [41] found significant differences ($p < 0.05$) between the treatments in the variables of crude protein (CP), ether extract (EE), crude fiber (CF), neutral detergent fiber (NDF), acid detergent fiber (ADF), and Nitrogen Free Extract (NFE). The increase in PC content as a plant response to foliar biofortification of NPsSe could be manifested because Se promotes the increase in the synthesis of sulfur amino acids (Cys and Met) and selenium-aminoacids such as SeCys and Semet, which are incorporated into proteins. However, there is also evidence that other non-protein amino acids are synthesized, such as γ -glutamyl methyl selenium-cysteine (γ -gluMetSeCys), methyl-SeCys and methyl-Semet ([32, 42], where Se was used as a fertilizer, and increased protein was obtained in the root and leaf of alfalfa (*Medicago sativa* L.). Similarly, [43] found that foliar spraying of selenate produced an increase in the levels of protein content and total root nitrogen in radish (*Raphanus sativus*). Other research [44] found that foliar supplementation of sodium selenate in wheat plants (*Triticum aestivum* variety BRS 264) increased total nitrogen (up to 20%) compared to the control. In addition [45] reported that fertilization with Se promotes the increase in fatty acids (oleic, linoleic, and linoleic), increasing the concentration of lipids; however, the metabolic pathway is unclear. Furthermore [46] found that foliar application of sodium selenite and selenate in four rice genotypes (Ariete, Albatros, OP1105, and OP1109) increased the content of total lipids, as well as increased the concentration of oleic acid (C18:1), linoleic (C18:2), and palmitic acid (C16:0). Also, [47] mention that the synthesis of compounds such as the content of starch, total soluble sugars, and reducing sugars are enhanced, caused by the possible increase in the activity of carbohydrate metabolism enzymes. The results obtained in this study are similar to those reported by [42], where Se was used to improve carbohydrate metabolism, finding an increase in these compounds in alfalfa root and leaf (*Medicago sativa* L). In the same way, [44] found that the carbohydrate content in wheat plants increased when biofortified with selenium, showing this trend in each crop.

Use of nanoelements in Animal Nutrition

Nanotechnology and its applications can revolutionize agricultural production. This technology has had various applications in animal production, including the use of new molecular and cellular tools for animal reproduction, preservation of the animal's identity from birth to the consumer's table ("traceability"), biosecurity of food from animal origin, a better understanding of the phenomena that govern animal nutrition from dietary ingestion to the uptake and use of nutrients, and others [48].

In animal nutrition, as in other areas, it is possible to apply nanotechnology to obtain information about a nutrient or bioactive component and its release at specific sites of action, greater availability and maintenance of adequate levels for more extended periods, greater use of food [49]. Minerals are one of the most widely used supplements in animal nutrition; However, the form in which these minerals are found influences their bioavailability, so if they have low bioavailability, the animal will not use them correctly. An example is Selenium (Se), an essential trace element with a narrow range of beneficial and toxic effects. Recently, nano selenium has attracted the attention of

many researchers due to its high bioavailability and low toxicity. Nonmetric particles exhibit novel characteristics, such as a larger contact surface, high surface activity, high absorption capacity, and low toxicity.

CONCLUSIONS

Biofortification with Se NPs benefits forage grass's morphological and physicochemical characteristics. Se NPs improve root development and synthesis, growth, and grass yield. Applying Se NPs increases the content of crude protein and ether extract.

Author contributions: "Conceptualization, ETK and ABL; software, MPG; validation, MRI; research, GLU; resources, PRS; writing—preparation of the original draft, ETK, and MPG; writing—review and editing. All authors have read and accepted the published version of the manuscript." Conflicts of interest: "The authors declare that they have no conflict of interest."

REFERENCES

1. Nations, F.a.A.O.o.t.U. Situación Alimentaria Mundial. Recuperado el 02 de 02 de 2017, de <http://www.fao.org/worldfoodsituation/foodpricesindex/es/>. 2007
2. Chen, L., *et al.*, Improved forage digestibility of tall fescue (*Festuca arundinacea*) by transgenic down regulation of cinnamyl alcohol dehydrogenase. *Plant Biotechnology Journal*, 2003. 1(6): p. 437-449.
3. Hartikainen, H., Biogeochemistry of selenium and its impact on food chain quality and human health. *Journal of Trace elements in Medicine and Biology*, 2005. 18(4): p. 309-318.
4. Germ, M., V. Stibilj, and I. Kreft, Metabolic importance of selenium for plants. *Eur. J. Plant Sci. Biotechnol*, 2007. 1(10): p. 91-97.
5. Ducsay, L. and O. Lozek, Effect of selenium foliar application on its content in winter wheat grain. *Plant Soil and Environment*, 2006. 52(2): p. 78.
6. Lira Saldivar, R.H., *et al.*, Potencial de la nanotecnología en la agricultura. *Acta universitaria*, 2018. 28(2): p. 9-24.
7. Bamberg, S.M., *et al.*, Effects of selenium (Se) application and arbuscular mycorrhizal (AMF) inoculation on soybean ('*Glycine max*') and forage grass ('*Urochloa decumbens*') development in oxisol. *Australian Journal of Crop Science*, 2019. 13(3): p. 380-385.
8. Moteszarezhadeh, B., S. Ghorbani, and H.A. Alikhani, The effect of selenium biofortification in alfalfa (*Medicago sativa*). *Journal of Plant Nutrition*, 2020. 43(2): p. 240-250.
9. Ros, G., *et al.*, Selenium fertilization strategies for bio-fortification of food: an agro-ecosystem approach. *Plant and Soil*, 2016. 404: p. 99-112.
10. Nedelkov, K., *et al.*, Alternative selenium supplement for sheep. *Animal Feed Science and Technology*, 2020. 261: p. 114390.
11. Meyer, U., *et al.*, Influence of various selenium sources on selenium concentration in the milk of dairy cows. *Journal für Verbraucherschutz und Lebensmittelsicherheit*, 2014. 9: p. 101-109.
12. Wang, Z., *et al.*, Effect of different levels of selenium yeast on the antioxidant status, nutrient digestibility, selenium balances and nitrogen metabolism of Tibetan sheep in the Qinghai-Tibetan Plateau. *Small ruminant research*, 2019. 180: p. 63-69.
13. Bawa, R., *et al.*, Protecting new ideas and inventions in nanomedicine with patents. *Nanomedicine: Nanotechnology, Biology and Medicine*, 2005. 1(2): p. 150-158.
14. Parisi, C., M. Vigani, and E. Rodríguez-Cerezo, Agricultural nanotechnologies: what are the current possibilities? *Nano Today*, 2015. 10(2): p. 124-127.
15. Han, D., Arsenic removal by novel nanoporous adsorbents-kinetics, equilibrium and regenerability. Report. Environmental and water Resources Division, Texas A&M University, 2007.
16. Stoimenov, P.K., *et al.*, Metal oxide nanoparticles as bactericidal agents. *Langmuir*, 2002. 18(17): p. 6679-6686.
17. Candel, J.J., Food security governance: A systematic literature review. *Food Security*, 2014. 6: p. 585-601.
18. Peters, R., *et al.*, Inventory of Nanotechnology applications in the agricultural, feed and food sector. *EFSA Supporting Publications*, 2014. 11(7): p. 621E.

19. Ibrahim, R.K., *et al.*, Environmental application of nanotechnology: air, soil, and water. *Environmental Science and Pollution Research*, 2016. 23: p. 13754-13788.
20. Smith, P., *et al.*, Policy and technological constraints to implementation of greenhouse gas mitigation options in agriculture. *Agriculture, Ecosystems & Environment*, 2007. 118(1-4): p. 6-28.
21. Adhikari, T., S. Kundu, and A.S. Rao, Zinc delivery to plants through seed coating with nano-zinc oxide particles. *Journal of Plant Nutrition*, 2016. 39(1): p. 136-146.
22. Hawthorne, J., *et al.*, Particle-size dependent accumulation and trophic transfer of cerium oxide through a terrestrial food chain. *Environmental science & technology*, 2014. 48(22): p. 13102-13109.
23. Fricker, M. and C. Willmer, Stomata. 2012: Springer Science & Business Media.
24. Perozo, A., Manejo de pastos y forrajes tropicales. 2013: Ali David Perozo Bravo.
25. Vera, R.R., The future for savannah and tropical grasslands: a Latin American perspective, in Grasslands. 2019, CRC Press. p. 516-539.
26. Wilkins, R., Forages and their role in animal systems. Forage Evaluation in Ruminant Nutrition, CAB International, 2000: p. 1-14.
27. Acosta, L., El selenio. Acosta L. Laboratorio Santa Elena, Uruguay: Disponible en: http://www.produccionanimal.com.ar/suplementacion_mineral/95-selenio.pdf. Fecha de consulta, 2007. 8(01): p. 2014.
28. Bentancur Arreseigor, S., L. Capdevielle Trindade, and A.C. Ferreira Deragon, Efecto de la suplementación con selenio sobre el desempeño reproductivo de vacas de cría pastoreando campo natural de basalto. 2015.
29. Barchielli, G., A. Capperucci, and D. Tanini, The role of selenium in pathologies: An updated review. *Antioxidants*, 2022. 11(2): p. 251.
30. Garduño, G.V., *et al.*, Role of selenium in sheep feeding: antioxidant effect and repercussion in meat shelf-life. *Nutritional strategies of animal feed additives*: p. 131.
31. Gaucin-Delgado, J.M., *et al.*, La biofortificación con selenio mejora los compuestos bioactivos y la actividad antioxidante en chile jalapeño. *Revista mexicana de ciencias agrícolas*, 2021. 12(8): p. 1339-1349.
32. Garza-García, J.J., *et al.*, The role of selenium nanoparticles in agriculture and food technology. *Biological Trace Element Research*, 2021: p. 1-21.
33. Hu, T., *et al.*, Absorption and bio-transformation of selenium nanoparticles by wheat seedlings (*Triticum aestivum* L.). *Frontiers in Plant Science*, 2018. 9: p. 597.
34. Aslani, F., *et al.*, Effects of engineered nanomaterials on plants growth: an overview. *The Scientific World Journal*, 2014. 2014.
35. Corredor, E., *et al.*, Nanoparticle penetration and transport in living pumpkin plants: in situ subcellular identification. *BMC plant biology*, 2009. 9: p. 1-11.
36. Rajae Behbahani, S., *et al.*, Red elemental selenium nanoparticles mediated substantial variations in growth, tissue differentiation, metabolism, gene transcription, epigenetic cytosine DNA methylation, and callogenesis in bittermelon (*Momordica charantia*); an *in vitro* experiment. *PloS one*, 2020. 15(7): p. e0235556.
37. González-Montiel, L., *et al.*, Bioactive Compounds, Antioxidant and Antimicrobial Activity of Propolis Extracts during *In Vitro* Digestion. *Applied Sciences*, 2022. 12(15): p. 7892.
38. Feng, T., *et al.*, Selenium improves photosynthesis and protects photosystem II in pear (*Pyrus bretschneideri*), grape (*Vitis vinifera*), and peach (*Prunus persica*). *Photosynthetica*, 2015. 53: p. 609-612.
39. Ragavan, P., A. Ananth, and M. Rajan, Impact of selenium nanoparticles on growth, biochemical characteristics and yield of cluster bean *Cyamopsis tetragonoloba*. *International Journal of Environment, Agriculture and Biotechnology*, 2017. 2(6): p. 238983.
40. Ikram, M., *et al.*, Foliar applications of bio-fabricated selenium nanoparticles to improve the growth of wheat plants under drought stress. *Green Processing and Synthesis*, 2020. 9(1): p. 706-714.
41. González-Lemus, U., *et al.*, Nutritional Parameters, Biomass Production, and Antioxidant Activity of *Festuca arundinacea* Schreb. Conditioned with Selenium Nanoparticles. *Plants*, 2022. 11(17): p. 2326.
42. Hajiboland, R., *et al.*, Selenium-induced enhancement in carbohydrate metabolism in nodulated alfalfa (*Medicago sativa* L.) as related to the glutathione redox state. *Soil science and plant nutrition*, 2015. 61(4): p. 676-687.
43. Schiavon, M., *et al.*, Selenium biofortification in radish enhances nutritional quality via accumulation of methyl-selenocysteine and promotion of transcripts and metabolites related to glucosinolates, phenolics, and amino acids. *Frontiers in Plant Science*, 2016. 7: p. 1371.
44. Lara, T.S., *et al.*, Selenium biofortification of wheat grain via foliar application and its effect on plant metabolism. *Journal of Food Composition and Analysis*, 2019. 81: p. 10-18.

45. García Márquez, V., *et al.*, Ionic selenium and nanoselenium as biofortifiers and stimulators of plant metabolism. *Agronomy*, 2020. 10(9): p. 1399.
46. Lidon, F.C., *et al.*, Selenium biofortification of rice grains and implications on macronutrients quality. *Journal of Cereal Science*, 2018. 81: p. 22-29.
47. Kaur, M., S. Sharma, and D. Singh, Influence of selenium on carbohydrate accumulation in developing wheat grains. *Communications in Soil Science and Plant Analysis*, 2018. 49(13): p. 1650-1659.
48. Scott, N., Nanotechnology and animal health. *Revue Scientifique Et Technique-Office International Des Epizooties*, 2005. 24(1): p. 425.
49. Ross, S.A., *et al.*, New technologies for nutrition research. *The Journal of nutrition*, 2004. 134(3): p. 681-685.

