



AgEcon SEARCH

RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

AGRO PRODUCTIVIDAD

Heliconias (Heliconiaceae) in rural landscapes

pág. 105

Año 17 • Volumen 17 • Número 7 • julio, 2024

- | | |
|---|----|
| <i>Spondias dulcis</i> propagation by seeds and stem cuttings | 3 |
| Variation of the nutritional content of different genotypes of <i>Lotus corniculatus</i> L. under optimum and sub-optimum soil moisture conditions throughout the seasons of the year | 13 |
| Kinetics of ammonium volatilization in the form of ammonia in soils through a textural gradient | 23 |
| Yield and nutritional value of <i>Cenchrus purpureus</i> VC Maralfalfa grass | 35 |
| Analysis of the natural regeneration of the white mangrove (<i>Laguncularia racemosa</i> (L.) C.F. Gaertn. with dasometric variables during the drought season | 43 |
| Evaluation of three forages as a source of fiber in diets of fattening rabbits in Aguascalientes, Mexico | 49 |

y más artículos de interés...



Colegio de
Postgraduados

Kinetics of ammonium volatilization in the form of ammonia in soils through a textural gradient

Perez-Rosales, Alejandro^{1*}

¹ Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias. Campo Experimental Zacatepec. Carr. Zacatepec-Galeana, km 0.5. Col. Centro. Zacatepec, Morelos. C.P. 62780.

* Correspondence: fitotecn@hotmail.com

ABSTRACT

Objective: To estimate the ammonia volatilization rate, using a texture gradient with increasing doses of ammonium in alkaline soils.

Design/Methodology/Approach: The experiment took place in the Análisis Químico de Suelos lab of the Soil Department of the Universidad Autónoma Chapingo. The incubations consisted of 25 g of soil from the former Texcoco lake, with 12.5% clay. The soil was air-dried and sieved with a no. 10 mesh. The soil was mixed with three different concentrations (17.5%, 22.5%, and 32.5%) of bentonite, in order to increase clay content. The mixtures were placed in plastic containers with airtight seals. Twelve-point five mL of an ammonium sulfate solution (with 150, 300, 450, 600, and 750 mg of nitrogen kg soil⁻¹) was added. The ammonia was recovered in a container with a boric acid solution. Volumetry was used to quantify ammonium. A completely randomized design with two factors (clay content and ammonium dose) was used. Data were analyzed with a regression analysis, analysis of variance, and Tukey Multiple Comparison Test, using the SAS OnDemand for Academics software.

Results: The ammonia volatilization has a linear trend, with the concentration of the applied nitrogen. The volatilization rate ranged from 0.02 to 0.03 mg of ammonium per milligram of the nitrogen applied per kg soil⁻¹. Significant statistical differences were recorded between the effect of the N dose and the clay content on the ammonia volatilization rate.

Study Limitations/Implications: Clay content in the soil and ammonia volatilization rate can be used as an indicator to estimate ammonia losses and the ammonia adsorption capacity of the soil.

Findings/Conclusions: Ammonia volatilization is independent from clay content and takes place immediately after its application. It has a linear trend regarding the ammonia dose applied. Ammonia volatilization rate decreases (asymptotic trend) as the clay content increases in the soil.

Keywords: clay, ammonium, alkaline, cationic exchange capacity.

Citation: Perez-Rosales, A. (2024). Kinetics of ammonium volatilization in the form of ammonia in soils through a textural gradient. *Agro Productividad*. <https://doi.org/10.32854/agrop.v17i7.2661>

Academic Editor: Jorge Cadena Iñiguez

Guest Editor: Juan Franciso Aguirre Medina

Received: August 07, 2023.

Accepted: June 21, 2024.

Published on-line: August 08, 2024.

Agro Productividad, 17(7). July. 2024. pp: 23-33.

This work is licensed under a Creative Commons Attribution-Non-Commercial 4.0 International license.



INTRODUCTION

The nitrogen (N) available for plants in the soil is the highest limiting factor (Martínez-Dalmau *et al.*, 2021). Consequently, during the last decade, the annual use of nitrogen fertilizers has increased by $\approx 1.4\%$ worldwide; however, their recovery efficiency is $< 50\%$ (Panday *et al.*, 2020; Mahmud *et al.*, 2021). Approximately 80% of the N used in agriculture is lost as a result of leaching, runoff, and gaseous emissions. The predominant gases are nitrous oxide (N₂O-N) and ammonia (NH₃-N), with 4 Tg N year⁻¹ and 12 Tg N year⁻¹, respectively (Krol *et al.*, 2020). The NH₃-N volatilization accounts for 10-60% of the total N applied as ammonium fertilizer. This process takes place a few days after the application (Chu *et al.*, 2020).

The $\text{NH}_3\text{-N}$ volatilization is the result of the ammonium ($\text{NH}_4\text{-N}$) adsorbed by the colloids and its interaction with the physical and chemical properties of the soil (Wang *et al.*, 2023). Volatilization is highly sensitive to pH and $\text{NH}_4\text{-N}$ concentration. The increases of pH promote the formation of $\text{NH}_3\text{-N}$ (Shi *et al.*, 2024) —*i.e.*, a >7 pH increases the formation of $\text{NH}_3\text{-N}$, as a consequence of the high concentration of the hydroxyl radical (HR) that reacts with the $\text{NH}_4\text{-N}$ (Skorupka and Nosalewicz, 2021). This dependency between pH and the $\text{NH}_4\text{-N}/\text{NH}_3\text{-N}$ balance (Pelster *et al.*, 2019) increases exponentially the formation of $\text{NH}_3\text{-N}$ (Ohnemus *et al.*, 2021). In calcareous soils, the losses account for 50% (Powlson and Dawson, 2021).

Clays, cation exchange capacity (CEC), texture, organic matter (Cassity-Duffey *et al.*, 2020), and damping capacity of soils determine the $\text{NH}_3\text{-N}$ volatilization rate, because they correlate with higher $\text{NH}_4\text{-N}$ adsorption rates. Clay soils have a lower volatilization than sandy soils (Pelster *et al.*, 2019; Mazloomi and Jalali, 2019): the amount of N lost in the volatilization is significantly influenced by soil texture and it is higher in coarse-textured soils than in fine-textured soils (Al-Badrani *et al.*, 2023). The $\text{NH}_4\text{-N}$ adsorption and desorption in the soil is highly dependent of the type and quantity of clays; these minerals have chemical reactivity and are grouped according to their silicon tetrahedra (1:1) and aluminum octahedra (2:1). The 2:1 type clays have a high CEC and a higher specific surface than the 1:1 type clays, increasing their $\text{NH}_4\text{-N}$ adsorption ($350\text{-}3,800 \text{ kg ha}^{-1}$ at ≤ 30 cm of depth); vermiculites, illites, and smectites predominate in these soils (Kome *et al.*, 2019). Consequently, the objective of this research was to estimate the ammonia volatilization rate, using a textural gradient and increasing doses of ammonium in alkaline soils.

MATERIALS AND METHODS

The experiment was carried out in the Análisis Químico lab of the Soil Department of the Universidad Autónoma Chapingo. To determine the volatilization rate, soils that had been previously air-dried and sieved with a no. 10 mesh were incubated.

Increasing concentrations of bentonite were added to the soil samples (texture gradient) to increase clay content. Bentonite is an expandible clay of the montmorillonite group (2:1) that has strong affinity and cation adsorption capacity in its internal and external layers (specific surface) (Alexander *et al.*, 2019). Bentonite has the following characteristics: $\text{pH}_{10:1}=9.42$, $\text{EC}_{10:1}=1.36 \text{ dSm}^{-1}$, $\text{CEC}=63.23 \text{ meq } 100 \text{ g}^{-1} \text{ S}$, carbonates $=1.5 \text{ mmol}_c \text{ L}^{-1}$, bicarbonates $=82.49 \text{ mmol}_c \text{ L}^{-1}$, and clay (particles $>0.002 \text{ mm}$) $=95.84\%$. Adding this mineral allowed the simulation of soils with higher contrasts regarding clay content (17.5%, 22.5%, and 32.5%). Soil without bentonite was used in the case of the 12.5% concentration. The soil samples collected from the former Texcoco lake had the following characteristics: 12.5% clay obtained with the pipette method (Van Reeuwijk, 2003); 9.6 pH, $389 \mu\text{Scm}^{-1}$ EC, and 32.0 mg kg^{-1} phosphorous (SEMARNAT, 2002); $8.78 \text{ Cmol kg S}^{-1}$ CEC (Pleyser *et al.*, 1986); and 1.60 g kg^{-1} soluble organic carbon (Harvey *et al.*, 2009). Once the sample and the bentonite were mixed, 25 g were weighted and placed in $15 \text{ cm} \times 20 \text{ cm}$ transparent polypropylene bags with airtight seals. An ammonium sulfate solution (12.5 mL) was added to the bags. This solution was

determined based on the effective porosity saturation (Wallach, 2019); the percentage used in this experiment was 50% (W/V).

The N doses applied were based on the proposals of Liu *et al.*, 2007 and Fan *et al.*, 2011. Ammonium sulfate was used as source of N (150, 300, 450, 600, and 750 mg N kg soil⁻¹). A modified version of the quantification methodology proposed by Ahmed *et al.* (2008), Chen *et al.* (2014) and Palanivell *et al.* (2015) for the recovery of ammonium was used: a 100 mL container with a 50 mL boric acid solution (20 g L⁻¹) was mixed with the bromocresol green and methyl red indicators and placed in the airtight sealed bag with the soil sample (previously treated with ammonium sulfate). Finally, the bag was airtight sealed to prevent water loss and gas and it was incubated at room temperature for three days. Volumetry was used to determine the amount of ammonium recovered from the boric acid solution (Shen *et al.*, 2020); in addition, the sulfuric acid concentration was adjusted to 0.005 N. The experiment was carried out using a completely randomized design with two factors (NH₄-N concentration and clay content of the soil) and three replicates. The analysis of the volatilization rate was carried out with a simple linear regression analysis. The NH₄-N doses were analyzed per clay level. The linear regression analysis determined the NH₃-N volatilization rate. Subsequently, a graph of these volatilization was developed per clay level, in order to evaluate the influence of clay content on the NH₃-N volatilization. The recovered NH₃ (mg) was subjected to an analysis of variance and to a Tukey Multiple Comparison Test, with the SAS OnDemand for Academics software.

RESULTS AND DISCUSSION

Effect of the clay content on the soil and the NH₄-N dose on the NH₃-N volatilization According to the analysis of variance and the regression analysis for the five NH₄-N levels in each clay layer, when $\alpha=0.01$ (Table 1), the behavior of the NH₃ volatilization has a proportional linear trend as the NH₄-N levels applied increase, given the positive correlation between NH₃-N emissions and NH₄-N concentration (Li *et al.*, 2020).

After the samples had been incubated for three days, a linear trend was identified between the concentration of NH₄-N applied and the amount of NH₃-N recovered, regardless of the clay content; this linear behavior is found in each clay layer evaluated, because high NH₄-N doses increase NH₃-N emissions (Wan *et al.*, 2021). The NH₃-N volatilization rate ranged from 0.02 to 0.03 mg per mg of N applied per kg of soil, with

Table 1. Analysis of variance of linear regression for four soil clay contents and five NH₄-N doses on NH₃-N volatilization.

Source	DF	Sum of squares			
		12.5	17.5	22.5	32.5
Clay (%)					
Reg	1	500.89**	430.56**	331.95**	282.87**
Error	13	0.48	0.57	0.61	0.69
Total	14				

DF: Degrees of Freedom; * $\alpha=0.05$; ** $\alpha=0.01$; NS: Not Significant.

>97% correlation coefficients (Figure 1). Minato *et al.* (2020) mention that the accumulated $\text{NH}_3\text{-N}$ volatilization has a sigmoid pattern with gradual increases. Its first stage had a high $\text{NH}_3\text{-N}$ emission rate, reaching its maximum point 72 h after the application ($0.4 \text{ kg ha}^{-1} \text{ day}^{-1} \text{ NH}_3\text{-N}$). It remained constant during the first 5 days.

The $\text{NH}_3\text{-N}$ losses recorded during the incubation with the various N doses (150, 300, 450, 600, and $750 \text{ mg}\cdot\text{kg soil}^{-1}$) suggest that, under experimental conditions, this situation takes place immediately after application and that, during the first stages, the soil lacks the capacity to absorb $\text{NH}_4\text{-N}$ from the clay. Viero *et al.* (2014) report a similar trend: a maximum loss of 0.40 and $0.53 \text{ kg ha}^{-1} \text{ day}^{-1}$ was registered 5-10 days after the application. Dari and Rogers (2022) determined that maximum volatilization takes place from 4 to 8 days after the application (8 to 11% of the total N applied). These emissions are generated by the reactivity of $\text{NH}_4\text{-N}$ to the OH^- content in the soil —*i.e.*, the post-application volatilization is mostly a chemical process, since the increase from 6 to 10 in pH increases the anionic association sites that react to $\text{NH}_4\text{-N}$, reducing the $\text{NH}_4\text{-N}$ concentration and, in turn, the adsorption capacity of soil clays (Fan *et al.*, 2021).

The analysis of variance found significant differences on the effect on the $\text{NH}_3\text{-N}$ volatilization of clay content, $\text{NH}_4\text{-N}$ doses, and the interaction between clay content and $\text{NH}_4\text{-N}$ doses (Table 2). Da Cruz Corrêa *et al.* (2021) established that an increase in the N doses results in a linear increase of volatilization, irrespective of the ammonium source. The $\text{NH}_4\text{-N}$ adsorption increases as clay content in the soil increases, given the negative correlation between the $\text{NH}_3\text{-N}$ volatilization, the clay content in the soil, and the CEC (Chuong *et al.*, 2020).

NH_3 volatilization is a chemical reaction determined by the pH of the soil; however, the clay content in the soil tends to diminish volatilization, since the ammonium applied in alkaline soils and soils with many clay layers or high CEC levels is absorbed by the sites where these clays are exchanged (Wester-Larsen *et al.*, 2022).




Figure 1. Linear regression model for the effect on ammonia volatilization of five levels of ammonium concentration and four levels of clay concentration found in the soil.

Table 2. Analysis of variance of the effect of four soil clay contents and five $\text{NH}_4\text{-N}$ doses on $\text{NH}_3\text{-N}$ volatilization.

Source	DF	SS (NH_3 mg)	F _c
Soil clay content (A)	3	98.9925	53.73**
Doses of ammonium sulphate (B)	4	1528.1210	622.03**
AB	12	23.0417	3.13**
Error	40	24.5667	
Total		1674.7218	

DF: Degrees of Freedom; SS: Sum of Squares; * $\alpha=0.05$; ** $\alpha=0.01$; NS: Not Significant.

The multiple comparison of means determined that a greater clay content in the soil diminished $\text{NH}_3\text{-N}$ volatilization. Likewise, statistical differences were found for each level evaluated (Table 3). An increase in the clay content of the soil increases cation exchange sites, enhancing the capacity of the soil to adsorb $\text{NH}_4\text{-N}$. Foerid *et al.* (2019) point out that the application of 10 t ha^{-1} zeolite increases NH_4 adsorption, without limiting the amount available for the crop.

Each level of the $\text{NH}_4\text{-N}$ doses applied had statistical differences regarding $\text{NH}_3\text{-N}$ volatilization. The $750 \text{ mg NH}_4\text{-N kg soil}^{-1}$ dose generates the largest volatilized $\text{NH}_3\text{-N}$; likewise, there is a proportional relation between the $\text{NH}_4\text{-N}$ dose applied and the $\text{NH}_3\text{-N}$ generated (Table 4). These results match the findings of Wang *et al.* (2021) who mentioned that $\text{NH}_3\text{-N}$ volatilization (4.3 mg, 5.27 mg, and 6.32 mg) gradually increases as the N dose applied increases (26.6 mg, 48.3 mg, and 69.0 mg).

The clay content of the soil and the application of ammonium sulfate showed statistical differences regarding $\text{NH}_3\text{-N}$ volatilization: 750 and 600 mg kg soil^{-1} doses recovered

Table 3. Effect of soil clay content on $\text{NH}_3\text{-N}$ volatilization.

Soil clay content (g kg S^{-1})	NH_3 (mg)
125.0	17.7333 a*
175.0	13.5467 b
225.0	12.6067 c
325.0	11.2267 d

* Means with the same letter are statistically equal (Tukey, $\alpha=0.05$). MSD=0.767.

Table 4. Effect of ammonium sulfate application on ammonia volatilization.

Doses of ammonium sulphate (mg kg S^{-1})	NH_3 (mg)
750	20.1417 a*
600	16.4167 b
450	13.3083 c
300	9.5167 d
150	5.7583 e

* Means with the same letter are statistically equal (Tukey, $\alpha=0.05$). MSD=0.9138.

more $\text{NH}_3\text{-N}$, an amount that gradually decreases as the N dose decreases (Table 5). Xie *et al.* (2019) reported a similar phenomenon: $\text{NH}_3\text{-N}$ volatilization increases as a result of the fertilization dose applied.

Effect of the clay content in the soil on the $\text{NH}_3\text{-N}$ volatilization rate

Figure 2 shows a slight decrease in volatilization speed as the clay content in the soil increases, based on the volatilization rates ($27.2, 25.3, 22.2,$ and $20.5 \mu\text{g NH}_3 \text{ mg NH}_4\text{-N kg soil}^{-1}$) with the clay content ($125, 175, 225,$ and $325 \text{ g kg soil}^{-1}$), which seemingly indicates an asymptotic trend for the study interval. This volatilization has a negative relation with the CEC and the clay content and its trend is adjusted to an exponential regression model (Hearn *et al.*, 2023), since bentonite is a 2:1 clay with a strong capacity to adsorb $\text{NH}_4\text{-N}$ ($19.01 \text{ mg NH}_4\text{-N g}^{-1}$ of clay) and high CEC ($40\text{-}130 \text{ Cmol kg}^{-1}$). $\text{NH}_4\text{-N}$ adsorption in a soil to which bentonite has been added is a result of its electrostatic interaction with the negative charges of the surface, resulting from isomorphous replacement (Han *et al.*, 2020). Mazloomi and Jalali (2019) studied the relation between adsorbed $\text{NH}_4\text{-N}$ and the concentration of the $\text{NH}_4\text{-N}$ applied in soils treated with vermiculite and determined that the $\text{NH}_4\text{-N}$ adsorption capacity of soils with 0% vermiculite is 285 mg kg^{-1} . It increases by 10.5%, 14%, and 24.6%, with

Table 5. Effect of soil clay content and ammonium sulfate application on ammonia volatilization.

Soil clay content (g kg S^{-1})	Doses of ammonium sulphate (mg kg S^{-1})	NH_3 (mg)
125	750	22.80 a
175	750	20.93 ab
225	750	19.40 bc
125	600	18.60 bc
325	750	17.43 cd
175	600	17.37 cde
225	600	15.80 def
125	450	15.00 efg
325	600	13.90 fgh
175	450	13.80 fgh
225	450	12.83 ghi
325	450	11.60 hij
125	300	11.20 ij
175	300	9.87 jk
225	300	8.60 kl
325	300	8.40 klm
225	150	6.40 lmn
125	150	6.07 mn
175	150	5.77 n
325	150	4.80 n

* Means with the same letter are statistically equal (Tukey, $\alpha=0.05$). MSD=2.4241.




Figure 2. Effect of the clay content on the ammonia volatilization rate.

the addition of 2%, 4%, and 8%, respectively. These results indicate that the adsorption capacity of the soil is directly dependent on clay, which can be interpreted through the CEC.

Significant differences were found as a result of the analysis of variance of the effect of clay content on NH₃-N volatilization. These results suggest that ammonia volatilization speed tends to decrease with a higher clay content; however, these losses continue (Table 6), since the adsorption capacity of bentonite behaves according to the Langmuir adsorption isotherm (Zaini *et al.*, 2021). Likewise, Linggi *et al.* (2020) mention that the capacity of various soils and clays (*e.g.*, bentonite) to adsorb ammonium can be explained with a non-linear model, using the Langmuir isotherm.

Significant differences were recorded in the effect of the clay content of the soil on the NH₃-N volatilization rate: the lowest clay content in the soil (125.0 g kg soil⁻¹) is more volatile (27.2409 µg NH₃ mg NH₄-N kg soil⁻¹) (Table 7). The decrease in the NH₃-N volatilization rate is the result of the NH₄-N adsorption in the exchange sites. This phenomenon prevents their reaction with the OH⁻ found in the soil solution, because bentonite has a wide specific surface, high CEC, and a high cation adsorption capacity (Barakan and Aghazadeh., 2020). In addition, it has negative charges generated through isomorphous replacement, which are not affected by pH (Jolin *et al.*, 2020).

Table 6. Analysis of variance of the effect of soil clay content on ammonium volatilization rates with three days of incubation with ammonium sulfate $\alpha=0.01$.

Source	DF	SS (µg NH ₃ mg NH ₄ -N kg S ⁻¹)	F _c
Tratamientos	3	83.03	12.26**
Error	8	18.07	
Total	11	101.10	

DF: Degrees of Freedom; SS: Sum of Squares; * $\alpha=0.05$; ** $\alpha=0.01$; NS: Not Significant.

Table 7. Effect of soil clay content and on the NH₃-N volatilization rate.

Soil clay content (g kg S ⁻¹)	Rate of ammonia volatilization ($\mu\text{g NH}_3$ mg NH ₄ -N kg S ⁻¹)
125.0	27.2409 a*
175.0	25.2560 ab
225.0	22.1760 bc
325.0	20.4711 c

* Means with the same letter are statistically equal (Tukey, $\alpha=0.05$). MSD=3.9292.

CONCLUSIONS

Ammonium losses are recorded immediately after the application of ammonium sulfate, regardless of the clay content in the soil and they are proportional to the amount of ammonium that was applied. Significant differences have been recorded regarding the amount of ammonia produced as a result of the increase in the clay content in the soil and the ammonium doses. Clay content in the soil reduces the speed of ammonia volatilization —*i.e.*, a greater clay content in the soil reduces the ammonia volatilization rate, as a consequence of the increase in the ion exchange sites in which the ammonium found in the solution is adsorbed. Ammonia volatilization takes place in alkaline soils and is more accentuated in sandy soils. Therefore, under these conditions, the use of ammonium sources must be limited and complemented with the application of soil improvers (organic carbon, fragments of recalcitrant carbon, and soil acidifiers) that increase exchange sites, in order to increase ammonium adsorption in the soil.

The ammonia volatilization rate helps to evaluate ammoniacal nitrogen sources, doses, and corrections, in order to increase the time that ammonium remains in the soil, as well as the efficiency of its recovery.

Clay content in the soil (measured as CEC) is an indicator of the ammonium adsorption capacity of alkaline soils. Although the NH₃-N volatilization is ruled by the pH, it is minimized by the type and amount of clay.

ACKNOWLEDGMENTS

The authors would like to thank CONAHCyT for funding this research, as part of the 2018 postdoctoral fellowship, to Departamento de Suelos of the Universidad Autónoma Chapingo and Génesis, Morfología y Clasificación Suelos lab of the Colegio de Postgraduados for the use of their experimental facilities.

REFERENCES

- Ahmed, O. H., Husin, A., & Hanif, A. H. M. (2008). Ammonia volatilization and ammonium accumulation from urea mixed with zeolite and triple superphosphate. *Acta Agriculturae Scandinavica. Section B, Soil and Plant Science*, 58(2), 182-186. <https://doi.org/10.1080/09064710701478271>
- Al-Badrani, W. A. A., Hassan, F. A., & Harbawi, M. A. (2023). Effect of Agricultural Sulfur and Poultry Manure on Ammonia Volatilization from Urea Fertilizer in two different Soil Textures. *International Journal of Agricultural and Statistics Sciences/International Journal of Agricultural and Statistical Sciences*, 19(01 (Supp)), 1153-1159. <https://doi.org/10.59467/ijass.2023.19.1153>
- Alexander, J. A., Zaini, M. A. A., Abdulsalam, S., Aliyu, E. U., & Omeiza, A. U. (2018). Surface modification of low-cost bentonite adsorbents—A review. *Particulate Science and Technology*, 37(5), 538-549. <https://doi.org/10.1080/02726351.2018.1438548>

- Barakan, S., & Aghazadeh, V. (2020). The advantages of clay mineral modification methods for enhancing adsorption efficiency in wastewater treatment: a review. *Environmental Science and Pollution Research International*, 28(3), 2572-2599. <https://doi.org/10.1007/s11356-020-10985-9>
- Cassidy Duffey, K., Cabrera, M. L., Franklin, D. A., Gaskin, J. W., & Kissel, D. E. (2020). Effect of soil texture on nitrogen mineralization from organic fertilizers in four common southeastern soils. *Soil Science Society of America Journal*, 84(2), 534-542. <https://doi.org/10.1002/saj2.20039>
- Chen, A., Lei, B., Hu, W., Lu, Y., Mao, Y., Zongyan, D., & Shi, Z. (2014). Characteristics of ammonia volatilization on rice grown under different nitrogen application rates and its quantitative predictions in Erhai Lake Watershed, China. *Nutrient Cycling in Agroecosystems*, 101(1), 139-152. <https://doi.org/10.1007/s10705-014-9660-7>
- Chu, Q., Xu, S., Xue, L., Liu, Y., Feng, Y., Yu, S., Li, Y., & Xing, B. (2020). Bentonite hydrochar composites mitigate ammonia volatilization from paddy soil and improve nitrogen use efficiency. *Science of the total environment*, 718, 137301. <https://doi.org/10.1016/j.scitotenv.2020.137301>
- Chuong, T., Plant, R. E., & Linquist, B. A. (2020). Fertilizer source and placement influence ammonia volatilization losses from water seeded rice systems. *Soil Science Society of America Journal*, 84(3), 784-797. <https://doi.org/10.1002/saj2.20074>
- Da Cruz Corrêa, D. C., Da Silva Cardoso, A., Ferreira, M. R., Simiscalchi, D., Gonçalves, P. H., Lumasini, R. N., Reis, R. A., & Ruggieri, A. C. (2021). Ammonia Volatilization, Forage Accumulation, and Nutritive Value of Marandu Palisade Grass Pastures in Different N Sources and Doses. *Atmosphere*, 12(9), 1179. <https://doi.org/10.3390/atmos12091179>
- Dari, B., & Rogers, C. W. (2021). Ammonia volatilization from fertilizer sources on a loam soil in Idaho. *Agrosystems, Geosciences & Environment*, 4(3). <https://doi.org/10.1002/agg2.20192>
- Fan, T., Wang, M., Wang, X., Chen, Y., Wang, S., Zhan, H., Chen, X., Lu, A., & Zha, S. (2021). Experimental Study of the Adsorption of Nitrogen and Phosphorus by Natural Clay Minerals. *Adsorption Science & Technology*, 2021, 1-14. <https://doi.org/10.1155/2021/4158151>
- Fan, X., Li, Y., & Alva, A. K. (2011). Effects of Temperature and Soil Type on Ammonia Volatilization from Slow-Release Nitrogen Fertilizers. *Communications In Soil Science and Plant Analysis*, 42(10), 1111-1122. <https://doi.org/10.1080/00103624.2011.566957>
- Foeroid, B., Alvarenga, E., Szocs, J., & Makádi, M. (2019). Ammonium Sorbed to Zeolite Is Partly Available to Wheat in the First Growth Cycle. *Agronomy*, 9(3), 122. <https://doi.org/10.3390/agronomy9030122>
- Han, B., Butterly, C., Zhang, W., He, J., & Chen, D. (2021). Adsorbent materials for ammonium and ammonia removal: A review. *Journal Of Cleaner Production*, 283, 124611. <https://doi.org/10.1016/j.jclepro.2020.124611>
- Harvey, O. R., Herbert, B., Harris, J., Stiffler, E. A., & Crenwelge, J. (2009). A New Spectrophotometric Method for Rapid Semiquantitative Determination of Soil Organic Carbon. *Soil Science Society of America Journal*, 73(3), 822-830. <https://doi.org/10.2136/sssaj2008.0268>
- Hearn, L., Barton, L., Schwenke, G., & Murphy, D. V. (2023). Ammonia volatilization losses from urea applied to acidic cropping soils is regulated by pH buffering capacity. *Soil Research*, 61(7), 685-696. <https://doi.org/10.1071/sr23044>
- Jolin, W. C., Richard, A., Vasudevan, D., Gascón, J. A., & MacKay, A. A. (2020). Aluminosilicate Mineralogy and the Sorption of Organic Cations: Interplay between Electrostatic Barriers and Compound Structural Features. *Environmental Science & Technology*, 54(3), 1623-1633. <https://doi.org/10.1021/acs.est.9b06121>
- Kome, G. K., Enang, R. K., Tabi, F. O., & Yerima, B. P. K. (2019). Influence of Clay Minerals on Some Soil Fertility Attributes: A Review. *Open Journal of Soil Science*, 09(09), 155-188. <https://doi.org/10.4236/ojss.2019.99010>
- Krol, D. J., Forrestal, P. J., Wall, D., Lanigan, G. J., Sanz-Gomez, J., & Richards, K. G. (2020). Nitrogen fertilizers with urease inhibitors reduce nitrous oxide and ammonia losses, while retaining yield in temperate grassland. *Science of the total environment*, 725, 138329. <https://doi.org/10.1016/j.scitotenv.2020.138329>
- Li, Y., Xu, J., Liu, S., Qi, Z., Li, A., Wei, Q., Gu, Z., Liu, X., & Hameed, F. (2020). Salinity-induced concomitant increases in soil ammonia volatilization and nitrous oxide emission. *Geoderma*, 361, 114053. <https://doi.org/10.1016/j.geoderma.2019.114053>
- Linggi, L., Berlin, M., Mallik, M., Roy, S., Natarajan, N., & Vasudevan, M. (2020). Feasibility investigation of adsorptive removal of NH_4^+ and NO_3^- species from clayey aquifer using special soils. *Environment, Development and Sustainability*, 23(5), 6749-6768. <https://doi.org/10.1007/s10668-020-00889-6>
- Liu, G., Li, Y., & Alva, A. K. (2007). High Water Regime Can Reduce Ammonia Volatilization from Soils under Potato Production. *Communications In Soil Science and Plant Analysis*, 38(9-10), 1203-1220. <https://doi.org/10.1080/00103620701328289>

- Mahmud, K., Panday, D., Mergoum, A., & Missaoui, A. (2021). Nitrogen Losses and Potential Mitigation Strategies for a Sustainable Agroecosystem. *Sustainability*, 13(4), 2400. <https://doi.org/10.3390/su13042400>
- Martínez-Dalmau, J., Berbel, J., & Ordóñez-Fernández, R. (2021). Nitrogen Fertilization. A Review of the Risks Associated with the Inefficiency of Its Use and Policy Responses. *Sustainability*, 13(10), 5625. <https://doi.org/10.3390/su13105625>
- Mazloomi, F., & Jalali, M. (2019). Effects of vermiculite, nano clay and zeolite on ammonium transport through saturated sandy loam soil: Column experiments and modeling approaches. *Catena*, 176, 170-180. <https://doi.org/10.1016/j.catena.2019.01.014>
- Minato, E. A., Cassim, B. M. A. R., Besen, M. R., Mazzi, F. L., Inoue, T. T., & Batista, M. A. (2020). Controlled-release nitrogen fertilizers: characterization, ammonia volatilization, and effects on second-season corn. *Revista Brasileira de Ciência Do Solo*, 44. <https://doi.org/10.36783/18069657rbc20190108>
- Ohnemus, T., Spott, O., & Thiel, E. (2021). Spatial distribution of urea induced ammonia loss potentials of German cropland soils. *Geoderma*, 394, 115025. <https://doi.org/10.1016/j.geoderma.2021.115025>
- Palanivell, P., O. H. Ahmed, K. Susilawati and Majid, N. M. A. (2015). Mitigating Ammonia Volatilization from Urea in Waterlogged Condition Using Clinoptilolite Zeolite. *International Journal of Agriculture & Biology*, 17(1), 1814-9596.
- Panday, D., Mikha, M. M., Collins, H. P., Jin, V. L., Kaiser, M., Cooper, J., Malakar, A., & Maharjan, B. (2020). Optimum rates of surface-applied coal char decreased soil ammonia volatilization loss. *Journal of environmental quality*, 49(2), 256–267. <https://doi.org/10.1002/jeq2.20023>
- Pelster, D. E., Watt, D., Strachan, I. B., Rochette, P., Bertrand, N., & Chantigny, M. H. (2019). Effects of Initial Soil Moisture, Clod Size, and Clay Content on Ammonia Volatilization after Subsurface Band Application of Urea. *Journal Of Environmental Quality*, 48(3), 549-558. <https://doi.org/10.2134/jeq2018.09.0344>
- Pleysier, J. L., Janssens, J. C. L., & Cremers, A. (1986). A Clay Suspension Stability End Point Titration Method for Measuring Cation Exchange Capacity of Soils. *Soil Science Society Of America Journal*, 50(4), 887-891. <https://doi.org/10.2136/sssaj1986.03615995005000040011x>
- Powlson, D. S., & Dawson, C. (2021). Use of ammonium sulphate as a Sulphur fertilizer: Implications for ammonia volatilization. *Soil Use And Management*, 38(1), 622-634. <https://doi.org/10.1111/sum.12733>
- SEMARNAT. Secretaria de Medio Ambiente y Recursos Naturales. (2002). Norma Oficial Mexicana NOM021-RECNAT-2000, que establece las especificaciones de fertilidad, salinidad y clasificación de suelos, estudios, muestreos y análisis. Diario Oficial de la Federación. Ciudad de México. México.
- Shen, Y., Hua, L., Gao, W., & Li, M. (2020). The effects of humic acid urea and polyaspartic acid urea on reducing nitrogen loss compared with urea. *Journal Of the Science of Food And Agriculture/Journal Of The Science Of Food And Agriculture*, 100(12), 4425-4432. <https://doi.org/10.1002/jsfa.10482>
- Shi, Z., She, D., Pan, Y., & Xia, Y. (2024). Promotion effects of salt-alkali on ammonia volatilization in a coastal soil. *Pedosphere*, 34(1), 191-199. <https://doi.org/10.1016/j.pedsph.2023.09.004>
- Skorupka, M., & Nosalewicz, A. (2021). Ammonia Volatilization from Fertilizer Urea—A New Challenge for Agriculture and Industry in View of Growing Global Demand for Food and Energy Crops. *Agriculture*, 11(9), 822. <https://doi.org/10.3390/agriculture11090822>
- Van Reeuwijk, R. (2003). Procedimientos para el análisis de suelos clasificación y correlación. Colegio de Postgraduados. Montecillo, Texcoco, México.
- Viero, F., Bayer, C., Fontoura, S. M. V., & De Moraes, R. P. (2014). Ammonia volatilization from nitrogen fertilizers in no-till wheat and maize in southern Brazil. *Revista Brasileira de Ciência Do Solo*, 38(5), 1515-1525. <https://doi.org/10.1590/s0100-06832014000500017>
- Wallach, R. (2008). Physical characteristics of soilless media. En Elsevier eBooks (pp. 41-116). <https://doi.org/10.1016/b978-044452975-6.50005-8>
- Wan, X., Wu, W., & Liao, Y. (2021). Mitigating ammonia volatilization and increasing nitrogen use efficiency through appropriate nitrogen management under supplemental irrigation and rain-fed condition in winter wheat. *Agricultural Water Management*, 255, 107050. <https://doi.org/10.1016/j.agwat.2021.107050>
- Wang, X., Zhu, H., Yan, B., Chen, L., Shutes, B., Wang, M., Lyu, J., & FuMan, Z. (2023). Ammonia volatilization, greenhouse gas emissions and microbiological mechanisms following the application of nitrogen fertilizers in a saline-alkali paddy ecosystem. *Geoderma*, 433, 116460. <https://doi.org/10.1016/j.geoderma.2023.116460>
- Wang, Y., Li, M., & Yan, H. (2021). Ammonia volatilization from urea in alfalfa field with different nitrogen application rates, methods and timing. *Agriculture, Ecosystems & Environment*, 312, 107344. <https://doi.org/10.1016/j.agee.2021.107344>

- Wester Larsen, L., Müller-Stöver, D. S., Salo, T., & Jensen, L. S. (2022). Potential ammonia volatilization from 39 different novel biobased fertilizers on the European market – A laboratory study using 5 European soils. *Journal Of Environmental Management*, 323, 116249. <https://doi.org/10.1016/j.jenvman.2022.116249>
- Xie, W., Li, S., Shi, W., Zhang, H., Fang, F., Wang, G., & Zhang, L. (2019). Quantitatively ranking the influencing factors of ammonia volatilization from paddy soils by grey relational entropy. *Environmental Science and Pollution Research International*, 27(2), 2319-2327. <https://doi.org/10.1007/s11356-019-06952-8>
- Zaini, N. S. M., Lenggono, I. W., Naim, M. N., Norihiro, Y., Man, H. C., Bakar, N. F. A., & Puasa, S. W. (2021). Adsorptive capacity of spray-dried pH-treated bentonite and kaolin powders for ammonium removal. *Advanced Powder Technology*, 32(6), 1833-1843. <https://doi.org/10.1016/j.ap.2021.02.036>

