AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

A Dynamic Principal-Agent Model of Human-Mediated Aquatic Species Invasions

Alexander J. Macpherson, Rebecca Moore, and Bill Provencher

This paper presents a dynamic principal-agent model of aquatic species invasions in which a manager, concerned about the spread of invasive species across lakes by boaters, sets *interseasonal* management controls on a lake-by-lake basis, and boaters make a series of *intraseasonal* trip decisions to maximize random utility during the course of the season, conditional on the controls imposed by the manager. The results of a simulated invasion of Eurasian watermilfoil (*Myriophyllum spicatum*) highlight interesting aspects of the optimal management policies under two different management objectives: maximizing boater welfare and minimizing milfoil spread.

Key Words: invasive species, random utility model, Eurasian watermilfoil, dynamic optimization, principal-agent

This paper presents a dynamic principal-agent model of aquatic species invasions in which a manager, concerned about the spread of invasive species across lakes by boaters, sets *interseasonal* management controls on a lake-by-lake basis, and boaters make a series of *intraseasonal* trip decisions to maximize random utility during the course of the season, conditional on the controls imposed by the manager. As such, this paper endogenizes resource user behavior in the management decisions related to a species invasion by allowing the lake manager, the principal, to anticipate boater reaction to management activity and the invasive species. This allows a more complete understanding of the impact of different

policy instruments on the lake system.

There are two fundamental reasons to endogenize boater movements in a model of the spread of aquatic invasions by boaters. The first is to provide a better forecast of the rate and direction of spread of the invader. Recent work concerning the spread of zebra mussel (*Dreissena polymorpha*) is instructive. The zebra mussel invasion has been well documented, is economically important, proceeds at a rapid but tractable pace, and has provided an opportunity to examine a variety of dispersal models, including human-based models, within the context of an ongoing invasion (Johnson and Carlton 1996). Zebra mussels are limited to aquatic environments with discrete boundaries and well-defined connections. This permits ecologists to make clear distinctions between dispersal modes within a lake, within a watershed, and between watersheds to determine the dispersal mechanism that best describes zebra mussel spread (Johnson and Carlton 1996). The human-mediated potentials for zebra mussel dispersal are plentiful: any submerged object or activity that moves water (like recreational boating) can transport the mussels (Johnson and Carlton 1996).

Much of the work on zebra mussel dispersal has focused on predictive and risk analysis models. Schneider, Ellis, and Cummings (1998) de-

Alexander J. Macpherson is a doctoral student and NSF-IGERT Fellow in the School of Forest Resources and Conservation at the University of Florida in Gainesville, Florida. Rebecca Moore and Bill Provencher are a doctoral student and Professor, respectively, in the Department of Agricultural and Applied Economics at the University of Wisconsin in Madison, Wisconsin.

This paper was presented at the Invasive Species Workshop, sponsored jointly by the Northeastern Agricultural and Resource Economics Association, the U.S. Environmental Protection Agency, USDA Economic Research Service (ERS), and the Farm Foundation, in Annapolis, Maryland, on June 14–15, 2005. The views expressed in this paper are the authors' and do not necessarily represent the policies or views of the sponsoring agencies.

This research was supported by an NSF biocomplexity grant administered by the Center for Limnology, University of Wisconsin-Madison. The authors also wish to thank Stephen R. Carpenter, Richard C. Bishop, and three anonymous reviewers for their valuable comments.

veloped a production-attraction transportation model of zebra mussel spread in Illinois streams to analyze the risks posed by the invasive species to native mussels in Illinois streams. The model was parameterized using Illinois boat ramp activity data to predict the frequency of boat movements between lakes and the likelihood that zebra mussels are transported. Bossenbroek, Kraft, and Nekola (2001) used a gravity model that allows for prediction of long-distance dispersal by considering the source populations and the spatial configuration and ecology of the potential colonization sites. Like Schneider, Ellis, and Cummings (1998), the gravity model predicted interlake trips and multiplied trips against a probability of colonization to predict the dispersal of zebra mussels. The model included a measure of the vulnerability (i.e., the attractiveness of the lake to the invader) of potentially invaded lakes to better model when and where long-distance dispersal will happen (Bossenbroek, Kraft, and Nekola 2001). Buchan and Padilla (1999) compared a standard diffusion model with patterns of recreational boater activities to find that boater behavior was a better predictor of the observed pattern of the zebra mussel invasions. The diffusion model approach underestimated the maximum rate and geographic extent of the invasions because it was less capable of incorporating the long-distance movements across unsuitable habitat that the mussels achieve via human-mediation. The authors concluded that management activities that slow the spread of the invasive species should focus on boater movements (Buchan and Padilla 1999).

None of the foregoing models account for the response of boaters to species invasions and management controls. The potential significance of this omission can be seen in a study of exotic lake trout (*Salvelinus namaycush*) in Yellowstone Lake, Wyoming, part of Yellowstone National Park (Settle, Crocker, and Shogren 2002, Settle and Shogren 2002). These papers present a joint economic-ecological model of a native trout species, an invasive trout species, park visitors, and park managers. Visitors to the park allocate their time between fishing for trout and other activities, such as sightseeing. In one version of the model, fishermen do not change their behavior in response to changing trout populations. A second version allows fishermen to adapt to decreased fish populations by spending less time fishing and

more time sightseeing. Simulations show that ignoring the response of the fishermen will yield significantly different trout populations and that the direction of this difference depends on how aggressively park managers can combat the invasive trout.

Finnoff et al. (2005) also use an endogenous risk framework to determine the importance of ecological-economic feedbacks when managing zebra mussels in a Midwest lake. In this study, two dimensions of feedbacks were identified: one between the biological system and a private economic agent damaged by zebra mussels (such as a power plant), and the second between the resource manager, the firm, and the biological system. The consequences of ignoring these feedbacks can be significant, but not in all cases, and they are sensitive to initial conditions and the parameters of the model. For example, ignoring the feedback between the biological system and the power plant resulted in significant economic consequences but only minor ecological consequences.

There are key differences between the single lake scenarios of these studies and the multi-lake system we address in this paper. For example, in the study of a single lake, the manager is concerned only with the growth of the single invasive population and is relatively unconcerned with agent activity outside of the lake. In a multi-lake system, the management of any individual lake could affect the state of all other lakes in the system. However, the results of these studies imply that for our problem, simple extrapolation of boater movements, while perhaps a better predictor of invasions than transportation and diffusion models, will prove unsatisfactory if boater reactions are significant. Suppose, for instance, that Lake A is uninfected by the invasive species, and the lake manager chooses to close the lake to cross-lake boat traffic (while perhaps leaving the lake open to boats that remain on the lake full-season) in order to protect it from an invasion. The boater trip pattern across the landscape may change as a result of the policy. Not all of the trips previously taken to Lake A are lost; some are diverted to other lakes. This raises the possibility that the lake closure effectively *advances* the spread of the invasive species by, for instance, causing boaters to “leapfrog” closed lakes in favor of more distant, “clean” lakes.

This example illustrates our second justification for endogenizing boater behavior in studies of aquatic invasions: it is necessary to accurately estimate welfare effects. The welfare effect of closing Lake A has an obvious *direct* welfare effect because now access to the lake requires keeping a boat on the lake, but it also has an *indirect* welfare effect because it shifts the spread of the invasive in a way that may leave society worse off overall.

The next section of the paper presents the integrated economic and ecological model of aquatic invasions. The third section presents a simulated Eurasian watermilfoil (*Myriophyllum spicatum*) invasion to highlight interesting aspects of the optimal management policies under two different management objectives: maximizing boater welfare and minimizing milfoil spread, both subject to a program budget. The fourth section discusses the specific challenges of applying this model to an actual lake system, and proposes techniques in survey research and dynamic programming that will address these challenges. The final section concludes.

The Model

A diagram of the model is presented in Figure 1. This schematic lays out the basic logic of the model: the sequence of decisions made by the manager and boaters, and the resulting transitions in the biological state of the lake system. Suppose there is a system of J lakes, $j = 1, \dots, J$, each with unique vectors of characteristics that make it attractive to resource users and vulnerable to species invasions. At the beginning of each season s , the principal—the lake system manager—chooses seasonal controls affecting boaters and therefore the spread of the invasive species. The lake manager is forward-looking: in setting controls, he anticipates the future spread of the invasive species and accounts for boater reactions to the controls.

There are T days in each season, indexed by $t = 1, \dots, T$. On each day, boaters, acting as the agents in the problem, decide whether to make a trip to a lake within the system. Because a boater might make multiple trips within a season, invasive propagules from a source lake can become attached to equipment and transported to uncolo-

nized lakes in the district. For our purposes, boaters are assumed to be the unique “carriers” of the invasive species via the accidental transport of propagules. Boater behavior is dependent on the current state of the invasion, lake-specific variables, and the controls chosen by the manager at the start of the season. As the season progresses, the rate and direction of the invasive species spread is determined by the controls chosen by the lake manager at the start of the season, the stochastic trip decisions of boaters, and the stochastic ecological processes underlying the invasion.

State of the Lake System and Management Controls

To make it easier to describe the essential features of the model, we present it in terms of a specific invasive species—Eurasian watermilfoil (*Myriophyllum spicatum*, hereafter called “milfoil”)—and two specific management controls relevant to the control of milfoil spread. These controls are the mandatory use of boat-cleaning equipment upon exit from a lake, to wash away milfoil propagules attached to the boat; and closing lakes to interlake boat traffic, so that access to a lake, although still possible—for instance, via an *in situ* boat livery—is nonetheless more expensive. The effects of a milfoil invasion, discussed in more detail in the following section, are predominantly negative impacts on lake ecology and recreational value, and so the lake manager does not face the possibility of a welfare-improving invasion, as might be the case when dealing with invasions of other species.

At the start of season s , the lake system is defined by three sets of dichotomous state variables relevant to the manager’s decision problem: i_{js} , taking a value of 1 when lake j is infected by milfoil, and 0 otherwise; e_{js} , taking a value of 1 if boat-cleaning equipment is already installed (with boater use required) on lake j , and 0 otherwise; and a_{js} , taking a value of 1 if a lake is already closed to interlake boat traffic, so that lake access is possible only via a costly alternative, and 0 otherwise. In the discussion below, the full set of state variables is defined by $\{\mathbf{i}_s, \mathbf{e}_s, \mathbf{a}_s\}$, where each element is a J -dimensioned vector of state values.

At the start of the season, the manager executes

Figure 1. Diagram of the Modeling Approach

two sets of decisions to control the spread of milfoil, conditional on the state of the system $\{\mathbf{i}_s, \mathbf{e}_s, \mathbf{a}_s\}$. The first pertains to the installation and required use of cleaning equipment, and the second pertains to the closure of lakes to interlake boat traffic. In the discussion below we treat

management controls implicitly, by using e'_{js} and a'_{js} to denote state variable values *after* the manager's control decisions. So, for instance, $a_{js} = 0$, $a'_{js} = 1$ indicates that at the start of season s lake j was open to interlake boat traffic, and the manager closed the lake; and $a_{js} = 1$, $a'_{js} = 1$ indicates

that at the start of season s lake j was closed, and the manager chose to keep it closed. The upshot is that whereas the manager chooses controls conditional on the state $\{\mathbf{i}_s, \mathbf{e}_s, \mathbf{a}_s\}$, boaters make trip decisions conditional on the state $\{\mathbf{i}_s, \mathbf{e}'_s, \mathbf{a}'_s\}$.

Boater Trip Decisions

In the model, a single representative boater decides on each day of the season whether to take a trip to a lake within the district.¹ Baseline utility (the utility associated with no trip) is zero, and the boater choosing to visit lake j on day t during season s receives money-measured utility

$$(1) \quad \begin{aligned} V_{jst} &= U_{js} + \varepsilon_{jst} \\ &= -C_j - \text{comply} \times E \times e'_{js} \\ &\quad - A \times a'_{js} + \beta Z_{js}(\mathbf{i}_{js}) + \varepsilon_{jst}, \end{aligned}$$

where C_j is the travel cost of a trip to lake j ; *comply* is the fraction of boaters who use the cleaning equipment; E denotes the cost of cleaning a boat, so $E \times e'_{js} = 0$ when boat-cleaning equipment is not installed on lake j in season s , and $E \times e'_{js} = E$ when boat-cleaning equipment is installed; A is the cost of a trip on a closed lake, so $A \times a'_{js} = 0$ when a lake is open to boat traffic, and $A \times a'_{js} = A$ when the lake is closed; Z_{js} is a vector of lake characteristics affecting trip decisions, such as lake size, and other lake attributes affected by whether the lake is infected by milfoil, such as the quality of fishing; and ε_{jst} is the component of the utility known to the user but unobserved by the analyst. We assume that this unobserved component is identically and independently Gumbel-distributed.

User welfare on day t during season s is measured in terms of expected utility, here corresponding to the well-documented *inclusive value*,

$$\ln \left(\sum_{j=0}^J \exp(U_{js}) \right).$$

Seasonal expected utility is then given by

$$(2) \quad IV_s(\mathbf{i}_s, \mathbf{e}'_s, \mathbf{a}'_s) = T \ln \left(\sum_{j=0}^J \exp(U_{js}) \right).$$

Invasive Species Dispersal

To fully develop the human-mediated dispersal model, we need to link the random utility model (RUM) to the ecological characteristics of the species and habitats in question. Under the assumption that boaters are the only means of interlake dispersal for milfoil, and invasive propagules remain viable for one day, the likelihood of an uninfected lake k being colonized at time $t+1$ can be represented as the probability that the representative boater visits an infected lake on day t , picks up a propagule from this lake, and visits lake k on day $t+1$, with the propagule then establishing a colony.

Let D_{st} be the event that the boater visits an infected lake and becomes a propagule carrier on day t of season s . We denote by J_s^* the set of lakes actually infected in season s , and we denote by $f(e'_{ks}, a'_{ks}, X_k)$ the probability that the boater becomes a propagule carrier upon a visit to lake $k \in J_s^*$; this probability depends on whether boat cleaning is required, whether the lake is closed to interlake boat traffic, and ecological characteristics of the lake, X_k . Keeping in mind that the unobserved component of utility is Gumbel-distributed, the probability of event D_{st} is given by

$$(3) \quad \Pr[D_{st}] = \sum_{k \in J_s^*} \left(f(e'_{ks}, a'_{ks}, X_k) \frac{\exp(U_{ks})}{1 + \sum_{j=1}^J \exp(U_{js})} \right).$$

We define $I_{ks,t+1}$ as the event that lake k is first colonized on day $t+1$ of season s .² The probability of this event occurring, *conditional* on the probability that the representative boater became a carrier on day t , is given by

¹ Nothing is gained by scaling up the model to N identical boaters.

² For the sake of simplicity, in this model a lake first colonized anytime during season s does not enter an infected state, thereby affecting boater and manager decisions, and becoming itself a source of propagules, until the start of season $s+1$. One might interpret this as representing the lag between the time at which an invasive species actually begins propagating at a new location, and the time at which its presence generates social costs.

$$(4) \quad \Pr[I_{ks,t+1} | D_{st}] = g(X_k) \frac{\exp(U_{ks})}{1 + \sum_{j=1}^J \exp(U_{js})},$$

where $g(X_k)$ relates the ecological characteristics of lake k to the probability of an introduced propagule establishing a colony. Combining equations (3) and (4) gives the *unconditional* probability of lake k being colonized on day $t+1$:

$$(5) \quad \Pr[I_{ks,t+1}] = \Pr[I_{ks,t+1} | D_{st}] \times \Pr[D_{st}]$$

$$= g(X_k) \frac{\exp(U_{ks})}{1 + \sum_{j=1}^J \exp(U_{js})}$$

$$\sum_{j \in J_s} \left(f(e'_{js}, a'_{js}, X_j) \frac{\exp(U_{js})}{1 + \sum_{r=1}^J \exp(U_{rs})} \right).$$

Left implicit in (5) is that $\Pr[I_{ks,t+1}]$ depends on the entire state of the system (through f and U).

Now we must extend the formulation of the colonization probability from a single day to the full season. Consider the probability that lake k is colonized by the end of season s , transitioning the lake into an infected state at the start of season $s+1$. On the first day of the season, there is no chance of colonization because there is no possibility of a trip on a previous day. On the second day of the season, the probability of an infection of lake k is given by (5). On the third day of the season, the probability of colonization of lake j is given by the probability of colonization on day three [from (5) again], multiplied by the probability of *not* being colonized on days one and two *plus* the probability of being colonized on days one and two. Letting t and q index the days within the season, the probability of colonization of lake k during season s assumes the following recursive structure:

$$(6) \quad \Pr[I_{ks} | \mathbf{i}, \mathbf{e}'_s, \mathbf{a}'_s] = \Pr[I_{ks2}] + \Pr[I_{ks3}](1 - \Pr[I_{ks2}]) +$$

$$\dots + \Pr[I_{ksT}] \prod_{t=2}^{T-1} (1 - \Pr[I_{kst}])$$

$$= \sum_{t=2}^T \left[\Pr[I_{kst}] \prod_{q=1}^{t-1} (1 - \Pr[I_{ksq}]) \right].$$

Management Objectives

At the beginning of each season the lake manager chooses which action to take on each lake. The options are to do nothing, mandate the use of cleaning equipment, or close a lake. We represent the cost of management controls in season s by $tc(\mathbf{e}_s, \mathbf{a}_s, \mathbf{e}'_s, \mathbf{a}'_s)$; this formulation recognizes that the cost of a control depends on the state of the management variables at the start of the season, and the manager's control choices. For instance, mandating that all boats exiting a lake must be cleaned is likely much less costly if cleaning equipment already exists at the lake. Seasonal net benefit, given that only boaters draw utility from changes in the state of the invasion (a modeling simplification, of course), is denoted by

$$(7) \quad B(\mathbf{i}_s, \mathbf{e}_s, \mathbf{a}_s, \mathbf{e}'_s, \mathbf{a}'_s) = IV(\mathbf{i}_s, \mathbf{e}'_s, \mathbf{a}'_s) - tc(\mathbf{e}_s, \mathbf{a}_s, \mathbf{e}'_s, \mathbf{a}'_s).$$

It is important to note that the value of the benefit function depends on the distribution of milfoil across the lake landscape, $\mathbf{i}_s$. The manager, as the principal of the model, anticipates how boaters will change their pattern of trip behavior in reaction to management activity and milfoil spread. This change in boater behavior will affect milfoil distribution in the next season, and the manager's benefit function implicitly recognizes this impact. This key element of the model allows the manager to truly appreciate the full effect of his management decisions and so identify the optimal management strategy.

In this simple model a lake's state of invasion is a binary variable, taking a value of 1 if milfoil is present in the lake and 0 otherwise. It follows that there exist 2^J possible states of invasion for the system as a whole. We denote this set of feasible invasion states by $\mathbf{I}$. Let $p_{i'}(\mathbf{i}, \mathbf{e}', \mathbf{a}')$ denote the probability that the system transitions to invasion state $\mathbf{i}' \in \mathbf{I}$ at the start of the next season, given the state of the invasion in the current season, and the manager's choice of controls for the current season (as indicated by the management state vectors $\mathbf{e}'$ and $\mathbf{a}'$). Then denoting the discount factor by δ , with $0 < \delta < 1$, and dropping the season subscript s to reduce notational clutter, the problem of a manager who wishes to maxi-

mize discounted utility, subject to a seasonal budget constraint Y , can be succinctly stated in Bellman's form as

$$(8) \quad V(\mathbf{i}, \mathbf{a}, \mathbf{e}) = \max_{\mathbf{a}', \mathbf{e}'} \left[B(\mathbf{i}, \mathbf{a}, \mathbf{e}, \mathbf{a}', \mathbf{e}') + \delta \sum_{\mathbf{i}' \in I} p_{\mathbf{i}'}(\mathbf{i}, \mathbf{a}', \mathbf{e}') V(\mathbf{i}', \mathbf{a}', \mathbf{e}') \right] \\ \text{s.t.} \quad tc(\mathbf{e}, \mathbf{a}, \mathbf{e}', \mathbf{a}') \leq Y.$$

The objective function in equation (8) represents a manager interested in maximizing the value of the lake ecosystem. Assuming that the value of the system is correctly measured, this objective is consistent with standard economic principles. However, invasive species management is a problem that crosses disciplinary lines, and assuming that the decision maker has such an objective could be incorrect. For example, from a purely ecological perspective, because milfoil has no ecological benefits, the goal of the manager might be to minimize the spread of milfoil across the landscape. In order to identify how optimal management strategies might differ under different management objectives, we compare the problem of the welfare-maximizing manager [given in equation (8)] with that of the spread-minimizing manager, who will choose seasonal values of $\mathbf{a}'$ and $\mathbf{e}'$ to minimize the rate of spread of the invasive species to new lakes, subject to a budget constraint:

$$(9) \quad V(\mathbf{i}, \mathbf{a}, \mathbf{e}) = \min_{\mathbf{a}', \mathbf{e}'} \left[\sum_{k \notin J^*} \Pr[I_k | \mathbf{i}, \mathbf{e}', \mathbf{a}'] + \delta \sum_{\mathbf{i}' \in I} p_{\mathbf{i}'}(\mathbf{i}, \mathbf{a}', \mathbf{e}') V(\mathbf{i}', \mathbf{a}', \mathbf{e}') \right] \\ \text{s.t.} \quad tc(\mathbf{e}, \mathbf{a}, \mathbf{e}', \mathbf{a}') \leq Y,$$

where

$$\sum_{k \notin J^*} \Pr[I_k | \mathbf{i}, \mathbf{e}', \mathbf{a}']$$

is effectively the expected number of newly infected lakes in the current season. It follows from a strictly positive discount rate (that is, the discount factor is strictly less than one) that even in the case where no control is foolproof—so that infection of the entire system is eventually ensured—it is worthwhile to control an infestation. Put another way, the timing of the invasion mat-

ters: the manager prefers to postpone an invasion for as long as possible.

Eurasian Watermilfoil Case Study

Eurasian watermilfoil is a perennial herbaceous freshwater submersed plant. The plant forms a dense canopy of branches floating at the surface (Madsen 1999) and grows best in fertile, fine-textured, inorganic sediments. An opportunistic species, milfoil establishes in highly disturbed lakes that receive significant nitrogen- and phosphorus-containing runoff (Wisconsin Department of Natural Resources 2004). Milfoil largely spreads asexually through vegetative fragments and not sexually by seed dispersal (Madsen, Eichler, and Boylen 1988). Because of this vegetative reproduction, milfoil spreads quickly via accidental carriage by boats, motors, trailers, bilges, live wells, or bait buckets (Madsen, Eichler, and Boylen 1988). Given damp conditions, milfoil fragments can remain viable for several days (Wisconsin Department of Natural Resources 2004). Between unconnected lakes, milfoil is spread primarily through accidental introductions by boats and boat trailers (Madsen, Eichler, and Boylen 1988).

Milfoil has several ecological effects once it establishes in a lake. First, the presence of significant submerged macrophyte biomass can contribute to nutrient enrichment in lakes (Carpenter 1980). This phenomenon occurs because the decaying shoots of the macrophyte release dissolved phosphorus and organic matter, which can make an important contribution to the pelagic production in a lake. This nutrient contribution can accelerate eutrophication processes (Carpenter 1980). Second, abundant milfoil can severely impact the diversity and density of native aquatic plant communities. Species richness and abundance decline with increased milfoil growth. These results are expected to extend to other components of the food web as well (Boylen, Eichler, and Madsen 1999). Third, high densities of submerged macrophytes can harm the quality of a fishery (Wiley et al. 1984, Bettoli et al. 1992). Feeding rates are reduced in lakes with dense beds of macrophytes by reducing predator efficiency by providing increased prey refuge (Olson et al. 1998). The lower mortality rates of the smaller fishes cause

greater population densities and stronger competitive interactions among forage fish (Mittelbach 1988). Experiments have shown that cutting channels through macrophyte beds may cause fish populations to increase, showing that harvesting may be a valuable management tool in infested lakes (Olson et al. 1998, Unmoth, Hansen, and Pellett 1999).

Results from a Four-Lake Simulation

Simulating the model requires a variety of data concerning boater preferences, probabilities of colonization, boater travel costs, management costs, and boater compliance rates. Given that little of this information is currently available, we simulated a minimal version of the model with realistic but mostly hypothetical data.

The lake system used in the simulations is comprised of four lakes, differentiated by the cost of travel to the lake and lake size. In the simulated model, the differential appeal of the lakes is straightforward, and supported by empirical evidence and/or economic theory: boaters prefer larger lakes to smaller ones, and closer lakes to more distant ones. The spatial distribution of lakes across the landscape is expected to have a large impact on boater trip behavior. For this reason, we explore two possible spatial arrangements of the four lakes. One arrangement places the largest lake farthest from the population center, and the smallest lake nearest. This arrangement presents the boater with a clear trade-off between lake size and trip cost. The second arrangement is one in which the largest lake is nearest the population center, and the smallest lake is farthest—a likely scenario in a region that develops around the use of its lakes. In the discussion below we call this the “endogenous development” scenario, and in the context of this scenario we refer to the large lake close to the population center as the “dominant” lake because it is preferable both in size and location.

A variety of parameters and management budgets were used in the simulations in an attempt to obtain a general sense of the role of boat traffic, and boater responses to management controls, on milfoil infestations. Several initial conditions were also examined, including scenarios in which all lakes are initially milfoil-free, one lake contains milfoil, three lakes contain milfoil, and all four

lakes contain milfoil. Though clearly minimal, the model is expected to reflect the key dynamics of managing an invasive aquatic plant species on a portfolio of lakes.³

Several observations can be made based on the patterns that emerge across the scenarios examined in the simulations. First, the results are consistent with Settle, Crocker, and Shogren (2002) in that the impact of explicitly recognizing boater reaction to milfoil spread can be quite large. For example, when the largest lake is farthest from the population center and boaters *do not* respond negatively to milfoil presence, the expected welfare value of a system entirely free of milfoil is about \$0.5 million higher than the value of the system where all lakes contain milfoil. When boaters *do* respond negatively to milfoil presence, the difference is about \$1.4 million. Even more, boaters’ negative response to environmental degradation caused by the presence of milfoil can play a significant role in controlling the dispersal of milfoil. Controls may cause a dramatic decrease in trips to regulated lakes, without a full redistribution of the trips to other lakes in the system. The expected size of this decline varies significantly with the spatial arrangement of the lakes. Overall, a policy that accounts for boater responses to controls is much more effective—no matter the objective—than one that does not.

Second, in models that incorporate a boater response, there is a significant difference in the optimal policies of a welfare-maximizing manager and a spread-minimizing manager. Under welfare maximization, management resources will often remain unspent, and the optimal policy is often to impose no controls, in order to avoid imposing excessive costs on boaters. This result obviously could change—perhaps drastically—in a more realistic model in which milfoil negatively affects non-boating shoreline property owners (Halsted et al. 2003). In contrast, managers interested in minimizing milfoil spread use controls more readily. Under this alternative objective, the budget is always binding and an increase in the budget can result in a *decrease* in welfare. This effect varies significantly with the spatial arrangement of the lakes.

³ A more detailed description of the mechanics of the simulation can be found in Macpherson (2004).

Finally, the difference in expected welfare value between a system entirely free of milfoil, and one where all lakes contain milfoil, appears to be greatest in the endogenous development scenario. In other words, the tendency for lake regions to develop around the largest lakes serves to increase the value of keeping the lake system free of an invasive. The explanation is that invasions tend to hit the most attractive lakes first, and endogenous development reinforces the stratification of lakes based on their “natural” appeal: the best lakes also become the closest lakes. A corollary to this result is that under welfare maximization in the endogenous development scenario, if the dominant lake is milfoil-free, the expected time to colonization of the entire system is relatively long, as significant resources are put into protecting the dominant lake, and trips to infected “satellite” lakes are relatively few; on the other hand, once the dominant lake is infected, the entire system quickly becomes infected.

Obstacles to a Large-Scale Application

The simulations described in the previous section illustrate the types of policy recommendations that can be derived from the model, but their usefulness from an applied perspective is limited by the small number of lakes considered. Though a larger, more complex system could be expected to behave similarly in many respects, it is not likely that an application to an actual lake system would generate exactly the same policy recommendations. Yet applying the model on a large scale is a daunting challenge, for two reasons.

The first and most obvious is that the dynamic programming problems (8) and (9) suffer from the well-known “curse of dimensionality”: as additional lakes are added to the system, the size of both the state space and the control space increases exponentially. The second reason is that estimation of a RUM model of boaters for a large-scale lake system is problematic. In particular, such estimation requires that the analyst define the choice sets of boaters at the lake level. Several authors have noted the bias in the estimation of RUMs when the choice set is misspecified (Haab and Hicks 1997, 1999). Most RUM studies use researcher-defined choice sets, such as including all lakes within a certain distance of the

individual’s residence, based on the maximum distance an individual would travel for a day trip (see Parsons 2003). Another possible option is to approximate larger choice sets by drawing a random subset of lakes. A series of studies (Parsons and Kealy 1992, Parsons and Needelman 1992, Feather 1994, Feather, Hellerstein, and Tomasi 1995, Lupi and Feather 1998) examine the bias and efficiency of aggregation and random selection.

It appears possible to surmount the challenges posed in applying our model to a large-scale aquatic system. With respect to the matter of solving large dynamic programming problems, Woodward, Wui, and Griffin (2005) present an innovative method of carrying out dynamic optimization in the context of a large simulation model of a red snapper fishery. With this method, an approximate optimization of an “aggregate” problem is solved using information from the full simulation model. While the application presented in their paper differs from our problem in a few key details—for instance, the fishery model control variables are continuous, while our model uses discrete control variables—we believe their “direct approach” to solving the dynamic programming simulation problem could be adapted to examine species invasions of large-scale aquatic systems.

Concerning the matter of identifying the lakes within a boater’s choice set, several authors argue persuasively that the best approach is to directly query decision makers about their choice sets (Peters, Adamowicz, and Boxall 1995, Hicks and Strand 2000). Yet this approach ultimately confronts the practical question of survey design: how does the analyst induce a respondent to identify all the lakes considered in the trip decision?

One possibility is to list all relevant lakes and request that survey respondents specify those they consider when making the trip decision. Another possibility is to pose the choice set as an open-ended question in which the respondent lists those lakes they contemplate visiting. With both approaches, respondent fatigue is likely to bias responses.

Recent advances in the use of Internet-based surveys offer a possible solution to this problem. Internet surveys allow for more flexibility and control over which questions the respondent actu-

ally sees. One could imagine, for instance, a clickable map of the study region. The respondent could click on the part of the region they most often visit. Zooming to this smaller region, the user could then simply click on all the lakes that should belong in the choice set—a much simpler task than writing down all of the lake names. The rest of the survey questions could then be tailored to those lakes identified by the respondent. In short, the Internet platform allows for relatively individualized survey design while still appearing concise and linear to the respondent.

Conclusions

This paper presents a general model of aquatic plant species invasions in which lake users are the primary vector of invasion and resource user behavior is explicitly incorporated into management decision. In the model, recreational boaters respond to both the presence of the invader and the actions taken by the lake manager. The lake manager acknowledges boater responses and makes management decisions accordingly. By incorporating a model of human behavior into a dynamic ecological system, we can better understand the complex interactions between human activity and the environment.

The simulated milfoil case study presented here illustrates the potential benefits of using this model to generate policy recommendations for lake managers. Though a large-scale application is desirable, there are several hurdles to its implementation. The discussion above argues that recent advances in the literature could allow us to overcome these obstacles. Doing so will allow simulations at a regional scale. With these results, managers will then be able to consider the full impacts of their policies, and so make more informed decisions.

References

- Bettoli, P.W., M.J. Maceina, R.L. Noble, and R.K. Betsill. 1992. "Piscivory in Largemouth Bass as a Function of Aquatic Vegetation Abundance." *North American Journal of Fisheries Management* 12: 509–516.
- Bossenbroek, J.M., C.E. Kraft, and J.C. Nekola. 2001. "Prediction of Long-Distance Dispersal Using Gravity Models: Zebra Mussel Invasion of Inland Lakes." *Ecological Applications* 11(6): 1778–1788.
- Boylen, C.W., L.W. Eichler, and J.D. Madsen. 1999. "Loss of Native Aquatic Plant Species in a Community Dominated by Eurasian Watermilfoil." *Hydrobiologia* 415: 207–211.
- Buchan, L.A.J., and D.K. Padilla. 1999. "Estimating the Probability of Long-Distance Overland Dispersal of Invading Aquatic Species." *Ecological Applications* 9(1): 254–265.
- Carpenter, S.R. 1980. "Enrichment of Lake Wingra, Wisconsin, by Submersed Macrophyte Decay." *Ecology* 61(5): 1145–1155.
- Feather, P.M. 1994. "Sampling and Aggregation Issues in Random Utility Model Estimation." *American Journal of Agricultural Economics* 76(4): 772–780.
- Feather, P., D. Hellerstein, and T. Tomasi. 1995. "A Discrete-Count Model of Recreational Demand." *Journal of Environmental Economics and Management* 29(2): 214–227.
- Finnoff, D., J.F. Shogren, G. Leung, and D. Lodge. 2005. "The Importance of Bioeconomic Feedback in Invasive Species Management." *Ecological Economics* 52(3): 367–381.
- Haab, T.C., and R.L. Hicks. 1997. "Accounting for Choice Set Endogeneity in Random Utility Models of Recreation Demand." *Journal of Environmental Economics and Management* 34(2): 127–147.
- . 1999. "Choice Set Considerations in Models of Recreation Demand: History and Current State of the Art." *Marine Resource Economics* 14(4): 271–281.
- Halsted, J.M., J. Michaud, S. Hallas-Burt, and J.P. Gibbs. 2003. "Hedonic Analysis of Effects of a Nonnative Invader (*Myriophyllum heterophyllum*) on New Hampshire (USA) Lakefront Properties." *Environmental Management* 32(3): 391–398.
- Hicks, R.L., and I. Strand. 2000. "The Extent of Information: Its Relevance for Random Utility Models." *Land Economics* 76(3): 374–385.
- Johnson, L.E., and J.T. Carlton. 1996. "Post-Establishment Spread in Large-Scale Invasions: Dispersal Mechanisms of the Zebra Mussel *Dreissena polymorpha*." *Ecology* 77(6): 1686–1690.
- Lupi, F., and P.M. Feather. 1998. "Using Partial Site Aggregation to Reduce Bias in Random Utility Travel Cost Models." *Water Resources Research* 34(12): 3595–3603.
- Macpherson, A.J. 2004. "A Model of Human-Mediated Aquatic Species Invasions: A Eurasian Watermilfoil Case Study." Master's thesis, Department of Agricultural and Applied Economics, University of Wisconsin-Madison.
- Madsen, J.D. 1999. "Predicting the Invasion of Eurasian Watermilfoil into Northern Lakes." Technical Report No. A-99-2, Waterways Experiment Station, U.S. Army Corps of Engineers, Vicksburg, MS.
- Madsen, J.D., L.W. Eichler, and C.W. Boylen. 1988. "Vegetative Spread of Eurasian Watermilfoil in Lake George, New York." *Journal of Aquatic Plant Management* 35: 63–68.
- Mittelbach, G.G. 1988. "Competition Among Refuging Sunfishes and Effects of Fish Density on Littoral Zone Invertebrates." *Ecology* 69(3): 614–623.
- Olson, M.H., S.R. Carpenter, P. Cunningham, S. Garny, B.R.

- Herwig, N.P., Nibbelink, T., Pellett, C., Storlie, A.S., Trebitz, and K.A. Wilson. 1998. "Managing Macrophytes to Improve Fish Growth: A Multi-lake Experiment." *Fisheries* 23(2): 6–12.
- Parsons, G.R. 2003. "The Travel Cost Model." In P.A. Champ, K.J. Boyle, and T.C. Brown, eds., *A Primer on Nonmarket Valuation*. Boston: Kluwer Academic Publishers.
- Parsons, G.R., and M.J. Kealy. 1992. "Randomly Drawn Opportunity Sets in a Random Utility Model of Lake Recreation." *Land Economics* 68(1): 93–106.
- Parsons, G.R., and M.S. Needelman. 1992. "Site Aggregation in a Random Utility Model of Recreation." *Land Economics* 68(4): 418–433.
- Peters, T., W.L. Adamowicz, and P.C. Boxall. 1995. "Influence of Choice Set Considerations in Modeling the Benefits from Improved Water Quality." *Water Resources Research* 31(7): 1781–1787.
- Schneider, D.W., C.D. Ellis, and K.S. Cummings. 1998. "A Transportation Model Assessment of the Risk to Native Mussel Communities from Zebra Mussel Spread." *Conservation Biology* 12(4): 788–800.
- Settle, C., T.D. Crocker, and J.F. Shogren. 2002. "On the Joint Determination of Biological and Economic Systems." *Ecological Economics* 42(1-2): 301–311.
- Settle, C., and J.F. Shogren. 2002. "Modeling Native-Exotic Species within Yellowstone Lake." *American Journal of Agricultural Economics* 84(5): 1323–1328.
- Unmoth, J.M.L., M.J. Hansen, and T.D. Pellett. 1999. "Effects of Mechanical Harvesting of Eurasian Watermilfoil on Largemouth Bass and Bluegill Populations in Fish Lake, Wisconsin." *North American Journal of Fisheries Management* 19(4): 1089–1098.
- Wiley, M.J., R.W. Gordon, S.W. Waite, and T. Powless. 1984. "The Relationship Between Aquatic Macrophytes and Sport-fish Production in Illinois Ponds: a Simple Model." *North American Journal of Fisheries Management* 4: 111–119.
- Wisconsin Department of Natural Resources. 2004. "The Facts of Eurasian Water Milfoil." Technical Publication No. PUB-WT-781, Wisconsin Department of Natural Resources, Madison, WI.
- Woodward, R.T., Y.S. Wui, and W.L. Griffin. 2005. "Living with the Curse of Dimensionality: Closed-Loop Optimization in a Large-Scale Fisheries Simulation Model." *American Journal of Agricultural Economics* (87)1: 48–60.