



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

Stata tip 1: The `eform()` option of `regress`

Roger Newson, King's College London, UK
roger.newson@kcl.ac.uk

Did you know about the `eform()` option of `regress`? It is very useful for calculating confidence intervals for geometric means and their ratios. These are frequently used with skewed Y -variables, such as house prices and serum viral loads in HIV patients, as approximations for medians and their ratios. In Stata, I usually do this by using the `regress` command on the logs of the Y -values, with the `eform()` and `noconstant` options. For instance, in the `auto` dataset, we might compare prices between non-US and US cars as follows:

```
. sysuse auto, clear
(1978 Automobile Data)
. generate logprice = log(price)
. generate byte baseline = 1
. regress logprice foreign baseline, noconstant eform(GM/Ratio) robust
Regression with robust standard errors
```

Number of obs =	74
F(2, 72) =	18043.56
Prob > F =	0.0000
R-squared =	0.9980
Root MSE =	.39332

logprice	GM/Ratio	Robust Std. Err.	t	P> t	[95% Conf. Interval]	
foreign	1.07697	.103165	0.77	0.441	.8897576	1.303573
baseline	5533.565	310.8747	153.41	0.000	4947.289	6189.316

We see from the `baseline` parameter that US-made cars had a geometric mean price of 5534 dollars (95% CI from 4947 to 6189 dollars), and we see from the `foreign` parameter that non-US cars were 108% as expensive (95% CI, 89% to 130% as expensive). An important point is that, if you want to see the baseline geometric mean, then you must define the constant variable, here `baseline`, and enter it into the model with the `noconstant` option. Stata usually suppresses the display of the intercept when we specify the `eform()` option, and this trick will fool Stata into thinking that there is no intercept for it to hide. The same trick can be used with `logit` using the `or` option, if you want to see the baseline odds as well as the odds ratios.

My nonstatistical colleagues understand regression models for log-transformed data a lot better this way than any other way. Continuous X -variables can also be included, in which case the parameter for each X -variable is a ratio of Y -values per unit change in X , assuming an exponential relationship—or assuming a power relationship, if X is itself log-transformed.