



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

THE STATA JOURNAL

Editor

H. Joseph Newton
Department of Statistics
Texas A & M University
College Station, Texas 77843
979-845-3142; FAX 979-845-3144
jnewton@stata-journal.com

Editor

Nicholas J. Cox
Geography Department
Durham University
South Road
Durham City DH1 3LE UK
n.j.cox@stata-journal.com

Associate Editors

Christopher Baum
Boston College
Rino Bellocco
Karolinska Institutet
David Clayton
Cambridge Inst. for Medical Research
Mario A. Cleves
Univ. of Arkansas for Medical Sciences
William D. Dupont
Vanderbilt University
Charles Franklin
University of Wisconsin, Madison
Joanne M. Garrett
University of North Carolina
Allan Gregory
Queen's University
James Hardin
University of South Carolina
Ben Jann
ETH Zurich, Switzerland
Stephen Jenkins
University of Essex
Ulrich Kohler
WZB, Berlin
Jens Lauritsen
Odense University Hospital

Stanley Lemeshow
Ohio State University
J. Scott Long
Indiana University
Thomas Lumley
University of Washington, Seattle
Roger Newson
King's College, London
Marcello Pagano
Harvard School of Public Health
Sophia Rabe-Hesketh
University of California, Berkeley
J. Patrick Royston
MRC Clinical Trials Unit, London
Philip Ryan
University of Adelaide
Mark E. Schaffer
Heriot-Watt University, Edinburgh
Jeroen Weesie
Utrecht University
Nicholas J. G. Winter
Cornell University
Jeffrey Wooldridge
Michigan State University

Stata Press Production Manager

Stata Press Copy Editors

Lisa Gilmore
Gabe Waggoner, John Williams

Copyright Statement: The Stata Journal and the contents of the supporting files (programs, datasets, and help files) are copyright © by StataCorp LP. The contents of the supporting files (programs, datasets, and help files) may be copied or reproduced by any means whatsoever, in whole or in part, as long as any copy or reproduction includes attribution to both (1) the author and (2) the Stata Journal.

The articles appearing in the Stata Journal may be copied or reproduced as printed copies, in whole or in part, as long as any copy or reproduction includes attribution to both (1) the author and (2) the Stata Journal.

Written permission must be obtained from StataCorp if you wish to make electronic copies of the insertions. This precludes placing electronic copies of the Stata Journal, in whole or in part, on publicly accessible web sites, file servers, or other locations where the copy may be accessed by anyone other than the subscriber.

Users of any of the software, ideas, data, or other materials published in the Stata Journal or the supporting files understand that such use is made without warranty of any kind, by either the Stata Journal, the author, or StataCorp. In particular, there is no warranty of fitness of purpose or merchantability, nor for special, incidental, or consequential damages such as loss of profits. The purpose of the Stata Journal is to promote free communication among Stata users.

The *Stata Journal*, electronic version (ISSN 1536-8734) is a publication of Stata Press, and Stata is a registered trademark of StataCorp LP.

Estimation and inference in dynamic unbalanced panel-data models with a small number of individuals

Giovanni S. F. Bruno
Istituto di Economia Politica, Bocconi University, Milan

Abstract. This article describes a new Stata routine, `xtlsdvc`, that computes bias-corrected least-squares dummy variable (LSDV) estimators and their bootstrap variance–covariance matrix for dynamic (possibly) unbalanced panel-data models with strictly exogenous regressors. A Monte Carlo analysis is carried out to evaluate the finite-sample performance of the bias-corrected LSDV estimators in comparison to the original LSDV estimator and three popular N -consistent estimators: Arellano–Bond, Anderson–Hsiao and Blundell–Bond. Results strongly support the bias-corrected LSDV estimators according to bias and root mean squared error criteria when the number of individuals is small.

Keywords: `st0091`, `xtlsdvc`, bias approximation, unbalanced panels, dynamic panel data, LSDV estimator, Monte Carlo experiment, bootstrap variance–covariance

1 Introduction

Situations in which past decisions have an impact on current behavior are ubiquitous in economics. To mention just one of the most familiar cases, in the presence of employment adjustment costs, the short-run labor demand of the firm will depend on past employment levels. Another crucial issue in empirical economics, strictly related to the modeling of dynamic relationships, is the presence of unobserved heterogeneity in individual behavior and characteristics. Panel datasets, where the behavior of N cross-sectional units is observed over T time periods, provide a solution to accommodating the joint occurrence of dynamics and unobserved individual heterogeneity in the phenomena of interest.

Since the seminal paper by Nickell (1981), where it is shown that the least-squares dummy variable (LSDV) estimator is not consistent for finite T in autoregressive panel-data models, a number of consistent instrumental variable (IV) and generalized method of moments (GMM) estimators have been proposed in the econometric literature as an alternative to LSDV. Anderson and Hsiao (1982) (AH) suggest two simple IV estimators that, upon transforming the model in first differences to eliminate the unobserved individual heterogeneity, use the second lags of the dependent variable, either differenced or in levels, as an instrument for the differenced one-time lagged dependent variable. Arellano and Bond (1991) (AB) propose a GMM estimator for the first-differenced model, which, relying on a greater number of internal instruments, is more efficient than AH. Blundell and Bond (1998) (BB) observe that with highly persistent data, first-

differenced IV or GMM estimators may suffer of a severe small-sample bias due to weak instruments. As a solution, they suggest a system GMM estimator with first-differenced instruments for the equation in levels and instrument in levels for the first-differenced equation.

A weakness of IV and GMM estimators is that their properties hold when N is large, so they can be severely biased and imprecise in panel data with a small number of cross-sectional units. This is often the case in most macro panels, but also in micro panels where heterogeneity concerns force the researcher not to use all information available, but rather to select a subsample of individuals from the original panel to estimate the parameters of interest. On the other hand, earlier Monte Carlo studies (Arellano and Bond 1991; Kiviet 1995; Judson and Owen 1999) demonstrate that LSDV, although inconsistent, has a relatively small variance compared to IV and GMM estimators.

Moving from the foregoing considerations, an alternative approach based upon the bias-correction of LSDV in dynamic panel-data models with strictly exogenous regressors has recently become popular in the econometric literature. Nickell (1981) derives an expression for the inconsistency of LSDV for $N \rightarrow +\infty$, which is bounded on the order T^{-1} . Kiviet (1995) uses higher-order asymptotic expansion techniques to approximate the small sample bias of the LSDV estimator to include terms of at most order $N^{-1}T^{-1}$. The approximations terms, however, all evaluated at the unobserved true parameter values, are of no direct use for estimation, so to make them operational, he suggests replacing the true parameters with the estimates from some consistent estimators. Monte Carlo evidence therein shows that the resulting bias-corrected LSDV estimator (LSDVC) often outperforms the IV–GMM estimators in terms of bias and root mean squared error (RMSE). Another piece of Monte Carlo evidence by Judson and Owen (1999) strongly supports LSDVC when N is small as in most macro panels. In Kiviet (1999), the bias expression is more accurate to include terms of at most order $N^{-1}T^{-2}$. Bun and Kiviet (2003), upon simplifying the approximations in Kiviet (1999), carry out Monte Carlo experiments showing that the first-order term of the approximation evaluated at the true parameter values is already capable to account for more than 90% of the actual bias.

None of the foregoing procedures to correct the LSDV estimator is feasible for unbalanced panels. This gap is partly filled in Bruno (2005), where the bias approximations in Bun and Kiviet (2003) are extended to accommodate unbalanced panels with a strictly exogenous selection rule. Monte Carlo evidence therein parallels that in Bun and Kiviet (2003).

This paper presents a new Stata program, `xtlsdvc`, which implements LSDVC building upon the theoretical approximation formulas in Bruno (2005) and estimates a bootstrap variance covariance matrix for the corrected estimator. Moreover, the relative performance of LSDVC is evaluated in comparison to LSDV, AB, AH, and BB for unbalanced panels with a small N (10 and 20 units) through various Monte Carlo experiments, thus extending the analysis by Judson and Owen (1999).

Monte Carlo results in this paper show that the three versions of LSDVC computed by `xtlsdvc` outperform all other estimators tried in terms of bias and RMSE. From this scenario, LSDVC clearly emerges as the preferred estimator for dynamic panel-data models with small N and strictly exogenous regressors. That said, practitioners should not forget an important limitation of the procedure: as opposed to IV-GMM estimators, in fact, no version of LSDVC is applicable in the presence of endogenous, or even only weakly exogenous, regressors. The results by Bun and Kiviet (2005) on the finite sample properties of LSDV and GMM estimators with weakly exogenous regressors, however, seem promising in the view of a general bias-correction procedure.

The paper is laid out as follows. The next section briefly reviews the theoretical results for corrected LSDV estimators. Section 3 describes the `xtlsdvc` routine. Section 4 contains the Monte Carlo analysis, and section 5 concludes the article. A demonstration of the code in the context of labor demand estimation is offered into an appendix.

2 Bias-corrected LSDV estimators

I consider the standard dynamic panel-data model

$$y_{it} = \gamma y_{i,t-1} + x'_{it}\beta + \eta_i + \epsilon_{it}; \quad |\gamma| < 1; \quad i = 1, \dots, N \text{ and } t = 1, \dots, T \quad (1)$$

where y_{it} is the dependent variable, x_{it} is the $\{(k-1) \times 1\}$ vector of strictly exogenous explanatory variables, η_i is an unobserved individual effect, and ϵ_{it} is an unobserved white-noise disturbance with constant variance σ_ϵ^2 .¹ Collecting observations over time and across individuals gives

$$y = D\eta + W\delta + \epsilon$$

where y and $W = (y_{-1} : X)$ are the $(NT \times 1)$ and $(NT \times k)$ matrices of stacked observations; $D = I_N \otimes \iota_T$ is the $(NT \times N)$ matrix of individual dummies (ι_T is the $(T \times 1)$ vector of all unity elements); η is the $(N \times 1)$ vector of individual effects; ϵ is the $(NT \times 1)$ vector of disturbances; and $\delta = (\gamma; \beta)'$ is the $(k \times 1)$ vector of coefficients.

It has been long recognized that the LSDV estimator for (1) is not consistent for finite T . Nickell (1981) derives an expression for the inconsistency for $N \rightarrow +\infty$, which is $O(T^{-1})$. Kiviet (1995) obtains a bias approximation that contains terms of higher order than T^{-1} . In Kiviet (1999), a more accurate bias approximation is derived. Bun and Kiviet (2003) reformulate the approximation in Kiviet (1999) with simpler formulas for each term.

Bruno (2005) extends Bun and Kiviet's (2003) formulas to unbalanced panels with a strictly exogenous selection rule. A more general version of (1) is considered, which allows missing observations in the interval $[0, T]$ for some individuals. Below I briefly present the approximation formulas for (possibly) unbalanced data as derived in Bruno (2005) and show their use to obtain LSDVC.

1. The bias-correction procedures of this paper can be applied to heteroskedastic cases after successful weighting of variables, although this may be admittedly difficult in practice.

Define a selection indicator r_{it} such that $r_{it} = 1$ if (y_{it}, x_{it}) is observed and $r_{it} = 0$ otherwise. From this, define the dynamic selection rule $s(r_{it}, r_{i,t-1})$, selecting only the observations that are usable for the dynamic model, namely, those for which both current values and one-time lagged values are observable:

$$s_{it} = \begin{cases} 1 & \text{if } (r_{i,t}, r_{i,t-1}) = (1, 1) \\ 0 & \text{otherwise} \end{cases} \quad i = 1, \dots, N \text{ and } t = 1, \dots, T$$

Thus for any i , the number of usable observations is given by $T_i = \sum_{t=1}^T s_{it}$. The total number of usable observations is given by $n = \sum_{i=1}^N T_i$; and $\bar{T} = n/N$ denotes the average group size. For each i , define the $(T \times 1)$ -vector $s_i = [s_{i1} \dots, s_{iT}]'$ and the $(T \times T)$ diagonal matrix S_i having the vector s_i on its diagonal. Define also the $(NT \times NT)$ block-diagonal matrix $S = \text{diag}(S_i)$. The (possibly) unbalanced dynamic model can then be written as

$$Sy = SD\eta + SW\delta + S\epsilon \quad (2)$$

The LSDV estimator is given by

$$\delta_{\text{LSDV}} = (W'M_s W)^{-1} W'M_s y$$

where

$$M_s = S \left\{ I - D(D'SD)^{-1} D' \right\} S$$

is the symmetric and idempotent $(NT \times NT)$ matrix, wiping out individual means and selecting usable observations.

Bias-approximation terms for unbalanced panels are the following:

$$c_1 \left(\bar{T}^{-1} \right) = \sigma_\epsilon^2 \text{tr}(\Pi) q_1 \quad (3)$$

$$c_2 \left(N^{-1} \bar{T}^{-1} \right) = -\sigma_\epsilon^2 \left\{ Q\bar{W}' \Pi M_s \bar{W} + \text{tr} \left(Q\bar{W}' \Pi M_s \bar{W} \right) I_{k+1} + 2\sigma_\epsilon^2 q_{11} \text{tr}(\Pi' \Pi \Pi) I_{k+1} \right\} q_1$$

$$c_3 \left(N^{-1} \bar{T}^{-2} \right) = \sigma_\epsilon^4 \text{tr}(\Pi) \left[2q_{11} Q\bar{W}' \Pi \Pi' \bar{W} q_1 + \left\{ \left(q_1' \bar{W}' \Pi \Pi' \bar{W} q_1 \right) + q_{11} \text{tr} \left(Q\bar{W}' \Pi \Pi' \bar{W} \right) + 2\text{tr}(\Pi' \Pi \Pi' \Pi) q_{11}^2 \right\} q_1 \right]$$

where $Q = \{E(W'M_s W)\}^{-1} = \left\{ \bar{W}' M_s \bar{W} + \sigma_\epsilon^2 \text{tr}(\Pi' \Pi) e_1 e_1' \right\}^{-1}$; $\bar{W} = E(W)$; $e_1 = (1, 0, \dots, 0)'$ is a $(k \times 1)$ vector; $q_1 = Qe_1$; $q_{11} = e_1' q_1$; L_T is the $(T \times T)$ matrix with unit first lower subdiagonal and all other elements equal to zero; $L = I_N \otimes L_T$; $\Gamma_T = (I_T - \gamma L_T)^{-1}$; $\Gamma = I_N \otimes \Gamma_T$; and $\Pi = M_s L \Gamma$. Clearly, in any balanced design $S \equiv I_{NT}$, so $M_s = I - D(D'D)^{-1} D'$, and the above terms reduce to those in Bun and Kiviet (2003).

With an increasing level of accuracy, the following three possible bias approximations emerge:

$$B_1 = c_1 \left(\overline{T}^{-1} \right); B_2 = B_1 + c_2 \left(N^{-1} \overline{T}^{-1} \right); B_3 = B_2 + c_3 \left(N^{-1} \overline{T}^{-2} \right) \quad (4)$$

In principle, bias-corrected LSDV estimators could be obtained by subtracting any of the above terms from LSDV. In practice, however, depending upon the unknown parameters σ_ϵ^2 and γ , approximations (4) are not feasible for bias correction. Nevertheless, consistent bias-corrected estimators can be obtained by finding consistent estimators for σ_ϵ^2 and γ , plugging them into the bias-approximations formulas, and then subtracting the resulting bias approximation estimates, \hat{B}_i , from LSDV as follows:

$$\text{LSDVC}_i = \text{LSDV} - \hat{B}_i, \quad i = 1, 2, \text{ and } 3 \quad (5)$$

Possible consistent estimators for γ are AH, AB, or BB, for example. Depending on the estimator of choice for γ , say h , a consistent estimator for σ_ϵ^2 is then given by

$$\hat{\sigma}_h^2 = \frac{e_h' M_s e_h}{(N - k - T)} \quad (6)$$

where $e_h = y - W\delta_h$, and $h = \text{AH, AB, and BB}$.

3 The xtlsdvc program

3.1 Syntax

The Stata program `xtlsdvc` written by the author calculates LSDVC for (1) using estimates for the bias approximations in (4). The basic syntax of `xtlsdvc` is the following:

```
xtlsdvc depvar [ indepvars ] [ if ], initial(estimator) [level(#) bias(#)
      vcov(#) first lsdv]
```

The routine automatically includes the lagged dependent variable as an explanatory variable and can fit the simple autoregressive model with no covariates.

3.2 Options

`initial(estimator)` is required and specifies the consistent estimator chosen to initialize the bias correction.

<i>estimator</i>	<i>description</i>
ah	AH estimator, with the dependent variable lagged two times, used as an instrument for the first-differenced model with no intercept ([R] ivreg)
ab	standard one-step AB estimator with no intercept ([XT] xtabond)
bb	standard BB estimator with no intercept, as implemented by the user-written Stata routine xtabond2 by Roodman (2003)
my	a row vector of initial values supplied directly by the user

To implement the last instance of this option, the user must create a $\{1 \times (k+1)\}$ matrix to be named `my`, the i th element of which serves as an initial value for the coefficient on the i th variable in *varlist* and the last, $(k+1)$ th, element as an estimate for the error variance. This may be useful in Monte Carlo simulations or if the user wishes to try initial estimators other than `ah`, `ab`, or `bb`.

`level(#)` specifies the confidence level, as a percentage, for confidence intervals of the coefficients. The default is `level(95)` or as set by `set level`; see [U] 20.6 **Specifying the width of confidence intervals.**

`bias(#)` determines the accuracy of the approximation: `# = 1` (default) forces an approximation up to $O(1/T)$; `# = 2` forces an approximation up to $O(1/NT)$; `# = 3` forces an approximation up to $O(N^{-1}T^{-2})$.

`vcov(#)` calculates a bootstrap variance–covariance matrix for LSDVC using `#` repetitions (`#` may not equal 1). The default is no bootstrap estimation of the variance–covariance matrix and standard errors. Notice that the bootstrap continues to work in the presence of gaps in the exogenous variables, although in this case, bootstrap samples for each unit are truncated to the first missing value encountered. Gaps in the dependent variable, instead, bear no consequence to the bootstrap sample size. This is explained in more detail in section 3.5. Also consider that bootstrap standard errors are downward biased when values for the unknown parameters are supplied through matrix `my` since the procedure in this case (keeping the values in `my` fixed over replications) neglects a source of variability for LSDVC.

`first` requests that the first-stage regression results be displayed.

`lsdv` requests that the original LSDV regression results be displayed.

To work out the approximations, `xtlsdvc` invokes the subroutine `xtlsdvc_1` that accomplishes the following tasks. First, `xtlsdvc_1` obtains the uncorrected LSDV estimates via a call to `xtreg ... , fe ([XT] xtreg)`.

Second, `xtlsdvc_1` obtains initial estimates for γ and β through one of the following instructions, depending on which *estimator* is specified in `initial`:

```
if "initial"=="ah" ivreg D.y D.x (LD.y=L2.y), noconstant
if "initial"=="ab" xtabond y x, noconstant
if "initial"=="bb" xtabond2 y L.y x, gmm(L.y) iv(x) noconstant
```

Then $\hat{\sigma}_h^2$, $h = \text{AH, AB, and BB}$, is computed as in (6).

Finally, `xtlsdvc_1` computes the bias approximations via the Stata `matrix` commands ([P] **matrix**) and corrects the LSDV estimates as indicated in (5).

3.3 Saved results

`xtlsdvc` saves in `e()`:

Scalars			
<code>e(N)</code>	number of observations	<code>e(sigma)</code>	estimates of σ from the first-stage regression
<code>e(Tbar)</code>	average number of time periods	<code>e(N_g)</code>	number of groups
Macros			
<code>e(cmd)</code>	<code>xtlsdvc</code>	<code>e(depvar)</code>	name of dependent variable
<code>e(ivar)</code>	panel variable	<code>e(predict)</code>	program used to implement <code>predict</code>
Matrices			
<code>e(b)</code>	<code>xtlsdvc</code> estimates	<code>e(V)</code>	variance-covariance matrix of the <code>xtlsdvc</code> estimator
<code>e(b_lsdv)</code>	<code>xtreg, fe</code> estimates		
<code>e(V_lsdv)</code>	variance-covariance matrix of the <code>xtreg, fe</code> estimator		
Functions			
<code>e(sample)</code>	marks estimation sample		

It is worth noting that the square root of the error variance estimate (6), saved in `e(sigma)`, uses residuals in levels computed via the first-stage coefficient estimates. As such, it need not coincide with the RMSE reported by Stata in the first-stage regression output, when AH is the chosen initialization, which is instead computed through first-differenced residuals. For the same reason, the squared value of `e(sigma)` does not coincide with the value of `e(sig2)` saved by `xtabond` when AB initializes `xtlsdvc`.

3.4 Syntax for predict

As with all Stata estimation commands, `xtlsdvc` supports the postestimation command `predict` ([R] `predict`) to compute fitted values and residuals. The syntax for `predict` following `xtlsdvc` is

```
predict [type] newvarname [if] [in] [, statistic]
```

where

<i>statistic</i>	description
<code>xb</code>	$\hat{\gamma}y_{i,t-1} + x'_{it}\hat{\beta}$, fitted values; the default
<code>ue</code>	$\hat{\eta}_i + \hat{\epsilon}_{it}$, the combined residuals
<code>*xbu</code>	$\hat{\gamma}y_{i,t-1} + x'_{it}\hat{\beta} + \hat{\eta}_i$, prediction, including fixed effect
<code>*u</code>	$\hat{\eta}_i$, the fixed effect
<code>*e</code>	$\hat{\epsilon}_{it}$, the observation-specific error component

Unstarred statistics are available both in and out of sample; type `predict ... if e(sample) ...` if wanted only for the estimation sample. Starred statistics are calculated only for the estimation sample, even when `if e(sample)` is not specified.

3.5 The bootstrap variance–covariance matrix

Kiviet and Bun (2001) show that LSDVC, however initialized, is asymptotically normal and derive the analytical expression for the asymptotic variance–covariance matrix of LSDVC in the version initialized by AH. Monte Carlo simulations therein, however, demonstrate that the analytical variance estimator performs poorly for a large γ , perhaps because of the unstable behavior of AH (documented also by the Monte Carlo analysis of this paper, see section 4). Alternatively, Kiviet and Bun (2001) suggest a parametric bootstrap procedure to estimate the asymptotic variance–covariance matrix of LSDVC, which seems superior to the analytical expression for at least three reasons: (1) it is simpler; (2) it always turns out to be relatively accurate; and (3) it can be applied to any version of LSDVC. Thus `xtlsdvc` adapts Kiviet and Bun’s (2001) bootstrap procedure for use with unbalanced panels, as described below.

A first difficulty here is brought about by the dependency in the data implied by the autoregressive data generation process (DGP), which does not permit us to adopt any of the official Stata bootstrap instructions, `bootstrap` and `bsample`. A parametric bootstrap is instead followed, which upon maintaining a normal distribution for the disturbances takes full account of the dependency in the DGP.

The subroutine `xtlsdvc_b` is called in `xtlsdvc` by the option `vcov()`. It is designed to yield a bootstrap sample and bootstrap LSDVC estimates and is iterated for `vcov(#)` times by `xtlsdvc`.

Let us focus on the generic iteration $(*)$ of `xtlsdvc_b`. It basically goes through the steps below.

1. Upon obtaining LSDVC estimates $\hat{\gamma}$ and $\hat{\beta}$ and $\hat{\sigma}^2$ from `xtlsdvc_1`, it calculates the N -vector of fixed-effect estimates $\hat{\eta} = \bar{y} - \hat{\gamma} \cdot \bar{y}_{-1} - \hat{\beta} \cdot \bar{x}$, where \bar{y} , \bar{y}_{-1} , and \bar{x} indicate N -vectors of group means.
2. It obtains bootstrap errors $\epsilon^{(*)}$ as a draw from $N(0, \hat{\sigma}^2)$.
3. Given x , S and y_0 , it obtains a bootstrap sample from $s_{it}y_{it}^{(*)} = s_{it}(\hat{\gamma} \cdot y_{i,t-1}^{(*)} + \hat{\beta} \cdot x_{it} + \hat{\eta}_i + \epsilon_{it}^{(*)})$, $i = 1, \dots, N$ and $t = 1, \dots, T$.
4. It applies LSDVC to $(y^{(*)}, S, x)$ to yield $\hat{\gamma}^{(*)}$ and $\hat{\beta}^{(*)}$.

While computational aspects of steps 1 and 2 are straightforward and step 4 only requires a call to `xtlsdvc_1` to calculate the corrected estimates from the generated bootstrap sample, step 3 is instructive and deserves some explanation. One possible way to implement step 3 would be to “manually” generate $y^{(*)}$ by recursion as a function of $\epsilon^{(*)}$, y_0 and x . But this is both computationally cumbersome and unnecessary in Stata. In fact, one can exploit the ability of `replace` ([D] **generate**) to work sequentially² to obtain $y^{(*)}$ in an effortless way:

2. I learned this by reading the messages by N. J. Cox and D. Kantor to Statalist on May 25, 2004, in response to a question of D. V. Masterov.

```
. by ivar: gen obs=_n
. replace y= GAMMA*L.y + BETA*x +THETA +EPSILON if obs>1.
```

Unbalancedness without gaps does not cause any trouble here, since different start-up dates can be dealt with very easily by the time-series operators in Stata. The presence of gaps, instead, may cause specific difficulty if they are found in any of the independent variables x 's, regardless of the way step 3 is implemented. In fact, since the recursion process generates $y^{(*)}$ from (y_0, S, x) , it must stop at the first missing value encountered in the x 's so that eventually a shorter sample is created at each replication. This decreases the accuracy of the estimates or even breaks down the identification of some coefficients in the shorter bootstrap sample and, consequently, of their standard errors. For example, if for all individuals there is a gap for a given time period, the coefficients on the time dummies corresponding and subsequent to the missing period would not be identified in each bootstrap sample so that their bootstrap standard errors could not be computed, too. To the opposite, gaps in the dependent variable are clearly immaterial for the size of the bootstrap samples since only the start-up values of y are used in the recursion process.

A `simulate` call ([R] **simulate**) in `xtlsdvc` replicates `xtlsdvc.b` for `vcov(#)` times, yielding a dataset of bootstrap LSDVC estimates $\hat{\delta}^*$, of dimension $(vcov \times k)$. Hence, `xtlsdvc` gets the bootstrap variance–covariance matrix V

$$V = \frac{\hat{\delta}^{*'} \hat{\delta}^*}{(vcov - 1)}$$

via `matrix accum ([P] matrix)`.

The bootstrap variance–covariance matrix V is then used to construct asymptotic t -ratio tests of parameter significance, as described in Kiviet and Bun (2001).

Attention should be paid when supplying the initial values through the matrix `my`. In this case, in fact, the bootstrap procedure would not be reliable since keeping the values in `my` fixed over replications neglects a source of variability for LSDVC so that the resulting bootstrap standard errors may be severely downward biased.

Finally, users should be warned that the bootstrap procedure may require a considerable amount of time. This tends to increase linearly with the number of replications. Also the procedure seems slightly faster if LSDVC is initialized by AH. Examples are given in the appendix.

4 Monte Carlo experiments

The Monte Carlo analyses in Kiviet (1995), Kiviet and Bun (2001), and especially, Judson and Owen (1999) provide support for LSDVC in balanced panels, compared to the traditional IV and GMM estimators. Moreover, Monte Carlo results in Bun and Kiviet (2003) for balanced panels and in Bruno (2005) for unbalanced panels demonstrate that the bias approximations (4), evaluated at the true γ and σ_e^2 , account for a significant

portion of the bias, never less than 90% and often virtually 100%. The relative merit of LSDVC in unbalanced panels is still to be explored, though. This is exactly what is accomplished here, where I evaluate the three versions of LSDVC as implemented by my code in a Monte Carlo study that extends Judson and Owen's (1999) under four respects. First, I evaluate LSDVC in the presence of various unbalanced designs; second, the performance of LSDVC is examined for the three different levels of accuracy; third, initial observations for the simulated data are generated following the procedure in McLeod and Hipel (1978), also adopted in Kiviet (1995) and Bruno (2005), which avoids wasting random numbers and small-sample nonstationary problems; finally, the comparison is extended to BB.

Data for y_{it} are generated by (1) with $k = 2$ and for x_{it} by

$$x_{it} = \rho x_{i,t-1} + \xi_{it}, \quad \xi_{it} \sim N(0, \sigma_\xi^2), \quad i = 1, \dots, N \text{ and } t = 1, \dots, T.$$

Initial observations y_{i0} and x_{i0} generated through the McLeod and Hipel (1978) procedure are kept fixed across replications. The long-run coefficient $\beta/(1 - \gamma)$ is kept fixed to unity, so $\beta = 1 - \gamma$; σ_ϵ^2 is normalized to unity; γ and ρ alternate between 0.2 and 0.8. The individual effects η_i are generated by assuming that $\eta_i \sim N(0, \sigma_\eta^2)$ and $\sigma_\eta = \sigma_\epsilon(1 - \gamma)$.

Two different sample sizes are considered: $(N, \bar{T}) = (20, 20)$ and $(N, \bar{T}) = (10, 40)$. Then following Baltagi and Chang (1995), I control for the extent of unbalancedness as measured by the Ahrens and Pincus index: $\omega = N / \left\{ \bar{T} \sum_{i=1}^N (1/T_i) \right\}$ ($0 < \omega \leq 1$, $\omega = 1$ when the panel is balanced). For each sample size, I analyze a case of mild unbalancedness ($\omega = 0.96$) and a case of severe unbalancedness ($\omega = 0.36$). Individuals are partitioned into two sets of equal dimension: one set contains the first $N/2$ individuals, each with the last h observations discarded, so $T_i = T - h$; the other contains the remaining $N/2$ individuals, each with $T_i = T$. I set T and h so that \bar{T} and ω take on the desired values (the four panel designs are summarized in table 1).

Table 1: Unbalanced designs

N	\bar{T}	T	T_i	ω
20	20	24	16 ($i \leq 10$), 24 ($i > 10$)	0.96
		36	4 ($i \leq 10$), 36 ($i > 10$)	0.36
10	40	48	32 ($i \leq 5$), 48 ($i > 5$)	0.96
		72	8 ($i \leq 5$), 72 ($i > 5$)	0.36

The simple AH estimator is the one chosen to initialize the correction procedure, based on the finding by Kiviet and Bun (2001) that differences in the initial estimators have only a marginal impact on the LSDVC performance. Then the LSDVC estimator is calculated for each of the three levels of accuracy in the estimated bias approximations.

4.1 Results

Results for γ are presented in figures 1 to 4, while results for β are presented in figures 5 to 8. In each figure, the first graph is for $\bar{T} = 20$ and the second for $\bar{T} = 40$. The bias and the RMSE are measured on the vertical axis, while the points on the horizontal axis always correspond to the eight possible combinations for γ , ρ , and ω . Since BB is specifically designed for highly persistent series, comparisons involving this estimator are restricted to $\gamma = 0.8$.

As a first general comment on the Monte Carlo results, I observe that according to a bias criterion, the three versions of LSDVC and, interestingly, AH have the best performances for both γ and β , with virtually zero bias in several cases. Turning to a RMSE criterion, the LSDVC estimators maintain the best performance, while AH shows the worst RMSE levels, also in comparison to LSDV, AB and, for highly persistent series, BB. This evidence highlights LSDVC as the preferred estimator for dynamic panel-data models with small N and strictly exogenous regressors, in line with that obtained by Kiviet (1995), Judson and Owen (1999), and Kiviet and Bun (2001) in similar Monte Carlo analyses.

That said, some interesting patterns seem to emerge when the behavior of each estimator is examined in more depth.

Estimating γ : bias

LSDVC₃ tends to perform slightly better than the other two LSDVC versions, especially when \bar{T} and γ increases. When $\gamma = 0.8$ and $\rho = 0.8$, however, all LSDVC estimators are slightly worse than AH (see figure 1).

After noting that the bias of LSDV and AB is always negative, confirming the findings by earlier studies (Kiviet and Bun 2001; Bond 2002; Bun and Kiviet 2003; Bruno 2005), I observe that LSDVC, LSDV, and AB estimators show similar patterns with respect to the degree of unbalancedness and average group size. As already shown in Bruno (2005) for LSDV, the biases of such estimators are decreasing in ω . This, always for AB and LSDV and often for LSDVC, brings with it an increase in the bias magnitude. When $\bar{T} = 20$, the AB estimator performs better than the LSDV estimator if ω is low but worse than the LSDV estimator when ω is high. When \bar{T} increases, however, besides observing an expected general tendency towards a smaller bias magnitude, I also notice an attenuation of the ω effect for all foregoing estimators. The bias of AH, instead, is always positive and increasing in ω , implying each time a worsening of the bias when unbalancedness reduces. The bias of BB is always positive and expected to be the largest in magnitude with lowly persistent series, but it dramatically improves when the persistence in y and x increases, reaching lower magnitudes than AB and LSDV when $\bar{T} = 20$ and comparable to AB and LSDV when $\bar{T} = 40$ (see figure 2).

Estimating γ : RMSE

The RMSE of the LSDVC estimators are almost coincident and always the smallest. On the other hand, AH almost always presents the highest RMSE, which hinders the attractiveness of such estimator in empirical work, despite its simplicity and good bias performance (see figure 3).

Except for BB, the RMSE for all estimators is increasing in γ and ρ , with the increase being especially large for AH. There is no apparent trend in the RMSE for BB. Similar to the previously discussed bias results, the RMSEs of the LSDVC, AB, and LSDV estimators are all increasing as the panel becomes closer to balanced. Again this effect is particularly strong for AB and when $\bar{T} = 20$. BB has a satisfactory RMSE in the presence of highly persistent series, performing generally better than AB and LSDV. In particular, when $\bar{T} = 40$ and $\omega = 0.96$, its RMSE gets very close to that of the LSDVC estimators (see figure 4).

The RMSE results for γ are summarized in table 2 ($\gamma = 0.2$) and table 3 ($\gamma = 0.8$), indicating for each case, the preferred estimator and its second best alternative.

Table 2: RMSE performance when $\gamma = 0.2$

$\omega \backslash \bar{T}$	20	40
0.36	1. LSDVC 2. AB (and LSDV if $\rho = 0.2$)	1. LSDVC 2. AB and LSDV
0.96	1. LSDVC 2. LSDV	1. LSDVC 2. AB and LSDV

Table 3: RMSE performance when $\gamma = 0.8$

$\omega \backslash \bar{T}$	20	40
0.36	1. LSDVC 2. BB	1. LSDVC 2. AB (and BB if $\rho = 0.8$)
0.96	1. LSDVC 2. BB	1. LSDVC and BB 2. LSDV

Estimating β : bias

LSDVC estimators and AH continue to show the best bias performance. While for $\rho = 0.2$ AB and LSDV also exhibit a negligible bias magnitude, for $\rho = 0.8$ their bias magnitude dramatically increases. With small \bar{T} , I notice a relatively bad performance of BB. When $\bar{T} = 40$ and $\omega = 0.36$, however, the bias attains acceptable levels and worsens when the degree of unbalancedness decreases (see figures 5 and 6).

Estimating β : RMSE

Results here parallel that evidenced for γ , with two differences: 1) There seems to be no clear role for the degree of unbalancedness. For example, when $\bar{T} = 20$, the RMSE of the LSDVC estimators benefits from a decreased unbalancedness, but when $\bar{T} = 40$ exactly the opposite occurs. 2) The RMSE for BB is now markedly increasing in ρ (see figures 7 and 8).

The documented evidence for a favorable impact of unbalancedness on bias and RMSE values in the estimation of γ , which is apparently surprising, can be explained by the fact that under investigation here is a notion of pure unbalancedness, not involving either gaps or any loss in degrees of freedom and average group size. Although more theoretical work, accompanied by broader Monte Carlo experiments, is needed to reach conclusive results on this issue, there is still a simple lesson to be learned from my Monte Carlo analysis; that is, smoothing unbalancedness at the cost of fewer time observations for the largest groups may be detrimental for estimation performance in dynamic panel-data models, especially if the average group size is small.

5 Conclusion

This paper has presented the new Stata code `xtlsdvc` implementing LSDVC estimators for dynamic (possibly) unbalanced panel-data models with a small N and strictly exogenous covariates. The procedure is based upon the bias approximations derived in Bruno (2005), who extends the result by Kiviet (1999) and Bun and Kiviet (2003) to unbalanced panels. The code also computes the bootstrap variance–covariance matrix of the estimators.

Monte Carlo experiments highlight the LSDVC estimators as the preferred ones in comparison to the original LSDV and widely used IV and GMM consistent estimators.

Future improvements of the code will enlarge the class of initial estimators, allowing more flexibility in defining the instrument set for the IV and GMM estimators.

6 Acknowledgments

I am grateful to an anonymous referee of this journal for helpful suggestions improving the presentation of the paper. I also benefited from useful discussions with Orietta Dessy and participants at the 10th UK Stata Users Group meeting, London 2004, and the 1st Italian Stata Users Group meeting, Rome 2004. Last but not least, I am grateful to all the Stata users who tested the `xtlsdvc` routine. In particular, Carl-Oskar Lindgren, Ivan Marinovic, and Clive Nicholas found bugs and provided helpful comments and suggestions. All remaining errors are my own. Financial support from Bocconi Ricerca di Base “Labor Demand, Production and Globalization” is gratefully acknowledged.

7 References

- Anderson, T. W. and C. Hsiao. 1982. Formulation and estimation of dynamic models using panel data. *Journal of Econometrics* 18: 570–606.
- Arellano, M. and S. Bond. 1991. Some tests of specification for panel data: Monte Carlo evidence and an application to employment equations. *Review of Economic Studies* 58: 277–297.
- Baltagi, B. H. and Y. J. Chang. 1995. Incomplete panels. *Journal of Econometrics* 62: 67–89.
- Blundell, R. and S. Bond. 1998. Initial conditions and moment restrictions in dynamic panel data models. *Journal of Econometrics* 87: 115–143.
- Bond, S. 2002. Dynamic panel data models: A guide to micro data methods and practice. Cemmap Working Paper CWP09/02.
- Bruno, G. S. F. 2005. Approximating the bias of the LSDV estimator for dynamic unbalanced panel data models. *Economics Letters* 87: 361–366.
- Bun, M. J. G. and J. F. Kiviet. 2003. On the diminishing returns of higher order terms in asymptotic expansions of bias. *Economics Letters* 79: 145–152.
- . 2005. The effects of dynamic feedbacks on LS and MM estimator accuracy in panel data models. *Journal of Econometrics*. Forthcoming.
- Judson, R. A. and A. L. Owen. 1999. Estimating dynamic panel data models: a guide for macroeconomists. *Economics Letters* 65: 9–15.
- Kiviet, J. F. 1995. On bias, inconsistency, and efficiency of various estimators in dynamic panel data models. *Journal of Econometrics* 68: 53–78.
- . 1999. Expectation of expansions for estimators in a dynamic panel data model; some results for weakly exogenous regressors. In *Analysis of Panels and Limited Dependent Variable Models*, ed. C. Hsiao, K. Lahiri, L.-F. Lee, and M. H. Pesaran, 199–225. Cambridge: Cambridge University Press.
- Kiviet, J. F. and M. J. G. Bun. 2001. The accuracy of inference in small samples of dynamic panel data models. Tinbergen Institute Discussion Paper TI 2001-006/4.
- McLeod, A. I. and K. W. Hipel. 1978. Simulation procedures for Box–Jenkins models. *Water Resources Research* 14: 969–975.
- Nickell, S. J. 1981. Biases in dynamic models with fixed effects. *Econometrica* 49: 1417–1426.
- Roodman, D. M. 2003. XTABOND2: Stata module to extend xtabond dynamic panel-data estimator. Statistical Software Components S435901, Boston College Department of Economics.

8 Appendix: Demonstrating xtlsdvc

I demonstrate the use of `xtlsdvc` in the context of labor demand estimation using the dataset `abdata.dta` (Arellano and Bond 1991), a typical micro panel of firm data with a moderately large N (140 firms). The labor demand of the firm is modeled according to specification (1), with the natural log of firm employment, n , as the dependent variable; the natural log of the real product wage, w ; the natural log of the gross capital stock, k ; and a set of time dummies as explanatory variables. The log of employment lagged one time is also included as a right-hand-side variable to allow costly employment adjustments.

Unlike in the customary approach, I do not use all information available to estimate the regression parameters. Instead, I follow a strategy that, exploiting the industry partition of the cross-sectional dimension as defined by the categorical variable `ind`, lets the slopes be industry-specific. This is easily accomplished by restricting the usable data to the panel of firms belonging to a given industry. While such a strategy leads to a less restrictive specification for the firm labor demand, it causes a reduced number of cross-sectional units for use in estimation so that the researcher must be prepared to deal with a potentially severe small-sample bias in any of the industry regressions. Clearly, `xtlsdvc` is the appropriate solution in this case.

The demonstration is kept as simple as possible by considering regressions for only one industry panel (`ind=4`).

Comparing two different initializations, AH and AB, I am able to confirm the feature found by Kiviet and Bun (2001) that differences in the initial estimators have only a marginal impact on the LSDVC estimates. Indeed, in this example, the evidence for the AB initialization is mixed. On the one hand, the one-step Sargan statistic suggests that the overidentifying restrictions used by AB are not satisfied. On the other hand, the second-order autocorrelation test does not reject the required lack of second-order autocorrelation in the differenced residuals. Be that as it may, the AB initialization has only negligible consequences on the resulting LSDVC estimates, as it clearly emerges upon comparing the latter with the LSDVC estimates initialized by AH.

The routine is reasonably fast when the bootstrap procedure is not invoked. Otherwise, the waiting time may be considerable, linearly increasing in the number of repetitions. To give you an idea of this, a message at the end of each execution displays the amount of time consumed by the code.

```
. use abdata, clear
. * Data description for industry 4
. xtides if ind==4
      id: 16, 18, ..., 133          n =          29
      year: 1976, 1977, ..., 1984    T =           9
      Delta(year) = 1; (1984-1976)+1 = 9
      (id*year uniquely identifies each observation)
Distribution of T_i:   min      5%      25%      50%      75%      95%      max
                     7         7         7         7         7         8         9
```

Freq.	Percent	Cum.	Pattern
18	62.07	62.07	1111111..
8	27.59	89.66	.1111111.
1	3.45	93.10	..1111111
1	3.45	96.55	.11111111
1	3.45	100.00	111111111
29	100.00		XXXXXXXXX

```
. set rmsg on
r; t=0.00 11:03:17
. * LSDVC initialized by AH.
. * Level 1 of accuracy.
. * AH and (uncorrected) LSDV estimates are also displayed.
. xtlsdvc n w k yr1977-yr1984 if ind==4, initial(ah) lsdv first
Note: Bias correction initialized by Anderson and Hsiao estimator
```

Instrumental variables (2SLS) regression

Source	SS	df	MS	Number of obs =	148
Model	1.35967485	10	.135967485	F(10, 138) =	.
Residual	.933924166	138	.006767566	Prob > F =	.
				R-squared =	.
				Adj R-squared =	.
Total	2.29359902	148	.015497291	Root MSE =	.08227

D.n	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
n						
LD.	.2204939	.4445225	0.50	0.621	-.658462	1.09945
w						
D1.	-.3771841	.134876	-2.80	0.006	-.643875	-.1104933
k						
D1.	.2204505	.0979079	2.25	0.026	.0268569	.4140442
yr1977						
D1.	.147631	.149344	0.99	0.325	-.1476674	.4429295
yr1978						
D1.	.1207165	.1386943	0.87	0.386	-.1535242	.3949572
yr1979						
D1.	.0977037	.1471064	0.66	0.508	-.1931704	.3885778
yr1980						
D1.	.0410339	.1448524	0.28	0.777	-.2453833	.3274512
yr1981						
D1.	-.0683895	.128972	-0.53	0.597	-.3234063	.1866273
yr1982						
D1.	-.1163022	.0788384	-1.48	0.142	-.2721896	.0395852
yr1983						
D1.	-.0512528	.0581115	-0.88	0.379	-.1661569	.0636513
yr1984						
D1.	(dropped)					

```
Instrumented: LD.n
Instruments: D.w D.k D.yr1977 D.yr1978 D.yr1979 D.yr1980 D.yr1981
              D.yr1982 D.yr1983 D.yr1984 L2.n
```

note: yr1984 dropped due to collinearity
in the LSDV regression
LSDV dynamic regression

	n	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
	n						
	L1.	.4056509	.0731424	5.55	0.000	.2622945	.5490074
	w	-.3541811	.1315442	-2.69	0.007	-.612003	-.0963593
	k	.2541555	.0525718	4.83	0.000	.1511167	.3571944
	yr1977	.0571224	.0614743	0.93	0.353	-.063365	.1776098
	yr1978	.0460914	.0619696	0.74	0.457	-.0753668	.1675497
	yr1979	.0147851	.0631942	0.23	0.815	-.1090733	.1386434
	yr1980	-.0403662	.0633203	-0.64	0.524	-.1644718	.0837394
	yr1981	-.1352945	.0620761	-2.18	0.029	-.2569615	-.0136275
	yr1982	-.1547943	.0570565	-2.71	0.007	-.266623	-.0429656
	yr1983	-.1019097	.0592481	-1.72	0.085	-.2180339	.0142145

note: Bias correction up to order $O(1/T)$

LSDVC dynamic regression
(SE not computed)

	n	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
	n						
	L1.	.5389829
	w	-.3375203
	k	.2218794
	yr1977	.030273
	yr1978	.0263007
	yr1979	-.005644
	yr1980	-.0604044
	yr1981	-.1508947
	yr1982	-.1562805
	yr1983	-.0928311

r; t=0.41 11:03:18

. * Level 2 of accuracy.
. xtlsdvc n w k yr1977-yr1984 if ind==4, initial(ah) bias(2)
Bias correction initialized by Anderson and Hsiao estimator

note: yr1984 dropped due to collinearity
in the LSDV regression

note: Bias correction up to order $O(1/NT)$

LSDVC dynamic regression
(SE not computed)

	n	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
	n						
	L1.	.5354691
	w	-.3380943
	k	.2226967
	yr1977	.0310655
	yr1978	.0269198
	yr1979	-.0050068
	yr1980	-.0597784
	yr1981	-.1503907
	yr1982	-.1561434
	yr1983	-.092829

```

r; t=0.39 11:03:18
. * Level 3 of accuracy
. xtlsdvc n w k yr1977-yr1984 if ind==4, initial(ah) bias(3)
Bias correction initialized by Anderson and Hsiao estimator
note: yr1984 dropped due to collinearity
      in the LSDV regression
note: Bias correction up to order  $O(1/NT^2)$ 
LSDVC dynamic regression
(SE not computed)

```

	n	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
n							
L1.		.6338054
w		-.3258186
k		.1988694
yr1977		.0112892
yr1978		.0123501
yr1979		-.0200475
yr1980		-.0745312
yr1981		-.1618727
yr1982		-.1572177
yr1983		-.0861093

```

r; t=0.39 11:03:18
. * LSDVC (level 3 of accuracy) initialized by AH, plus bootstrap SE
. * 100 replications
. xtlsdvc n w k yr1977-yr1984 if ind==4, initial(ah) bias(3) vcov(100)
Bias correction initialized by Anderson and Hsiao estimator
note: yr1984 dropped due to collinearity
      in the LSDV regression
note: Bias correction up to order  $O(1/NT^2)$ 
LSDVC dynamic regression
(bootstrapped SE)

```

	n	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
n							
L1.		.6338054	.2384333	2.66	0.008	.1664848	1.101126
w		-.3258186	.1624866	-2.01	0.045	-.6442865	-.0073507
k		.1988694	.0652599	3.05	0.002	.0709623	.3267765
yr1977		.0112892	.0908366	0.12	0.901	-.1667472	.1893257
yr1978		.0123501	.0928353	0.13	0.894	-.1696038	.194304
yr1979		-.0200475	.0956793	-0.21	0.834	-.2075756	.1674805
yr1980		-.0745312	.0988459	-0.75	0.451	-.2682655	.1192032
yr1981		-.1618727	.0885033	-1.83	0.067	-.335336	.0115906
yr1982		-.1572177	.0651537	-2.41	0.016	-.2849166	-.0295188
yr1983		-.0861093	.0703664	-1.22	0.221	-.224025	.0518064

```

r; t=38.47 11:03:57
. * 200 replications
. xtlsdvc n w k yr1977-yr1984 if ind==4, initial(ah) bias(3) vcov(200)
Bias correction initialized by Anderson and Hsiao estimator
note: yr1984 dropped due to collinearity
      in the LSDV regression
note: Bias correction up to order  $O(1/NT^2)$ 
LSDVC dynamic regression

```

(bootstrapped SE)

n	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
n						
L1.	.6338054	.2366395	2.68	0.007	.1700005	1.09761
w	-.3258186	.1740695	-1.87	0.061	-.6669885	.0153514
k	.1988694	.082856	2.40	0.016	.0364747	.3612641
yr1977	.0112892	.091363	0.12	0.902	-.1677789	.1903574
yr1978	.0123501	.0935808	0.13	0.895	-.1710649	.1957652
yr1979	-.0200475	.09732	-0.21	0.837	-.2107912	.1706962
yr1980	-.0745312	.0977652	-0.76	0.446	-.2661475	.1170852
yr1981	-.1618727	.088817	-1.82	0.068	-.335951	.0122055
yr1982	-.1572177	.0666269	-2.36	0.018	-.2878041	-.0266313
yr1983	-.0861093	.0714245	-1.21	0.228	-.2260987	.05388

r; t=76.44 11:05:13

. * LSDVC (level 3 of accuracy) initialized by AB,
. * plus bootstrap SE (100 replications).
. * AB estimates are also displayed.
. xtlsdvc n w k yr1977-yr1984 if ind==4, initial(ab) first bias(3) vcov(100)
Note: Bias correction initialized by Arellano and Bond estimator
note: yr1977 dropped due to collinearity

Arellano-Bond dynamic panel-data estimation Number of obs = 148
Group variable (i): id Number of groups = 29
Wald chi2(.) = .
Time variable (t): year Obs per group: min = 5
avg = 5.103448
max = 7

One-step results

D.n	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
n						
LD.	.2721012	.0875276	3.11	0.002	.1005503	.4436521
w						
D1.	-.4926766	.1138765	-4.33	0.000	-.7158704	-.2694828
k						
D1.	.2026031	.0527761	3.84	0.000	.0991637	.3060424
yr1978						
D1.	-.0219591	.0198588	-1.11	0.269	-.0608816	.0169633
yr1979						
D1.	-.0509516	.0202053	-2.52	0.012	-.0905533	-.01135
yr1980						
D1.	-.1080377	.0204241	-5.29	0.000	-.1480682	-.0680073
yr1981						
D1.	-.2176279	.0214474	-10.15	0.000	-.2596641	-.1755918
yr1982						
D1.	-.2527341	.0260614	-9.70	0.000	-.3038136	-.2016546
yr1983						
D1.	-.1992322	.0387691	-5.14	0.000	-.2752182	-.1232462
yr1984						
D1.	-.0629971	.0509664	-1.24	0.216	-.1628893	.0368952

Sargan test of over-identifying restrictions:

chi2(27) = 81.60 Prob > chi2 = 0.0000

Arellano-Bond test that average autocovariance in residuals of order 1 is 0:

H0: no autocorrelation z = -1.09 Pr > z = 0.2748

Arellano-Bond test that average autocovariance in residuals of order 2 is 0:

H0: no autocorrelation $z = -1.25$ $\Pr > z = 0.2129$

note: yr1984 dropped due to collinearity
in the LSDV regression

note: Bias correction up to order $O(1/NT^2)$

LSDVC dynamic regression
(bootstrapped SE)

n	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
n						
L1.	.6360273	.0912651	6.97	0.000	.4571509	.8149037
w	-.3256377	.143472	-2.27	0.023	-.6068377	-.0444377
k	.1988754	.0537594	3.70	0.000	.0935089	.304242
yr1977	.0080108	.0625058	0.13	0.898	-.1144982	.1305199
yr1978	.0097372	.0659415	0.15	0.883	-.1195059	.1389802
yr1979	-.0238944	.0678355	-0.35	0.725	-.1568495	.1090607
yr1980	-.0778375	.0684853	-1.14	0.256	-.2120662	.0563912
yr1981	-.1649284	.0675663	-2.44	0.015	-.2973559	-.032501
yr1982	-.1599435	.0592059	-2.70	0.007	-.2759848	-.0439021
yr1983	-.088907	.0644108	-1.38	0.167	-.2151498	.0373357

r; t=43.81 11:05:57

. set rmsg off

(Continued on next page)

9 Appendix: Figures





Figure 1: Biases of $LSDVC_1$, $LSDVC_2$, $LSDVC_3$, and AH for γ (γ, ρ, ω).

Figure 2: Biases of all estimators for γ (γ, ρ, ω).









Figure 3: RMSEs of $LSDVC_1$, $LSDVC_2$, $LSDVC_3$, and BB for γ (γ, ρ, ω).

Figure 4: RMSEs of all estimators for γ (γ, ρ, ω).

Figure 5: Biases of LSDVC₁, LSDVC₂, LSDVC₃, and AH for $\beta(\gamma, \rho, \omega)$.

Figure 6: Biases of all estimators for β (γ , ρ , ω).

Figure 7: RMSEs of LSDVC₁, LSDVC₂, LSDVC₃, and BB for $\beta(\gamma, \rho, \omega)$.

Figure 8: RMSEs of all estimators for β (γ , ρ , ω).