



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

THE STATA JOURNAL

Editor

H. Joseph Newton
Department of Statistics
Texas A&M University
College Station, Texas 77843
979-845-8817; fax 979-845-6077
jnewton@stata-journal.com

Editor

Nicholas J. Cox
Department of Geography
Durham University
South Road
Durham City DH1 3LE UK
n.j.cox@stata-journal.com

Associate Editors

Christopher F. Baum
Boston College

Nathaniel Beck
New York University

Rino Bellocco
Karolinska Institutet, Sweden, and
Univ. degli Studi di Milano-Bicocca, Italy

Maarten L. Buis
Vrije Universiteit, Amsterdam

A. Colin Cameron
University of California–Davis

Mario A. Cleves
Univ. of Arkansas for Medical Sciences

William D. Dupont
Vanderbilt University

David Epstein
Columbia University

Allan Gregory
Queen's University

James Hardin
University of South Carolina

Ben Jann
ETH Zürich, Switzerland

Stephen Jenkins
University of Essex

Ulrich Kohler
WZB, Berlin

Frauke Kreuter
University of Maryland–College Park

Jens Lauritsen
Odense University Hospital

Stanley Lemeshow
Ohio State University

J. Scott Long
Indiana University

Thomas Lumley
University of Washington–Seattle

Roger Newson
Imperial College, London

Austin Nichols
Urban Institute, Washington DC

Marcello Pagano
Harvard School of Public Health

Sophia Rabe-Hesketh
University of California–Berkeley

J. Patrick Royston
MRC Clinical Trials Unit, London

Philip Ryan
University of Adelaide

Mark E. Schaffer
Heriot-Watt University, Edinburgh

Jeroen Weesie
Utrecht University

Nicholas J. G. Winter
University of Virginia

Jeffrey Wooldridge
Michigan State University

Stata Press Editorial Manager

Stata Press Copy Editors

Lisa Gilmore

Jennifer Neve and Deirdre Patterson

The *Stata Journal* publishes reviewed papers together with shorter notes or comments, regular columns, book reviews, and other material of interest to Stata users. Examples of the types of papers include 1) expository papers that link the use of Stata commands or programs to associated principles, such as those that will serve as tutorials for users first encountering a new field of statistics or a major new technique; 2) papers that go “beyond the Stata manual” in explaining key features or uses of Stata that are of interest to intermediate or advanced users of Stata; 3) papers that discuss new commands or Stata programs of interest either to a wide spectrum of users (e.g., in data management or graphics) or to some large segment of Stata users (e.g., in survey statistics, survival analysis, panel analysis, or limited dependent variable modeling); 4) papers analyzing the statistical properties of new or existing estimators and tests in Stata; 5) papers that could be of interest or usefulness to researchers, especially in fields that are of practical importance but are not often included in texts or other journals, such as the use of Stata in managing datasets, especially large datasets, with advice from hard-won experience; and 6) papers of interest to those who teach, including Stata with topics such as extended examples of techniques and interpretation of results, simulations of statistical concepts, and overviews of subject areas.

For more information on the *Stata Journal*, including information for authors, see the web page

<http://www.stata-journal.com>

The *Stata Journal* is indexed and abstracted in the following:

- Science Citation Index Expanded (also known as SciSearch®)
- CompuMath Citation Index®

Copyright Statement: The *Stata Journal* and the contents of the supporting files (programs, datasets, and help files) are copyright © by StataCorp LP. The contents of the supporting files (programs, datasets, and help files) may be copied or reproduced by any means whatsoever, in whole or in part, as long as any copy or reproduction includes attribution to both (1) the author and (2) the *Stata Journal*.

The articles appearing in the *Stata Journal* may be copied or reproduced as printed copies, in whole or in part, as long as any copy or reproduction includes attribution to both (1) the author and (2) the *Stata Journal*.

Written permission must be obtained from StataCorp if you wish to make electronic copies of the insertions. This precludes placing electronic copies of the *Stata Journal*, in whole or in part, on publicly accessible web sites, file servers, or other locations where the copy may be accessed by anyone other than the subscriber.

Users of any of the software, ideas, data, or other materials published in the *Stata Journal* or the supporting files understand that such use is made without warranty of any kind, by either the *Stata Journal*, the author, or StataCorp. In particular, there is no warranty of fitness of purpose or merchantability, nor for special, incidental, or consequential damages such as loss of profits. The purpose of the *Stata Journal* is to promote free communication among Stata users.

The *Stata Journal*, electronic version (ISSN 1536-8734) is a publication of Stata Press. Stata and Mata are registered trademarks of StataCorp LP.

Tests for unbalanced error-components models under local misspecification

Walter Sosa-Escudero
Department of Economics
Universidad de San Andrés
Buenos Aires, Argentina
wsosa@udesa.edu.ar

Anil K. Bera
Department of Economics
University of Illinois at Urbana–Champaign
Champaign, IL

Abstract. This paper derives *unbalanced* versions of the test statistics for first-order serial correlation and random individual effects summarized in Sosa-Escudero and Bera (2001, *Stata Technical Bulletin Reprints*, vol. 10, pp. 307–311), and updates their `xttest1` routine. The derived test statistics should be useful for applied researchers faced with the increasing availability of panel information where not every individual or country is observed for the full time span. The test statistics proposed here are based on ordinary least-squares residuals and hence are computationally very simple.

Keywords: sg164_1, `xttest1`, error-components model, unbalanced panel data, testing, misspecification

1 Introduction

A standard specification check that accompanies the output of almost every estimated error-components model is a simple test for the presence of random individual effects. The well-known Breusch–Pagan statistic (Breusch and Pagan 1980), based on the Rao-score (RS) principle, is a frequent choice. Bera, Sosa-Escudero, and Yoon (2001) demonstrated that, in the presence of first-order serial correlation, the test too often rejects the correct null hypothesis of no random effects. Consequently, they propose a modified version that is not affected by the presence of local serial correlation. A similar concern affects the standard test for first-order serial correlation derived by Baltagi and Li (1991), which overrejects the true null hypothesis when random effects are present. For this case, an adjusted RS test was also derived by Bera, Sosa-Escudero, and Yoon (2001). These test statistics, along with their `xttest1` routine in Stata and some empirical illustrations, are presented in Sosa-Escudero and Bera (2001). For a textbook exposition, see Baltagi (2005, 96–97).

These test procedures were originally derived for the *balanced* case, that is, in the panel-data terminology, the case where all individuals are observed for the same number of periods, and in every period all individuals are observed. On the other hand, in applied work the availability of *unbalanced* panels is far from being an uncommon situation. Though in some cases statistical procedures designed for the balanced case can be straightforwardly extended to accommodate unbalanced panels, many estimation or test procedures require less trivial modifications.

Baltagi and Li (1990) derived an unbalanced version of the Breusch–Pagan statistic. The purpose of this paper is to derive unbalanced versions of the test for first-order serial correlation originally proposed by Baltagi and Li (1991) and of the modified tests proposed by Bera, Sosa-Escudero, and Yoon (2001). As a simple extension, we also derive an unbalanced version of the joint test of serial correlation and random effects proposed by Baltagi and Li (1991). The derived test statistics, being based on ordinary least-squares residuals after pooled estimation, are computationally very simple. Finally, the Sosa-Escudero and Bera (2001) `xttest1` routine is appropriately updated to handle unbalanced panels.

2 Tests for the unbalanced case

Consider a simple linear model for panel data allowing for the presence of random individual effects and first-order serial correlation:

$$\begin{aligned} y_{it} &= x'_{it}\beta + u_{it} \\ u_{it} &= \mu_i + \nu_{it} \\ \nu_{it} &= \lambda\nu_{i,t-1} + \epsilon_{it}, \quad |\lambda| < 1 \end{aligned}$$

where x_{it} is a $k \times 1$ vector of explanatory variables with 1 in its first position, β is a $k \times 1$ vector of parameters including an intercept, $\mu_i \sim N(0, \sigma_\mu^2)$, and $\epsilon_{it} \sim N(0, \sigma_\epsilon^2)$. We will assume $\nu_{i,0} \sim N\{0, \sigma_\epsilon^2/(1 - \lambda^2)\}$.

We will be interested in testing for the absence of random effects ($H_0 : \sigma_\mu^2 = 0$) and/or first-order serial correlation ($H_0 : \lambda = 0$). The panel will be unbalanced in the sense that for every individual $i = 1, \dots, N$ we will observe, possibly, a different number of time observations T_i . We will restrict the analysis to the cases where missing observations occur either at the beginning or at the end of the sample period for each individual (that is, there are no “gaps” in the series), and the starting and final periods are determined randomly. Hence, without loss of generality and to avoid complicating the notation too much, we can safely assume that the series for each individual starts at the same period ($t = 1$) and finish randomly at period $t = T_i$.

Let $m = \sum_{i=1}^N T_i$ be the total number of observations. Let u be an $m \times 1$ vector with typical element u_{it} where observations are sorted first by individuals and then by time, so the time index is the faster one. Then in our setup, $V(u) \equiv \Omega$ can be written as

$$V(u) = \sigma_\mu^2 \tilde{H} + \sigma_\epsilon^2 \tilde{V}$$

where \tilde{H} is an $m \times m$ block diagonal matrix with blocks H_i equal to matrices of ones, each with dimensions $T_i \times T_i$. Similarly, \tilde{V} will be a block diagonal $m \times m$ matrix with blocks V_i equal to

$$V_i = \begin{bmatrix} 1 & \lambda & \lambda^2 & \dots & \lambda^{T_i-1} \\ \lambda & 1 & \lambda & \dots & \lambda^{T_i-2} \\ \vdots & \vdots & \vdots & & \vdots \\ \lambda^{T_i-1} & \lambda^{T_i-2} & \lambda^{T_i-3} & \dots & 1 \end{bmatrix}$$

For the purpose of deriving the test statistics, the log-likelihood function will be

$$L(\beta, \lambda, \sigma_\epsilon^2, \sigma_\mu^2) = \text{constant} - \frac{1}{2} \log |\Omega| - \frac{1}{2} u' \Omega^{-1} u$$

The information matrix for this problem is known to be block diagonal between β and the remaining parameters. Therefore, for the purposes of this paper, we will concentrate only on the parameters λ, σ_μ^2 , and σ_ϵ^2 . Under a more general setup, suppose the log likelihood can be characterized by a three-parameter vector $\theta = (\psi, \phi, \gamma)'$. Let $d(\theta)$ be the score vector and $J(\theta)$ the information matrix. If it can be assumed that $\phi = 0$, the standard Rao-score (RS) test statistic for the null hypothesis $H_0: \psi = 0$ is given by

$$\text{RS}_\psi = d_\psi(\hat{\theta}) J_{\psi \cdot \gamma}^{-1}(\hat{\theta}) d_\psi(\hat{\theta}) \quad (1)$$

where d_ψ is the element of the score corresponding to the parameter ψ , $J_{\psi \cdot \gamma}(\theta) = J_\psi - J_{\psi \gamma} J_\gamma^{-1} J_{\gamma \psi}$, and $\hat{\theta}$ is the maximum likelihood estimator (MLE) of θ under the restriction implied by the null hypothesis and the assumption $\phi = 0$. Asymptotically, this test statistic under the null hypothesis $H_0: \psi = 0$ is known to have a central chi-squared distribution. In the context of our error-components model, if $\gamma = \sigma_\epsilon^2$ and if we set $\psi = \sigma_\mu^2$ and $\phi = \lambda$, (1) is a test for random effects assuming no serial correlation; and if we set $\psi = \lambda$ and $\phi = \sigma_\mu^2$, (1) gives a test for serial correlation assuming no random effects. The standard Breusch–Pagan test for random effects (assuming no serial correlation) and the Baltagi–Li test for first-order serial correlation (assuming no random effects) are derived from this principle.

Bera and Yoon (1993) showed that the test statistic (1) is invalid when $\phi \neq 0$, in the sense that the test tends to reject the null hypothesis too frequently even when it is correct. More specifically, the RS_ψ statistic is found to have an asymptotic *noncentral* chi-squared distribution under $H_0: \psi = 0$, when $\phi = \delta/\sqrt{n}$, that is, when the alternative is *locally misspecified*. In particular, this implies that when the null is correct, the Breusch–Pagan test tends to reject the true null of absence of random effects if the error term is serially correlated, even in a local sense. A similar situation arises for the test for serial correlation of Baltagi and Li (1991) in the local presence of random effects. In order to remedy this problem, Bera and Yoon (1993) proposed the following modified RS statistic:

$$\begin{aligned}
\text{RS}_\psi^* &= \frac{1}{n} \left\{ d_\psi(\hat{\theta}) - J_{\psi\phi\cdot\gamma}(\hat{\theta}) J_{\phi\cdot\gamma}^{-1}(\hat{\theta}) d_\phi(\hat{\theta}) \right\}' \\
&\quad \left\{ J_{\psi\cdot\gamma}(\hat{\theta}) - J_{\psi\phi\cdot\gamma}(\hat{\theta}) J_{\phi\cdot\gamma}^{-1}(\hat{\theta}) J_{\phi\psi\cdot\gamma}(\hat{\theta}) \right\}^{-1} \\
&\quad \left\{ d_\psi(\hat{\theta}) - J_{\psi\phi\cdot\gamma}(\hat{\theta}) J_{\phi\cdot\gamma}^{-1}(\hat{\theta}) d_\phi(\hat{\theta}) \right\}
\end{aligned} \tag{2}$$

where $\hat{\theta}$ is the MLE of θ under the joint null $\psi = \phi = 0$. This modified test statistic has an asymptotic *central* χ_1^2 distribution under the null hypothesis $H_0: \psi = 0$ and when $\phi = \delta/\sqrt{n}$, that is, the modified test statistic has the correct size even when the underlying model is locally misspecified. Based on this principle, [Bera, Sosa-Escudero, and Yoon \(2001\)](#) derived modified tests for random effects (serial correlation), which are valid in the presence of local first-order serial correlation (random effects) assuming that the panel is balanced.

To derive tests for the unbalanced case, let $\theta = (\lambda, \sigma_\mu^2, \sigma_\epsilon^2)'$ and $\hat{\theta} = (0, 0, \hat{\sigma}_\epsilon^2)'$ be the MLE of θ under the joint null hypothesis $H_0: \lambda = \sigma_\mu^2 = 0$. The following formula by [Hemmerle and Hartley \(1973\)](#) will be useful to derive the score vector for the problem:

$$d_{\theta_r} \equiv \frac{\partial L}{\partial \theta_r} = -\frac{1}{2} \text{tr} \left(\Omega^{-1} \frac{\partial \Omega}{\partial \theta_r} \right) + \frac{1}{2} \left(u' \Omega^{-1} \frac{\partial \Omega}{\partial \theta_r} \Omega^{-1} u \right) \tag{3}$$

where θ_r denotes the r th element of θ , $r = 1, 2, 3$. Note that $\partial \Omega / \partial \sigma_\mu^2 = \tilde{H}$ with $\text{tr}(\tilde{H}) = m$. Similarly, $\partial \Omega / \partial \sigma_\epsilon^2 = \tilde{V}$, which under the restricted MLE is an $m \times m$ identity matrix with trace equal to m . Also $\partial \Omega / \partial \lambda = \sigma_\epsilon^2 \tilde{G}$, where \tilde{G} is a block diagonal matrix with blocks equal to G_i , with $G_i = \partial V_i / \partial \lambda$ given by

$$G_i = \begin{bmatrix} 0 & 1 & 2\lambda & \cdots & (T_i - 1)\lambda^{T_i-2} \\ 1 & 0 & 1 & \cdots & (T_i - 2)\lambda^{T_i-3} \\ \vdots & \vdots & \vdots & & \vdots \\ \vdots & \vdots & 1 & 0 & 1 \\ (T_i - 1)\lambda^{T_i-2} & \cdots & \cdots & 1 & 0 \end{bmatrix}$$

Under the restricted MLE, G_i is a bidiagonal matrix as follows:

$$G_i(\hat{\theta}) = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 1 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & \cdots & 1 & 0 & 1 \\ 0 & \cdots & \cdots & 1 & 0 \end{bmatrix}$$

Hence, $\text{tr} \{ G_i(\hat{\theta}) \} = 0$. Replacing these results in (3) and evaluating the expression under the restricted MLE, we obtain

$$\begin{aligned}
d_{\sigma_\mu^2}(\hat{\theta}) &= -\frac{1}{2}\text{tr}\left(\frac{1}{\hat{\sigma}_\epsilon^2}I_m\tilde{H}\right) + \frac{1}{2}e'\frac{1}{\hat{\sigma}_\epsilon^2}I_m\tilde{H}\frac{1}{\hat{\sigma}_\epsilon^2}e \\
&= -\frac{1}{2}\frac{1}{\hat{\sigma}_\epsilon^2}m + \frac{1}{2}\frac{1}{\hat{\sigma}_\epsilon^4}e'\tilde{H}e = -\frac{m}{\hat{\sigma}_\epsilon^2}A
\end{aligned}$$

where e is an $m \times 1$ vector with typical element $e_{it} = x'_{it}\hat{\beta}$, and $\hat{\beta}$ is the restricted MLE of β . Similarly, $\hat{\sigma}_\epsilon^2 = e'e/m$ is the restricted MLE of σ_ϵ^2 , and $A \equiv 1 - e'\tilde{H}e/(e'e)$. In a similar fashion,

$$\begin{aligned}
d_\lambda(\hat{\theta}) &= -\frac{1}{2}\text{tr}\left\{\frac{1}{\hat{\sigma}_\epsilon^2}\hat{\sigma}_\epsilon^2\tilde{G}(\hat{\theta})\right\} + \frac{1}{2}2\frac{1}{\hat{\sigma}_\epsilon^2}e'\tilde{G}(\hat{\theta})e \\
&= \frac{1}{\hat{\sigma}_\epsilon^2}e'\tilde{G}(\hat{\theta})e = mB
\end{aligned}$$

where $B \equiv e'\tilde{G}e/e'e$.

To derive the elements of the information matrix, we will use the following formula from Baltagi (2005, 59–60):

$$J_{r,s}(\theta) = E\left(-\frac{\partial^2 L}{\partial \theta_r \partial \theta_s}\right) = \frac{1}{2}\text{tr}\left(\Omega^{-1}\frac{\partial \Omega}{\partial \theta_r}\Omega^{-1}\frac{\partial \Omega}{\partial \theta_s}\right)$$

Then

$$\begin{aligned}
J_{\sigma_\epsilon^2, \sigma_\epsilon^2}(\hat{\theta}) &= \frac{1}{2}\text{tr}\left\{\frac{1}{\hat{\sigma}_\epsilon^2}\tilde{V}(\hat{\theta})\frac{1}{\hat{\sigma}_\epsilon^2}\tilde{V}(\hat{\theta})\right\} = \frac{1}{2}\text{tr}\left(\frac{1}{\hat{\sigma}_\epsilon^4}I_m\right) = \frac{m}{2\hat{\sigma}_\epsilon^4} \\
J_{\hat{\sigma}_\mu^2, \hat{\sigma}_\mu^2}(\hat{\theta}) &= \frac{1}{2}\text{tr}\left(\frac{1}{\hat{\sigma}_\epsilon^2}\tilde{H}\frac{1}{\hat{\sigma}_\epsilon^2}\tilde{H}\right) = \frac{1}{2}\frac{1}{\hat{\sigma}_\epsilon^4}\text{tr}(\tilde{H}\tilde{H}) = \frac{\sum_{i=1}^N T_i^2}{2\hat{\sigma}_\epsilon^4} \\
J_{\lambda, \lambda}(\hat{\theta}) &= \frac{1}{2}\text{tr}\left\{\frac{1}{\hat{\sigma}_\epsilon^2}\hat{\sigma}_\epsilon^2\tilde{G}(\hat{\theta})\frac{1}{\hat{\sigma}_\epsilon^2}\hat{\sigma}_\epsilon^2\tilde{G}(\hat{\theta})\right\} = \frac{1}{2}\text{tr}\left\{\tilde{G}(\hat{\theta})\tilde{G}(\hat{\theta})\right\} \\
&= \frac{1}{2}\sum_{i=1}^N 2(T_i - 1) = m - N \\
J_{\hat{\sigma}_\epsilon^2, \hat{\sigma}_\mu^2}(\hat{\theta}) &= \frac{1}{2}\text{tr}\left\{\frac{1}{\hat{\sigma}_\epsilon^2}\tilde{V}(\hat{\theta})\frac{1}{\hat{\sigma}_\epsilon^2}\tilde{V}(\hat{\theta})\right\} = \frac{1}{2}\frac{1}{\hat{\sigma}_\epsilon^4}\text{tr}(\tilde{H}) = \frac{m}{2\hat{\sigma}_\epsilon^4} \\
J_{\hat{\sigma}_\epsilon^2, \lambda}(\hat{\theta}) &= \frac{1}{2}\text{tr}\left\{\frac{1}{\hat{\sigma}_\epsilon^2}\tilde{V}(\hat{\theta})\frac{1}{\hat{\sigma}_\epsilon^2}\tilde{G}(\hat{\theta})\right\} = \frac{1}{2}\frac{1}{\hat{\sigma}_\epsilon^4}\text{tr}\left\{\tilde{G}(\hat{\theta})\right\} = 0 \\
J_{\lambda, \hat{\sigma}_\mu^2}(\hat{\theta}) &= \frac{1}{2}\text{tr}\left\{\frac{1}{\hat{\sigma}_\epsilon^2}\hat{\sigma}_\epsilon^2\tilde{G}(\hat{\theta})\frac{1}{\hat{\sigma}_\epsilon^2}\tilde{H}\right\} = \frac{1}{2}\frac{1}{\hat{\sigma}_\epsilon^2}\text{tr}\left\{\tilde{G}(\hat{\theta})\tilde{H}\right\} \\
&= \frac{1}{2}\frac{2}{\hat{\sigma}_\epsilon^2}\left(\sum_{i=1}^N T_i - N\right) = \frac{1}{\hat{\sigma}_\epsilon^2}(m - N)
\end{aligned}$$

where we have used the facts that $\text{tr} \left\{ \tilde{G}_i(\hat{\theta}) \tilde{G}_i(\hat{\theta}) \right\} = \text{tr} \left\{ \tilde{G}_i(\hat{\theta}) \tilde{H}_i \right\} = 2(T_i - 1)$, and $\text{tr}(\tilde{H}_i \tilde{H}_i) = T_i^2$.

Collecting all the elements, the information matrix evaluated at the restricted MLE under the joint null can be expressed as

$$J(\hat{\theta}) = \frac{1}{2\hat{\sigma}_\epsilon^4} \begin{bmatrix} m & m & 0 \\ m & a & 2\hat{\sigma}_\epsilon^2(m - N) \\ 0 & 2\hat{\sigma}_\epsilon^2(m - N) & 2\hat{\sigma}_\epsilon^4(m - N) \end{bmatrix}$$

where $a \equiv \sum_{i=1}^N T_i^2$. For the balanced case $T_i = T$, we get exactly the same expression for $J(\hat{\theta})$ as in [Baltagi and Li \(1991, 279\)](#). From the above expression of $J(\hat{\theta})$, we can show that

$$\begin{aligned} J_{\mu\lambda \cdot \sigma_\epsilon^2} &= \frac{m - N}{\hat{\sigma}_\epsilon^2} \\ J_{\mu \cdot \sigma_\epsilon^2} &= \frac{a - m}{2\hat{\sigma}_\epsilon^4} \\ J_{\lambda \cdot \sigma_\epsilon^2} &= m - N \end{aligned}$$

Substituting these results in (2), we obtain the unbalanced version of the modified test for random effects as

$$\text{RS}_\mu^* = \frac{m^2 (A + 2B)^2}{2(a - 3m + 2N)}$$

When $T_i = T$ (the balanced case), the above expression boils down to

$$\text{RS}_\mu^* = \frac{NT (A + 2B)^2}{2(T - 1) \{1 - (2/T)\}}$$

as in [Bera, Sosa-Escudero, and Yoon \(2001\)](#) for the balanced case.

Similarly, the modified test statistic for serial correlation is

$$\text{RS}_\lambda^* = \left(B + \frac{m - N}{a - m} A \right)^2 \frac{(a - m)m^2}{(m - N)(a - 3m + 2N)}$$

and when $T_i = T$, we get

$$\text{RS}_\lambda^* = \left(B + \frac{A}{T} \right)^2 \frac{NT^2}{(T - 1)(1 - 2/T)}$$

which is the expression in [Bera, Sosa-Escudero, and Yoon \(2001\)](#) for the balanced case.

For computational purposes, it is interesting to see that

$$A = 1 - \frac{\sum_{i=1}^N \left(\sum_{t=1}^{T_i} e_{it}^2 \right)^2}{\sum_{i=1}^N \sum_{t=1}^{T_i} e_{it}^2}$$

and

$$B = \frac{\sum_{i=1}^N \sum_{t=2}^{T_i} e_{i,t} e_{i,t-1}}{\sum_{i=1}^N \sum_{t=1}^{T_i} e_{it}^2}$$

and, therefore, there is no need to construct the \tilde{G} or \tilde{H} matrices; hence, the test statistics can be easily computed right after ordinary least-squares estimation without constructing any matrices.

The previous derivations allow us to obtain the unbalanced version of the test for serial correlation assuming no random effects:

$$\text{RS}_\lambda = \frac{m^2 B^2}{m - N}$$

which again reduces to $NT^2 B^2 / (T - 1)$, originally derived by Baltagi and Li (1991) for balanced panels. Also, for completeness, the unbalanced version of the test for random effects assuming no serial correlation is given by

$$\text{RS}_\mu = \frac{\frac{1}{2} m^2 A^2}{a - m}$$

This test statistic is a particular case of the Baltagi–Li test for the two-way error-components model.

Suppose that we are interested in the joint null hypothesis of no random effects and no first-order serial correlation. Let $\text{RS}_{\phi, \psi}$ be the RS test statistic for the joint null hypothesis $H_0 : \phi = \psi = 0$. Bera and Yoon (2001) show that the following identities hold:

$$\text{RS}_{\phi\psi} = \text{RS}_\psi^* + \text{RS}_\phi = \text{RS}_\phi^* + \text{RS}_\psi$$

This simplifies computations, as illustrated in Sosa-Escudero and Bera (2001). Then, as a simple byproduct of the previous derivations, we can obtain a statistic for jointly testing serial correlation and random effects, as

$$\text{RS}_{\lambda\mu} = m^2 \left\{ \frac{A^2 + 4AB + 4B^2}{2(a - 3m + 2N)} + \frac{B^2}{m - N} \right\}$$

When $T_i = T$, $RS_{\lambda\mu}$ simplifies to

$$RS_{\lambda\mu} = \frac{NT^2}{2(T-1)(T-2)} (A^2 + 4AB + 2TB^2)$$

which is the original joint test statistic of Baltagi and Li (1991).

Finally, because $\sigma_\mu^2 \geq 0$, it is natural to consider one-sided versions of the tests for the null $H_0 : \sigma_\mu^2 = 0$. As in Bera, Sosa-Escudero, and Yoon (2001), appropriate test statistics can be readily constructed by taking the signed square roots of the original two-sided tests RS_μ and RS_μ^* . Denoting their one-sided versions, respectively, as RSO_μ and RSO_μ^* , we have

$$RSO_\mu = -\sqrt{\frac{\frac{1}{2} m^2}{a - m}} A$$

and

$$RSO_\mu^* = -\sqrt{\frac{m^2}{2(a - 3m + 2N)}} (A + 2B)$$

3 Empirical illustration

As an illustration of these procedures, we provide an empirical exercise that is based on Gasparini, Marchionni, and Sosa-Escudero (2001). It consists of a simple linear panel-data model where the dependent variable is the Gini coefficient for 17 regions of Argentina. The vector of explanatory variables includes mean income and its square (**ie** and **ie2**); proportion of the population employed in the manufacturing industry (**indus**) and in public administration, health, or education (**adpubedsal**); unemployment rate (**desempleo**); activity rate (**tactiv**); public investment as percentage of GDP (**invipib**); degree of openness (**apertura**); social assistance (**pyas4**); proportion of population older than 64 (**e64**); proportion of population that completed high school (**supc**); and average family size (**tamfam**); for details see Gasparini, Marchionni, and Sosa-Escudero (2001). Models of this type have been used extensively in the literature exploring the links between inequality and development, usually to study the so-called “Kuznets hypothesis”, which postulates an inverted *U*-shaped relationship between these two variables (for example, see Anand and Kanbur [1993] and Gustafsson and Johansson [1999]).

Income-related variables, including the Gini coefficients, are constructed using Argentina’s *Permanent Household Survey* (Encuesta Permanente de Hogares), which surveys several socioeconomic variables at the household level for several regions of the country. Because of certain administrative deficiencies, the panel is largely unbalanced, so the number of available temporal observations ranges from 5 to 8 years in the period 1992–2000.

First, we `tsset` the data and then use `xtreg` to estimate the parameters of a one-way error-components model with region-specific random effects:

```
. use ginipanel5
. tsset naglo ano
    panel variable:  naglo (unbalanced)
    time variable:  ano, 1992 to 2000, but with a gap
                delta:  1 unit
. xtreg gini ie ie2 indus adpubedsal desempleo tactiv invipib apertura pyas4
> e64 supc tamfam, re i(naglo)

Random-effects GLS regression                Number of obs   =       128
Group variable: naglo                      Number of groups  =        17

R-sq:  within = 0.5096                     Obs per group: min =         6
        between = 0.6153                             avg   =       7.5
        overall = 0.5344                             max   =         8

Random effects u_i ~ Gaussian              Wald chi2(12)     =    121.30
corr(u_i, X)      = 0 (assumed)           Prob > chi2      =     0.0000
```

| gini | Coef. | Std. Err. | z | P> z | [95% Conf. Interval] | |
|------------|-----------|-----------------------------------|-------|-------|----------------------|----------|
| ie | -.0000995 | .0001823 | -0.55 | 0.585 | -.0004568 | .0002578 |
| ie2 | 1.64e-08 | 2.19e-07 | 0.08 | 0.940 | -4.12e-07 | 4.45e-07 |
| indus | -.041974 | .0704982 | -0.60 | 0.552 | -.1801478 | .0961999 |
| adpubedsal | -.0635789 | .0531777 | -1.20 | 0.232 | -.1678053 | .0406475 |
| desempleo | -.1177452 | .0638999 | -1.84 | 0.065 | -.2429868 | .0074963 |
| tactiv | .0999584 | .0737997 | 1.35 | 0.176 | -.0446864 | .2446031 |
| invipib | -.3307239 | .1912258 | -1.73 | 0.084 | -.7055197 | .0440718 |
| apertura | .4289793 | .0768693 | 5.58 | 0.000 | .2783183 | .5796404 |
| pyas4 | 2.884162 | 1.626136 | 1.77 | 0.076 | -.3030061 | 6.071331 |
| e64 | -.1339182 | .1505384 | -0.89 | 0.374 | -.4289681 | .1611316 |
| supc | .2417907 | .0946423 | 2.55 | 0.011 | .0562952 | .4272861 |
| tamfam | .0169905 | .0174328 | 0.97 | 0.330 | -.0171771 | .0511581 |
| _cons | .3084864 | .1031351 | 2.99 | 0.003 | .1063453 | .5106274 |
| sigma_u | .01370805 | | | | | |
| sigma_e | .01377936 | | | | | |
| rho | .49740589 | (fraction of variance due to u_i) | | | | |

Next the command `xttest1` with the `unadjusted` option presents the following output:

```
. xttest1, unadjusted
Tests for the error component model:
      gini[naglo,t] = Xb + u[naglo] + v[naglo,t]
      v[naglo,t] = lambda v[naglo,(t-1)] + e[naglo,t]
Estimated results:

```

| | Var | sd = sqrt(Var) |
|------|----------|----------------|
| gini | .0006167 | .0248335 |
| e | .0001899 | .01377936 |
| u | .0001879 | .01370805 |

```

Tests:
Random Effects, Two Sided:
LM(Var(u)=0)      =    13.50 Pr>chi2(1) =   0.0002
ALM(Var(u)=0)     =     6.03 Pr>chi2(1) =   0.0141
Random Effects, One Sided:
LM(Var(u)=0)      =     3.67 Pr>N(0,1) =   0.0001
ALM(Var(u)=0)     =     2.46 Pr>N(0,1) =   0.0070
Serial Correlation:
LM(lambda=0)      =     9.32 Pr>chi2(1) =   0.0023
ALM(lambda=0)     =     1.86 Pr>chi2(1) =   0.1732
Joint Test:
LM(Var(u)=0,lambda=0) =   15.35 Pr>chi2(2) =   0.0005

```

The unadjusted version of the tests for random effects ($\text{LM}(\text{Var}(u)=0)$) and serial correlation ($\text{LM}(\text{lambda}=0)$), and the test for the joint null ($\text{LM}(\text{Var}(u)=0, \text{lambda}=0)$) suggest rejecting their nulls at the 5% significance level. Care must be taken in deriving conclusions about the direction of the misspecification because, in light of the results in [Bera, Sosa-Escudero, and Yoon \(2001\)](#), rejections may arise because of the presence of random effects, serial correlation, or both. To explore the possible nature of the misspecification, we restore the modified versions of the test. The adjusted version of the test for serial correlation $\text{ALM}(\text{lambda}=0)$ now fails to reject the null hypothesis while the adjusted version of the test for random effects $\text{ALM}(\text{Var}(u)=0)$ still does. This suggests that the possible misspecification is likely due to the presence of random effects rather than the serial correlation. Consequently, and to stress the main usefulness of these procedures, in this example the presence of the random effects seems to confound the unadjusted test for serial correlation, making it spuriously reject its null.

4 Acknowledgments

We thank an anonymous referee for useful suggestions that helped improve the routine considerably, and we thank Javier Alejo for excellent research assistance.

5 References

- Anand, S., and R. Kanbur. 1993. Inequality and development: A critique. *Journal of Development Economics* 41: 19–43.
- Baltagi, B., and Q. Li. 1990. A Lagrange multiplier test for the error components model with incomplete panels. *Econometric Reviews* 9: 103–107.
- . 1991. A joint test for serial correlation and random individual effects. *Statistics and Probability Letters* 11: 277–280.
- Baltagi, B. H. 2005. *Econometric Analysis of Panel Data*. 3rd ed. New York: Wiley.
- Bera, A., W. Sosa-Escudero, and M. Yoon. 2001. Tests for the error component model in the presence of local misspecification. *Journal of Econometrics* 101: 1–23.
- Bera, A., and M. Yoon. 1993. Specification testing with locally misspecified alternatives. *Econometric Theory* 9: 649–658.
- . 2001. Adjustments of Rao's score test for distributional and local parametric misspecifications. Mimeo: University of Illinois at Urbana–Champaign.
- Breusch, T., and A. Pagan. 1980. The Lagrange multiplier test and its applications to model specification in econometrics. *Review of Economic Studies* 47: 239–253.
- Gasparini, L., M. Marchionni, and W. Sosa-Escudero. 2001. *Distribucion del Ingreso en la Argentina: Perspectivas y Efectos sobre el Bienestar*. Cordoba: Fundacion Arcor-Triunfar.
- Gustafsson, B., and M. Johansson. 1999. In search of smoking guns: what makes income inequality vary over time in different countries? *American Sociological Review* 64: 585–605.
- Hemmerle, W. J., and H. Hartley. 1973. Computing maximum likelihood estimates for the mixed AOV model using the W transformation. *Technometrics* 15: 819–831.
- Sosa-Escudero, W., and A. Bera. 2001. sg164: Specification tests for linear panel data models. *Stata Technical Bulletin* 61: 18–21. Reprinted in *Stata Technical Bulletin Reprints*, vol. 10, pp. 307–311. College Station, TX: Stata Press.

About the authors

Walter Sosa-Escudero is an associate professor and the chairman of the Department of Economics, Universidad de San Andrés, Buenos Aires, Argentina.

Anil K. Bera is a professor in the Department of Economics, University of Illinois at Urbana–Champaign.