



**AgEcon** SEARCH  
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

# A Farm Level Evaluation of a New Twinning Technology in Beef Cattle

R.J. Farquharson\*

This article describes a farm level case study evaluation of a new beef production technology. The evaluation was undertaken as input into a technical research project aimed at developing farm level management strategies to facilitate use of the twinning technology when it becomes commercially available. The evaluation includes budgeting methods and a Linear Programming analysis. A risk analysis was also conducted in which key factors are varied stochastically and the resulting distributions of activity gross margins compared. The analysis shows that twinning could potentially improve the returns to intensive beef production systems, depending importantly on the level and variability of weaning percentage.

## 1. Introduction

There are a number of research and development (R&D) institutions in Australia which are conducting and/or funding agricultural projects and programs. These institutions are interested in maximising the extent of uptake or eventual use of technologies by a target group or audience. There are a number of strategies which can be used to achieve this aim. One of these is to determine the potential payoffs from alternative R&D projects and programs as an aid in the project planning process and in making resource allocation decisions. Is the new technology likely to be profitable at the farm level and what are the aggregate industry impacts?

At the farm level the assessment is aimed at establishing whether a new technology really will 'improve the farmer's lot' and whether the anticipated improvements are reasonably substantial. This assessment can be undertaken while the technology is still being developed as an aid to the scientists involved in the development project (Anderson and Hardaker 1979). It was for this purpose that the present study was undertaken.

### 1.1 Twinning Technologies for Beef Cattle

New technologies are being developed with the

potential to revolutionise cattle breeding in Australia. These technologies include:

- (a) *in vitro* fertilisation;
- (b) cloning of individual cells from early embryos;
- (c) embryo splitting;
- (d) embryo transfer; and
- (e) fertility vaccines.

The objective of the research into these techniques is to develop methods which allow multiple pregnancies with the aim, through twinning, of raising the annual reproductive rate of cows (Anon 1990). The success of such programs will depend largely on applying appropriate levels of management and on catering adequately for the increased nutritional requirements of the herd.

Three methods of non-genetically improving reproductive technology are currently being investigated (Piper and Bindon 1990). One involves the embryo transfer of eggs that have been fertilised *in vitro*, the second involves vaccination of the cow against the hormone Inhibin (which normally restricts ovaries to shedding one egg at each oestrus) and the third involves a hormone injection to increase ovulation rate (Follicle Stimulating Hormone). It is assumed that either the anti-Inhibin vaccine approach or the *in vitro* fertilisation/embryo transfer twinning method will be available commercially in the near future (Piper and Bindon 1990). The embryo transfer method has the benefit of allowing control over features such as day of conception and therefore date of birth, genotype

\* Division of Rural and Resource Economics, NSW Agriculture, Agricultural Research and Advisory Station, Grafton. This is a revised version of a paper presented at the 35th Annual Conference of the Australian Agricultural Economics Society, 11-14 February 1991. The contributions of Louise Patten and Garry Griffith, and the comments of anonymous referees, are gratefully acknowledged.

Review coordinated by Blair Bartholomew.

and, perhaps eventually, sex of calf. However, the anti-Inhibin vaccination method is simpler and probably cheaper as it only requires the annual injection of cows prior to mating. Therefore, despite the fact that this method provides control over nothing more than the shedding of an extra egg, it may be more appealing to beef producers.

Twinning in beef cattle through anti-Inhibin vaccination would require all breeding cows in the target herd to be injected prior to mating. After mating all cows would be scanned ultrasonically to identify those with twins, and the diagnosed twin-bearing cows would then be subjected to a greater degree of management attention. It is only in the more productive agricultural areas with intensive management that twinning is considered to be potentially successful. Important management factors for successful twinning might include use of restricted mating seasons, supplementary feeding, pregnancy testing and culling strategies for infertile and aged cows. It is considered that country capable of producing prime lambs is likely to be suitable for twinning in beef cattle (Dr B. Bindon, CSIRO, pers. comm.). The twinning technology could also potentially be used in the dairy industry. With more intensive feeding and management, dairy herds may be particularly suited to twinning. However, no assessment of twinning in the dairy industry has been included in this analysis.

Research studies are being conducted by the New South Wales and Victorian Departments of Agriculture in a coordinated program into various aspects of nutrition and management required for successful use of twinning at the farm level (Drs D.W. Hennessy, J.F. Wilkins and L.J. Cummins, pers. comm.). These studies will determine the extra feed demand of twin-bearing beef cows and investigate alternative farm management strategies for accommodating twinning herds.

## 1.2 Problem Definition and Hypothesis to be Tested

The problem addressed in this analysis is the *ex ante* evaluation of potential economic gains from twinning by vaccination at the farm level. This type of assessment can be used either as an input into aggregate industry-level assessments or to identify

areas where the technology needs fine tuning prior to commercial release. If aspects of a technology are identified as constraining its use in normal farm operation, or if the potential financial benefits are not large enough, then the technology can be reviewed and changed if necessary to improve it for the target group.

Important aspects of the appeal a twinning enterprise may have at the farm level include:

- (a) whether a twinning enterprise is complementary or competitive with existing enterprises in terms of resource requirements;
- (b) the expected level of financial return from twinning compared to that of other enterprises; and
- (c) the impact of a twinning enterprise on beef production risk, including the variability in return from twinning as well as the covariability in return between twinning and other enterprises.

The assessment will need to be conducted in a manner that will account for these questions.

The null and alternative hypotheses to be tested are  $H_0$ : Twinning is not economically appealing to beef producers in the target group; and  $H_a$ : Twinning provides an economically appealing alternative enterprise to beef producers. The null hypothesis will be rejected if twinning is among the group of activities selected as optimal in a farm plan where twinning is an option.

## 2. Representing the Target Group

The aim of this type of analysis is to undertake an assessment for a 'representative farm' from which can be drawn general conclusions for the target group. Three main approaches to this problem were outlined by Anderson and Hardaker (1979) - a case study approach, a representative farm approach and a sample survey. In this study a case study approach was used, but the farm chosen was also considered to be fairly representative of the likely target population.

A mixed-enterprise (sheep-beef) property situated between Glen Innes and Inverell in the Northern Tablelands of New South Wales was used as the case study farm. It was located on fertile soils with

improved pastures and was subject to relatively intensive management. Furthermore, the farmer kept good records and was willing to discuss the farm operation. Information was collected by personal interview on the property.

### 3. Methods of Assessment

#### 3.1 Budgeting methods

At the farm level different activities (technologies)

can be initially compared using budgeting methods (Longworth and Menz 1980). Gross margins, cash flow and partial budgets can be used for comparative analysis of activities and to examine the effects of changes over a number of years (Makeham and Malcolm 1988, Dillon and Hardaker 1984). A partial budget for the introduction of twinning to a beef store/vealer activity is shown in Table 1. The impacts of increased weaning percentage and reduced weaning weight are that a net benefit of \$33 to \$55 per herd cow is demonstrated.

**Table 1: Partial Budget of Twinning in a 100 Cow Beef Activity**

	\$ per cow	\$ per 100 cows
Returns foregone		
Single calf	360 (300 kg vealer at \$1.20) (a)	32400
Extra returns		
Twin Calves	610-660 (2 x 255 kg (b) at \$1.20 at \$1.30(c))	43650 45900
NET CHANGE IN REVENUE	250-300 per twin-bearing cow	11250-13500
Extra costs		
Induction (d)	20	2000
Diagnosis (d)	15	1500
Supervision at Calving (e,f)	30	1350
Extra feed: (e,g)		
- pregnancy	38	1710
- lactation	32	1440
TOTAL EXTRA COSTS		8000
NET BENEFITS : - without feed costs		6400-8650
- with feed costs		3250-5500
(a) Normal weaning rate is 90 per cent (b) Assumes 10 cows dry, 45 with single calves, 45 with twins and 15 per cent lower liveweight per twin weaner (c) Beef advisory officers indicate that liveweight price (\$/kg) is often greater for lighter animals (d) All cows (e) Twin-bearing cows only (f) Includes veterinary costs at calving (g) Grain at \$200/tonne fed		

Budgeting methods are a useful first step in comparing farm activities, but they do have a number of disadvantages. One of these is that in a whole-farm context, with other competing activities, the limited availability of resources means that the process of choosing a farm plan can be more complex. In this respect the methods of whole-farm analysis discussed in Section 3.3 are more appropriate for major changes to farm plans.

Another disadvantage of budgeting approaches is that they generally do not account for risk. Here risk is conceptualised as variability in outcome with consequences for human choice (Anderson 1988). The inclusion of aspects of risk in a gross margin results in a stochastic gross margin (Anderson 1976). If a stochastic variable is to be introduced into a budget, information is required on the type and parameters of the distribution of that variable. This implies that the probability of different outcomes can be estimated. This type of information may be available from historical records or experimental data but often it may not. The introduction of risk into a gross margin budget requires that stochastic distributions for the most important variables be known or capable of estimation and that covariability of those distributions be included. If it is possible to estimate the probability of the risky outcome (e.g. gross margin per hectare) being less than a range of values then the comparison of risky gross margins is possible.

### 3.2 Stochastic efficiency rules

If risk is incorporated into a gross margin, the result can be expressed as a probability density function (PDF) or as a (less than) cumulative distribution function (CDF) - see Figure 1. In comparing technologies the problem then becomes one of comparing PDFs or CDFs.

Some attempts have been made to measure in aggregate the risk attitudes of Australian farmers. Bond and Wonder (1980) measured risk attitudes using a risk attitude questionnaire and Bardsley and Harris (1987) measured risk aversion coefficients by observing the actual behaviour of farmers in a realistic economic environment. Both studies concluded that Australian farmers were generally risk averse.

The utility or preference function shows the willingness to trade off extra expected income against increased variability in income. However, the utility functions for individual farmers or small groups are unknown and the formal specification of such functions is a difficult and costly process. Concepts of stochastic efficiency are useful in comparisons of activities or technologies when utility functions are unknown. Anderson (1974) and Anderson, Dillon and Hardaker (1977, Ch. 9) have reviewed these rules which search for efficient sets of distributions that are not dominated given certain assumptions about the behaviour or preferences of decision-makers. CDFs can be compared using first-, second- and third-degree stochastic dominance tests.

The concept of stochastic dominance with respect to a function (SDWRF) enhances the stochastic efficiency rules. It was developed by Meyer (1977a, b) based on Pratt's (1964) coefficient of absolute risk aversion,  $r(x)$ , which relates to the curvature of the utility function,  $U(x)$

$$r(x) = -U''(x)/U'(x)$$

The use of SDWRF requires specifying a preference interval bounded by upper and lower values of  $r(x)$ . Once the preference interval is narrowed the SDWRF criteria become more powerful in distinguishing between distributions. This approach allows different sets of decision-makers to be distinguished for different bounds (values of  $r(x)$ ).

Pandey (1990) identified risk-efficient irrigation strategies for wheat in India using four ranges of risk aversion coefficients between 0 and 0.04. Australian farmers might be expected to be less risk averse because of their greater wealth (including human capital). Bardsley and Harris (1987) estimated values for median income and partial risk aversion coefficient for the Pastoral, Wheat-Sheep and High Rainfall Zones of Australia. From their results the value of absolute risk aversion for Australian farmers was  $1 \times 10^{-5}$ ,  $6 \times 10^{-5}$  and  $1 \times 10^{-6}$  for the High Rainfall, Wheat-Sheep and Pastoral Zones respectively. These figures are at the lower end of the range of values used by Pandey (1990).




Figure 1 An Equivalent Probability Density Function (PDF) and Cumulative Distribution Function (CDF)

A microcomputer program developed by Goh, Raskin and Cochrane (1987) was used to conduct the stochastic efficiency comparisons using SDWRF. The distributions in this analysis were compared pairwise over a range of values for absolute risk aversion intervals including those estimated from the Bardsley and Harris (1987) results.

### 3.3 Whole-farm approach

To overcome the disadvantages of partial budgeting, the whole-farm approach allows a more holistic or systematic view of technology testing and adoption (Dillon 1976). It requires the setting up of a model of a farm that is representative in some sense of the population of farms (see Section 2).

The model can be used to test the effect of including the new technology as an alternative activity. In effect the model results are compared with and without the new technology to show how successful it might be and to indicate the 'size' of the effects of the new activity. This information can be used to make estimates of the wider socio-economic impact of the technology.

Farquharson (1991) reviewed some methods of whole-farm analysis. These included gross margins analysis, simplified programming, linear programming (LP) and quadratic risk programming (QRP). Ghodake and Hardaker (1981) listed other methods that extend the LP approach to incorporate risk. These include linear risk programming,

stochastic programming and Monte Carlo programming, as well as systems simulation.

### 3.4 Alternative analytical approaches

The discussion in the preceding sections has indicated the desirability of using a whole-farm approach and of accounting for risk in comparing alternative farm activities. LP was considered the most useful method of undertaking the basic whole-farm analysis because it provides good information as a basis for comparing technologies, it is relatively straightforward to use and computer programs are readily available and user friendly.

For the risk analysis a spreadsheet-based program, @RISK (Palisade Corporation 1989), was available which enabled the definition of uncertain spreadsheet cell values as probability distributions. A considerable number of distribution types can be specified and cells within the spreadsheet can be designated as the output range which contains the distribution of possible results.

A number of alternative approaches to conducting a risk analysis as an extension of LP are possible. Two particular approaches were considered for this analysis. The first was to use a QRP model which accounts for variability and covariability in farm activity net revenues. The alternative was to place the LP results for key activities back into a spreadsheet format and use @RISK to generate CDFs for comparison using the stochastic efficiency concepts mentioned in Section 3.2.

The latter approach was preferred in this analysis for a number of reasons. The main advantages are that @RISK allows the incorporation of stochastic variability into a number of parameters within the beef activity rather than just net revenue, as in QRP. In terms of technology assessments in conjunction with scientific R&D projects, @RISK allows physical parameters to be varied which the scientists can directly relate to and which are important for the analysis. Also @RISK has a great deal of flexibility in specifying types of distributions and accounting for different types of distributional information.

The main disadvantage of this latter approach is

that the selection of the optimal farm plan in the whole-farm analysis only allows for risk in some activities and does not account for the influence of that variability in the whole-farm plan. However, an analysis using QRP would require much more information on the net revenue variability and covariability of all potential farm activities which might be very difficult to determine. For the purposes of the particular technology assessment in this analysis the use of LP and @RISK was considered the best approach.

## 4. Results

### 4.1 Linear programming model

In constructing an LP model of the case study farm a set of gross margin budgets for the existing and proposed activities was constructed. First the energy requirements for livestock were derived on a herd or flock basis (Rickards and Passmore 1977, Agricultural Research Council 1980, Ministry of Agriculture, Fisheries and Food 1984, Animal Production Committee 1990). The estimated metabolisable energy (ME) requirements per breeding female (including followers) throughout the year in units of megajoules of metabolisable energy per female per day are shown in Table 2.

Then the energy supplied by pastures throughout the year was specified in the same units. Published estimates of improved pasture production from basaltic (self-mulching) soils at the Glen Innes Agricultural Research and Advisory Station were used (Hennessy and Robinson 1974, 1979). Carryover of pasture from one time period (season) to the next was incorporated in the model, with the amounts of energy in feed carried over estimated to be 33 per cent from Summer to Autumn, 20 per cent from Autumn to Winter, zero from Winter to Spring and 28 per cent from Spring to Summer (Dr P. Mears, NSW Agriculture, pers. comm.).

Because of the winter feed gap, supplementary feeding of breeding stock through winter is often undertaken in the tablelands areas of New South Wales. Oddy (1983) set out the basis for use of the ME system for drought feeding of sheep and cattle including the average ME and dry matter content of feeds. The relationship between animal liveweight

**Table 2: Estimated Metabolisable Energy Requirements per Breeding Female (including followers) throughout the year**

Breeding female	Quarter			
	Spring	Summer	Autumn	Winter
	MJ of ME/female/day			
Cows				
- single	153	146	166	161
- twin	170	166	186	282
Ewes				
- Merino	13	13	15	10
- First-cross	23	26	11	14

and ME content of feed can be used to determine the quantity of feed required per day for sheep and cattle. This allows the amount of different types of feed to be estimated for specific types of animals, and was used in developing the supplementary feeding activities of the LP model.

Gross margin budgets were drawn up for the beef (store/vealer production) and sheep (Merino wool growing, first- and second-cross lamb production) activities of the case study farm. The beef twinning activity budgets were set up based on three alternative scenarios: cows first calve at 2 years, cast-for-age (cfa) at 10 years (the same as the normal beef activity); cows first twin calve at 4 years (after 2 years of singles), cfa at 10 years; and cows first twin calve at 4 years, cfa at 9 years. The implications of these alternative assumptions for herd structure and turnoff were derived from use of a herd model (Holmes 1988) as shown in Table 3.

Labour budgets were also constructed for each activity. In discussions about labour requirements for farm activities, the case study farmer indicated that he considered the labour budgets in Turvey (1988) to be adequate. The labour requirements for twinning activities were set as pro-rata increases over the normal beef herd according to the increased calf weaning rates.

In discussing supplementary feeding the case study

farmer indicated that due to the shortage of pasture breeding stock carried through winter are fed supplements rather than winter fodder crops or conserved fodder (hay or silage). Breeding ewes are fed grain and cows are fed molasses and cottonseed meal. The ME content and average daily requirements, given liveweight and pregnancy status, were estimated (Oddy 1983) and the costs per feed unit determined. This information was used in the development of winter supplementary feeding activities for both cattle and sheep in the LP model.

As shown in Table 4 sheep activities were more profitable than cattle activities and twinning activities were more profitable than the normal beef activities. The profitability of sheep was partly due to the buoyant wool prices in the early part of 1990. Since the lowering and then removal of the floor price for wool the relative advantage of wool activities has been reduced, but no investigation of this outcome was undertaken here. However, the relative profitability of sheep over cattle activities had implications for construction of the LP model.

The LP model was based on the land, labour and feed resources of the case study farm. The treatment of resource constraints was considered in line with the discussion by Dent, Harrison and Woodford (1986). In particular the personal attitudes and longer term goals of the case study farmer were important determinants of the balance of livestock



**Table 3: Herd Structures for Single- and Twin-Bearing Herds Producing Vealers and Store Weaners**

Cattle type	Singles	Twins(a) (2-10)	Twins(b) (4-10)	Twins(c) (4-9)
Calves born start of year	92	138	125	124
Heifers aged 1 start of year (heifers retained)	17	15	15	16
Heifers aged 2 start year	15	14	14	15
Cows aged 3 years plus	68	71	71	69
Bulls	3	3	3	3
<b>Total</b>	<b>195</b>	<b>241</b>	<b>229</b>	<b>229</b>
Cows and heifers mated	100	100	100	100
Calves branded	92	138	125	124
Breeder deaths	1	3	3	3
Sales - cull cows	8	6	6	7
- cfa cows	7	7	8	9
- surplus heifers	29	53	47	45
- steers	45	68	62	61

(a) First calving at 2, last calving at 10, 9 opportunities to calve as twins.  
(b) 2 opportunities to calve as singles, 7 opportunities to calve as twins.  
(c) 2 opportunities to calve as singles, 6 opportunities to calve as twins.

activities on the farm and in the LP model. The LP matrix for the livestock activities was based on the model structure of Muir and Vere (1987). In setting up the base model the relatively greater

**Table 4: Gross Margin Comparisons for Livestock Activities**

Activity	GM/cow	GM/ewe	GM/ME unit (a)	GM/labour unit (b)
	\$	\$	\$	\$
<b>Beef</b>				
- normal	279	-	1397	140
- twins (2-10)	373	-	1863	186
- twins (4-10)	343	-	1716	172
- twins (4-9)	345	-	1724	172
<b>Sheep</b>				
- Merino wool	-	67	3354	335
- First-cross lamb	-	66	3321	332
- Second-cross lamb	-	52	2613	261

(a) 20 MJ/day  
(b) 200 hours/season

**Table 5: Base results of Linear Programming Model**

Activity	Unit	Value
Merino ewes (mated to Merino rams)	no.	4622
Merino ewe replacements	no.	1294
Sell Merino ewe hoggets	no.	893
Sell Merino wether lambs	no.	2265
Transfer old Merino ewes	no.	1063
Sell Merino wool	kg	37355
Merino ewes (mated to Border Leicester rams)	no.	1063
Sell first-cross ewe hoggets	no.	592
Sell first-cross wether lambs	no.	627
Sell old Merino ewe culls	no.	999
Sell first-cross wool	kg	1183
Sell Border Leicester wool	kg	191
Breeding cows	no.	239
Cow replacements	no.	41
Surplus heifers	no.	68
Steer vealers	no.	55
Steer stores	no.	55
Sell vealers	kg live	15931
Sell stores	kg live	13733
Sell heifers	kg live	17129
Pasture for cattle	ha	650
Pasture for sheep	ha	1500
Purchase molasses	kg	0
Purchase grain	kg	230
Livestock selling costs	\$	15019
Wool selling costs	\$	42014
Sheep production costs	\$	64484
Beef production costs	\$	1890
Cattle pasture carryover Summer to Autumn	MJ/ha/day	158639
Cattle pasture carryover Autumn to Winter	MJ/ha/day	76683
Cattle pasture carryover Spring to Summer	MJ/ha/day	55308
Sheep pasture carryover Summer to Autumn	MJ/ha/day	324442
Sheep pasture carryover Autumn to Winter	MJ/ha/day	150162
Sheep pasture carryover Spring to Summer	MJ/ha/day	109120
Objective Function Value	\$	383 666

profitability of merino woolgrowing meant that it was the only activity selected. This result was considered to be unrealistic in light of the management strategy of diversifying activities undertaken by the case study farmer. To overcome this problem the land resource was separated into two categories (one for cattle and one for sheep) based on the case study farm area and each land category was

provided with a feed supply. This structure reduced the potential value of the objective function and partly determined the outcome of the model. The approach was justified on the grounds that it was a more realistic outcome in comparison with the actual case study farm plan.

The main results of the base LP model are shown in

**Table 6: Key Binding and Slack Constraints in the Linear Programming Base Solution**

Resource	Unit	Binding or Slack	Amount of Slack	Shadow Price
Labour spring	hr	S	117	-
Labour summer	hr	S	658	-
Labour autumn	hr	S	740	-
Labour winter	hr	S	1264	-
Grain	kg	B	0	63.75

**Table 7: Results of Introducing Beef Twinning Activity**

Activity	Unit	Base solution	Twinning included
Breeding cows	no.	239	216
Replacement cows	no.	41	32
Surplus heifers	no.	68	116
Steer vealers	no.	55	-
Steer stores	no.	55	149
Sell vealers	kg live	15931	-
Sell stores	kg live	13733	36028
Sell heifers	kg live	17129	24748
Cattle pasture	ha	650	650
Purchase molasses	kg	0	450
Selling costs(a)	\$	15019	15714
Beef production costs	\$	1890	6007
Objective Function(a)	\$	383 666	387 681

(a) Includes sheep activities

**Table 8: Key Binding and Slack Constraints under Twinning**

Resource	Unit	Binding or Slack	Amount of Slack	Shadow Price
Labour spring	hr	S	17	-
Labour summer	hr	S	591	-
Labour autumn	hr	S	628	-
Labour winter	hr	S	1127	-
Molasses	kg	B	-	16.23

Table 5. The Merino wool growing activity was the major contributor to the objective function. Merino cfa ewes were transferred to the first-cross lamb activity rather than being sold. However, the second-cross lamb activity did not enter the final optimal solution. The beef herd consisted of 239 breeding cows.

The names and amounts of slack resources for this base solution are shown in Table 6. Excess labour was available in all four quarters. Pasture supply for sheep in winter was limiting the objective function. Also shown in Table 6 is the shadow price of the binding constraint. The shadow price is the marginal value of this resource - the objective function would increase by \$63.75 if one further unit of grain for sheep could be supplied (Lee, Moore and Taylor 1985). The shadow price of grain is greater than the cost of purchase (\$11.70). Therefore it would be profitable to increase the supplementary feed supply for sheep, and hence the stocking rate, but this would be at the expense of increased risk in a bad season.

When the beef twinning activity was included in the matrix the optimal solution was as shown in Table 7. Since the sheep activities were unchanged only the beef results are presented. When the twinning activity was included in the model it replaced the single-breeding cow activity completely. There were fewer cows, increased progeny (numbers and weight), higher costs and an increase of \$4015 in the value of the objective function.

The main resource constraints under twinning are shown in Table 8. In comparison with Table 7, inclusion of the twinning activity resulted in more labour being used in each quarter. Cattle winter feed supply was limiting and the shadow price of an extra unit (MJ/day) of molasses in winter was \$16.23.

Two further analyses with this twinning model were undertaken. To determine the impact of reducing labour availability, the use of only 3.5 full time labour units (compared to 4 people) was tested. When this change was made the twinning activity did not enter the optimal solution - the normal single vealer/store activity was selected and the objective function value was reduced to

\$363 716. Labour in spring was a binding constraint and the shadow price of spring labour was \$170 per hour.

To determine the impact of changed feed supplies on twinning the supplementary feeding of molasses and cottonseed meal to breeding cows in winter was deleted from the model by setting the amount of molasses available for feeding to zero. The resulting solution was that 197 twinning cows were selected and the objective function value was \$373 005.

The twinning activity added to the base model was for cows first calving at 2 years and being culled at 10 years. The two additional twinning scenarios identified in Table 3 were added individually to the base model (no twinning) and to the twinning (join year 2, cfa year 10) model. In no case were these other twinning activities selected in the optimal farm plan. Therefore the second and third twinning activities do not appear to be profitable in the whole-farm context. None of these further results are reported in detail here but they are discussed in Farquharson (1991).

#### 4.2 Risk analysis

In this section the LP results are extended through a risk analysis. A more realistic analysis will include some of the inherent biological and economic variability in at least some of the important factors. The methods used in this section allows more information about key variables to be incorporated, including the range of possible values for a variable and the likelihood of occurrence of each value. The resulting analysis is a simulation in which a large number of 'what if' scenarios are presented together. 'Simulation' can be defined as 'the use of an analogue to study the properties of a system' where an analogue 'pertains to any device which represents a variable by a continuously moving or varying entity' (The Macquarie Dictionary 1985). The output of the simulation can be presented graphically and comparisons made between beef production with and without the new technology.

The risk analysis was conducted using the @RISK program. The distribution types used in this analysis were normal (@NORMAL), triangular

(@TRIANG) and truncated normal (@TNORMAL). The arguments for these functions in the @RISK program are (mean, standard deviation (SD)) for @NORMAL, (minimum, most likely, maximum) for @TRIANG and (mean, SD, minimum, maximum) for @TNORMAL.

The @TRIANG distribution is useful for rough modelling when actual data are not available. The @TNORMAL is used in this analysis where the market destination for steers changes at a certain weight or where a certain proportion of the heaviest females must be retained as replacement cows. For steers, the vealer trade (local butchers or supermarket) commences at about 160 kg dressed weight (or 290 kg liveweight at 55 percent dressing) (Mr P. Doyle, pers. comm.). Below this weight steers are generally store quality.

The risk analysis was conducted only for the beef cattle activities within the LP model. The main results from the model for the beef activity with and without the twinning technology were shown in Table 7. Two spreadsheets were developed from these results and these are presented in Tables 9 and 10. These spreadsheets contain the same technical coefficients as in the LP model. The main differences between these tables and the original budgets are that Tables 9 and 10 include actual livestock numbers from the LP results and the supplementary feeding and superphosphate costs.

The variables subject to risk simulation are weaning percentage, weaning weight and saleyard price for beef. The level and distribution of weaning percentage may be the most important factor in determining the success of twinning in beef cattle. This factor determines the number of calves born and weaned. On the case study farm the distribution of weaning percentage was estimated to be @TRIANG (90,92,94). For twinning the weaning percentage was assumed to rise by 50 per cent (Farquharson and Griffith 1991) so the mean weaning percentage became 138 per cent. Optimistic and pessimistic scenarios for twinning were derived by setting different distributions for weaning percentage. These scenarios were analysed in an attempt to determine whether the distribution of this variable could influence the final choice of technology, but there were no experimental data

available as a guide to the parameters of the optimistic and pessimistic distributions.

As purely hypothetical cases the distributions of weaning percentage were set as @TRIANG (126,138,150) and @TRIANG (80,155,179) for the optimistic and pessimistic scenarios respectively. The pessimistic scenario has a wider distribution and a longer tail to the left, but both of these distributions have expected values of 138 per cent so that the comparisons with the LP results are valid.

The weaning weight of progeny will vary within a herd in any year. On the case study farm all heifer weaners are weighed as part of the process of selecting heifer replacements. These weights in the last season ranged from 310 kg to 200 kg with an 'average' of 251 kg. By allocating the maximum and minimum weights to probabilities of 0.995 and 0.005 respectively, an estimate for SD of heifer weaning weight of 16.5 kg was derived from the standard normal distribution. Heifer weights were assumed to be distributed normally with mean 251 kg and SD 16.5 kg. However, because a certain percentage of the heaviest need to be kept as replacements the normal distribution of heifers for sale was truncated at a maximum weight (estimated as 266 kg) that allowed the correct number of replacements to be retained.

Steer weaners are not individually weighed on the case study farm, although the average weight was estimated to be 285 kg. Without any information on the distribution of steer weights the SD was estimated to be in the same ratio to the mean as for heifers. Thus the SD of steer weights was 18.7 kg. Given this distribution, the proportion of steers above and below 290 kg liveweight was estimated to determine the number of steers going to the vealer and store markets. The weight distributions for steer progeny were truncated at 290 kg for the vealer and store markets. The distributions of weights for steers and heifers used in the risk analysis are shown in Table 11.

For twinning the mean weaning weight is assumed to be 15 per cent lower than for single calves (i.e. 213 kg for heifers and 242 kg for steers). It can be assumed that in comparison with single calves,

**Table 9: Beef Vealer Spreadsheet for Risk Analysis**

Cattle pasture :	650 ha	Weaning percentage :	92		
Herd size :	239 cows	Steer progeny :	110		
		Heifer progeny :	110		
<b>INCOME</b>					
Sale of	No.	@ kg	\$/kg	\$/beast	\$
Vealers	55	290	1.35	-	21532.50
Store steers	55	250	1.45	-	19937.50
Surplus heifers	68	251	1.20	-	20481.60
Cull cows	36	-	-	530	19080.00
Cull bulls	2	-	-	800	1600.00
<b>TOTAL INCOME</b>					<b>82631.60</b>
<b>VARIABLE COSTS</b>					
<b>Bull replacements</b>					
Purchase	2	-	-	2000	4000.00
Transport	2	-	-	5.7	11.40
<b>Beef production</b>					
Cows	239	-	-	7.23	1727.97
Heifer replacements	41	-	-	3.39	138.99
Surplus heifers	68	251	.005	-	85.34
Vealers	55	290	.006	-	95.70
Steer stores	55	250	.008	-	110.00
Bulls	8	-	-	3.19	25.52
<b>Selling costs</b>					
Vealers	55	290	.083	-	1323.85
Steer stores	55	250	.075	-	1031.25
Heifers	68	250	.075	-	1280.10
Cows	36	-	-	43.3	1558.80
Bulls	2	-	-	54.1	108.20
<b>Pasture costs</b>					
Super	650 ha	@	\$34/ ha		22100.00
<b>Supplementary feed costs</b>					
Molasses and cottonseed meal		0 kg	@	\$ 16.2/ kg	0.00
<b>TOTAL COSTS</b>					<b>33597.12</b>
<b>TOTAL GROSS MARGIN</b>					<b>49034.48</b>

**Table 10: Beef Twinning Spreadsheet for Risk Analysis**

Cattle pasture :	650 ha	Weaning % :	138		
Herd size :	216 cows	Steer progeny :	149		
		Heifer progeny :	149		
<b>INCOME</b>					
Sale of	No.	@ kg	\$/kg	\$/beast	\$
Vealers	0	290	1.35	-	0.00
Store steers	149	242	1.45	-	52284.10
Surplus heifers	116	213	1.20	-	29649.60
Cull cows	28	-	-	530	14840.00
Cull bulls	2	-	-	800	1600.00
<b>TOTAL INCOME</b>					<b>98373.70</b>
<b>VARIABLE COSTS</b>					
<b>Bull replacements</b>					
Purchase	2	-	-	2000	4000.00
Transport	2	-	-	5.7	11.40
<b>Beef production</b>					
Cows	216	-	-	27.23	5881.68
Heifer replacements	32	-	-	3.39	108.48
Surplus heifers	116	213	.005	-	123.54
Vealers	0	290	.006	-	0.00
Steer stores	149	242	.008	-	288.46
Bulls	6	-	-	3.19	19.14
<b>Selling costs</b>					
Vealers	0	290	.083	-	0.00
Steer stores	149	242	.075	-	2704.35
Heifers	116	213	.075	-	1853.10
Cows	28	-	-	43.3	1212.40
Bulls	2	-	-	54.1	108.20
<b>Pasture costs</b>					
Super	650 ha	@	\$34/ ha		22100.00
<b>Supplementary feed costs</b>					
Molasses and cottonseed meal		450 kg	@	\$ 16.2/ kg	7290.00
<b>TOTAL COSTS</b>					<b>45700.75</b>
<b>TOTAL GROSS MARGIN</b>					<b>52672.95</b>

weights for twin calves are 20 per cent lower at birth, 10-15 per cent lower at weaning and similar at yearling (Dr B. Bindon, CSIRO, pers. comm.). The distribution of weaning weights is suspected to be wider than for single calves, especially if heifers and young cows are in the twinning herd (Dr B. Bindon, CSIRO, pers. comm.). For twin heifers and steers the SD of weaning weight was set at 16.4 and 17.5 kg respectively.

The level and distributions of prices and their relationships with liveweight for different beef types is an area where little information is available. It is considered (Mr P. Doyle, NSW Agriculture, pers. comm.) that the relationship between price and liveweight is weakly negative for vealers (the local butcher trade may pay more \$/kg for lighter vealer carcasses than the supermarket trade pays for heavier carcasses), strongly negative for store steers and strongly positive for heifers (the heavier heifers are in more demand as replacements). Todd and Cowell (1981) estimated the regression coefficient for weight explaining beef auction prices (c/kg basis) at a domestic trade type of cattle auction to be -0.18. Park (1979) found that an increase in liveweight was correlated with a decrease in unit price.

The @RISK program uses dependency coefficients to represent the degree of correlation between dependent and independent variables when sampling, but these are not the equivalent of correlation coefficients. The dependency coefficient can be set between 1 and -1 to determine the type of correlation between two variables in the sampling process. The closer to 1 or -1 the stronger the (positive or negative) correlation. The levels of dependency coefficients between liveweight and price (\$/kg) were set as shown in Table 12.

The Livestock Marketing Reporting Service (LMRS) (NSW Meat Industry Authority 1990) provides information on prices for slaughter cattle at regional markets for cattle categorised by age, sex, liveweight, muscle score and fat score. No individual weight data are recorded by the LMRS. Price data from the Armidale and Inverell sales from January 1988 to June 1990 for vealers in the weight range 280-370 kg were analysed to determine the mean and SD. Altogether 12316 vealers

were marketed through these two saleyards over the 18 month period with a mean price of \$1.29/kg liveweight and a SD of \$0.21. Information on store cattle prices (\$/kg liveweight basis) and weights is not collected by the LMRS or any other agency. For store steers and heifers triangular distributions were used based on current market experience. All price distributions are shown in Table 11.

Three beef scenarios were analysed - normal vealer production (single-bearing cows) and twin-bearing cows with an optimistic and a pessimistic distribution for weaning percentage. The main @RISK results are shown in Table 13 as total gross margin (TGM) of beef activities from 650 ha of land. In comparison with normal beef the optimistic twinning scenario indicated a greater mean and a greater range of expected results. The pessimistic twinning scenario resulted in a higher mean than normal and a much wider range of possible outcomes. These results are shown graphically as CDFs in Figure 2.

From Figure 2 the normal beef CDF intersects once with each of the twinning CDFs. In comparison with the normal beef CDF the optimistic twinning CDF (which is also normally distributed) has a higher mean and a higher variance. Therefore the (E-V) comparison using means and variances (Anderson, Dillon and Hardaker 1977, p.287) cannot determine stochastic dominance. Anderson, Dillon and Hardaker (1977, p.288) describe distributions which intersect once as 'simply related' and define second-degree stochastic dominance also in terms of proneness to low outcomes. However, the optimistic twinning CDF is slightly more prone to low outcomes and so this criterion cannot be used to separate it from the normal beef CDFs. And because the pessimistic twinning CDF is not normally distributed it cannot be compared with the other two CDFs using these rules.

The distributions in Figure 2 were compared using SDWRF as noted in Section 3.2. The results of these comparisons are shown in Table 14. At the levels of risk aversion measured by Bardsley and Harris (1987) the optimistic twinning CDF was stochastically efficient and dominated the other distributions. At the higher level of risk aversion the small possibility of a lower outcome under



**Table 11: Types and Parameters of Distributions of Key Variables**

Variable	Distribution type	Distribution parameters (a)	Source of information
<b>Weaning percentage</b>			
- normal	@TRIANG	(90,92,94)	CS farm
- twins			
pessimistic	@TRIANG	(80,155,179)	Estimate
optimistic	@TRIANG	(126,138,150)	Estimate
<b>Weaning weight</b>			
- normal			
heifers	@TNORMAL	(251,16.5,0,266)	CS farm
vealer steers	@TNORMAL	(285,18.7,290,1000)	CS farm
store steers	@TNORMAL	(285,18.7,0,290)	CS farm
- twins			
heifers	@TNORMAL	(213,16.4,0,226)	Estimate
vealer steers	@TNORMAL	(242,17.5,290,500)	Estimate
store steers	@TNORMAL	(242,17.5,0,290)	Estimate
<b>Prices</b>			
heifers	@TRIANG	(1.05,1.11,1.20)	CS Farm
vealer steers	@NORMAL	(1.29,0.21)	LMRS
store steers	@TRIANG	(1.40,1.45,1.50)	Estimate
<p>CS farm = Case Study farm  LMRS = Livestock Market Reporting Service  (a) @TRIANG (minimum, most likely, maximum)  @NORMAL (mean, SD)  @TNORMAL (mean, SD, minimum, maximum)</p>			

**Table 12: Correlations between Weights and Prices**

Beef Type	Independent Variable	Dependent Variable	Dependency Coefficient	Comment
Heifer	Live weight	Price	+0.75	Strongly positive
Store steer	Live weight	Price	-0.75	Strongly negative
Vealer steer	Live weight	Price	-0.25	Weakly negative

<b>Table 13: Results of Risk Analysis</b>			
Distribution measure	Scenario		
	Normal vealer production	Twinning optimistic	Twinning pessimistic
	\$(a)	\$(a)	\$(a)
Mean	48295	50573	50534
Maximum	55371	62762	79002
Minimum	40275	36405	13647
Range	15096	26357	65355
SD	2790	4621	13032

(a) Total Gross Margin from beef activities on 650 ha.

<b>Table 14: Dominant Beef Production Strategies for Ranges of Absolute Risk Aversion</b>	
Risk Aversion Range (a)	Dominant Technology
$1 \times 10^{-6} - 1 \times 10^{-5}$	Optimistic twinning
$1 \times 10^{-5} - 1 \times 10^{-4}$	Optimistic twinning
$1 \times 10^{-4} - 1 \times 10^{-3}$	Optimistic twinning, normal beef production

(a) Measured by the absolute risk aversion coefficient.

optimistic twinning (Figure 2) has become more important for that class of decision-maker.

Therefore under the assumptions made about the distributions of key variables, the optimistic twinning scenario presented here would be selected by a 'normally' risk averse Australian beef producer, but the pessimistic twinning scenario would not.

## 5. Summary and Conclusions

The null hypothesis is that twinning is not economically appealing to beef producers in the target

group. This hypothesis has been tested using one principal methodology (LP) and an extension (risk analysis). Both of these were based on detailed financial, labour and energy budgets. The LP model results presented above have shown that the twinning technology (given various assumptions) is superior to the conventional beef activity in some circumstances. If cows are joined and cfa at the normal ages, twinning is more profitable. Under twinning the number of cows is lower than normal for the same given land area or feed supply. Although the quality of the progeny is lower (lighter weights and no vealers) and costs are higher, the




Figure 2 CDF's for normal beef and two twinning activities

greater turnover increases the value of the TGM for twinning.

The increase in value of the objective function is \$4015. Whether this is sufficient to convince beef producers to adopt the twinning technology depends on the minimum rate of return they would require on their investment in breeding stock. Under twinning the number of breeders is reduced from 239 to 216 (Table 7) and it is likely that the funds invested in breeding stock would be reduced.

Examination of the main LP results clearly indicates that, assuming profit maximisation, beef production using twinning can provide an increased return in a whole-farm context, and therefore the null hypothesis must be rejected. The alternative hypothesis, that twinning provides an economically appealing alternative enterprise to beef producers, is therefore accepted.

However, some qualifications of this conclusion have been indicated from the LP analysis. The number of calves weaned per cow joined is an important factor. The two alternative twinning scenarios under which the weaning percentage was reduced through cows being joined later and/or culled earlier than normal were not economically superior and so this question of overall herd fecundity is an important one for technology developers. Although this result should not be surprising its confirmation by the analysis for two not-realistic possibilities is of interest. This result should also be compared with the gross margin budgets summarised in Table 4. The implications of those results are that all three twinning activities would be economically superior to normal beef production, but the LP analysis has shown that by accounting for all resource availabilities the latter two twinning scenarios are not superior. Thus the advantage of using a whole-farm analysis over simple budgeting is demonstrated.

Another important result was that although twinning was still selected when supplementary winter feeding was reduced, when the labour supply was reduced normal beef production proved to be more profitable. Extra feed and labour requirements are both expected *a priori* to be important, but this result indicates that labour availability is a neces-

sary requirement for twinning. This highlights an aspect of the twinning technology - that is is expected to be quite labour-intensive and beef producers would need to have adequate labour available to undertake twinning.

The risk analysis required further assumptions about the behaviour of twinning in a commercial context. In particular assumptions about the distributions of key variables needed to be made. The approach taken was to use the best available information or opinions of experts in the field, and to apply a sensitivity analysis to investigate the impact of different distributional assumptions for the key variables.

When the risk analysis results were compared for a risk averse beef producer, an optimistic twinning scenario proved superior to normal beef production and the null hypothesis must be rejected. However, under the pessimistic scenario of the distribution of weaning percentage, the probability of a lower financial outcome for much of the time means that the beef producer would not consider twinning given the range of other risks in livestock production and farming generally.

These results can be compared with two other economic studies of twinning in beef cattle. Farquharson and Griffith (1991) used a partial budget to estimate the potential economic benefits from twinning at the farm level. The LP results from this analysis agree with their conclusion that twinning provided economic benefits. Herd *et al.* (1991) used an LP approach to look at a number of different beef cattle production systems at the farm level (including twinning). They concluded that the beef system based on twinning would be the most profitable to run.

In drawing conclusions from this type of analysis the crucial dependence on assumptions about potential impacts of the new technology must always be remembered. Close consultation with scientists and advisory officers has been maintained in attempting to set up a model that is as realistic as possible. The aim should be to do as good a job as possible with the resources and knowledge available at the time, but it must be recognised that the final outcome may be different for unforeseen rea-

sons (Anderson, Dillon and Hardaker 1977).

Overall the results of this analysis have indicated that, on the basis of the profit maximisation criterion, the twinning technology for beef cattle might be expected to be appealing to at least some beef producers. When aspects of potential increasing variability in returns associated with twinning are included, risk averse beef producers might still be attracted to twinning depending on the level of variability in the final weaning percentage. The reliability of the commercial product is thus indicated to be of primary importance.

## References

- AGRICULTURAL RESEARCH COUNCIL (1980), *The Nutrient Requirements of Ruminant Livestock*, Commonwealth Agricultural Bureaux, Farnham Royal.
- ANDERSON, J.R. (1974), 'Risk efficiency in the interpretation of agricultural production research', *Review of Marketing and Agricultural Economics* 42(3), 131-84.
- ANDERSON, J.R. (1976), *Methods and Programs for Analysis of Risky Gross Margins*, Miscellaneous Publication No. 5, Department of Agricultural Economics and Business Management, University of New England, Armidale.
- ANDERSON, J.R. (1988), 'Accounting for risk in livestock improvement programs', in *Economics of Livestock Improvements*, Proceedings of the Seventh Conference, Australian Association of Animal Breeding and Genetics, University of New England, Armidale, September.
- ANDERSON, J.R., DILLON, J.L. and HARDAKER, J.B. (1977), *Agricultural Decision Analysis*, Iowa State University Press, Ames.
- ANDERSON, J.R. and HARDAKER, J.B. (1979), 'Economic analysis in design of new technologies for small farmers', in VALDES, A., SCOBIE, G.M. and DILLON, J.L. (eds), *Economics and the Design of Small-farmer Technology*, Iowa State University Press, Ames, Iowa.
- ANIMAL PRODUCTION COMMITTEE (1990), *Feeding Standards for Australian Livestock, Ruminants*, edited by J.L. Corbett, CSIRO, East Melbourne.
- ANON (1990), 'ANU artificial breeding technologies "revolutionise" livestock production', *Laboratory News*, January.
- BARDSLEY, P. and HARRIS, M. (1987), 'An approach to the econometric estimation of attitudes to risk in agriculture', *Australian Journal of Agricultural Economics* 31(2), 112-126.
- BOND, G. and WONDER, B. (1980), 'Risk attitudes amongst Australian farmers', *Australian Journal of Agricultural Economics* 24(1), 16-34.
- DENT, J.B., HARRISON, S.R. and WOODFORD, K.B. (1986), *Farm Planning with Linear Programming: Concept and Practice*, Butterworths, Sydney.
- DILLON, J.L. (1976), 'The economics of systems research', *Agricultural Systems* 1, 5-22.
- DILLON, J.L. and HARDAKER, J.B. (1984), *Farm Management Research for Small Farmer Development*, FAO Agricultural Services Bulletin 41, Rome.
- FARQUHARSON, R.J. (1991), *An Ex Ante Economic Assessment at the Farm Level of a New Beef Production Technology*, Agricultural Economics Bulletin No. 39, Department of Agricultural Economics and Business Management, University of New England, Armidale.
- FARQUHARSON, R.J. and GRIFFITH, G.R. (1991), 'The potential economic returns to the Australian beef industry from a new twinning technology', *Agricultural Systems* 36, 79-103.
- GHODAKE, R.D. and HARDAKER, J.B. (1981), 'Whole-farm modelling for assessment of dryland technology', Economics Program Progress Report No. 29, International Crops Research Institute for the Semi-Arid Tropics, Patancheru, India.
- GOH, S., RASKIN, R. and COCHRANE, M.J. (1987), 'A generalised stochastic dominance program for the IBM-PC', University of Arkansas.
- HENNESSY, D.W. and ROBINSON, G.G. (1974), 'Problem: feeding Northern Tablelands beef cattle all year round', *Agricultural Gazette of New South Wales* 85(1), 4-5.
- HENNESSY, D.W. and ROBINSON, G.G. (1979), 'The herbage intake, eating behaviour and calf production of beef cows grazing improved pastures on the Northern Tablelands of New South Wales', *Australian Journal of Experimental Agriculture and Animal Husbandry* 19, 261-8.
- HERD, R.M., BOOTLE, B.W. and PARFETT, D.C. (1991), "An economic evaluation of traditional and novel systems of beef production", in *Genetics for Profit and Profits for Genetics*, Proceedings of the Ninth Conference, Australian Association of Animal Breeding and Genetics, University of Melbourne.
- HOLMES, W.E. (1988), 'Instructions for Breedcow and Dynama', Department of Primary Industries, Townsville, December.
- LEE, S.M., MOORE, L.J. and TAYLOR, B.W. (1985), *Management Science*, Second Edition, W.C. Brown, Dubuque, Iowa.

- LONGWORTH, J.W. and MENZ, K.M. (1980), 'Activity analysis: bridging the gap between production economics theory and practical farm management procedures', *Review of Marketing and Agricultural Economics* 48(1), 7-20.
- MAKEHAM, J.P. and MALCOLM, L.R. (1988), *The Farming Game*, Gill Publications, Armidale.
- MEYER, J. (1977a), 'Choice among distributions', *Journal of Economic Theory* 14, 326-36.
- MEYER, J. (1977b), 'Second degree stochastic dominance with respect to a function', *International Economic Review* 18, 477-87.
- MINISTRY OF AGRICULTURE, FISHERIES and FOOD (1984), *Energy Allowances and Feeding Systems for Ruminants*, 2nd edn, Reference Book 433, HMSO, London.
- MUIR, A.M. and VERE, D.T. (1987), 'The economics of pasture improvement on the New South Wales tablelands: a linear programming approach', Research Workpaper No. 8/87, Division of Marketing and Economic Services, New South Wales Department of Agriculture, July.
- NSW MEAT INDUSTRY AUTHORITY (1990), 'Understanding cattle and sheep market reports', Livestock Market Reporting Service, Sydney.
- ODDY, V.H. (1983), *Feed Requirements of Sheep and Cattle during Drought using a Metabolisable Energy System*, AGbulletin 3, Department of Agriculture New South Wales, Sydney.
- PALISADE CORPORATION (1989), '@RISK: Risk analysis and modelling for the PC', New York.
- PANDEY, S. (1990), 'Risk-efficient irrigation strategies for wheat', *Agricultural Economics* 4, 59-71.
- PARK, S.W. (1979), 'Variability in the appraisal of beef carcass characteristics and their relationship with prices in Australia', Unpublished M. Ec. dissertation, University of New England, Armidale.
- PIPER, L.R. and BINDON, B.M. (1990), 'Genetic and non-genetic approaches to increasing prolificacy in beef cattle', in *Technology Prospects and Transfer*, Proceedings of Eighth Conference, Australian Association of Animal Breeding and Genetics, Hamilton and Palmerston North, New Zealand.
- PRATT, J. (1964), 'Risk aversion in the small and the large', *Econometrica* 3, 122-36.
- RICKARDS, P.A. and PASSMORE, A.L. (1977), *Planning for Profit in Livestock Grazing Systems*, Professional Farm Management Guidebook No. 7, ABRI, University of New England, Armidale.
- THE MACQUARIE DICTIONARY (1985), Revised edition, Macquarie Library, Dee Why.
- TODD, M.C. and COWELL, M.D. (1981), 'Within-sale price variation at cattle and carcass auctions', *Australian Journal of Agricultural Economics* 25(1), 30-47.
- TURVEY, Ross (1988), 'Enterprise budgets for the North-West of NSW', COMPLAN Handbook No. 8, Agricultural Business Research Institute, University of New England, Armidale.