



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

Papers downloaded from AgEcon Search may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

ANALYSIS OF DIFFUSION STRATEGIES IN NORTHEAST BRAZIL FOR NEW CASSAVA VARIETIES WITH IMPROVED NUTRITIONAL QUALITY

Carolina González

International Center for Tropical Agriculture (CIAT) - International Food Policy Research Institute (IFPRI)
A.A. 6713, Cali, Colombia
E-mail: c.gonzalez@cgiar.org

Salomón Pérez

Socioeconomic and impact analysis, International Center for Tropical Agriculture (CIAT), Cali,
Colombia
E-mail: s.p.suarez@cgiar.org

Carlos Estevao Cardoso

Mandioca e Fruticultura, Embrapa, Brazil
E-mail: estevao@cnpmf.embrapa.br

Robert Andrade

Socioeconomic and impact analysis, International Center for Tropical Agriculture (CIAT), Cali,
Colombia
E-mail: r.s.andrade@cgiar.org

Nancy Johnson

Economist, team leader, International Livestock Research Institute (ILRI)
E-mail: n.johnson@cgiar.org

Selected Poster prepared for presentation at the International Association of Agricultural Economists (IAAE) Triennial Conference, Foz do Iguaçu, Brazil, 18-24 August, 2012.

Copyright 2012 by Carolina González, Salomón Pérez, Carlos Estevao Cardoso, Robert Andrade and Nancy Johnson. All rights reserved. Readers may make verbatim copies of this document for non-commercial purposes by any means, provided that this copyright notice appears on all such copies.

Carolina González¹, Salomón Pérez², Carlos Estevao Cardoso³, Robert Andrade⁴ and Nancy Johnson⁵

1 International Center for Tropical Agriculture (CIAT), Cali, Colombia -International Food Policy Research Institute (IFPRI), Washington, D.C., United States of America; 2,4 Socioeconomic and Impact Analysis, International Center for Tropical Agriculture (CIAT), Cali, Colombia; 3 Mandioca e Fruticultura, Embrapa, Brazil; 5 Economist, Team Leader, International Livestock Research Institute (ILRI)

Introduction

More than 130 million pre-school children suffer from vitamin A deficiency (Meenakshi *et al.*, 2010), which increases the prevalence and severity of infectious diseases (morbidity and mortality) and may cause severe eye problems, including permanent blindness. In addition, VAD results in high costs for the health system and the economy as a whole (Qaim *et al.*, 2007).

In 2006, EMBRAPA supported by the HarvestPlus Program and CIAT, released in Northeast of Brazil four yellow cassava varieties with improved levels of provitamin A: BRS Dourada, BRS Gema de Ovo, Amarelo I and Amarelo II (Fukuda *et al.*, 2008). The provitamin A levels in the new yellow cassava varieties varied between levels of 4 and 12 µg per g. of fresh weight.



Source: www.texbrasil.com.br

Methods

Two comprehensive surveys were carried out in Northeast Brazil, the first with producers involved in a participatory research process (Group 1 – sample of 359 farmers) and the second with producers who requested stakes (seeds) following a launch event via telephone or internet (Group 2 – sample of 40 farmers).

For Group 1, in addition to descriptive analysis, a logit model was implemented in data analysis, using a dichotomous dependent variable of the potential adoption rate (whether or not producers would continue planting the new cassava varieties based on their experiences). This variable provided an indication of the success of the transfer process.

Results Group 1

- ✓ 62% of the sample group responded affirmative to the question of whether they intended to plant the new varieties (potential adoption rate).
- ✓ The main reasons for adopting the new varieties were: nutritional content (90%), family preference for their flavor (7%) and acceptance in the market (3%).
- ✓ The reasons given for not adopting the new varieties were: not knowing how to obtain the seed (43%), dislike of the taste (27%), lack of tradition in the region for planting yellow cassava varieties (21%) and low productivity compared to traditional varieties in combination with low resistance to diseases (10%).

Table 1. Improved yellow varieties versus conventional varieties (% affirmative answers)

Variable		Group 1 (n=359) %	Group 2 (n=40) %
Culinary quality	More than six months	15.6	15.0
	Less than six months	3.6	2.5
	Same as conventional varieties	5.3	2.5
	Don't know or no answer	75.5	80.0
Perishable*	Slower	10.3	5.0
	Faster	6.1	2.5
	Same as conventional varieties	7.2	10.0
	Don't know or no answer	76.3	82.5
Ease of harvesting	Easier	35.1	42.5
	More difficult	1.7	2.5
	Same as conventional varieties	8.4	17.5
	Don't know or no answer	54.8	38.0
Ease of peeling	Easier	33.7	47.5
	More difficult	1.4	0
	Same as conventional varieties	8.9	10.0
	Don't know or no answer	56.0	42.5

*Ref: Conventional varieties used in their region

Results Group 2

- ✓ 75% of the producers remembered having received seed of the two varieties from EMBRAPA, and 62.5% planted the seeds in 2007.
- ✓ The reasons given for not planting generally related to a lack of seed availability and adaptability of the new varieties to the region.
- ✓ 37% professed to have given away their new variety stakes to neighbors and friends.
- ✓ 17.5% used the new varieties in purposes other than direct consumption, such as cassava flour and starch consumption.
- ✓ The main reasons for replanting yellow varieties were: a high level of acceptance by family members (40%) and their improved nutritional value (26%).

Results of the binary model – Group1


Variable	Coef.	s.e.
Income > 2 minimum monthly wage	-1.52 **	0.40
Farm size (hectares)	0.02***	0.01
Pulp color preference (white=1)	-0.68 **	0.34
Involved in participatory research activities (yes=1)	0.59 *	0.35
Main sources of information (Private)	-0.88***	0.30
Does not look for information	-1.30***	0.39
Perceives differences in crop management (yes =1)	-1.65***	0.54
Perceives differences in culinary quality (yes=1)	2.36***	0.53
Aware of nutritional advantages (yes=1)	0.68 *	0.38

*,**,*** Statistically significant at the 0.10, 0.05 and 0.01 level, respectively

- ✓ Producers with low income levels were found to have more probability of adopting the new varieties, while producers who preferred the white flesh cassava tended not to adopt these.
- ✓ Variables related to the diffusion process that increased the probability of adoption included: involvement in participatory research, knowledge about the nutritional advantages of new cassava varieties and access to information from a public source (i.e. EMBRAPA).

Conclusions

- ❖ Awareness of the nutritional advantages of new varieties, to promote participatory research activities and to socialize projects among producers are strong determinants in the success of the adoption process.
- ❖ The lack of seed availability has been the main factor limiting the adoption of the new cassava varieties.
- ❖ Seed production and distribution systems should be improved.
- ❖ It is important to continue exploring other low-cost diffusion strategies such as agricultural products stores, launch events, web pages and publications, although they need to be complemented with field work given that many poor producers typically do not have access to these information sources.
- ❖ The overarching recommendation is that future studies about cost-effectiveness should be undertaken, in combination with the development of an impact evaluation of different diffusion strategies for providing information, so as to facilitate future decisions about the diffusion programs.



Source: Welsch R et al. Plant Cell 2010;22:3348-3356

References

1. Meenakshi, J. V., Johnson, N. L., Manyong, V., de Groot, H., Javelosa, J., Yanggen, D., Naher, F., Gonzalez, C., Garcia, J. and Meng, E. (2010). How cost-effective is biofortification in combating micronutrient malnutrition? An ex ante assessment. *World Development* 38: 64–75.
2. Qaim, M., Stein, A. J., and Meenakshi, J. V. (2007). Economics of biofortification. *Agricultural Economics* 37(S1): 119–133.
3. Fukuda, M. W. G., Souza, L., Fukuda, C.H., Santos, V., Borromeu, C., Silva, M. N., Coreolano, J. W., Pinho, J. L. And Santos, A. R. (2008). Adoção de variedades de mandioca de polpa amarela para mesa no nordeste brasileiro. *Embrapa Mandioca e Fruticultura Tropical, Cruz das Almas, Brasil. Boletim de Pesquisa e Desenvolvimento* no. 41.