



*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

*No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.*

## Stata tip 76: Separating seasonal time series

Nicholas J. Cox  
Department of Geography  
Durham University  
Durham, UK  
n.j.cox@durham.ac.uk

Many researchers in various sciences deal with seasonally varying time series. The part rhythmic, part random character of much seasonal variation poses several graphical challenges for them. People usually want to see both the broad pattern and the fine structure of trends, seasonality, and any other components of variation. The very common practice of using just one plot versus date typically yields a saw-tooth or roller-coaster pattern as the seasons repeat. That method is often good for showing broad trends, but not so good for showing the details of seasonality. I reviewed several alternative graphical methods in a *Speaking Stata* column (Cox 2006). Here is yet another method, which is widely used in economics. Examples of this method can be found in Hylleberg (1986, 1992), Ghysels and Osborn (2001), and Franses and Paap (2004).

The main idea is remarkably simple: plot separate traces for each part of the year. Thus, for each series, there would be 2 traces for half-yearly data, 4 traces for quarterly data, 12 traces for monthly data, and so on. The idea seems unlikely to work well for finer subdivisions of the year, because there would be too many traces to compare. However, quarterly and monthly series in particular are so common in many fields that the idea deserves some exploration.

One of the examples in Franses and Paap (2004) concerns variations in an index of food and tobacco production for the United States for 1947–2000. I downloaded the data from <http://people.few.eur.nl/paap/pbook.htm> (this URL evidently supersedes those specified by Franses and Paap [2004, 12]) and named it `ftp`. For what follows, year and quarter variables are required, as well as a variable holding quarterly dates.

```
. egen year = seq(), from(1947) to(2000) block(4)
. egen quarter = seq(), to(4)
. gen date = yq(year, quarter)
. format date %tq
. tsset date
. gen growth = D1.ftp/ftp
```

Although a line plot is clearly possible, a scatterplot with marker labels is often worth trying first (figure 1). See an earlier tip by Cox (2005) for more examples.

```
. scatter growth year, ms(none) mla(quarter) mlabpos(0)
```




Figure 1. Year-on-year growth by quarter for food and tobacco production in the United States: separate series

Immediately, we see some intriguing features in the data. There seems to be a discontinuity in the early 1960s, which may reflect some change in the basis of calculating the index, rather than a structural shift in the economy or the climate. Note also that the style and the magnitude of seasonality change: look in detail at traces for quarters 1 and 4. No legend is needed for the graph, because the marker labels are self-explanatory. Compare this graph with the corresponding line plot given by [Franses and Paap \(2004, 15\)](#).

In contrast, only some of the same features are evident in more standard graphs. The traditional all-in-one line plot (figure 2) puts seasonality in context but is useless for studying detailed changes in its nature.

```
. tsline ftp
```




Figure 2. Quarterly food and tobacco production in the United States

The apparent discontinuity in the early 1960s is, however, clear in a plot of growth rate versus date (figure 3).

```
. tsline growth
```




Figure 3. Year-on-year growth by quarter for food and tobacco production in the United States: combined series

An example with monthly data will push harder at the limits of this device. Grubb and Mason (2001) examined monthly data on air passengers in the United Kingdom for 1947–1999. The data can be found at <http://people.bath.ac.uk/mascc/Grubb.TS>; also see Chatfield (2004, 289–290). We will look at seasonality as expressed in monthly shares of annual totals (figure 4). The graph clearly shows how seasonality is steadily becoming more subdued.

```
. egen total = total(passengers), by(year)
. gen percent = 100 * passengers / total
. gen symbol = substr("123456789OND", month, 1)
. scatter percent year, ms(none) mla(symbol) mlabpos(0) mlabsize(*.8) xtitle("")
> ytitle(% in each month) yla(5(5)15)
```




Figure 4. Monthly shares of UK air passengers, 1947–1999 (digits 1–9 indicate January–September; O, N, and D indicate October–December)

Because some users will undoubtedly want line plots, how is that to be done? The `separate` command is useful here: see Cox (2005), [D] `separate`, or the online help. Once we have separate variables, they can be used with the `line` command (figure 5).

```

. separate percent, by(month) veryshortlabel
. line percent1-percent12 year, xtitle("") ytitle(% in each month) yla(5(5)15)
> legend(pos(3) col(1))

```




Figure 5. Monthly shares of UK air passengers, 1947–1999

You may think that the graph needs more work on the line patterns (and thus the legend), although perhaps now the scatterplot with marker labels seems a better possibility.

If graphs with 12 monthly traces seem too busy, one trick worth exploring is subdividing the year into two, three, or four parts and using separate panels in a `by()` option. Then each panel would have only six, four, or three traces.

## References

- Chatfield, C. 2004. *The Analysis of Time Series: An Introduction*. 6th ed. Boca Raton, FL: Chapman & Hall/CRC.
- Cox, N. J. 2005. Stata tip 27: Classifying data points on scatter plots. *Stata Journal* 5: 604–606.
- . 2006. Speaking Stata: Graphs for all seasons. *Stata Journal* 6: 397–419.
- Franses, P. H., and R. Paap. 2004. *Periodic Time Series Models*. Oxford: Oxford University Press.
- Ghysels, E., and D. R. Osborn. 2001. *The Econometric Analysis of Seasonal Time Series*. Cambridge: Cambridge University Press.

Grubb, H., and A. Mason. 2001. Long lead-time forecasting of UK air passengers by Holt–Winters methods with damped trend. *International Journal of Forecasting* 17: 71–82.

Hylleberg, S. 1986. *Seasonality in Regression*. Orlando, FL: Academic Press.

Hylleberg, S., ed. 1992. *Modelling Seasonality*. Oxford: Oxford University Press.