AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

Agricultural Adaptation to a Changing Climate

Economic and Environmental Implications Vary by U.S. Region

Scott Malcolm, Elizabeth Marshall, Marcel Aillery,
Paul Heisey, Michael Livingston,
and Kelly Day-Rubenstein

www.ers.usda.gov

Visit Our Website To Learn More!

www.ers.usda.gov

Recommended citation format for this publication:

Malcolm, Scott, Elizabeth Marshall, Marcel Aillery, Paul Heisey, Michael Livingston, and Kelly Day-Rubenstein. *Agricultural Adaptation to a Changing Climate: Economic and Environmental Implications Vary by U.S. Region*, ERR-136, U.S. Department of Agriculture, Economic Research Service, July 2012.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and, where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

United States
Department
of Agriculture

Economic
Research
Report
Number 136

July 2012

A Report from the Economic Research Service

www.ers.usda.gov

Agricultural Adaptation to a Changing Climate

Economic and Environmental Implications Vary by U.S. Region

Scott Malcolm, smalcolm@ers.usda.gov

Elizabeth Marshall, Marcel Aillery, Paul Heisey,
Michael Livingston, and Kelly Day-Rubenstein

Abstract

Global climate models predict increases over time in average temperature worldwide, with significant impacts on local patterns of temperature and precipitation. The extent to which such changes present a risk to food supplies, farmer livelihoods, and rural communities depends in part on the direction, magnitude, and rate of such changes, but equally importantly on the ability of the agricultural sector to adapt to changing patterns of yield and productivity, production cost, and resource availability. Study findings suggest that, while impacts are highly sensitive to uncertain climate projections, farmers have considerable flexibility to adapt to changes in local weather, resource conditions, and price signals by adjusting crops, rotations, and production practices. Such adaptation, using existing crop production technologies, can partially mitigate the impacts of climate change on national agricultural markets. Adaptive redistribution of production, however, may have significant implications for both regional land use and environmental quality.

Keywords: climate change, adaptation, water resources, agricultural pests, Regional Environment and Agriculture Programming (REAP) model, regional crop mix, regional environmental effects, drought tolerance, pest management

Acknowledgments

The authors gratefully acknowledge the assistance of Gerald Nelson from the International Food Policy Research Institute and USDA Economic Research Service analysts Ryan Williams, Vince Breneman, and Paul Westcott for their contributions. The authors appreciate the thoughtful reviews from Allan Auclair, Noel Gollehon, Jerry Hatfield, Greg Johnson, Ron Sands, Roger Sedjo, along with Bill Hohenstein and the staff of USDA's Climate Change Program Office. The authors also thank Dale Simms, Angela Anderson, and Wynnice Pointer-Napper for editorial and design assistance.

Contents

Summary	iii
Introduction	1
Scope of the Research	5
Climate Change and Agricultural Impact Analysis	8
Regional Environment and Agriculture Programming Model	8
Characterizing the Study's Climate Projections	10
Quantifying Climate Change Impacts on Crop Yields	12
Quantifying the Impacts of Climate Change and Adaptation Behavior Using REAP	14
Quantifying the Additional Impacts of Pest Distribution Change	15
Quantifying the Impacts of Drought-Tolerant Varieties	17
Quantifying the Agricultural Impact of Climate Change and the Potential for Agricultural Adaptation	18
Climate Impact Analysis: No Adaptation	18
Climate Change Impacts When Farmers Adapt Crop Rotations, Tillage, and Land Use	21
How Farmer Adaptation Affects Measured Impacts of Climate Change	27
Impact of Changes in Pest Prevalence on Crop Production Under Climate Change	32
Lessening the Impact of Climate Change Through Technical Change: Targeting Crop Breeding for Adaptation to Climate Stress	41
Adoption of Plant Varieties With Greater Tolerance to Climate Change-Related Stresses	41
Impact of Adopting Drought-Tolerant Crop Varieties	44
Conclusions	53
Analysis Limitations and Future Research	55
References	59
Appendix A: The REAP Model	67
Appendix B: Climate Projections for 2030	68
Appendix C: Sensitivity of Analysis Results to Climate Change Elements	70
Appendix D: An Econometric Model of Regional Pesticide Expenditures	73

Summary

What Is the Issue?

Agricultural production has always been affected by variability in weather, and U.S. farmers have adopted production practices and strategies appropriate to their local climate. The weather that shapes the structure of U.S. agricultural production, however, is changing along with world climatic conditions. Climate models predict increases in average temperatures worldwide, with wide-ranging impacts on local temperature and rainfall. Whether such changes present a risk to food supplies, farmer livelihoods, and rural communities depends partly on the direction, magnitude, and rate of such changes, but also on the agricultural sector's responsiveness to changing yield and productivity patterns, production costs, and resource availability. Adaptive behaviors will allow producers to mitigate costs of climate change and even to capitalize on new opportunities. The introduction of crop varieties better adapted to new growing conditions could facilitate this transition.

What Did the Study Find?

The projected impacts of climate change in 2030 vary widely both across climate scenarios and across regions within a single scenario, primarily due to the direction and magnitude of precipitation changes. Farmers' ability to alter crops, rotations, and production practices enables them to lessen the impact of changes in local weather, resource conditions, and price signals. Redistributing production across regions can greatly mitigate the impact of climate change on national agricultural markets. Such redistribution, however, will alter land use and environmental quality. Key findings (with ranges expressed across different climate scenarios) include:

- National acreage changes when farmers adapt are relatively small across climate change scenarios (from 0.2 to 1.0 percent compared with the baseline), although acreage changes vary considerably by region. Crop acreage and planting patterns in the Corn Belt and Northern regions, in general, are less sensitive to climate change than in Southern regions, where yield changes have a wider range across crops (for example, acreage changes in the Delta region range from -9.8 to 5.0 percent). Acreage changes indicate considerable capacity in the agricultural system to reallocate crop production in response to shifting conditions.
- Although climate change leads to higher prices for corn and soybeans under hotter, drier scenarios as a result of considerably lower national yields, adaptation to climate change dampens the rise in prices for most commodities.
- Aggregate national returns to crop production decline with the increasing severity of the climate change scenario. The same trend holds for the Corn Belt, which accounts for over half of all returns to U.S. field crop production. The complex interaction between regional yield changes, markets, and production options—combined with the Corn Belt's large production—creates a larger absolute impact than in other regions, although the percentage decline in returns is smaller than in other regions. Changes in returns vary in the other regions, however, with no direct correspondence to the magnitude of the scenario's temperature and

precipitation change. This is due to shifts in the economic attractiveness of crops in regions other than the Corn Belt.

- Aggregate impacts of climate change on net returns to crop farmers range from an estimated increase of \$3.6 billion to a loss of \$1.5 billion per year, under the four climate change scenarios. Spread and redistribution of agricultural pests may reduce these returns by \$1.5 billion to \$3.0 billion.
- Regionally, crop sector impacts from climate change are likely to be greatest in the Corn Belt, with annual losses ranging from \$1.1 billion to \$4.1 billion across scenarios. Heightened damage from crop pests could lead to additional losses of \$400 million to \$600 million in that region. Economic effects in other regions may be positive or negative, depending on how well crop rotation and tillage practices accommodate changes in temperature and precipitation and how market-mediated prices change for predominant regional crops. Drought-tolerant varieties increase returns nationally and in regions that plant them, indicating that further development of drought-tolerant varieties could be beneficial under a wide range of adverse climate changes.
- Changes in crop production result in and reflect changes in crop prices. Soybean markets may be particularly sensitive, with estimated price effects ranging from -4 to 22 percent. Corn prices are estimated to change between -2 and 6 percent, while wheat prices are estimated to decline across all four scenarios. Shifting agricultural pest populations cause the price range to widen and crop prices to increase for all crops except cotton. The availability of drought-tolerant crop varieties is estimated to reduce prices.
- Climate change is projected to slightly increase aggregate natural resource and environmental impacts from U.S. agricultural production, although local effects may be more significant. Cropland area is projected to expand 0.2-1.0 percent, while nitrogen fertilizer losses are projected to grow 1.4-5.0 percent. Rainfall-related soil erosion changes range from -0.9 to 1.2 percent above baseline levels. The disproportionate change in nitrogen loss to water relative to acreage expansion reflects changes in regional crop distribution, input use, and the varying impacts of changes in production practices.

This report focuses on how crop farmers will adapt to changing climate conditions and how extensively changing pest pressures and emergent technologies such as drought-resistant crops might alter the benefits of adaptation. While interactions between the crop and livestock sectors are included in the analysis, changes in the livestock sector are not the focus of the report. Consumers will likewise be affected by adjustments in both the crop and livestock sectors. Livestock producers will see changes in the prices they pay for feed, and retail food prices will adjust to commodity price changes.

Our climate change analysis focused on the yield-related impacts associated with increased average temperatures, regional changes in average precipitation, increased carbon dioxide concentration in the atmosphere, the expanded incidence of pests, and the market-mediated price impacts that arise from regional shifts in crops and practices. Model limitations precluded analysis of yield impacts from the potential increase in extreme weather events, nor could

the analysis address the potential for, and constraints to, expanding irrigated acreage and water use, which is particularly important in the Western United States where there is already significant competition for water resources.

How Was the Study Conducted?

Downscaled climate projections from four different general circulation models—based on the Intergovernmental Panel on Climate Change’s (IPCC) Special Report on Emissions Scenarios (SRES) A1B emissions scenario—represent possible climate futures in the United States. A crop-growth simulator—the Environmental Productivity and Integrated Climate (EPIC) model—is used to estimate the effect on crop yields of associated weather patterns resulting from each climate projection and a suite of environmental indicators associated with each regional production enterprise, which consists of a single crop rotation/tillage/fertilizer regime. Climate projections, historical climate data, and Agricultural Resource Management Survey (ARMS) data are also used to estimate cost and yield impacts associated with potential changes in the geographic distribution and severity of pest and disease outbreaks resulting from climate change. The Regional Environment and Agriculture Programming (REAP) model—a mathematical programming model of the U.S. agricultural sector—is then used to project shifts in regional agricultural production given climate-induced changes in crop productivity patterns and price/demand feedback from national commodity and livestock markets. REAP also allows researchers to estimate the impact on national agricultural production, crop prices, regional farmer income, and—in combination with EPIC results—regional indicators of environmental quality.

Introduction

Agricultural production has always been closely linked with, and vulnerable to, trends in weather. As a result, agricultural production enterprises and practices have adapted to local climatic conditions, and farmers have developed strategies for responding to local weather variability. Corn farmers in the Corn Belt push back planting dates in response to a wet spring, for example, and may switch to soybean production if persistent wet weather delays corn planting excessively. During extremely dry periods, farmers in the Plains States may increase moisture-conserving tillage practices, such as no-till, ridge-till, and mulch-till (Ding, 2009). Local strategies for weather adaptation are based on years of producer experience and farming-system research specific to regional conditions.

The range of local weather conditions that has shaped the current structure of domestic agricultural production, however, is changing in response to broad shifts in general climatic conditions across the country and around the world. General climatic conditions have adjusted slowly throughout the 20th century, with global average temperature increasing 1.3 degrees Fahrenheit (°F) (IPCC, 2007). As atmospheric concentrations of carbon dioxide (CO₂) have increased, the rate of temperature increase appears to be accelerating, and recent climate models predict further warming trends over time that may have a significant impact on local temperature and precipitation patterns.

Agricultural productivity, and the degree to which other inputs (such as fertilizer, pesticides, and irrigation) are needed to augment production, depend a great deal on local climate conditions. Increases in average temperature, changes in precipitation patterns, and increases in the frequency of extreme weather events would significantly alter the local production environment through the distribution of crop yields, crop acreage planted to different crops, reliance on dryland and irrigated production systems, and the geographic range and severity of pest outbreaks. Changes in water availability for crop production will be an important factor affecting regional agricultural production. Shifting precipitation patterns in combination with warming temperatures may increase water scarcity in some regions, intensifying competition for water currently used in agriculture. In other areas, increased soil-moisture availability may increase opportunities for agricultural production.

Agricultural systems respond to the changing production environment associated with climate change through the process of adaptation. The Intergovernmental Panel on Climate Change (IPCC) defines adaptation as the “adjustment in natural or human systems in response to actual or expected climatic stimuli or their effects, which moderates harm or exploits beneficial opportunities” (IPCC, 2007). Agricultural systems adapt to climate change at a number of levels, from national-level investments in agricultural research and development, climate forecasting, or infrastructure to behavioral adjustments of individual farm households. Smit and Skinner (2002) organize agricultural adaptation options within four interdependent categories (table 1):

- Technological developments,
- Government programs and insurance,
- Farm production practices, and
- Farm financial management.

Table 1

Types and examples of agricultural adaptation options**Technological developments**

Crop development:

- Develop new crop varieties, including hybrids, to increase the tolerance of and suitability of plants to temperature, moisture, and other relevant climatic conditions.

Weather and climate information systems:

- Develop early warning systems that provide daily weather predictions and seasonal forecasts.

Resource management innovations:

- Develop water management innovations, including irrigation, to address the risk of moisture deficiencies and the increasing frequency of droughts.
- Develop farm-level resource management innovations to address the risk associated with changing temperature, moisture, and other relevant climatic conditions.

Government programs and insurance

Agricultural subsidy and support programs:

- Modify crop insurance programs to influence farm-level risk management strategies with respect to climate-related loss of crop yields.
- Modify subsidy, support, and incentive programs to influence farm-level production practices and financial management.

Private insurance:

- Develop private insurance to reduce climate-related risks to farm-level production, infrastructure, and income.

Resource management programs:

- Develop and implement policies and programs to influence farm-level land and water resource use and management practices in light of changing climate conditions.

Farm production practices

Farm production:

- Diversify crop and livestock types and varieties to address environmental variations and economic risks associated with climate change.
- Change production intensity to address environmental variations and economic risks associated with climate change.

Land use:

- Use alternative fallow and tillage practices to address climate-related moisture and nutrient deficiencies.

Irrigation:

- Implement irrigation practices to address the moisture deficiencies associated with climate change and reduce the risk of income loss due to recurring drought.

Timing of operations:

- Change farm operation timing to address the changing duration of growing seasons and associated changes in temperature and moisture.

Farm financial management

Crop insurance:

- Purchase crop insurance to reduce the risks of climate-related income loss.

Crop shares and futures:

- Invest in crop shares and futures to reduce the risks of climate-related income loss.

Household income:

- Diversify household income to address the risk of climate-related income loss.

Source: Adapted from Smit and Skinner, 2002

While adaptation may take many forms throughout the farm economy, this report focuses specifically on the potential for adaptation at the first level of response—farmer behavior. Adapting to changing conditions is nothing new for farmers; they regularly adapt to changes in crop demand, new technological developments, farm policy provisions, land development pressure, and, most significantly, weather variability.

Changes in individual farmer behavior in response to climate change may include, but are not limited to, growing different crops or crop varieties; adjusting planting and harvest dates; altering input use, such as applied fertilizers, pesticides, and water; adopting new production methods; expanding planted acreage; or abandoning farming altogether. How individual farmers respond to changing conditions is a function of each farmer's location, resource endowment, economic incentives, and knowledge of alternatives. While farming enterprises are likely to adapt in some way to shifting climate conditions, the costs and benefits of adaptation may vary considerably depending on the farm's location, the crops grown, and other factors that differ across operations.

Similarly, regional impacts of changing climate will not be homogeneous; some regions may see an improvement in crop growth potential, while others may face declining productivity (see box, "Climate Change Impact on Yields"). An analysis that focuses exclusively on the average effect of climate change on national production, commodity and food prices, and agricultural trade would mask important regional differences. Climate change that alters the relative profitability of regional crop production may redistribute production and resource allocations across regions, with potentially significant implications for producer income, resource use, and environmental quality. Assessing the potential impact of climate change on the U.S. agricultural sector requires the ability to differentiate among regional impacts and allow for adaptive behavior that results in shifts within and across production regions in response to changing climate regimes.

Historically, genetic enhancement—the combination of biological research, plant breeding, and genetic resources—has played a key role in maintaining and improving agricultural productivity. As agriculture adapts to global climate change, however, genetic combinations that are optimal for *current* growing environments are unlikely to be optimal for *future* growing environments. Adaptive genetic enhancement of traits, such as drought and heat tolerance, may offer critical assistance to producers' long-term response to the challenges of climate change.

The extent to which changing weather patterns will impact the distribution and severity of pests (Hatfield et al., 2008) and invasive species (USDA, 2010a) may also influence adaptive decisionmaking strategies in the short and long term. Increased pesticide and herbicide use is one possible response (Bridges, 1992; Joyce et al., 2008). Genetic manipulation of crops to better resist pest and disease infestation is another. Crop distribution may also change, with production of vulnerable crops moving to less risky regions. Fully characterizing the potential agricultural impact of climate change means assessing how crop distributions, yield impacts, and the costs of prevention and control might be affected by regional temperature changes and precipitation levels and associated shifts in agroecological systems.

There are several pathways through which the changing conditions associated with climate change are likely to influence crop growth and development.

Increasing Temperatures

The impact of increasing temperatures on crop growth will depend on how climate change shifts local temperatures relative to the optimal temperature range for the crop varieties growing in that region. Research suggests that crops may be particularly sensitive to temperature extremes during the reproductive phase, when pollen viability and seed setting are vulnerable to high temperatures (USCCSP, 2008). Higher average temperatures may also result in accelerated crop maturity, as optimal air temperatures for growth occur earlier in the season, which can result in less seasonal growth and lower yield potential.

Temperature also has an important effect on crop water demand. Increased crop water requirements under a warming climate may place greater demands on available soil moisture and irrigation water supplies. Actual water demand will depend on other climatic factors as well, including field humidity and shifts in solar radiation caused by changing cloud cover and aerosol concentrations.

Changes in Local Precipitation Patterns

A significant body of research has addressed the impact of climate change on water resources (NWAG, 2000; Thomson et al., 2005; IPCC, 2007; USCCSP, 2008; USDOL, 2011). While General Circulation Models (GCMs) predict a wide range of future precipitation patterns for the United States, some projected precipitation trends have emerged more consistently than others from the modeling literature. Annual precipitation has been projected to increase over much of the Eastern United States and across the middle-to-high latitudes of the Central and Western United States.¹ In contrast, potential precipitation declines are projected for the Southwest, Central Mountain region, Southern Plains, and Delta region, with the direction of precipitation change less evident across the Southeastern United States and Central Plains. Changes in total precipitation are also projected to be accompanied by interseasonal shifts in the timing of precipitation, with a larger share of precipitation falling in the winter months and smaller amounts in the summer.

¹These projections are generally consistent with higher levels of recorded precipitation over the latter half of the 20th century (USCCSP, 2008).

²The quantitative impacts of increased CO₂ on yield are still being discussed in the literature. Most results come from greenhouse or open-top chambers in the field, with only a few experiments conducted through FACE (free-air carbon dioxide enrichment) methods, which may provide results more representative of actual field conditions (Lobell and Burke, 2010).

Increasing Atmospheric Carbon Dioxide (CO₂)

Crop yields have been observed to increase with increasing levels of atmospheric CO₂, though yield response differs by crop. Yield increases associated with increasing CO₂ arise through two pathways: increased rates of photosynthesis and reduced water loss through transpiration. Research suggests that rising CO₂ concentrations that limit plant transpiration through the stomata could help mitigate the increase in crop water stress experienced as a result of higher temperatures (Izaurralde et al., 2003). The transpiration effect (the magnitude of which depends largely on soil moisture levels) operates in all crops. Impacts on crop yields via the photosynthetic pathway, however, operate only in a subset of plants. Plants have two different metabolic pathways for photosynthesis—C₃ and C₄—but only the C₃ photosynthetic pathway responds to increased atmospheric CO₂. C₃ crops are therefore projected to have a higher yield response to increased atmospheric CO₂ than are C₄ crops.

Among the REAP model's major field crops, only corn and sorghum are C₄ plants; other major crops, such as wheat, soybeans, and cotton, are classified as C₃ crops and therefore are more likely to respond positively to increased atmospheric CO₂.² The U.S. Climate Change Science Program (USCCSP, 2008) reported that a doubling of CO₂ increased estimated yields by approximately 4 percent for corn, 0-8 percent for sorghum, 44 percent for cotton, and 34-38 percent for soybeans. Actual responses to increasing atmospheric CO₂ will depend upon whether crop growth is constrained by other stressors, such as nitrogen or water limitations.

Changing Patterns of Pests and Disease

Changes in the geographic distribution of crop pests and diseases (Hatfield et al., 2008) and invasive species (USDA, 2010a; Mooney and Hobbs, 2000; Ziska et al., 2010) as a result of climate change are expected to increase yield losses and management costs. In general, weed species are expected to benefit more than crop species from increasing temperatures and CO₂ concentration levels, and crop species less able to adapt to changing climatic conditions are expected to be more susceptible to attack by pests. Although the shift in range of particular pests and invasive species will vary, with some expanding and others contracting (Bradley et al., 2009), climate change is expected to lead to a northward expansion

Continued on page 5

of many damaging pests and diseases. Furthermore, herbicide use and associated costs are expected to increase, not only because of increases in pest pressure, but also because herbicides generally become less effective as temperatures and CO₂ levels rise (Kiely et al., 2004). Control costs and crop losses as a result of weeds, insects, and diseases are therefore expected to increase, especially in northern regions where U.S. field crop production is concentrated.

Changes in Soil Fertility and Erosion Rates

For many years, researchers have speculated that the higher temperatures associated with climate change could accelerate the decomposition of organic matter in the soil, making soil less fertile and quicker to release CO₂ and nitrous oxide (N₂O) from the soil. As far back as 1938, scientists recognized the importance of soil organic matter to maintain soil productivity and access to plant-available nitrogen and observed an empirical relationship between decreased soil organic matter and increased temperature and/or decreased precipitation (Albrecht, 1938). Recent field research has corroborated that differential impacts of temperature on soil organic compounds may mean that warmer temperatures shift soil molecular carbon composition toward forms of

carbon less accessible to plants (Feng et al., 2008). Increased soil erosion may also contribute to soil fertility losses in a warming world. Climate change may impact erosion rates through a number of possible pathways, including increased intensity of rainfall events, shifting incidence of precipitation from snowfall to rainfall, changing soil organic structure, and changes in residue or litter cover due to changing yields, cultivation practices, and decomposition rates.

Changes in Climatic Variability and the Incidence of Extreme Events

Climate modeling assessments also point to an increase in precipitation variability, including increases in extreme weather events, as a potential result of a warming climate. Risk of flood damages are likely to rise in basins projected for higher annual runoff or rapid early-season snowmelt runoff (USDOI, 2011). Greater frequency and intensity of storm events would likely increase the potential for pollutant runoff. Evidence also points to increasing drought frequency and severity, particularly across the central and southern tier regions of the United States (USCCSP, 2008; Strzepek et al., 2010).

Individual farmer decisions, when aggregated to the national level, will have consequences on agricultural markets through production levels, trade, and prices and on resource use that affects environmental quality. Production adjustments may involve changes in aggregate land under cultivation; regional cropping pattern shifts, including movement of crops into areas not historically cultivated; changes in the distribution of regional crop rotations; and changes in tillage practices and fertilizer use. This study explores the regional and national implications of such farm-level adjustments for agricultural markets and environmental quality and the net effect of such adjustments on the projected impact of climate change on U.S. crop production.

Scope of the Research

Our research focused on how the crop sector might respond to climate change, specifically:

- How might farmers adjust land-use and land-management decisions when faced with a new production regime shaped by climate change, and what are the implications for regional expansion and contraction of cropland?
- How might negative impacts, such as higher prices to consumers, lower incomes for farmers, and intensification of environmental consequences, be reduced or eliminated through adaptation to climate change?
- How might changes in climate and the geographic distribution and severity of pest and disease outbreaks affect crop production and prices for major U.S. field crops?

- What impact might advances in crop research and development have on the farm production environment?
- How might the response of crop production to climate change impact soil and water quality?

We explored climate change's impacts on crop production and the potential for, and possible constraints to, adaptive behavior that addresses those impacts. Our quantitative modeling analysis then empirically examined the implications of altered climate regimes on production patterns and market conditions projected to occur by the year 2030. The quantitative methodology consisted of three phases:

- **PHASE I:** The Environmental Productivity and Integrated Climate (EPIC) model linked projections of future climate conditions to crop yields and other biophysical indicators. Since future climate projections are highly uncertain, we employed climate projections from several models to capture a range of possible climate outcomes.
- **PHASE II:** The Regional Environment and Agriculture Programming (REAP) model was first used to examine regional crop and livestock production, input use, cropping practices, economic returns, and environmental quality based on projected USDA market and production conditions under constant climate conditions (i.e., no climate change). REAP was then used to examine how production, market, and environmental measures behave under the climate change scenarios. To isolate the effect of adaptation, we examined two initial cases assuming climate change: one where farmers are not allowed to adapt by adjusting crop acreage or production practices, and a second where farmers can choose crop acreage, rotation, and tillage in response to climate-induced changes in crop yields.
- **PHASE III:** We refined the adaptation case explored in phase two to consider two important issues that may affect behavioral outcomes under climate change. One case estimated possible changes in costs and yields due to a shift in the geographic distribution and severity of pest outbreaks. A second case introduced changes to yields that might result from research and development supporting crop genetic resources for drought tolerance.

This research focused on agricultural production in the United States. Our purpose was to illustrate possible regional impacts of climate change within the United States and to explore the potential for existing and proposed production technologies both to mitigate negative impacts and to take advantage of beneficial impacts of regional shifts in relative crop yield. We recognize, however, that ecosystems across the globe will be affected by climate change, putting additional pressure on international markets and agricultural production systems worldwide through changes in commodity demand, trade patterns, and broader economic conditions. This study does not explicitly incorporate international market changes caused by climate change into its analysis.

While REAP's strength lies in the specification of crop production detail for major commodity crops, the model's structure also allows for a limited set of adaptation behaviors within the livestock sector. The model permits livestock

producers to change what they feed livestock in order to minimize costs under the new price regimes associated with climate change. The flexibility of such changes, however, is limited to a pre-existing spectrum of historically observed diet options. New grains, feed meals, and feed combinations that historically have not been used are not included in the set of livestock diet options. Furthermore, the impacts of climate change on the livestock sector are limited to those experienced indirectly through feed markets and increased competition for pasture land; the model does not capture the direct impacts of climate change on livestock productivity and production costs, such as those associated with climate control costs for confined livestock production (Key and Sneeringer, 2011). While livestock supply and demand is included in the model, results for the sector are not reported.

We first established a baseline scenario that assumes the current climate will prevail through 2030. The “no climate change scenario” establishes a baseline pattern of rotation acreages that meets a projected set of yield, production, price, and acreage measures under a set of “current” weather conditions, as measured by an average of weather conditions between 1950 and 2000. To measure the impact of climate change on those production patterns, we considered four analytical cases that reflect differing scopes of potential climate change impact and behavioral response, as shown in table 2. These cases will be described more fully as they are introduced in the report. While not exhaustive, the cases illustrate the implications of different elements of climate change impact and potential opportunities for adaptation across a range of climate change scenarios.

Table 2

Description of analysis cases

Cases: U.S. agriculture in 2030	Climate change scenarios	Adaptation allowed	Additional pest effects	Drought-tolerant varieties available
Baseline: Without climate change but with anticipated changes in export demand ¹ for agricultural commodities and continued historical rates of growth in U.S. crop yields.	No	-	-	-
No adaptation: With direct effects of climate change in temperature and precipitation but no farmer adaptation in crops, rotations, tillage, or land use.	Yes	No	No	No
Farmer adaptation: With direct effects of climate change but farmers adapt through changes in crops, rotations, tillage, and land use using existing technical options.	Yes	Yes	No	No
Additional pest damage: In addition to direct effects, climate change also alters the distribution of agricultural crop pests, leading to reduced yields and increased pesticide use and management costs.	Yes	Yes	Yes	No
Drought-tolerant varieties: Assumes farmers may adopt new, drought-tolerant varieties for some crops.	Yes	Yes	No	Yes

Note: Adaptation refers to changes by farmers in crops, crop rotations, tillage, and land use in response to the direct effects of climate change on temperature and precipitation.

¹The trend in projected future demand for U.S. farm exports does not vary across the cases, or with climate change. Although climate change may also affect foreign demand for U.S. agricultural commodities, modeling these effects is beyond the scope of this report.

Regional Environment and Agriculture Programming Model

Figure 1
Regional Environment and Agriculture Programming (REAP) model regions and USDA Farm Production Regions

Source: USDA Farm Production Regions and Natural Resources Conservation Service (NRCS) Land Resource Regions.

Each REAP model region includes a set of available crop rotations that are implemented using one of up to five tillage practices. The combination of region, rotation, and tillage practice is referred to as a production enterprise and represents the basic unit of crop production economic activity in the REAP model. A selection of regionally appropriate production enterprises was derived for each REAP region from 1997 National Resources Inventory (NRI) data. When REAP solves for agricultural production patterns under changed climate, technology, or policy conditions, acreage in each region is distributed among production enterprises based on an assessment of relative rates of return arising from differences in yields, costs, and returns and is further constrained by acreage distribution parameters that capture historically observed patterns of production. (For more information on the REAP modeling framework, see Appendix A.)

REAP Baseline

To construct a baseline against which to compare the impacts of climate change, REAP's pattern of production enterprises was calibrated to projected agricultural production conditions for 2030, assuming constant climate conditions based on climate variables calculated as a monthly average over 1950-2000 (Hijmans et al., 2005). This "no climate change" baseline scenario assumes that technology and market conditions will continue to change at historical rates and holds the suitability of a given region to produce crops constant according to a baseline set of weather conditions.

Technology and market condition projections in 2030 were extrapolated from USDA's annual agricultural production and market indicator projections (USDA, 2010b). The USDA projections include estimates of planted and harvested acreage, anticipated crop yields, trade volumes, and market prices to 2030 (table 3). The projections assume that agricultural policies remain constant and that improvements in crop yields grow at a fixed rate. Yield estimates for the baseline case were calculated by first running EPIC under the "current" weather conditions. EPIC yields were then adjusted, using

Table 3

Baseline production and market projections for 2030

Crop	Planted acres (million)	Harvested acres (million)	Production (million bushels, except as noted)	Harvested yield (bu/acre)	Price (dollars)
Corn	89	81.8	16,400	200.4	3.65
Sorghum	6.7	5.8	370	63.4	3.45
Barley	3.3	2.9	252	78.2	3.93
Oats	3.1	1.3	192	71.5	2.25
Wheat	52.5	44.6	2,371	50	4.7
Rice ¹	3.075	3.057	296.9	84.35	16.76
Soybeans	76	75	3,829	51	9.3
Cotton ²	11.5	10.2	21.495	2.125	333.6

¹Rice units measured in million cwt, or hundredweight.

²Cotton units measured in million 480-pound bales.

Source: USDA, Economic Research Service, Paul Westcott, personal communication, 2011.

a crop-specific adjustment factor, to meet projected average yields for 2030 to capture assumptions about exogenous increases in crop productivity. The baseline projections do not consider the likelihood of shocks to agricultural production from extreme weather or changes over time in other economic conditions, such as energy prices, incomes, or exchange rates that might affect exports, imports, or input prices.

Characterizing the Study's Climate Projections

There are various sources of uncertainty associated with generating estimates of future local weather conditions suitable for agricultural production impact analysis. Most significant among these are:

- The rate at which carbon and other greenhouse gases (GHG) are expected to be emitted into the atmosphere in the coming decades;
- The effects of that GHG accumulation on climate dynamics and core average climate variables, such as temperature, precipitation, and relative humidity; and
- Our capacity to downscale geographically coarse average climate projections into local projections for temperature, precipitation, and other variables that reflect daily weather changes across a finer spatial and temporal scale.

The IPCC Special Report on Emissions Scenarios (SRES) addressed the first source of uncertainty. The report defined several emissions scenarios that reflect different sets of assumptions about global population change, technology adoption, energy use, and macroeconomic conditions. The uncertainty surrounding carbon dynamics and climate response is reflected by a wide array of models that attempt to project future climate trends. These models—general circulation models (GCMs)—differ from one another in the numerical methods used, as well as in the spatial resolution at which climate projections are made. As a result, different models may vary considerably with respect to predictions of the magnitude and direction of precipitation and temperature change for given points or regions.

Each of the IPCC scenarios represented an estimated future path of CO₂ emissions that can be used as emissions input data into a GCM. Atmospheric CO₂ levels are an important driver of many long-term climate phenomena, so the various emissions scenarios result in different long-term climate projections. This analysis used climate projections derived from a single emissions scenario—the SRES A1B emissions scenario—which was designed to reflect “very rapid economic growth,” “the rapid introduction of new and more efficient technologies,” and a balanced portfolio of energy sources that included both fossil fuels and renewable energy technologies (IPCC, 2007). The SRES A1B emissions scenario represented a middle ground between other illustrative scenarios. Because this research focuses on an analysis year (2030) that occurs prior to significant divergence in emissions levels across the different scenarios, sensitivity analysis of results across the different emissions scenarios was not considered a high priority. In contrast, the significant variability across GCM results using a single emissions scenario suggested a need to explore climate projections across a number of models to capture a range of possible climate impacts arising from a single projected path of emissions.

“Downscaling” refers to the process of translating the large-scale climate information that emerges from GCMs into finer temporal and spatial resolution. There are several methods available for downscaling GCM output; the downscaled data used in this study were generated by Jones, Thornton, and Heinke (2009). Each of the four datasets represents output from a different GCM running the SRES A1B emissions scenario (table 4). For more information on the climate projections and how the downscaled data were aggregated to regions suitable for the REAP analysis, see Appendix B.

Figure 2 shows the variation in mean annual maximum temperature and precipitation change between the base period and 2030 for each of the four climate projections across REAP regions. The Model for Interdisciplinary Research on Climate (MIROC) represented the most extreme change from the base period, in that it demonstrated the largest temperature increase and the most negative precipitation change. The Commonwealth Scientific and Industrial Research Organisation (CSIRO) and Max Planck Institute’s ECH scenarios represented the mildest change projections; both predicted a similar range of temperature increases across REAP regions, but the ECH scenario predicted slightly wetter conditions relative to the CSIRO projections. The Centre National de Recherches (CNR) projection was highly variable in both temperature increase and precipitation change across the REAP regions and represented a moderate national projection in terms of the severity of climate change predicted among the scenarios considered.

While the magnitudes of shifts in maximum temperatures differed across GCMs, regional patterns of temperature impact were somewhat similar (fig. 3). The models generally projected the most moderate temperature increases in the West and Southeast and more significant temperature impacts in the Midwest and Northeast. That pattern of impact was roughly consistent with a composite of multiple IPCC model simulations generated for North America, which projected temperature increases from approximately 1 degree Celsius (°C) in the Southeast to more than 2 °C in northern Canada, with intermediate values over the rest of the contiguous United States (USCCSP, 2008).

In contrast to the consistent pattern of relative temperature impacts, there was little consistency in precipitation-change projections across GCMs

Table 4
General circulation models adopted for use in this study

Model name	Label	Institution	Reference
CNRM-CM3	CNR	Centre National de Recherches Meteorologiques (CNRM), Meteo France, France	Déqué et al. (1994)
CSIRO-Mark 3.0	CSIRO	Commonwealth Scientific and Industrial Research Organisation (CSIRO) Atmospheric Research, Australia	Gordon et al. (2002)
ECHam5	ECH	Max Planck Institute for Meteorology, Germany	Roeckner et al. (2003)
MIROC 3.2	MIROC	Center for Climate System Research (University of Tokyo), National Institute for Environmental Studies, and Frontier Research Center for Global Change (JAMSTEC), Japan	K-1 Developers (2004)

CM3 = Climate Model, version 3.
ECHam5 = European Centre – Hamburg.
MIROC 3.2 = Model for Interdisciplinary Research on Climate.

Figure 2

Estimated change in mean annual maximum temperature and precipitation for each REAP production region under the four climate change scenarios

Temperature change, 2000-2030 (degrees Celsius)

REAP = Regional Environment and Agriculture Programming model.

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations and Jones, Thornton, and Heinke, 2010.

(fig. 4). Some regions exhibited an increase in precipitation according to some models and a decrease in precipitation according to others. A few regions demonstrated a consistent direction of impact across models; the Pacific Northwest exhibited an increase in precipitation across all models, while the Texas/Louisiana region exhibited a decline in precipitation across all models. Nevertheless, even regions with a consistent direction of impact exhibited a wide range of estimated magnitudes across GCMs. Because there is no basis with which to assign probabilities to climate outcomes predicted across GCMs, crop yield impacts, adaptation potential, and aggregate system impacts were calculated and presented independently for each of the illustrative climate projections.

Quantifying Climate Change Impacts on Crop Yields

Climate change is expected to impact crop growth and development through a number of pathways (see box, “Climate Change Impacts on Crop Yields,” p. 4). In this analysis, climate change impacts on crop yields were estimated using EPIC—a field-scale biophysical model that uses a daily time step to simulate crop growth, soil impacts, hydrology, nutrient cycling, and pesticide fate under various cropping systems (e.g., tillage, crop rotation, soil and nutrient management) and weather scenarios. A random weather generator built into EPIC uses the average monthly climate information derived from the GCMs—minimum daily temperature (TMIN), maximum daily temperature (TMAX), and precipitation (PRCP)—to generate daily temperature and precipitation patterns for simulated crop growth in each REAP region.

Figure 3

Change in mean annual maximum temperature (degrees Celsius), from the baseline under the four climate change scenarios

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations and Jones, Thornton, and Heinke, 2010.

To represent a range of possible weather scenarios associated with each GCM's set of average monthly estimates, simulation results were run 10 times for 20 years, using a different random weather seed for each run. Results from the first 10 years of each run (a total of 100 years) were discarded to minimize the impact of initial soil conditions on yield and environmental impact estimates. Results from the remaining 100 years were used to calculate the average yield and environmental impact results associated with each production enterprise. Because variability estimates for future weather cannot be derived from either the original or the downscaled GCM climate output, weather variability—and therefore the incidence of extreme weather events—was held constant in this analysis across the baseline and future weather scenarios.

For each production enterprise, EPIC was used to calculate a set of yield and environmental impact measures associated with region-specific weather assumptions and four sets of regional soils differentiated by highly erodible, non-highly erodible, with tile drainage, and without tile drainage. To calculate the impact of the climate change scenarios on crop growth in each region, the crop growth parameters and geophysical process parameters used

Figure 4

Change in annual precipitation (millimeters), from the baseline under the four climate change scenarios

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations and Jones, Thornton, and Heinke, 2010.

in EPIC's simulations were held constant across the estimates generated using the baseline climate conditions and the projected climate conditions emerging from the GCMs. In moving from the baseline to the climate change projection scenarios, however, we assumed that ground-level CO₂ concentrations increased from 381 parts per million (ppm) in the baseline to 450 ppm across GCM projections. EPIC calculated the impact of the increased atmospheric CO₂ effect using a nonlinear plant response equation with crop-specific parameters (see Appendix C for a more detailed discussion of the impact of increased atmospheric CO₂ levels in this analysis). The only other variables that differed between the baseline and the climate change yield estimates were the TMIN, TMAX, and PRCP variables.

Quantifying the Impacts of Climate Change and Adaptation Behavior Using REAP

Climate-induced changes in agricultural production were assessed by substituting into REAP the yield and cost estimates for production enterprises that were estimated in EPIC using new, regionally variable climate conditions

associated with climate projections. Yield estimates were again adjusted (by the same crop-specific adjustment factor) to account for exogenous increases in productivity up to 2030. Because climate change affects crops and regions differently, the relative productivity and economic value of regional production enterprises will change under the projected climate scenarios. Production enterprises that are economical under one climate regime may not be economical under another.

Our analysis showed that in each climate change scenario, several historically established crop rotations were no longer economical to employ, possibly because the enterprise was marginally economical in the baseline and was sensitive to climate-induced changes in yield or price. Another economic driver may be that changes in other crop yields and/or prices make them more favorable to produce and force out crop rotations that do not experience improved yields or increased prices. Optimizing agricultural production levels and patterns under projected climate conditions produced a new pattern of production enterprises that reflected changes in regional production levels, including shifts among crops, crop rotations, tillage used, and expansion or contraction of cropland.

Quantifying the Additional Impacts of Pest Distribution Change

Pests and diseases reduce crop yields through several means. Weeds often reduce yields by competing for external resources. Generally, there are few genetic sources of host plant resistance to weeds, with the exception of parasitic weeds that invade the roots of crop plants. Host plant resistance to insects is more common in field crops, and host plant resistance is even more important in field crops for many plant diseases, particularly fungal diseases. Whether measured in pounds of active ingredient or in pesticide costs, herbicides are the most widely used chemicals in U.S. field crop production. Insecticide use has been common in corn and cotton. Fungicide use is quite low in field crops, although it is much higher in fruits and vegetables (Osteen and Livingston, 2006; Padgitt et al., 2000).

Global climate change, at least in terms of average annual temperature, will have the effect of making production conditions in northern U.S. regions more similar to production conditions in southern regions. In southern regions, problems with pests, especially weeds—the most important type of crop pest in terms of pesticide expenditures and yield losses—are much more severe (Bridges, 1992). In this analysis, the assumption of a temperature change-induced migration of pest costs and impacts was used to estimate the potential additional pest-related impacts on yield losses and production patterns associated with changing climate conditions. Our analysis assumed that temperature is an important factor driving changes in the geographical distributions of pests and invasive species. This was a reasonable assumption for many biological organisms because average temperatures during winter months are important determinants of overwintering survival rates and because average annual temperatures are correlated with average temperatures during winter months (Hatfield et al., 2008). Changes in precipitation patterns also guide changes in the distribution of pests and invasive species. Estimating the direct impact of precipitation changes with consistency,

however, was beyond current modeling capacity. We instead focused on estimating the direct effect of average annual temperature changes, while accounting for annual weather variation over time and space.

We began by estimating quadratic relationships between average annual temperature and latitude for the REAP crop production regions for 2000 and 2030 based on CNR, CSIRO, ECH, and MIROC temperature projections. These estimates were then used to estimate the percentage shift in each region's latitude centroid—the latitude at the center of the region—associated with the respective temperature change projections. Those percentage shifts were used to characterize the extent of each region's southward movement in latitude-temperature space (see Appendix D for more information).

We then used Agricultural Resource Management Survey (ARMS)¹ data for barley (2003), corn (1996, 2001, 2005), cotton (1997, 2003, 2007), oat (2005), rice (2000, 2006), sorghum (2003), soybean (1997, 2002, 2006), and wheat (2004) producers to estimate linear relationships between real pesticide expenditures per acre and latitude for each crop, while accounting for weather variation over time and 10 USDA National Agricultural Statistics Service (NASS) crop production regions.² The estimates from this model were combined with the estimates used to characterize the southward movement of the REAP regions in latitude-temperature space (Appendix D, table 5) to obtain percentage changes in pesticide expenditures and yield losses for each crop and for each region under climate change in 2030 relative to 2000. The percentage changes in pesticide expenditures and yield losses were then input into the REAP model to examine the pest-impact scenarios associated with climate change.

Generally, both yield loss and pesticide use increase with pest pressure (Livingston, Carlson, and Fackler, 2004; Hatfield et al., 2008). Because observations on yield loss due to pests are not available in the ARMS data, percentage yield-loss impacts were specified as a constant multiple of the percentage pesticide-expenditure impacts. The constant multiple was the elasticity of yield loss with respect to pesticide applications (0.97), which was based on estimates of a yield-loss function for cotton reported by Livingston et al. (2007). While the elasticity of yield loss with respect to pesticide applications likely varies by crop—as well as over time, as new pesticides and genetically engineered (GE) crop varieties become available—we used the estimate for cotton to specify yield-loss impacts for all crops because cotton was the only U.S. crop for which reliable yield-loss estimates due to pests were reported.

This method implicitly assumes that the shares of pesticide expenditures and yield losses associated with domestic and invasive pests, and the rate of introduction of new invasive species, would not change during 2000-30 relative to 1996-2007. These assumptions were necessary because information was not available to determine the allocation of pesticide expenditures across both pest categories in the ARMS data and because impacts of climate change on invasive species introductions are difficult to predict.

Estimates suggest that with changes in the distribution of pest populations for each climate-change scenario, pesticide expenditures and yield losses would increase for barley, corn, oats, sorghum, soybeans, and wheat; decline for cotton; and remain fairly constant for rice (table 5). The estimates

¹ARMS is a joint project of USDA's Economic Research Service and the National Agricultural Statistics Service. Visit <http://www.ers.usda.gov/Briefing/ARMS/> for more information and to download summary statistics based on these data.

²Observations on average annual temperature and precipitation were not available in the ARMS data; therefore, we used a time index and NASS production-region fixed effects to account for annual weather variation over time and space. We used the results reported in appendix table 6 to specify the pesticide-expenditure and yield-loss impacts used in the analysis.

Table 5

Average percentage change in pesticide expenditures and yield losses across the REAP (Regional Environment and Agriculture Programming model) regions for each climate change scenario, by crop

Crop	ECH		CSIRO		CNR		MIROC	
	Cost	Yield loss	Cost	Yield loss	Cost	Yield loss	Cost	Yield loss
<i>Percent</i>								
Barley	4.9	4.7	3.9	3.8	6.0	5.8	8.5	8.2
Corn	2.9	2.8	2.3	2.3	3.5	3.4	5.0	4.8
Cotton	-1.0	-1.0	-0.8	-0.8	-1.2	-1.2	-1.7	-1.7
Oats	7.4	7.1	5.9	5.7	9.0	8.7	12.9	12.4
Rice	-0.5	-0.4	-0.4	-0.4	-0.6	-0.5	-0.8	-0.8
Sorghum	3.9	3.8	3.2	3.1	4.8	4.6	6.8	6.6
Soybeans	4.2	4.1	3.4	3.3	5.1	4.9	7.3	7.0
Wheat	3.9	3.7	3.1	3.0	4.7	4.5	6.7	6.4

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Notes: Estimates based on the maximum likelihood estimates relating pesticide expenditures to latitude and the southward movements of the latitude centroids of each REAP region (see Appendix D). Percentage changes in yield losses are a constant multiple (0.9661) of the percentage changes in pesticide expenditures. The constant multiple is the elasticity of yield loss with respect to pesticide applications and is based on estimates and data reported by Livingston et al. (2007).

suggest that the largest impacts would occur in oats, which could experience increases in pesticide expenditures between 6 and 13 percent (for CSIRO and MIROC, respectively). The least deleterious impacts would occur in corn, which might experience increases in pesticide expenditures between 2 and 5 percent (for CSIRO and MIROC, respectively).

Quantifying the Impacts of Drought-Tolerant Varieties

To complement the analysis of adaptive behavior of agricultural producers, we analyzed a case that represents the potential for technical change to provide additional adaptation opportunities and the implications of those opportunities on the magnitude or pattern of climate change impacts. There are many promising avenues of research on plant genetics, soil management, and inputs to production that may lead to advances that mitigate climate change impacts. We did not attempt to describe all the possible benefits of such research. We considered an illustrative case that introduces varieties for selected crops that can maintain yields under conditions of reduced precipitation, so-called “drought-tolerant” varieties, thereby reducing yield losses due to climate change for some crops in regions with low precipitation.

Quantifying the Agricultural Impact of Climate Change and the Potential for Agricultural Adaptation

Applying information about the yield impact of climate change across projected production patterns in the United States is a straightforward, if naïve, approach to estimating the potential economic impact of climate change. This method assumes an unrealistic future in which farmers' yields and returns are affected by climate change, but farmers fail to adapt their production decisions to changing climate conditions. Nevertheless, illustrating just such a scenario allows us to visualize regional and crop differences in the biophysical impacts of climate change and, as described later, to differentiate changes that take place in the agricultural sector due to biophysical impacts from those resulting from dynamic behavioral adaptation.

We first illustrate a “no adaptation” case and then provide the results of a more comprehensive assessment of climate change impacts, incorporating the impacts of farmer adaptation decisions in determining production and price patterns under changed climate conditions. A comparison between the “no adaptation” and “adaptation” cases illustrates the benefits of adaptation.

Climate Impact Analysis: No Adaptation

“No adaptation” case results were derived by putting new yield numbers into REAP but prohibiting the model from adjusting projected 2030 base-line acreage, tillage, or rotation allocations to crop production activities in response to the new climate-adjusted yields. REAP then used the baseline (i.e., no climate change) production patterns, together with the adjusted yield and environmental impact information, to calculate crop production, farmer income, price impacts, and environmental impacts under each future climate scenario. Note that, for this case, REAP eliminated farmer adaptation but retained the flexibility to adjust the livestock sector, most notably with respect to its demand for feed grains, in response to changed production and price patterns. Changes in national productivity by crop due to climate change, assuming no adaptive behavior on the part of farmers, are shown in figure 5.

These crop yield averages reflect the average impact of climate change on individual production enterprises (i.e., region/rotation/tillage combinations) weighted by the amount of crop acreage in that production enterprise, which remained constant across the climate projections in this case. Several interesting climate effects on crop productivity are evident here. Climate change impacts were most negative for corn and soybean productivity, though the least extreme scenario (ECH) produced an increase in crop yields for both. Several other crops experienced crop productivity increases for some or all of the scenarios, though the yield increases associated with the more extreme climate change scenarios (MIROC and CNR) were generally lower than those associated with the milder scenarios (ECH and CSIRO). While the impact of any temperature increase was generally negative, positive crop productivity impacts can arise both from beneficial precipitation changes (increases in water-constrained regions) or from the CO₂ fertilization effect projected when atmospheric carbon dioxide concentrations increase from 381 to 450 parts per million (ppm). For several crops, the latter positive effects

Figure 5

Changes in national average crop yield without adaptation, by climate change scenario

Percent change in yield from baseline

Cwt=Hundredweight.

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations.

outweighed the negative temperature-related losses for some or all of the projected climate projections.³

Aggregating crop productivity impacts at the national level, however, masked considerable variability in both crop productivity by region under the baseline and in regional impacts on productivity under the climate change scenarios. Disaggregating the results for corn to the level of the farm production region produced the results shown in figure 6. While the productivity results for the Corn Belt drove the pattern of national averages shown in figure 5 (because the Corn Belt accounts for 53-56 percent of U.S. corn production under these scenarios), there were regional differences in corn's response under a given climate scenario. In some regions, one or both of the milder climate change scenarios actually increased corn yields. Furthermore, corn production increased under even the extreme climate scenarios in the minor corn-producing regions of the Pacific and Mountain States and the Southern Plains region. A portion of corn production in those regions is irrigated and, therefore, less sensitive to precipitation losses from climate change but also mildly responsive to carbon dioxide fertilization gains.

In a situation with fewer interacting parts, price results might mirror productivity results; when national average crop productivity decreases (increases), the price of that crop increases (decreases) (fig. 7). Since crop and livestock markets are integrated, however, this simple dynamic did not play out for several crops and climate scenarios. Because corn prices go up in every scenario, there was always an incentive to substitute away from corn in livestock diets, which has implications for the price of other grains and feed meals. The significant corn price increases in the CNR and MIROC

³For further information on the sensitivity of EPIC's results to different elements of climate change, see Appendix C.

Figure 6

Regional differences in national average corn yield without adaptation, by farm production region and climate change scenario

Percent change from baseline

AP = Appalachia, CB = Corn Belt, DL = Delta, LA = Lake States, MN = Mountain, NP = Northern Plains, NT = Northeast, PA = Pacific, SE = Southeast, SP = Southern Plains.

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations.

Figure 7

Price changes relative to the “no climate change baseline,” by climate change scenario

Percent change in crop price from baseline

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations.

scenarios, in particular, appeared to pull up soybean, oat, sorghum, and barley prices, despite field productivity increases for some of those crops. Productivity, and therefore supply, of oats increased under every scenario, but prices also increased under all but the ECH scenario. Similarly, barley’s substitutability with the other feed grains in livestock diets led to increased

barley demand and a demand-induced price increase under the MIROC scenario that persisted despite increasing supply.

The productivity impacts illustrated in this section reflect only EPIC's yield-change calculations based on changing climate conditions, while the price impacts reflect a limited set of interacting demand-and-supply forces across agricultural sectors. Projecting potential climate change impacts on the crop sector, however, requires a more comprehensive analysis to capture how farmers may respond to biophysical impacts in their production decisions. In the following sections, we discuss how farmers might adapt to the biophysical impacts of climate change and the implications of such production adjustments for aggregate crop production, prices, agricultural acreage, and a suite of environmental indicators under changing climate regimes.

Climate Change Impacts When Farmers Adapt Crop Rotations, Tillage, and Land Use

The economic value of planting specific crops in each region changed in response to the new production conditions, since yields and costs did not change uniformly in magnitude or direction for all regions in our scenarios. In some regions, adaptation to climate change resulted in reduced planted acreage, while planted acreage in other regions increased. Regional production effects reflected both changes in yield and planted acreage. Differences in production levels, coupled with demand response to substitute crops, in turn drove changes in crop prices. The combination of changes in acreage, yield, and price influenced the degree to which farm revenues responded in a region. In this section, we report results from the REAP model that describe the economic and environmental impacts of the climate change scenarios on U.S. crop production, taking into account how farmers may adjust their crop and tillage decisions. The results are shown relative to the baseline projection (assuming no climate change) for U.S. agriculture in 2030.

Regional Shifts in Planted Acreage for Selected Crops

Each of the four climate change scenarios demonstrated a small increase in total planted acreage compared with baseline acreage levels, though there is variability in the direction of acreage change by region (fig. 8). The total acreage change, though relatively small compared with the baseline acreage, was composed of changes in the acreage planted to individual crops. The individual crops showed a much wider range across climate change scenarios, following differences in productivity and regional redistribution (table 6). Corn acres increased in all scenarios, reflecting the decline in corn yields (see fig. 5) and the need for additional acreage to compensate. Response of other crops varied by scenario, with the ECH and CSIRO scenarios showing a reduction in wheat acres that corresponds to the larger wheat-yield increases in these scenarios. Soybean acres declined in the CNR and MIROC scenarios; despite higher soybean prices, acreage decline likely reflects a decrease in the relative returns to soybean production arising from the significant yield decline under the warmer climate scenarios. The corresponding corn price increase in the MIROC scenario keeps soybean acres from declining further as soybeans are often produced in rotation with corn, particularly in the Corn Belt.

Figure 8
Changes in total regional planted acres from “no climate change baseline”

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.
 Source: USDA, Economic Research Service calculations.

Table 6
Total U.S. acreage change, by climate change scenario

Crop	ECH	CSIRO	CNR	MIROC
<i>Percent change</i>				
Total	0.6	0.6	0.2	1.0
Corn	1.7	2.8	3.0	4.2
Wheat	-1.1	-0.2	1.0	0.8
Soybeans	1.4	1.0	-2.8	-1.8
Other crops	-0.1	-1.5	-0.2	0.5

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Even though total U.S. acreage planted increased in all scenarios, no individual region showed an increase in planted acreage across all scenarios. While the variation in total acreage change was small across scenarios from a national perspective, the regions showed different degrees of response. Acreage in the Corn Belt was the least sensitive to climate change, with total changes ranging from -2.7 to 1.4 percent. In contrast, the Delta region showed a range of change between -9.8 and 5.0 percent. The sensitivity to change reflects the capacity of the region to economically shift to a different crop mix, indicating crop and production practice substitution possibilities and the larger regional yield changes relative to the national changes for each crop.

Figures 9-11 illustrate the change in planted acreage by climate change scenario for the three major field crops (corn, wheat, and soybeans). As with the national total, corn acreage in the Corn Belt was the least sensitive (-1.3 to 1.1 percent). For each of the other regions, at least one scenario demonstrated at least a 10-percent change in acreage from the baseline. The Southern Plains increased corn acreage in all scenarios, with the Delta and Pacific regions reducing corn acres in all scenarios.

Figure 9
Changes in regional planted corn acres from “no climate change baseline”

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.
 Source: USDA, Economic Research Service calculations.

Figure 10

Changes in regional planted wheat acres from “no climate change baseline”

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations.

The Corn Belt also showed the smallest change in wheat acres (0.4 to 2.4 percent). The Lake States and Corn Belt increased wheat acres in all scenarios; all other regions showed both increases and decreases across scenarios. Climate change induced soybean acres to move into the Northern and Southern Plains (see fig. 11). Once again, the Corn Belt showed the smallest range in soybean acreage change (-5.6 to 0.8 percent); by contrast, change in soybean acres in the Delta region ranged from -19.2 to 11.4 percent.

Corn acreage increased in the combined Northern regions (Corn Belt, Lake States, Northern Plains, and Northeast) in all scenarios, while corn acres in the combined Southern regions (Appalachian, Delta, Southeast, and Southern Plains) declined in the ECH and CSIRO scenarios and increased in the CNR and MIROC scenarios. The same relationship holds for wheat acres. Taken as a group, the Southern regions were more sensitive to weather-induced yield change than were the Northern regions. This trend was indicative of the larger range in EPIC-derived yields in the Southern regions under the climate change scenarios.

Figure 11

Changes in regional planted soybean acres from “no climate change baseline”

Note: Soybeans not cultivated in Pacific and Mountain regions.

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations.

Changes in Crop Prices and Regional Farm Revenue

Crop price changes were highly scenario-specific, although there were some features common to the results. The CNR and MIROC scenarios led to higher corn and soybean prices than under the baseline, whereas the CSIRO and ECH scenarios generally resulted in lower prices (except for a small price increase in soybeans in CSIRO) (table 7). Corn and soybean yields were lower in the CNR and MIROC scenarios in all regions, while the CSIRO and ECH scenarios showed increased yields in some regions. Wheat prices declined in all scenarios. Wheat and corn prices were generally less sensitive to changes in climate than soybean prices, mainly as a result of smaller projected sensitivity of wheat yields to temperature and rainfall and the poor substitutability between wheat and corn in the diets of livestock within REAP. While REAP’s existing diets reflect feed combinations that historically have been used in the livestock sector, significant feed price changes may induce changes within the livestock sector in popular diet combinations. Our analysis did not capture that dynamic.

Climate change reduced returns to corn in all scenarios relative to the baseline, although less so in the higher temperature/precipitation change scenario, where price increases helped support producer revenue (table 8). Because of the positive effect of climate change on cotton yields, cotton

returns increased significantly in all scenarios. Returns to other crops varied by scenario. Returns in the Corn Belt mimicked changes in precipitation and temperature, whereas returns in other regions did not necessarily mirror the same trend (table 9). The more extreme scenarios generally led to lower returns, but not for all regions. Nationally, the milder scenarios led to an increase in returns, whereas the more extreme scenarios led to a decline in returns. From the perspective of crop returns, the Southern Plains was the most robust, even though the regional change in acreage was large compared with other regions. This result reflects the relatively higher returns to cotton production under climate change, and the ability of farmers in the Southern Plains region to reallocate production resources to minimize the regional impact on profitability.

Table 7

Change in crop prices with adaptation relative to the “no climate change” baseline

Crop	ECH	CSIRO	CNR	MIROC
<i>Percent change</i>				
Corn	-2.2	-2.1	3.7	6.0
Wheat	-1.6	-5.9	-0.8	-1.0
Soybeans	-3.5	0.3	7.6	22.1

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Table 8

Change in annual returns to crop production from “no climate change baseline”

Crop	ECH	CSIRO	CNR	MIROC
<i>\$ Million</i>				
Corn	-742	-839	-33	-223
Wheat	-10	332	-265	-456
Soybeans	1,361	-180	-2,772	-3412
Cotton	1,135	1,081	1,474	1,266

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Table 9

Change in annual returns to crop production from “no climate change baseline”

Region	ECH	CSIRO	CNR	MIROC
<i>\$ Million</i>				
Corn Belt	-1,114	-2,165	-2,112	-4,053
Delta	904	167	-521	-146
Lake States	41	902	1,001	-37
Northern Plains	1,256	1,671	-914	255
Southern Plains	418	322	7	681
U.S. total	3,619	2,165	-332	-1,465

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Changes in Environmental Outcomes

Shifts in crops and production practices as a result of climate-induced changes in crop yields will have an influence on the environmental impacts associated with agricultural production. Changes in total acreage, in tandem with the redistribution of crop rotation/tillage practices, will affect regional soil and water quality through nutrient loss and soil erosion. Increases in environmental impacts caused by acreage expansion in one region may not be offset by acreage decreases in other regions, resulting in aggregate national increases in water quality impacts. Table 10 depicts the changes in nitrogen loss (leaching and runoff to ground and surface water) and rainfall-related soil erosion (specifically sheet and rill erosion) compared with the baseline, along with the change in U.S. planted acres. Total nitrogen lost to water (measured as nitrogen deep percolation and runoff at the field edge) increased in all scenarios, which follows from the general increase in acreage nationally. Increased nitrogen loss was not uniform across the country, with the MIROC scenario resulting in the most widespread changes (fig. 12). The Corn Belt, Northern Plains, and Southeast regions all increased nitrogen lost to water in all scenarios.

Soil lost to water erosion also increased in all but the ECH scenario, and this measure was generally less sensitive to climate scenario than the measure for nitrogen loss to water (table 10). Changes in the Corn Belt and Northern Plains dominated the national erosion change, as a result of shifts in crop rotation and tillage and an increase in planted acres (fig. 13).

How Farmer Adaptation Affects Measured Impacts of Climate Change

In the no-adaptation case, farmers did not adjust production decisions in response to changes in expected yields across crops. As a result, regions may overplant crops whose returns have declined relative to other potential crops, while underplanting crops that have become relatively more profitable. In the adaptation case, in contrast, farmers adjusted their land use, crop rotations, and tillage regimes in response both to changes in climate and adjustments in market conditions resulting from climate effects. Below, we examine how incorporating farmer adaptive response to climate change affects measured economic and environmental impacts.

Table 10

Change in total U.S. planted acreage and select environmental measures with adaptation relative to the “no climate change” baseline

Environmental measure	ECH	CSIRO	CNR	MIROC
	Percent			
Total acreage	0.6	0.6	0.2	1.0
Nitrogen loss to water	1.4	1.5	2.1	5.0
Sheet and rill erosion	-0.9	0.6	0.9	1.2

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Figure 12

Regional change in total nitrogen loss to water from “no climate change baseline”

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations.

Regional Shifts in Planted Acreage for Selected Crops

Adaptation includes the various strategies farmers use to adjust their production decisions (e.g., shifts in crops or crop rotations) in response to absolute and relative changes in yield and management costs. Figure 14 illustrates the redistribution of planted acreage that resulted from adaptation for this analysis. In most region/scenario combinations, adaptation led to an increase in corn acreage (fig. 15). Failing to adapt (no adaptation case) restricted the supply of corn, which led to higher consumer prices. When farmers are permitted to adapt, they respond to high corn prices by changing production in favor of corn. Thus, farmer behavior can moderate climate change impacts on production by diverting productive resources to crops whose loss most negatively impacts other farm-sector stakeholders.

Changes in Crop Prices and Regional Farm Revenue

Although national welfare (the sum of economic benefits to consumers plus benefits to producers) increases when farmers have the flexibility to adapt, the benefits of adaptation differ across regions and between consumers and

Figure 13

Regional change in sheet and rill erosion from “no climate change baseline”

Table 11
Price difference from “no adaptation case” to “adaptation case,”
by crop

	ECH	CSIRO	CNR	MIROC
Percent change				
Corn	-3.2	-3.2	-3.6	-3.9
Sorghum	-1.1	-1.8	-1.1	-1.2
Barley	2.1	2.6	-1.6	-2.9
Oats	-0.6	-0.4	-8.7	-7.1
Wheat	0.2	-3.0	-0.7	-0.6
Rice	0.0	-0.3	-0.4	-0.3
Soybeans	-0.6	-0.1	0.7	1.9
Cotton	-8.2	-5.5	-5.7	-9.0
Silage	0.9	0.7	0.5	0.3
Hay	0.3	0.6	0.5	0.8

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Figure 14

Regional change in total acreage, from “no adaptation case” to “adaptation case”

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Note: Figure 14 is identical to figure 8 because production acreage is fixed in the “no adaptation case” at the level of production acreage in the “no climate change baseline”.

Source: USDA, Economic Research Service calculations.

producers. Consumers generally benefited from the process of adaptation. Prices (a measure of consumer benefit) were generally lower if all farmers adapted (table 11), which moderated consumer expenditure impacts of climate change and increased consumer welfare measures relative to the case where adaptation was not permitted.

Producers were not necessarily better off as a result of adaptation, however. For some regions, nationwide adaptation led to lower returns relative to not adapting by, for instance, driving down the price of a major crop in that region. Returns to crop production in the Corn Belt did not benefit from nationwide adaptation (table 12); this is the consequence of smaller acreage changes in the Corn Belt relative to other regions combined with the lower prices that resulted from adaptation.

Changes in Environmental Outcomes

Table 13 illustrates the consequences of adaptation with respect to selected environmental measures when compared with the fixed-acreage, no adaptation case. The percent increase in nitrogen loss was greater than the change

Figure 15

Regional change in corn acreage from “no adaptation case” to “adaptation case”

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Note: Figure 15 is identical to Figure 9 because corn acreage is fixed in the “no adaptation case” at the level of corn acreage in the “no climate change baseline”.

Source: USDA, Economic Research Service calculations.

Table 12

Direction of change in returns to crop production from “no adaption case” to “adaptation case,” by region

	ECH	CSIRO	CNR	MIROC
Appalachian	↑	↑	↑	↑
Corn Belt	↓	↓	↓	↓
Delta	↑	↑	↑	↑
Lake States	↑	↑	↓	↓
Mountain	↑	↑	↑	↑
Northern Plains	↑	↑	↑	↑
Northeast	↓	↓	↓	↓
Pacific	↑	↑	↑	↑
Southeast	↑	↑	↑	↑
Southern Plains	↑	↑	↑	↑
U.S. total	↑	↑	↓	↓

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

in acreage in all scenarios, suggesting that climate change adaptation induced crop production to shift into areas and production practices with greater water-quality impacts. Climate change-induced crop reallocation, however, significantly reduced nitrogen loss in the Pacific and Mountain regions while resulting in substantial increases in the Southern Plains and mixed impacts in other regions (fig. 16). Even the more extreme scenarios showed regions where nitrogen loss declined, despite a 1.8- to 3.6-percent increase in losses nationally (table 13). Regional impacts on soil loss to water erosion followed a similar pattern to that of nitrogen loss (fig. 17), although the national change in the erosion measure was similar to, not greater than, the national acreage change.

Impact of Changes in Pest Prevalence on Crop Production Under Climate Change

Our initial REAP analyses of the climate change scenarios considered only the crop-yield differences resulting from changes in prevailing precipitation, temperature, and carbon dioxide concentration. Additional cost and yield implications of changes in pest prevalence (weed, insect, and plant disease) are likely to occur with climate change. Costs and yields were modified for each climate change

Figure 16
Regional change in nitrogen loss to water from “no adaptation case” to “adaptation case”

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.
Source: USDA, Economic Research Service calculations.

scenario, and the results were compared with the climate change scenario results for the case where additional pest impacts were not considered, allowing us to examine the additional contribution of climate change-induced pest pressure to potential changes in agricultural production and commodity markets.

Table 13

Change in total U.S. planted acreage and select environmental measures from “no adaptation case” to “adaptation case”

Environmental measure	ECH	CSIRO	CNR	MIROC
<i>Million acres</i>				
Total acreage	1.8	1.8	0.8	3.2
<i>Percent change</i>				
Total acreage	0.6	0.6	0.2	1.0
Nitrogen loss to water	4.3	4.0	1.8	3.6
Sheet and rill erosion	5.4	4.3	4.3	1.4

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Figure 17

Regional change in sheet and rill erosion from “no adaptation case” to “adaptation case”

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations.

An increase in pest prevalence uniformly reduced yields across all crops (except cotton and, to a lesser extent, rice) in all regions. Thus, while the climate change scenarios described in the previous section showed some regions with yield increases for some crops, increased pest presence eliminated some, if not all, of those yield increases and reduced even further the projected yields that declined under climate change. The geographic and crop-specific nature of pest impacts led to differing implications for regional crop production.

Regional Shifts in Planted Acreage for Selected Crops

Table 14 shows the national outcome on total acreage for select crops when farmers account for yield and cost effects from increased pests. All scenarios showed an increase in total acreage planted to compensate for the yield reductions from increased pests. The two higher temperature-change scenarios—CNR and MIROC—exhibited the largest acreage impact. For soybeans, there was a general decrease in acreage, except in the highest temperature-change scenario. Regional acreage changes varied for select crops. Pest changes reduced acreage for all crops in the Corn Belt, whereas other regions had a mixed response (fig. 18). Corn acreage increased in all regions except the Corn Belt, where acreage remained fairly constant (fig. 19). Wheat acreage changes varied regionally, with acreage declining generally in the North Central United States and increasing in the West and Southwest (fig. 20). Wheat became more important in the Southeast in the CNR and MIROC scenarios. Soybeans shifted away from the Corn Belt and into the Northern and Southern Plains in all scenarios (fig. 21).

Changes in Crop Prices and Regional Farm Revenue

Additional pest pressure resulted in price increases compared with the “adaptation case” without pest effects for all crops (except cotton) (table 15). Corn and soybean price changes were greater than the changes in wheat price in all scenarios. Regional and national returns to crop production were also affected (table 16). Additional increased pest pressure reduced returns from the adaptation-only level in all scenarios. The magnitude of the changes in returns associated with additional pests increased as the scenarios became more extreme, indicating that additional pests compound the temperature- and precipitation-induced yield changes as climate diverged from current conditions. In the Corn Belt, the impact of additional pests was insensitive

Table 14
National change in planted acres from 2030 climate change scenarios as a result of changes in pest distribution

Crop	ECH	CSIRO	CNR	MIROC
<i>Million acres</i>				
Total	2.1	0.9	4.2	5.6
Corn	1.0	0.6	2.2	2.0
Wheat	1.3	0.7	1.7	2.4
Soybeans	-0.9	-1.2	-0.2	0.4
Cotton	0.1	0.0	0.0	0.1

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Figure 18
Regional change in total planted acres from 2030 climate change scenarios as a result of changes in pest distribution

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.
 Source: USDA, Economic Research Service calculations.

Table 15
Price difference from 2030 climate change scenarios as a result of changes in pest distribution

Crop	ECH	CSIRO	CNR	MIROC
<i>Percent change</i>				
Corn	4.4	3.9	4.3	6.2
Sorghum	1.2	1.6	0.3	0.7
Barley	4.5	3.2	4.7	7.6
Oats	6.7	4.1	9.5	10.2
Wheat	1.5	1.7	1.7	2.6
Rice	0.0	0.0	0.0	0.0
Soybeans	3.1	2.6	5.1	9.1
Cotton	-0.8	-0.6	-0.9	-1.4
Silage	1.0	0.8	1.5	2.3
Hay	2.2	1.7	2.6	3.7

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Figure 19

Regional change in planted corn acres from 2030 climate change scenarios as a result of changes in pest distribution

Note: The Pacific region's acreage change was greater than 10 percent in ECH, CSIRO, and MIROC. See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models. Source: USDA, Economic Research Service calculations.

Table 16
Change in annual returns to crop production from 2030 climate change scenarios as a result of changes in pest distribution

Region	ECH	CSIRO	CNR	MIROC
\$ Million				
Corn Belt	-462	-444	-568	-510
Delta	-61	-38	-10	52
Lake States	-322	-218	-502	-618
Northern Plains	-416	-280	-760	-997
Southern Plains	-114	-117	-181	-184
U.S. total	-1,903	-1,471	-2,604	-3,007

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Figure 20

Regional change in planted wheat acres from 2030 climate change scenarios as a result of changes in pest distribution

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations.

to the climate conditions because of the relatively smaller temperature range across climate scenarios compared with other regions.

Changes in Environmental Outcomes

Increased pest pressure on yields generally resulted in an increase in nitrogen losses to water (fig. 22), a consequence of the increase in planted acreage required to make up for the lower yields. Soil lost to water erosion also increased in response to the increased acreage planted (fig. 23). The impact of pests on soil erosion is generally less than on nitrogen loss, evident by the range of change over the scenarios. Nitrogen loss ranged from -15 to 15 percent over the climate change scenarios, whereas the range of soil erosion impacts was less (-3 to 10 percent), with the Northern Plains and Mountain regions showing an increase in all scenarios.

Figure 21

Regional change in planted soybean acres from 2030 climate change scenarios as a result of change in pest distribution

Note: Soybeans not cultivated in the Pacific and Mountain regions.
 See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.
 Source: USDA, Economic Research Service calculations.

Figure 22

Regional change in nitrogen lost to water from 2030 climate change scenarios as a result of changes in pest distribution

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.
Source: USDA, Economic Research Service calculations.

Figure 23

Regional change in sheet and rill erosion from 2030 climate scenarios as a result of changes in pest distribution

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.
Source: USDA, Economic Research Service calculations.

Lessening the Impact of Climate Change Through Technical Change: Targeting Crop Breeding for Adaptation to Climate Stress

We previously considered the regional effects of temperature and precipitation changes on crop yields, and then examined how potential shifts in the geographic distribution of pests and diseases might further impact crop yields and farmer adaptation decisions. We now consider how technical changes provide additional adaptation opportunities and the implications of those opportunities on land use, crop production, commodity markets, and environmental quality. We first analyzed current crop genetic research addressing particular production constraints accompanying climate change, particularly heat, drought, and pests/diseases. We then designed and analyzed a case exploring the potential effects of new drought-tolerant crop varieties on the likely impacts of climate change.

Adoption of Plant Varieties With Greater Tolerance to Climate Change-Related Stresses

Plant genetic resources might be used as an adaptation strategy in a number of ways:

- One crop species may simply replace another in response to changing growing conditions within a region. Existing crop varieties may also be used in new locations as conditions change.
- New varieties can be developed by plant breeders to improve tolerance to heat, drought, and pests/diseases, using classical breeding techniques, molecular methods (e.g., genetic engineering), or some combination of the two. These new plant varieties may also respond better to other adaptive management technologies, such as those related to irrigation and tillage (Rosenzweig and Hillel, 1995).

Some changes in genetic resource use may come from individual farmers using existing resources and technologies. Development of new crop varieties and many other technological advances will require the involvement of the formal agricultural research system. Various institutions, both public and private (including gene banks and plant breeding research stations), are pioneering the use of existing varieties in new locations, developing new varieties, and testing the efficacy of new varieties in conjunction with other adaptive strategies.

Potential Adaptive Benefits From Genetic Resource Use

Heat Tolerance

Tolerance to temperatures (extreme heat or cold) is a complex trait governed by multiple genes that involve a number of physiological traits and metabolic pathways. At present, few public and private sector research resources have been devoted to developing heat-tolerant crop cultivars. Hatfield et al. (2008) discussed the role of “failure temperatures” above which yields for different crops fall to zero. In a recent review, Wahid et al. (2007) concluded that “there are a few examples of plants with improved heat tolerance through the

use of traditional breeding protocols,” but that “the success of [the] genetic transformation approach has been thus far limited. The latter is due to limited knowledge and availability of genes with known effects on plant heat-stress tolerance, though these may not be insurmountable in the future.”

Advances are being made in heat-tolerance research, however. A number of studies supported genetic variability for heat tolerance in important field crops and suggested particular selection criteria that might enhance breeding for heat tolerance (e.g., Maestri et al., 2002; Martineau et al., 1979; Radin et al., 1994; ur Rahman et al., 2004; Reynolds et al., 1994). Other researchers have focused on targeted changes in plant metabolism, such as improved photosynthetic performance at higher temperatures (Parry et al., 2011). Nevertheless, the combination of long research lags and limited research investment suggests that the probability of substantial development of commercially viable heat-tolerant crops by 2030 is relatively low.

Carbon Dioxide Responsiveness

Evidence suggests that a portion of crop yield gains in the past may be attributed to yield response to increasing atmospheric CO₂ (McGrath and Lobell, 2011). Plant breeders may have inadvertently selected for response to CO₂ fertilization, which raised the question of whether breeders might wish to select for greater response deliberately. Ziska et al. (Ziska and Blumenthal, 2007; Ziska, 2008; Ziska and McClung, 2008) suggested that:

- In some cases, newer varieties did not appear to be more responsive to atmospheric carbon dioxide fertilization than older varieties of the same crop;
- Some genetic variability for this trait—response to atmospheric CO₂—did exist within cultivated species or wild relatives; and
- The relative ease of incorporating such a trait may vary by crop.

Pests and Diseases

The first generation of genetically engineered (GE) crops for commercial use was bred specifically to control for pests (particularly insects) and disease as well as to improve herbicide tolerance. Herbicide-tolerant crops introduce a genetic element into weed control by making the crop tolerant of herbicides, such as glyphosate, which are then used to control weeds. As a result, using GE crops has reduced pesticide use (Fernandez-Cornejo and Caswell, 2006). Genetic engineering for disease resistance has also been explored (Mourges et al., 1998; Punja, 2001) but, to date, a disease-resistant GE crop has not been commercialized.

A great deal of money has been invested in both public and private research for disease/pest-tolerant or resistant crops. As a result, creating an adaptive genetic resource may reduce some of the negative impacts of the disease/pest landscape likely to accompany climate change. For example, different forms of herbicide tolerance incorporated into crop varieties may be part of the management response if weeds become a more significant problem (Duke and Powles, 2008).

New threats may also emerge with climate change. Weeds may compete increasingly with crops if their growth is more responsive to CO₂ fertilization than the crop, or diseases may expand their ranges as alternate hosts also expand their range (e.g., soybean rust). Violent storms may have aided the spread of rust spores, first from Asia to Africa, then to South America, and finally to the United States. The farm sector's first line of defense typically has been fungicides, with potentially large increases in fungicide use tempered by changes in cropping patterns (Livingston et al., 2004).⁴ USDA coordinated early soybean rust surveillance and warning networks that also reduced the impact of the disease (Roberts et al., 2006).

Drought Tolerance

Drought tolerance, like heat tolerance, is a complex genetic trait. Over the past half century or more, "selection for high yield in stress-free conditions has, to a certain extent, indirectly improved yield in many water-limited conditions" (Cattivelli et al., 2008). U.S.-grown corn is one example of this phenomenon, where more than 50 years of selection in multi-environment trials has increased grain yield under drought conditions (Campos et al., 2004). Selection for yield stability may have played some role in improving performance under stress conditions. Attempts to improve drought tolerance directly through conventional plant breeding date back to the 1970s, if not earlier. More recently, molecular and genomic analyses identified gene networks that may be important in drought stress (Shinozaki and Yamaguchi-Shinozaki, 2007). In applied breeding, seed companies have increased their efforts to develop drought-tolerant crops, particularly corn, and several companies have placed new, conventionally bred drought-tolerant hybrids on the market. Large seed companies are also using transgenic⁵ means to develop drought-tolerant crops.

The first transgenic drought-tolerant corn hybrids received regulatory approval in 2011, which will be followed by extensive field testing in late 2012; these hybrids may be marketed as early as 2013, pending regulatory approvals.⁶ These hybrids may have been altered with genes from bacteria that regulate metabolic pathways for cold shock response; in plants, they may regulate pathways for other stresses, such as drought (Castiglioni et al., 2008). Potential yields from transgenic drought-tolerant corn were shown to increase by approximately 10 percent under managed stress environmental testing, and up to 15 percent under dryland conditions in environments where the average control yield was about 78 bushels/acre (Castiglioni et al., 2008).

Alternatively, Edmeades (2008) hypothesized that drought-tolerant corn had the potential to improve every 5 years over a 20-year period based on a combination of conventional selection, marker-aided selection, and a new transgenic event. Edmeades calculated yield improvement above a base yield of 48 bushels/acre to reflect conditions in some developing countries. These research assumptions projected a tripling of yield over 20 years. Such estimates might be used as the rough basis for lower and upper bounds on achievable improvements in drought tolerance over a given period.

Current Research Emphases

Several of the physical factors that accompany climate change can influence crop yields, and both genetic and management adaptations to these physical

⁴Historically, fungicide use in soybean production has been very low.

⁵A gene or genetic material transferred naturally or via genetic engineering from one organism to another.

⁶See http://www.aphis.usda.gov/newsroom/2011/05/ea_corn.shtml and <http://www.monsanto.com/products/Pages/drought-tolerant-corn.aspx>.

factors have the potential to change yield response. The impacts of increasing atmospheric CO₂, increasing temperatures, and variable precipitation on pest/disease incidence and temperature- and water stress-related yield impacts are both complex and uncertain. A number of public agricultural research projects have addressed crop tolerance to heat, drought, and pest/disease infestations, as reported in USDA's Current Research Information System (CRIS) (table 17).

Based on project counts from USDA's CRIS, relatively little research has been done on crop adaptation to higher temperatures.⁷ A great deal of research has focused on pests and diseases and the potential for host plant resistance.

Private-sector companies have invested a great deal of resources into GE crops, particularly for insect resistance or herbicide tolerance (HT). Genetically engineered HT, particularly to Roundup, is widely used in U.S.-grown soybeans, corn, and cotton, and the bacterium *Bacillus thuringiensis*, or Bt, is used for insect resistance in corn and cotton. Genetically engineering crops for drought tolerance is still at various phases of research development, depending on the crop, but several companies do have conventionally bred drought-tolerant corn varieties on the market. The level of heat tolerance research by private seed-biotechnology companies is unknown, since at present they do not include this trait in reported research pipelines.

Impact of Adopting Drought-Tolerant Crop Varieties

Drought-tolerant crop varieties are a good example of an adaptive genetic response to climate change with likely impact by 2030. We assumed patterns for the potential impact of drought-tolerant varieties for four major crops on nonirrigated land:

- Corn:** Drought-tolerant varieties increased yields on nonirrigated land by 15 percent in all areas where annual precipitation was less than 700 millimeters (mm) and nonirrigated yields in 2010 were less than 80 bushels per acre (bu/acre).

Table 17
U.S. public sector agricultural research projects, by crop stressor, as reported in 2010*

Stress	Projects that focus on crop stressor	Projects that partially focus on crop stressor
	<i>Number (percent)</i>	
Heat	17 (3.7)	59 (5.8)
Drought	49 (10.7)	149 (14.7)
Pests and diseases	390 (85.5)	808 (79.5)
Total	456 (100)	1,016 (100)

*Ninety-five percent of these projects were initiated during 2001-10.
 Source: USDA, Current Research Information System (CRIS) database, November 2010.

⁷Total financial resources devoted to each project were not, in general, publicly available.

- **Wheat:** Drought-tolerant varieties increased yields on nonirrigated land by 10 percent in all areas where annual precipitation was less than 625 mm and nonirrigated yields in 2010 were less than 50 bu/acre.
- **Soybeans:** Drought-tolerant varieties increased yields on nonirrigated land by 10 percent in all areas where annual precipitation was less than 700 mm.
- **Cotton:** Drought-tolerant varieties increased yields on nonirrigated land by 10 percent in all areas where annual precipitation was less than 900 mm.

The relative yield increases for wheat, soybeans, and cotton were assumed to be less than that for corn, since a drought-tolerant trait will soon be deployed in corn. Drought-tolerant varieties of wheat, soybeans, and cotton are in earlier stages of development. Also, no advances in drought tolerance were assumed for crops other than these four. Yield increase assumptions for drought-tolerant corn varieties were based on current experimental results reported by Castiglioni et al. (2008). In the absence of similar experimental results for wheat, soybeans, and cotton, lower percentage yield increases were assumed for REAP regions where lower yields and precipitation totals were both operative. Figure 24 shows the regions where the drought-tolerant varieties are planted in the scenario analyzed.

Figure 24

Regions assumed to adopt drought-tolerant corn, wheat, soybeans, and cotton

Note: The crops and regions using drought-tolerant varieties vary by scenario.

Source: USDA, Economic Research Service calculations.

Several factors suggest that these yield increments might be conservative if improved drought tolerance is commercialized within the next few years and then widely adopted. First, additional research might enhance potential yields beyond 10-15 percent in low-moisture environments. Second, depending on how well drought-tolerant varieties perform relative to other varieties under better moisture conditions, some drought-tolerance traits might be introduced into varieties for higher rainfall areas, as insurance against less favorable production conditions in some years. The threshold assumption currently defining water-stressed regions concerns annual precipitation; a more sophisticated analysis would consider precipitation and soil moisture conditions during crucial periods of the growing cycle.

Our analysis compared variables of interest, such as crop area and crop prices, between scenarios in which farmers had adapted crops, rotations, and production practices with the availability of drought-tolerant varieties and scenarios in which they had made these adaptations without drought-tolerant varieties.⁸ Tables 18 and 19 indicate aggregate acreage and price changes within each climate change scenario. In each instance, estimates from the introduction of drought-tolerant varieties were compared with results without drought-tolerant varieties within the context of that particular climate change

⁸In other words, the starting points for the analysis were the adaptation scenarios from earlier sections of the report. Drought-tolerant varieties were added as an additional adaptation option for farmers.

Table 18

U.S. acreage change if drought-tolerant varieties are available relative to adaptation case with current varieties

Crop	ECH	CSIRO	CNR	MIROC
<i>Million acres</i>				
Total	-0.5	-1.3	-1.2	-1.4
Corn	0.5	0.1	0.2	0.1
Wheat	-1.1	-1.3	-1.3	-1.1
Soybeans	0.4	0.2	0.1	-0.2
Cotton	0.0	0.1	0.0	0.0

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Table 19

Crop price impacts of drought-tolerant varieties compared with baseline prices

Crop	Adaptation	ECH	CSIRO	CNR	MIROC
<i>Percent change</i>					
Corn	Normal	-2.2	-2.1	3.7	6.1
	DT	-3.0	-2.9	2.8	5.3
Soybeans	Normal	-3.5	0.3	7.6	22.1
	DT	-3.6	0.1	7.4	21.9
Wheat	Normal	-1.5	-5.9	-0.8	-1.0
	DT	-4.3	-9.4	-4.0	-4.3
Cotton	Normal	-19.8	-14.6	-17.8	-22.8
	DT	-20.6	-15.6	-18.7	-23.6

DT = Drought-tolerant corn, soybeans, wheat, and cotton planted in some regions.

Normal = No drought-tolerant varieties planted.

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models

scenario. Including drought-tolerant varieties reduced total planted acreage across all the scenarios. Corn acreage increased in all scenarios, as did soybeans, except for in the MIROC scenario; wheat acreage decreased in all scenarios due to improved productivity in major wheat-producing regions (table 18). The change in acreage was less sensitive to the climate scenario than in other cases examined.

The introduction of drought-tolerant varieties reduced prices for corn, soybeans, wheat and cotton in all scenarios, with wheat showing the biggest reduction (table 19). Prices changed for these crops in the same direction from the baseline in all scenarios; prices that went up in the adaptation case went up less with drought tolerance and prices that declined were further reduced with the adoption of drought-tolerant varieties. Price changes for other crops varied by crop and scenario.

Net returns for all scenarios (table 20) increased for the United States as a whole compared with the adaptation scenario without drought-tolerant varieties. The Northern and Southern Plains regions benefited from the introduction of drought-tolerant varieties, but returns were reduced in the Corn Belt, which does not plant drought-tolerant varieties under any scenario. The change in returns was consistent across climate change scenarios, indicating that drought tolerance could be beneficial under a wide range of adverse climate changes.

Impacts on acreage from the introduction of drought-tolerant varieties were more distinct at the regional level than at the national aggregate level. The only region that showed an increase in total acreage under all scenarios was the Northern Plains. Acreage moved from other western regions; regions east of the Mississippi River showed little change in acreage (fig. 25).

The impacts of introducing drought-tolerant varieties were most apparent when examining crop production by region. With a few exceptions, corn, wheat, and soybean production shifted from the eastern half of the country to the drier Plains and Mountain regions. These changes were seen only at the margin; the Corn Belt region, for example, remained the largest producer of

Table 20
Change in annual net returns to crop production from 2030 climate change scenarios with introduction of drought-tolerant varieties

Region	ECH	CSIRO	CNR	MIROC
\$ Million				
Corn Belt	-369	-374	-391	-327
Delta	-36	-31	-27	-31
Lake States	-86	-91	-97	-84
Northern Plains	479	638	610	684
Southern Plains	273	307	275	296
Other U.S. regions	146	89	158	115
U.S. total	372	506	487	623

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models. Results describe the change relative to the 2030 climate change scenario without the addition of drought-tolerant varieties.

Figure 25

Regional change in total planted acres from 2030 climate change scenarios with introduction of drought-tolerant varieties

Results describe the change relative to the 2030 climate change scenario without the addition of drought tolerant varieties. See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations.

corn and soybeans. The addition of drought-tolerant varieties does not lead to significant corn production redistribution, even under the more extreme scenarios (fig. 26); corn is not a major crop in the regions where drought-tolerant varieties are planted, and the yield benefit in those regions is not sufficient to allow corn to compete significantly with other crops, or other regions. Wheat production increased in the Northern Plains with the introduction of drought-tolerant varieties, but decreased in the Southern Plains as a result of declining prices (fig. 27). Soybean production increased slightly in the Northern Plains and Southern Plains with the introduction of drought-tolerant varieties (fig. 28).

One environmental variable—nitrogen loss to water—showed little change at the national level. Nitrogen loss improved in the Northern Plains, even as total acreage increased (fig. 29). Production in the region moved from continuous to multi-crop rotations, which tend to have lower nutrient loss. Slightly reduced acreage led to a decrease in nitrogen lost to water in the eastern regions. Acreage in the Southern Plains increased in crop rotations that had higher nitrogen losses, leading to increased overall nitrogen losses for the region. The pattern of regional change in sheet and rill erosion is similar to that of nitrogen losses to water.

Figure 26

Regional change in corn production from 2030 climate change scenarios with introduction of drought-tolerant varieties

Results describe the change relative to the 2030 climate change scenario without the addition of drought tolerant varieties.

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations.

Figure 27

Regional change in wheat production from 2030 climate change scenarios with introduction of drought-tolerant varieties

Results describe the change relative to the 2030 climate change scenario without the addition of drought tolerant varieties. See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models. Source: USDA, Economic Research Service calculations.

Figure 28

Regional change in soybean production from 2030 climate change scenarios with introduction of drought-tolerant varieties

Note: Soybeans not cultivated in the Pacific and Mountain regions.
Results describe the change relative to the 2030 climate change scenario without the addition of drought tolerant varieties.
See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.
Source: USDA, Economic Research Service calculations.

Figure 29
Regional change in nitrogen loss to water from 2030 climate change scenarios with introduction of drought-tolerant varieties

■ -2.2% to -1.5% ■ -1.5% to -0.9% ■ -0.9% to -0.2% ■ -0.2% to 0.4% ■ 0.4% to 1.1% ■ 1.1% to 1.8% ■ 1.8% to 2.4%
 Results describe the change relative to the 2030 climate change scenario without the addition of drought tolerant varieties.
 See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.
 Source: USDA, Economic Research Service calculations.

Conclusions

The biophysical impacts from climate change on crop growth are both complex and uncertain. Although temperature increases associated with climate change are widely expected to lower crop yields, shifting regional precipitation patterns may either increase or decrease yields. There has also been considerable debate over the likely impacts of increasing atmospheric CO₂ concentrations on plant growth. Our analysis suggests that the possible negative implications of climate-induced yield effects on crop prices, farm revenue, and food supply can be mitigated somewhat by farmers' ability to adapt to changing climatic conditions through crop production decisions, technologies used, and the regional allocation of land. There also may be increased environmental impacts associated with expansion of cropland. Changes in pest distribution exacerbate production and revenue losses.

Nationally, acreage in crop production was fairly robust to climate change in the sense that aggregate acreage changes across scenarios (compared with the baseline projection) tended to be relatively small (less than 1 percent), whereas the range for individual regions was typically greater than the national range. This finding reflects the flexibility of the U.S. farm sector to respond to changing resource and market forces, resulting in production reallocations that minimize the aggregate disturbance to commodity supply and demand. In this analysis, available adaptation strategies included changing crops, crop rotations, and tillage types, as well as expansion or contraction of crop production acreage. Without additional pest impacts, mild climate change led to a reduction in crop prices; more extreme climate change resulted in price increases in some crops, most notably for soybeans and corn. Adaptation led to an increase in crop production, in general, though even with adaptation corn and soybean production declined under the more extreme climate projections.

Total acreage in U.S. field crop production increased in all climate change scenarios, although variations in yield changes across scenarios did not necessarily translate into acreage expansion in all regions. Corn and cotton were the only crops that increased acreage in all scenarios. Higher temperature and atmospheric carbon dioxide concentrations raised cotton yields considerably in every region where cotton was grown. The increase in supply, together with relatively elastic demand, caused cotton prices to drop precipitously, facilitating enough increase in demand to actually expand acreage even though existing acreage was already more productive. The corn yield effect was small relative to other crops across the scenarios, but high demand for corn raised corn acreage to compensate for lost yields. Corn acreage increased relatively more in regions where corn was not the predominant crop. Crop distributions in the Corn Belt and the Northern regions, in general, were less sensitive to climate change than in the Southern regions.

Price and production changes contributed to shifting regional returns to crop production. Returns in the Corn Belt, where much of the Nation's field crop production is concentrated, declined under all climate change scenarios in proportion to the severity of projected change. Farm returns nationally, however, increased under the two milder climate projections, with losses in the Corn Belt compensated for by increased returns in other regions as a

result of changes in price and production patterns. Our findings suggest that, for regions outside the Corn Belt, changes in crop returns did not necessarily correspond to the general magnitude of the temperature and precipitation change of the scenario. Changes in relative productivity both across crops and within crops across regions, together with the resulting market-mediated price impacts, appeared to be the primary determinant of how any given region fared under changing climate conditions.

Extending the climate impact analysis to consider projected increases in pest-related yield and associated control costs showed that such impacts led to expanded planted acreage in all crops except for soybeans. Increases in pest prevalence reduced yields across all crops (except cotton and rice), but the differentiation of impacts by region and crop led to regional differences in acreage response. Corn production expanded in all regions except the Corn Belt, while wheat became relatively more important in the Southeast and soybeans shifted into the Northern and Southern Plains. Increased pest management costs meant a decline in farmer returns, though higher crop prices under the extreme climate scenarios partially offset increased control costs, with a greater impact on farmer returns in the hottest, driest scenarios. Since prices go up, producer losses are not offset by consumer gains, implying that consumers would be worse off.

Research is underway to generate adaptation options for climate change by introducing crop varieties that are tolerant to environmental stresses such as drought and high temperatures. Such developments will depend on technological opportunities, as well as on investments in research over time. We modeled one particular example—the introduction of drought-tolerant varieties of corn, wheat, soybeans, and cotton—that resulted in 10- to 15-percent higher yields in drier but nonirrigated environments. Within each climate change scenario, introduction of drought-tolerant varieties resulted in a slight reduction in crop prices. With a few exceptions, drought-tolerant varieties led to small shifts in corn, wheat, and soybean acreages from the eastern half of the United States into the Plains or Mountain States, where drought-tolerant varieties were introduced. In contrast to the additional pest-pressure case, lower producer returns in many regions were complemented by lower prices, benefiting consumers.

Under most scenarios, changes in environmental indicators of soil and water quality generally followed acreage changes. Environmental impacts were also sensitive, however, to changes in production practices. Under several scenarios, the shift in nitrogen loss to water was proportionately larger than the total increase in cropped acreage, indicating an increase in production intensity for regions where more severe environmental impacts were observed, although soil loss to water erosion was commensurate with acreage change.

The results for the pest-pressure and drought-tolerance cases have been presented as changes relative to the climate change adaptation scenarios, rather than as changes from the baseline. This was done to show that the range of climate projections had an influence in determining how factors, such as biotic impacts (i.e., pests) and technological advancements, will ultimately affect U.S. crop production patterns and the environmental consequences associated with them. An alternative way to consider the results

is to compare the scenario results for each case directly to the “no climate change baseline.” Rather than capturing the contribution of differences in crop yields between climate change scenarios, comparison with the baseline provides a measure of the composite effects of climate change plus the changes in production environment introduced in each case. Table 21 illustrates the changes in prices and returns from the climate change scenarios with respect to the “no climate change baseline.” The table indicates a consistent pattern corresponding to the climate change scenarios. In all cases, the ECH and CSIRO scenarios led to lower aggregate crop prices than under the “no climate change baseline,” whereas the more extreme CNR and MIROC scenarios led to higher aggregate crop prices. Similarly, crop returns followed the same trend. In the more extreme scenarios (CNR and MIROC), higher crop prices were generally paired with lower returns (compared with the “no climate change” baseline). Changes in relative profitability within and across regions resulted in lower returns in the Corn Belt across all scenarios in the adaptation, pest-pressure, and drought-tolerance cases.

Environmental consequences generally increased across all scenarios for all cases (table 22). Increases in the environmental measures were largely the result of an expansion in total planted acres under the climate change scenarios, with some contribution from redistribution of crop production between regions with different soil and water characteristics. For nitrogen loss to water, the percent change in the indicator value was generally greater than the percent change in acreage, suggesting both intensification of agricultural production through increased fertilizer use and redistribution of crop production to regions and rotations more susceptible to nutrient losses. In aggregate, changes in soil lost to water erosion were more moderate than those for nitrogen loss, with the percentage change in erosion impacts often less than the change in acreage overall. Regional erosion impacts varied widely, however.

Analysis Limitations and Future Research

In exploring the implications of climate change for U.S. field crop production, our analysis focused on the yield-related impacts associated with increased regional average temperatures, varied regional changes in average precipitation, increased carbon dioxide concentration in the atmosphere, the expanded incidence of pests, and the market-mediated price impacts that arise as a result of decentralized impacts and adaptive responses. The GCM results used to drive our climate change projections did not allow us to estimate changes in the variability of daily temperature and precipitation, but other studies suggest that changes in volatility of weather parameters—including increased incidence of extreme weather events—may also be a significant driver of yield and impact changes (Isik and Devadoss, 2006). By capturing only changes in average maximum daily temperature, minimum daily temperature, and precipitation, our EPIC results may underestimate the full yield impacts expected from changes in those climate conditions.

Furthermore, direct yield impacts represent only a partial, if important, subset of the climate change elements likely to impact farmers. Other atmospheric factors—such as ground-level ozone, which is expected to increase in tandem with increasing CO₂ emissions—may have significant agricultural

Table 21

**Economic effects of climate change scenarios on agriculture:
Changes from “no climate change” baseline**

REAP Cases	Changes in crop prices (aggregated)			
	ECH	CSIRO	CNR	MIROC
	<i>Percent</i>			
No adaptation	-2.1	-0.9	3.9	8.3
Adaptation	-4.0	-2.9	1.9	6.4
Pest damage	-1.5	-0.8	5.1	11.9
Drought-tolerance	-4.7	-3.6	1.1	5.6
	Changes in annual net returns to crop production (United States)			
	ECH	CSIRO	CNR	MIROC
	<i>\$ Million</i>			
No adaptation	4,185	2,664	493	-6
Adaptation	3,619	2,165	-332	-1,465
Pest damage	1,716	694	-2,936	-4,473
Drought-tolerance	3,992	2,671	155	-843
	Changes in annual net returns to crop production (Corn Belt)			
	ECH	CSIRO	CNR	MIROC
	<i>\$ Million</i>			
No adaptation	299	-680	-747	-2,592
Adaptation	-1,114	-2,165	-2,112	-4,053
Pest damage	-1,576	-2,608	-2,680	-4,564
Drought-tolerance	-1,482	-2,538	-2,503	-4,381
	Changes in annual net returns to crop production (all regions except Corn Belt)			
	ECH	CSIRO	CNR	MIROC
	<i>\$ Million</i>			
No adaptation	3,886	3,344	1,240	2,586
Adaptation	4,733	4,330	1,780	2,588
Pest damage	3,292	3,303	-256	91
Drought-tolerance	5,474	5,210	2,658	3,538

Notes: ECH, CSIRO, CNR, and MIROC are climate change scenarios (see table 4 for sources). The “no climate change” “...baseline crop returns in REAP were \$43 billion. The “no adaptation” case assumes climate change affects crop yield but that farmers’ land allocations and cropping practices do not adjust. The “adaptation” case allows farmers to adjust their crop rotations, tillage practices, and land allocation decisions in response to how the climate change affects crop yield and resulting market conditions. The “pest damage” case considers the additional effects of changes in the distribution of crop pests resulting from climate change and how this affects crop yields and pest control costs. The “drought-tolerance” case assumes that drought-tolerant crop varieties of corn, wheat, soybeans, and cotton become available and provide another option for how farmers can adapt to climate change. It does not include damages from changes in pest populations. See table 2 for a complete discussion of the REAP analysis cases.

Table 22

**Environmental effects of climate change scenarios on agriculture:
Changes from “no climate change baseline”**

REAP Cases	Changes in total crop acreage			
	ECH	CSIRO	CNR	MIROC
	<i>Million acres</i>			
No adaptation	-	-	-	-
Adaptation	1.9	1.8	0.8	3.2
Pest damage	4.0	2.7	5.0	8.8
Drought-tolerance	1.4	0.6	-0.3	1.9
	Changes in total crop acreage			
	ECH	CSIRO	CNR	MIROC
	<i>Percent</i>			
No adaptation	-	-	-	-
Adaptation	0.6	0.6	0.2	1.0
Pest damage	1.3	0.9	1.6	2.8
Drought-tolerance	0.5	0.2	-0.1	0.6
	Changes in nitrogen losses to water			
	ECH	CSIRO	CNR	MIROC
	<i>Percent</i>			
No adaptation	-2.7	-2.4	0.4	1.4
Adaptation	1.4	1.5	2.1	5.0
Pest damage	5.7	5.0	5.2	6.8
Drought-tolerance	1.5	1.4	2.0	4.7
	Changes in sheet and rill erosion			
	ECH	CSIRO	CNR	MIROC
	<i>Percent</i>			
No adaptation	-6.0	-3.5	-3.2	-0.2
Adaptation	-0.9	0.6	0.9	1.2
Pest damage	-0.8	1.0	3.3	4.9
Drought-tolerance	-1.6	0.1	0	0.3

Notes: ECH, CSIRO, CNR, and MIROC are climate change scenarios (see table 4 for sources). The “no adaptation” case assumes climate change affects crop yield but that farmers’ land allocations and cropping practices do not adjust. The “adaptation” case allows farmers to adjust their crop rotations, tillage practices, and land allocation decisions in response to how the climate change affects crop yield and resulting market conditions. The “pest damage” case considers the additional effects of changes in the distribution of crop pests resulting from climate change and how this affects crop yields and pest control costs. The “drought-tolerance” case assumes that drought-tolerant crop varieties of corn, wheat, soybeans, and cotton become available and provide another option for how farmers can adapt to climate change. It does not include damages from changes in pest populations. See Table 2 for a complete discussion of the REAP analysis cases.

impacts and may influence the incentives and constraints that farmers face in responding to a changing climate. The potential impacts of climate change on the supply and costs of agricultural inputs—such as land, energy, fertilizer, water, and labor—would also affect relative returns to different types of production and would possibly create region-specific constraints on the adaptive strategies available to farmers. Such considerations were beyond the scope of this analysis. Moreover, our research focuses on the potential for adaptation within the U.S. crop sector, with a particular emphasis on major field crops. We did not consider specialty crops—fruits, vegetables, nursery crops, and other specialty crops—which account for an important share of the value of U.S. agricultural production. We recognize, however, that climate change can have an important bearing on resources supporting specialty crop production as well. The model did consider livestock sector impacts through changes in feedgrain markets, although we did not capture the full range of substitution of feed ingredients that may occur due to climate change. Nor did we explicitly model the effects of climate change on animal productivity and other aspects of livestock management costs.

Our analysis also was limited in the range of adaptation strategies available to farmers that could be examined within the existing modeling framework. REAP evaluates adaptation strategies related to changing crop patterns and practices, but existing production enterprises in the model did not allow for other farm-level adaptation strategies, such as changing harvesting and planting dates or the timing and magnitude of applied irrigation or fertilizer. Such strategies are common responses to weather variability and will be an important element in farmers' adaptation responses. The potential for, and constraints to, expanding irrigated acreage and water use may be a particularly significant factor in adaptation strategies where there is already significant competition for water resources, such as in the Western United States. REAP does not currently allow for an analysis of shifting irrigation patterns within U.S. agriculture; however, such modifications are underway and will inform ongoing ERS research related to climate adaptation.

References

- Adams, R.M., and D.E. Peck. "Climate Change and Agriculture: Economic Impacts," *Choices*, 1st Quarter, 23(1), 2008.
- Adusei, E.O., and G.W. Norton. "The Magnitude of Agricultural Maintenance Research in the USA," *Journal of Production Agriculture* 3: pp. 1-6, 1990.
- Albrecht, W.A. "Loss of soil organic matter and its restoration," *Soils and Men, USDA Yearbook of Agriculture*, U.S. Government Printing Office: Washington, DC, pp. 347-60, 1938.
- Alston, J.M., M.A. Andersen, J.S. James, and P.G. Pardey. *Persistence Pays: U.S. Agricultural Productivity Growth and the Benefits from Public R&D Spending*, Springer: New York, 2010.
- Alston, J.M., G.W. Norton, and P.G. Pardey. *Science Under Scarcity: Principles and Practice for Agricultural Research Evaluation and Priority Setting*, Cornell University Press: Ithaca, NY, 1995.
- Bradley, B.A., M. Oppenheimer, and D.S. Wilcove. "Climate Change and Plant Invasions: Restoration Opportunities Ahead?" *Global Change Biology* 15: pp. 1511-21, 2009.
- Brennan, J.P. "An analytical model of a wheat breeding program," *Agricultural Systems* 31: pp. 349-66, 1989.
- Bridges, D.C. (Ed.). *Crop Losses Due to Weeds in the United States*, Weed Society of America: Champaign, IL, p. 403, 1992.
- Campos, H., M. Cooper, J.E. Habben, G.O. Edmeades, and J.R. Schussler. "Improving drought tolerance in maize: A view from industry," *Field Crops Research* 90: pp. 19-34, 2004.
- Castiglioni, P., D. Warner, R.J. Bensen, D.C. Anstrom, J. Harrison, M. Stoecker, M. Abad, G. Kumar, S. Salvador, R. D'Ordine, S. Navarro, S. Back, M. Fernandes, J. Targolli, S. Dasgupta, C. Bonin, M.H. Luethy, and J.E. Heard. "Bacterial RNA chaperones confer abiotic stress tolerance in plants and improved grain yield in maize under water-limited conditions," *Plant Physiology* 147: pp. 446-55, 2008.
- Cattivelli, L., F. Rizza, F.W. Badeck, E. Mazzucotelli, A.M. Mastrangelo, E. Francia, C. Marè, A. Tondelli, and A.M. Stanca. "Drought tolerance improvement in crop plants: An integrated view from breeding to genomics," *Field Crops Research* 105: pp. 1-14, 2008.
- Dettinger, M., and S. Earman. "Western Ground Water and Climate Change—Pivotal to Supply Sustainability or Vulnerability in Its Own Right?" *Ground Water: News and Views* 4 (1): pp. 4–5, June 2007.
- Déqué, M., C. Drevet, A. Braun, and D. Cariolle. "The ARPEGE/IFS atmosphere model: a contribution to the French community climate modeling," *Climate Dynamics* 10 (4-5): pp. 249–66, 1994.

- Ding, Ya, and Karina Schoengold. "The Impact of Weather Extremes on Agricultural Production Methods: Does Drought Increase Adoption of Conservation Tillage Practices?" *Journal of Agricultural and Resource Economics* 34(3): pp. 395-411, December 2009.
- Duke, S.O., and S.B. Powles. "Glyphosate: A Once-In-A-Century Herbicide," *Pest Management Science* 64(4): pp. 319-25, 2008.
- Eathington, S.R., T.M. Crosbie, M.D. Edwards, R.S. Reiter, and J.K. Bull. "Molecular markers in a commercial breeding program," *Crop Science* 47(S3): pp. S154-63, 2007.
- Edmeades, G.O. "Drought tolerance in maize: an emerging reality," *Global Status of Commercialized Biotech/GM Crops: 2008*, International Service for the Acquisition of Agri-biotech Applications Brief No. 39, 2008.
- Feng, X., A.J. Simpson, K.P. Wilson, D.D. Williams, and M.J. Simpson. "Increased cuticular carbon sequestration and lignin oxidation in response to soil warming," *Nature Geoscience* 1, pp. 836-39, 2008.
- Fernandez-Cornejo, J., and M. Caswell. *The First Decade of Genetically Engineered Crops in the United States*, EIB-11, L. Mitchell, E. Golan, and F. Kuchler (Contrib.), U.S. Department of Agriculture, Economic Research Service, 2006.
- Frumhoff, P.C., J.J. McCarthy, J.M. Melillo, S.C. Moser, and D.J. Wuebbles. *Confronting Climate Change in the U.S. Northeast: Science, Impacts, and Solutions*, Northeast Climate Impacts Assessment, Union of Concerned Scientists: Cambridge, MA, p. 146, 2007.
- Fuglie, K.O., and P.W. Heisey. *Economic Returns to Public Agricultural Research*, EB-10, U.S. Department of Agriculture, Economic Research Service, 2007.
- Fuglie, K.O., P.W. Heisey, J.L. King, C.E. Pray, K. Day-Rubenstein, D. Schimmelpfennig, S.L. Wang, and R. Karmarkar-Deshmukh. "Private Research and Development for Crop Genetic Improvement," *Research Investments and Market Structure in the Food Processing, Agricultural Input, and Biofuel Industries Worldwide*, ERR-130, December 2011, <http://ers.usda.gov/publications/err130/>.
- Gleick, Peter. *Water: The Potential Consequences of Climate Variability and Change for the Water Resources of the United States*, Water Sector Assessment Team of the National Assessment of the Potential Consequences of Climate Variability and Change, U.S. Global Change Research Program, p. 160, September 2000.
- Gordon, H.B., L.D. Rotstain, J.L. McGregor, M.R. Dix, E.A. Kowalczyk, S.P. O'Farrell, L.J. Waterman, A.C. Hirst, S.G. Wilson, M.A. Collier, I.G. Watterson, and T.I. Elliott. *The CSIRO Mk3 Climate System Model*, CSIRO Atmospheric Research Technical Paper No. 60, Commonwealth Scientific and Industrial Research Organisation (CSIRO): Aspendale,

- Victoria, Australia, p. 130, 2002, http://www.cmar.csiro.au/e-print/open/gordon_2002a.pdf.
- Hatch, U., S. Jagtap, J. Jones, and M. Lamb. "Potential Effects of Climate Change on Agricultural Water Use in the Southeast U.S.," *Journal of the American Water Resources Association* Vol. 35, No. 6, December 1999.
- Hatfield, J., K. Boote, P. Fay, L. Hahn, C. Izaurralde, B.A. Kimball, T. Mader, J. Morgan, D. Ort, W. Polley, A. Thomson, and D. Wolfe. "Agriculture," *The Effects of Climate Change on Agriculture, Land Resources, Water Resources, and Biodiversity in the United States*, U.S. Climate Change Science Program and the Subcommittee on Global Change Research: Washington, DC, p. 362, 2008.
- Heap, I. *International Survey of Herbicide Resistant Weeds – Glycines (G/9) Resistant Weeds by Species and Country*, <http://www.weedscience.org/Summary/UspeciesMOA.asp?lstMOAID=12>.
- Hijmans, R.J., S.E. Cameron, J.L. Parra, P.G. Jones, and A. Jarvis. "Very high resolution interpolated climate surfaces for global land areas," *International Journal of Climatology* 25: pp. 1965-78, 2005.
- Homer, C., C. Huang, L. Yang, B. Wylie, and M. Coan. "Development of a 2001 National Landcover Database for the United States," *Photogrammetric Engineering and Remote Sensing* Vol. 70, No. 7, pp. 829-40, July 2004.
- Howden, S.M., J.F. Soussana, F.N. Tubiello, N. Chhetri, M. Dunlop, and H. Meinke. "Adapting agriculture to climate change," *Proceedings of the National Academy of Sciences* 104, 50: pp. 19691-96, 2007.
- Howitt, R., J. Medellin-Azuara, and D. MacEwan. "Climate Change, Markets, and Technology," *Choices*, 3rd Quarter 25(3), 2010.
- Howitt, R., and K. Hansen. "The Evolving Western Water Markets," *Choices*, 1st Quarter 20(1), 2005.
- Intergovernmental Panel on Climate Change (IPCC Report). *Climate Change 2007: Impacts, Adaptation and Vulnerability*, Fourth Assessment Report of the Intergovernmental Panel on Climate Change, M.L. Parry, O.F. Canziani, J.P. Palutikof, J.P. van der Linden, and C.E. Hanson (Eds.), Cambridge University Press: Cambridge, UK, p. 976, 2007.
- Isik, Murat, and Stephen Devadoss. "An analysis of the impact of climate change on crop yields and yield variability," *Applied Economics* 38: pp. 835-44, 2006.
- Izaurralde, R. César, Norman J. Rosenberg, Robert A. Brown, and Allison M. Thomson. "Integrated assessment of Hadley Center (HadCM2) climate-change impacts on agricultural productivity and irrigation water supply in the conterminous United States. Part II. Regional agricultural production in 2030 and 2095," *Agricultural and Forest Meteorology* 117, pp. 97-122, 2003.

- Jones, P.G., P.K. Thornton, and J. Heinke. *Generating characteristic daily weather data using downscaled climate model data from the IPCC's Fourth Assessment*, project report, International Livestock Research Institute, 2009.
- Joyce, L.A., G.M. Blate, J.S. Littell, S.G. McNulty, C.I. Millar, S.C. Moser, R.P. Neilson, K.A. O'Halloran, and D.L. Peterson. "National Forests," *Preliminary Review of Adaptation Options for Climate-Sensitive Ecosystems and Resources, Synthesis and Assessment Product 4.4*, U.S. Environmental Protection Agency, pp. 3-1–3-127, 2008.
- K-1 Model Developers. *K-1 Coupled GCM (MIROC) Description*, K-1 Technical Report No. 1, H. Hasumi and S. Emori (Eds.), Center for Climate System Research, University of Tokyo, National Institute for Environmental Studies, Frontier Research Center for Global Change, p. 34, 2004, <http://www.ccsr.u-tokyo.ac.jp/kyosei/hasumi/MIROC/techrepo.pdf>.
- Karl, T.R., J. M. Melillo, and T. C. Peterson, (Eds.). *Global Climate Change Impacts in the United States*, Cambridge University Press, 2009.
- Kenny, J.F., N.L. Barber, S.S. Hutson, K.S. Linsey, J.K. Lovelace, and M.A. Maupin. *Estimated Use of Water in the United States in 2005*, U.S. Geological Survey Circular 1344, p. 52, 2009.
- Key, N. and S. Sneeringer. "The Potential Effects of Climate Change on the Productivity, Costs, and Returns of U.S. Dairy Production," selected paper at the AAEE Annual Meeting, Pittsburgh, PA, 2011.
- Kiely, T., D. Donaldson, and A. Grube. *Pesticides Industry Sales and Usage: 2000 and 2001 Market Estimate*, U.S. Environmental Protection Agency, 2004.
- Knowles, N., M.D. Dettinger, and D.R. Cayan. "Trends in snowfall versus rainfall in the western United States," *Journal of Climate* 19 (18): pp. 4545–59, September 2006.
- Livingston, M.J., G.A. Carlson, and P.L. Fackler. "Managing Resistance Evolution in Two Pests to Two Toxins with Refugia," *American Journal of Agricultural Economics* 86(1): pp. 1-13, 2004.
- Livingston, M., R. Johansson, S. Daberkow, M. Roberts, M. Ash, and V. Breneman. *Economic and Policy Implications of Wind-Borne Entry of Asian Soybean Rust into the United States*, OCS-04D-02, U.S. Department of Agriculture, Economic Research Service, 2004, <http://www.ers.usda.gov/Publications/OCS/Apr04/OCS04D02/>.
- Livingston, M.J., N.P. Storer, J.W. Van Duyn, and G.G. Kennedy. "Do Refuge Requirements for Biotechnology Crops Promote Economic Efficiency? Some Evidence for Bt Cotton," *Journal of Agricultural and Applied Economics* 39(1): pp. 171–85, 2007.

- Lobell, D., and M. Burke. "Economic impacts of climate change on agriculture to 2030," *Climate Change & Crop Production*, M.P. Reynolds (Ed.), CAB International: Wallingford, UK, pp. 38-49, 2010.
- López-Pereira, M.A., and M.P. Filippello. "Maize Seed Industries, Revisited: Emerging Roles of the Public and Private Sectors," *CIMMYT 1993-94 World Maize Facts and Trends*, Centro Internacional de Mejoramiento de Maíz y Trigo (CIMMYT): Mexico, D.F., 1994.
- Maestri, E., N. Klueva, C. Perrotta, M. Gulli, H.T. Nguyen, and N. Marmioli. "Molecular genetics of heat tolerance and heat shock proteins in cereals," *Plant Molecular Biology* 48: pp. 667-81, 2002.
- Martineau, J.R., J.H. Williams, and J.E. Specht. "Temperature Tolerance in Soybeans. II. Evaluation of Segregating Populations for Membrane Thermostability," *Crop Science* 19(1): pp. 79-81, 1979.
- McGrath, J.M., and D.B. Lobell. "An independent method of deriving the carbon dioxide fertilization effect in dry conditions using historical yield data from wet and dry years," *Global Change Biology* 17(8): pp. 2689-96, 2011.
- Monsanto. *Research and Development Pipeline*, 2011, <http://www.monsanto.com/products/Documents/pipeline-flash/pdfs/TechPipelineALL.pdf>.
- Mooney, H.A., and R.J. Hobbs (Eds.). *Invasive Species in a Changing World*, Island Press: Washington, 2000.
- Mote, P.W., A.F. Hamlet, M.P. Clark, and D.P. Lettenmaier. "Declining mountain snowpack in western North America," *Bulletin of the American Meteorological Society* 86: pp. 39-49. 2005.
- Mourgues, F., M.N. Brisset, and E. Chevreau. "Strategies to improve plant resistance to bacterial diseases through genetic engineering," *Trends in Biotechnology* 16: pp. 203-10, 1998.
- National Water Assessment Group (NWAG). *Water: The Potential Consequences of Climate Variability and Change for the Water Resources of the United States*. A report of the National Water Assessment Group for the U.S. Global Change Research Program. Lead Author: Peter Gleick. Pacific Institute for Studies in Development, Environment, and Security, Oakland, CA, Sept. 2000.
- Ojima, D., L. Garcia, E. Elgaali, K. Miller, T. Kittel, and J. Lockett. "Potential Climate Change Impacts on Water Resources in the Great Plains," *Journal of the American Water Resources Association* Vol. 35, No. 6, December 1999.
- Osteen, C., and M. Livingston. "Pest management practices," *Agricultural Resources and Environmental Indicators 2006 Edition*, EIB-16, K. Wiebe and N. Gollehon (Eds.), U.S. Department of Agriculture, Economic Research Service, 2006, <http://www.ers.usda.gov/Publications/AREI/EIB16/>.

- Owen, M.D.K. "Weed Species Shifts in Glyphosate-Resistant Crops," *Pest Management Science* 64(4): pp. 377-87, 2008.
- Padgitt, M., D. Newton, R. Penn, and C. Sandretto. *Production Practices for Major Crops in U.S. Agriculture, 1990-97*, SB-969, U.S. Department of Agriculture, Economic Research Service, 2000, <http://www.ers.usda.gov/Publications/SB969/>.
- Parry, M.A.J., M. Reynolds, M.E. Salvucci, C. Raines, P.J. Andralojc, X.G. Zhu, G.D. Price, A.G. Condon, and R.T. Furbank. "Raising yield potential of wheat. II. Increasing photosynthetic capacity and efficiency," *Journal of Experimental Botany* 62(2): pp. 453-67, 2011.
- Punja, Z.K. "Genetic engineering of plants to enhance resistance to fungal pathogens—a review of progress and future prospects," *Canadian Journal of Plant Pathology* 23: pp. 216-35, 2001.
- Radin, J.W., Z. Lu, R.G. Percy, and E. Zeiger. "Genetic variability for stomatal conductance in Pima cotton and its relation to improvements of heat adaptation," *Proceedings of the National Academy of Sciences* 91: pp. 7217-21, 1994.
- Reilly, J.M., and D. Schimmelpfennig. "Agricultural impact assessment, vulnerability, and the scope for adaptation," *Climatic Change* 43: pp. 745-88, 1999.
- Reynolds, M.P., M. Balota, M.I.B. Delgado, I. Amani, and R.A. Fischer. "Physiological and morphological traits associated with spring wheat yield under hot, irrigated conditions," *Australian Journal of Plant Physiology* 21: pp. 717-30, 1994.
- Roberts, M.J., D. Schimmelpfennig, E. Ashley, M. Livingston, M. Ash, and U. Vasavada. *The Value of Plant Disease Early Warning Systems: A Case Study of USDA's Soybean Rust Coordinated Framework*, ERR-18, U.S. Department of Agriculture, Economic Research Service, 2006, <http://www.ers.usda.gov/Publications/ERR18/>.
- Roeckner, E., G. Bäuml, L. Bonaventura, R. Brokopf, M. Esch, M. Giorgetta, S. Hagemann, I. Kirchner, L. Kornblueh, E. Manzini, A. Rhodin, U. Schlese, U. Schulzweida, and A. Tompkins. *The Atmospheric General Circulation Model ECHAM5. Part I: Model Description*, MPI Report 349, Max Planck Institute for Meteorology: Hamburg, Germany, p. 127, 2003.
- Rosenzweig, C., and D. Hillel. "Potential Impacts of Climate Change on Agriculture and Food Supply," *Consequences* 1(2): pp. 23-32, 1995.
- Shinozaki, K., and K. Yamaguchi-Shinozaki. "Gene networks involved in drought stress response and tolerance," *Journal of Experimental Botany* 58(2): pp. 221-27, 2007.
- Smit, Barry, and Mark W. Skinner. "Adaptation options in agriculture to climate change: A typology," *Mitigation and Adaptation Strategies for Global Change* 7: pp. 85-114, 2002.

Soil and Water Conservation Society (SWCS). *Conservation Implications of Climate Change: Soil Erosion and Runoff from Cropland*, pp.1-24, January 2003.

Strzepek, Kenneth, Gary Yohe, James Neumann, and Brent Boehlert. "Characterizing changes in drought risk for the United States from climate change," *Environmental Research Letters*, 5 044012, 2010, <http://iopscience.iop.org/1748-9326/5/4/044012>.

Thomson, A.M., R.A. Brown, N.J. Rosenberg, R. Srinivasan, and R.C. Izaurralde. "Climate change impacts for the conterminous USA: An integrated assessment," *Climatic Change* 69: pp. 43-105, 2005.

ur Rahman, H., S.A. Malik, and M. Saleem. "Heat tolerance of upland cotton during the fruiting stage evaluated using cellular membrane thermostability," *Field Crops Research* 85: pp. 149-58, 2004.

U.S. Climate Change Science Program (USCCSP). *The Effects of Climate Change on Agriculture, Land Resources, Water Resources, and Biodiversity in the United States*, Subcommittee on Global Change Research, P. Backlund, A. Janetos, and D. Schimel (Lead authors), M. Walsh (Managing editor), U.S. Department of Agriculture, pp. 362, 2008.

U.S. Department of Agriculture. "New Germplasm Helps Assure Food for the Future," *USDA News Feature, News Division*, Office of Public Affairs, December 6, 1990.

U.S. Department of Agriculture, National Agricultural Statistics Service (USDA/NASS). *Census of Agriculture, 2007*, <http://www.agcensus.usda.gov/>.

U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Center for Plant Health Science and Technology (CPHST). *A System to Integrate Climate Change Scenarios into Response Programs for Invasive Plant Pests: Proposal Submitted to the National Institute of Food and Agriculture*, 2010a.

U.S. Department of Agriculture, Interagency Agricultural Projections Committee. *Agricultural Projections to 2019*, OCE 2010-1, 2010b, <http://usda.mannlib.cornell.edu/usda/ers/94005/2010/OCE101.pdf>.

U.S. Department of the Interior (DOI), Bureau of Reclamation. *SECURE Water Act Section 9503(c) - Reclamation Climate Change and Water*, Report to Congress, April 2011.

U.S. Department of Labor, Bureau of Labor Statistics. *Producer Price Index for Pesticides and Other Agricultural Chemicals*, PCU325320325320, 2010, <http://www.bls.gov/ppi/>.

U.S. Government Accountability Office. *USDA: Information on Classical Plant and Animal Breeding Activities*, GAO-07-1171R, October 2007.

- Wahid, A., S. Gelani, M. Ashraf, and M.R. Foolad. "Heat tolerance in plants: An overview," *Environmental and Experimental Botany* 61: pp. 199-223, 2007.
- Wolfe, D.W., L. Ziska, C. Petzoldt, A. Seaman, L. Chase, and K. Hayhoe. "Projected Change in Climate Thresholds in the Northeastern U.S.: Implications for Crops, Pests, Livestock, and Farmers," *Mitigation and Adaptation Strategies for Global Change* 13(5-6): pp. 555-75, 2007.
- Ziska, L.H., and D.M. Blumenthal. "Empirical selection of cultivated oat in response to rising atmospheric carbon dioxide," *Crop Science* 47(4): pp. 1547-52, 2007.
- Ziska, L.H., D.M. Blumenthal, P. Teal, G.B. Runion, E.R. Hunt, Jr., and H. Diaz-Soltero. "Invasive Species and Climate Change: An Agronomic Perspective," *Climatic Change* 105: pp. 13-42, 2010.
- Ziska, L.H. "Three-year field evaluation of early and late 20th century spring wheat cultivars to projected increases in atmospheric carbon dioxide," *Field Crops Research* 108(1): pp. 54-9, 2008.
- Ziska, L.H., and A. McClung. "Differential response of cultivated and weedy (red) rice to recent and projected increases in atmospheric carbon dioxide," *Agronomy Journal* 100(5): pp. 1259-63, 2008.

Appendix A: The REAP Model

The Regional Environment and Agriculture Programming (REAP) model is a static, partial equilibrium optimization model of the agricultural sector that quantifies agricultural production and its associated environmental outcomes for 48 regions in the United States (see fig. 1). The regions are defined by the intersection of USDA's Farm Production Regions (defined by State boundaries) and Land Resource Regions (defined by predominant soil type and geography).

REAP employs survey data (from USDA's Agricultural Resource Management Survey (ARMS)) and simulated input data (from the Environmental Productivity and Integrated Climate (EPIC) model) at the regional level on crop yields, input requirements, costs and returns, and environmental parameters to estimate long-run equilibrium outcomes. Regional production levels are determined for 10 crops and 13 livestock categories, and national production levels are determined for 20 processed products. For each REAP region, land use, crop mix, and acreage allocations by multi-year crop rotation and tillage practice are endogenously determined by REAP's constrained optimization process. Input use and national-level prices are also determined endogenously. The model has been applied to address a wide range of agri-environmental issues, such as soil conservation and environmental policy design, environmental credit trading, climate change mitigation policy, and regional effects of trade agreements.

REAP is implemented as a nonlinear mathematical program using the General Algebraic Modeling System (GAMS) programming environment. The model determines a welfare-maximizing set of crop, livestock, and processed product production levels subject to land constraints and processing and production balance requirements. Production activities for crops within a region (defined by crop rotation and tillage) are allocated in the model solution based on a constant elasticity of transformation (CET) relationship. The CET specification helps to avoid unrealistic "corner point" solutions by accounting for cost/return and risk considerations embedded in observed acreage allocations but not explicitly included in the model. The model is calibrated to USDA baseline production levels over a multi-year timeframe using the Positive Mathematical Programming (PMP) method.

Production "shocks" under policy, technical, or environmental scenarios can be introduced via changes or additions to constraints, modifications of baseline data assumptions, adjustments in objective function terms, or some combination of approaches. Changes in policy, commodity demand, or production/processing technology can be imposed on the model and the results examined to determine their effects on:

- Regional supply of crops and livestock;
- Commodity prices;
- Crop management and production input use;
- Farm income; and
- Environmental indicators, such as nutrient and pesticide runoff, soil loss, GHG emissions, soil carbon fluxes, and energy use.

For more information on REAP and its applications, see model documentation at <http://www.ers.usda.gov/publications/tb1916/tb1916fm.pdf>.

Appendix B: Climate Projections for 2030

Using Climate Projections in Agro-Economic Analysis

Interest in climate change and weather processes, in general, has resulted in a wide array of General Circulation Models (GCMs) that attempt to project climate trends. GCMs are highly complex models that use assumptions about future emissions paths and the evolution of atmospheric concentration of various greenhouse gases (GHGs) to estimate spatial patterns of temperature and precipitation. Because of their numerical complexity, GCMs simulate climate conditions at a very coarse spatial and temporal resolution. They simulate how atmospheric behavior that is averaged over relatively long periods of time (climate) will change, but they do not model weather or localized atmospheric conditions, such as storm events, drought, or temperature extremes over short periods of time. Also, the spatial resolution of GCMs is not sufficient to capture the wide variation in temperature, precipitation, and other meaningful weather conditions influenced by detailed ground topography that, in turn, influence crop production. A grid cell in a GCM may cover approximately 10,000 square kilometers, resulting in an average of only 15 cells per REAP region.

Downscaling

Spatial downscaling is a process that provides finer resolution climate information from the lower resolution data emerging from GCMs. Downscaling methods generally apply assumptions about regional climate dynamics to disaggregate the GCM averages down into region-specific forecasts. One of the major assumptions of spatial downscaling is that it is possible to determine significant relationships between local and large-scale climate that will remain valid under future climate conditions. There are several methods, often involving simulation or statistics, to spatially downscale GCM output; the method used to generate the data used in this report is described in Jones et al., 2009.

Choice of Models and Data

For this analysis, we focused on the SRES A1B emissions scenario. To choose GCM outputs, we considered whether the outputs:

- Were downscaled by the same method to the same resolution across our entire U.S. study area;
- Included the appropriate set of climate variables for the EPIC crop-growth simulator analysis—monthly minimum temperature and maximum temperature and precipitation; and
- Covered the 2030 projection year used in the REAP analysis, using an average of weather projections over a period of at least 5 years.

The downscaled GCM output we employed was generated by Jones et al. (2009). The four models and their sources are listed in table 4. The data included the following disclaimer, “These downscaled climate data are NOT

predictions of what the future climate will be like in any place. They are projections of possible future climate, and should be treated with considerable caution. There is a great deal of variability between different climate models, between different greenhouse gas emission scenarios, and between different downscaling methods.”

Aggregating Climate Data to REAP Regions

The downscaled climate data covered the coterminous United States. Most of the points corresponded to land unsuitable to agriculture, such as mountain ranges, urban areas, or bodies of water. Much of the United States is range-land that is not in use and is unlikely to be used in the near future for crop agriculture. Since we required a single average weather value representative of each region to derive crop yield estimates, it was important that the values reflected conditions that were representative of conditions in agricultural areas. Nonagricultural data points can bias weather characteristics and alter the performance of the crop-growth simulation. The downscaled weather data were confined to agricultural production areas by overlaying National Land Cover Database (NLCD) results for cultivated crops and pasture/hay categories onto the REAP regions (Homer et al., 2004). Average monthly values for maximum temperature and precipitation and minimum temperature were computed from agricultural land within each REAP region (see fig. 1 for a map of the REAP regions).

Appendix C: Sensitivity of Analysis Results to Climate Change Elements

The Environmental Productivity and Integrated Climate (EPIC) simulation of the yield impacts of simultaneously changing values of temperature, precipitation, and carbon dioxide concentration drives REAP’s analysis of the impacts of future climate scenarios relative to a baseline scenario. EPIC’s results are in turn driven by a large set of technical parameter assumptions that are held constant across climate scenarios but that, through their influence on the relative impact of temperature, precipitation, and carbon dioxide concentration on crop yields, can subsequently influence differences in impact across future climate scenarios. Examples of such assumptions include the minimal and optimal growth temperatures for each crop, the parameters of the relationship between carbon dioxide concentration and crop growth, water-related parameters, such as maximum stomatal conductance, and assumptions about the rate of decline in radiation use efficiency with increasing vapor pressure deficits.¹

Because there is ongoing debate about the expected magnitude of impacts from factors such as carbon dioxide concentration (i.e. carbon dioxide fertilization), and to understand how each element of the climate change impact behaves individually in EPIC’s results, it is helpful to present disaggregated climate change impact results. EPIC scenarios in which temperature, precipitation, and carbon dioxide concentrations are varied independently of one another are described in appendix table 1. Note that because of interaction effects, the impact of the combined changes is not a strict sum of the impact of individual effects. Carbon dioxide fertilization’s impact on transpiration, for example, could alter the sensitivity of impact results to precipitation changes.

To isolate the biophysical impacts from the behavioral impacts in this analysis, results are presented for a series of “Impact without Adaptation” cases where production acreage is fixed across all scenarios (Appendix fig. 1). Productivity changes are due exclusively to changes in biophysical impact. As mentioned in the section on “Climate Impact Analysis: No Adaptation,” regional yield changes reflect yield changes at the crop rotation level (as measured by EPIC) that are then weighted by rotation acreage in aggregating up to the regional level.

¹For a complete list of EPIC’s parameters, see the EPIC documentation at <http://epicapex.brc.tamus.edu/media/23015/epic0509usermanualupdated.pdf>.

Appendix table 1
Climate scenarios used to explore yield impact sensitivity to climate change elements

Measure	Baseline	CNR	CNR_No_CO ₂	CNR_JUST_CO ₂	CNR_Base_T = CNR	CNR_Base_P = CNR
Temperature (year)	2000	2030	2030	2000	2000	2030
Precipitation (year)	2000	2030	2030	2000	2030	2000
Carbon dioxide concentration (ppm)	381	450	381	450	450	450

CNR = Centre National de Recherches climate scenario.
CNR_No_CO₂ = CNR temperature and precipitation, baseline CO₂ concentration.
CNR_JUST_CO₂ = Baseline temperature and precipitation, elevated CO₂ concentration.
CNR_Base_T = CNR climate scenario precipitation and elevated CO₂ concentration, with baseline temperature.
CNR_Base_P= CNR climate scenario temperature and elevated CO₂ concentration, with baseline precipitation.
Ppm = Parts per million.

Appendix figure 1

U.S. average crop yields under various climate sensitivity scenarios

Average national crop yield
(units/acre)

Units/acre for silage, cotton, and hay

CNR = Centre National de Recherches climate scenario.

CNR_No_CO₂ = CNR temperature and precipitation, baseline CO₂ concentration.

CNR_JUST_CO₂ = Baseline temperature and precipitation, elevated CO₂ concentration.

CNR_Base_T = CNR climate scenario precipitation and elevated CO₂ concentration, with baseline temperature.

CNR_Base_P = CNR climate scenario temperature and elevated CO₂ concentration, with baseline precipitation.

Cwt = Hundredweight.

Source: USDA, Economic Research Service calculations

Appendix figure 1 reveals the magnitude of carbon dioxide fertilization impact by crop, as well as the relative impact of temperature change versus precipitation change in the presence of carbon dioxide fertilization. As expected, carbon dioxide fertilization alone (CNR_JUST_CO₂) resulted in crop productivity increases for all crops relative to the baseline case (appendix table 2). The aggregate numbers masked differences in carbon dioxide fertilization impacts across regions. Appendix table 3 illustrates those differences for corn and soybeans.

A comparison of CNR_T_Base_P and CNR_P_Base_T results illustrates the relative impact of temperature change versus precipitation change on biophysical impacts, when averaged nationwide and in the presence of the carbon dioxide fertilization effect. For some crops, the precipitation change associated with the CNR scenario actually increased productivity relative to carbon dioxide fertilization alone; that dynamic was seen for rice and silage, for instance. Silage, with concentrations of production in the Northeast and Lake States regions, benefited from the increased precipitation in those regions associated with the CNR projection.²

Applying the temperature changes associated with the CNR model (CNR_T_Base_P) almost always decreased productivity relative to the carbon dioxide fertilization only case (CNR_JUST_CO₂). There was very little temperature impact on barley and hay, and cotton yields increased significantly with the temperature increase. Cotton yield's positive response to temperature change was not consistent with the literature and possibly was a result of a disparity between the optimal temperature for leaf and

²Rice reacts counterintuitively; with production concentrated in the Delta region, rice thrives despite lower precipitation levels. Because rice is heavily irrigated, one could imagine little reaction to precipitation change, but a positive yield change seems an unlikely result. We are currently improving our treatment of rice irrigation and its interaction with REAP's soil types for future analyses with a more sophisticated focus on water resources.

Appendix table 2

Change, by crop, in nationwide average yield as a result of carbon dioxide fertilization in the CNR scenario

Crop	Percent yield change
Barley	8.1
Corn	2.9
Oats	6.9
Rice	5.3
Sorghum	4.7
Soybeans	10.0
Wheat	5.8
Silage	4.7
Cotton	9.7
Hay	15.7

CNR - Centre National de Recherches general circulation model.

Source: USDA, Economic Research Service calculations based on Regional Environment and Agriculture Programming (REAP) model data.

Appendix table 3

Regional differences in carbon dioxide fertilization effect for corn and soybeans

Region	Corn yield	Soybean yield
<i>Percent change</i>		
Appalachia	1.5	9.7
Corn Belt	1.8	10.3
Delta	1.8	8.1
Lake States	5.3	9.1
Mountain	5.1	N/A
Northern Plains	4.4	12.1
Northeast	3.5	10.0
Pacific	3.4	N/A
Southeast	2.3	11.0
Southern Plains	3.0	11.0

N/A – Not applicable.

Source: USDA, Economic Research Service calculations based on Regional Environment and Agriculture Programming (REAP) model data.

vegetative growth (37° C) and the optimal temperature for cotton boll growth (25-26° C). The EPIC analysis did not capture a decline in harvest index at the higher temperature and therefore translated higher biomass production at higher temperatures into higher yields.

The aggregate crop productivity impacts at both the regional and national level are weighted averages of what is occurring at the field scale for each of REAP's production enterprises, so the magnitude of change is not necessarily representative of what happens for any single rotation. The results for corn production in the Corn Belt, for instance, are an average of what would happen to corn yields in a continuous corn rotation and in a corn/soybean rotation (among others). Because the yield impacts of any single element of climate change are dependent on other factors in the crop production system, particularly water and nutrient constraints, those impacts vary significantly across production enterprises for the same crop within a single region.

Appendix D: An Econometric Model of Regional Pesticide Expenditures

Movement of Crop Production Regions in Latitude-Temperature Space

The premise of the pest impact analysis in our study was based on the assumption that rising temperatures with climate change will make production conditions—and potential pest infestations—more comparable with observed conditions farther south. Quadratic relationships between average annual temperature and latitude centroid for REAP crop production regions for 2000 and 2030 were estimated using temperature projections from the ECH, CSIRO, CNR, and MIROC climate models applied to the Special Report on Emissions Scenarios (SRES) A1B emissions scenario. These relationships were used to estimate the effective southward percentage shift in each region's latitude centroid in temperature space for each climate-model projection.

Average annual temperature was regressed on a constant latitude centroid, and latitude centroid squared for current weather conditions and for each model 2030 projection using ordinary least squares (OLS) (appendix table 4). As shown, latitude accounted for 93 percent of the variation in average annual temperatures for each of the climate models. Additionally, coefficient estimates were statistically different from zero at the 1-, 5-, and 10-percent levels, except for the square of the latitude centroid in the CSIRO ($p=0.195$) and MIROC ($p=0.116$) climate models.

The relationships between average annual temperature and latitude centroid are displayed graphically for the base year (2000) and for each climate model for 2030 in appendix figure 2. Except for the lower latitudes (below 32°N), the largest increases in average annual temperature occurred using the MIROC temperature projections, and the lowest increases in average annual

Appendix table 4

Ordinary least-squares estimates for the quadratic relationships between average annual temperature and REAP region latitude centroids

Variable	Base year ^a	ECH ^b	CSIRO ^c	CNR ^d	MIROC ^e
Intercept	60.9399***	61.097784***	56.707154***	61.776466***	58.6598***
Latitude centroid	-1.6851***	-1.623285***	-1.407831***	-1.631187***	-1.486675***
Latitude centroid ²	0.0107*	0.009895	0.007144*	0.009859	0.008593***

*=Statistically different at the 1-percent level.

**=Statistically different at the 5-percent level.

***=Statistically different at the 10-percent level.

Notes: Average annual temperatures in each of the 48 REAP regions for a 2000 base year and 2030 temperature projections based on the ECH, CSIRO, CNR, and MIROC climate models applied to the SRES A1B emissions scenario were regressed on a vector of ones and the REAP regions' latitude centroids and latitude centroids squared. The least-squares estimates are reported in this table. The standard errors do not account for the use of temperature projections for 2030. Forty-eight observations were used in each regression.

^aThe standard error of the estimate, σ , is 1.28, the coefficient of determination, R^2 , is 0.93, and the hypothesis that all coefficient estimates are zero can be rejected at the 1-percent level, $F=320.63$ ***.

^b $\sigma=1.16$, $R^2=0.95$, and $F=391.12$ ***.

^c $\sigma=1.20$, $R^2=0.94$, and $F=361.97$ ***.

^d $\sigma=1.25$, $R^2=0.94$, and $F=347.42$ ***.

^e $\sigma=1.18$, $R^2=0.94$, and $F=344.70$ ***.

Quadratic relationships between average annual temperature and latitude centroid for the base year and for the climate models in 2030

Average annual temperature (degrees Celsius)

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Source: USDA, Economic Research Service calculations

temperature occurred using the CSIRO temperature projections. Except for the lower latitudes, average annual temperatures for the CNR, CSIRO, and ECH climate projections were between base year levels and MIROC temperature projections. The largest increases in average annual temperature below roughly 29°N occurred using the CNR temperature projections.

The simulated southward movement of each REAP region's latitude centroid implied by the OLS estimates is reported in appendix table 5. These figures represent the estimated southward shift of each region in latitude-temperature space. Empirical estimates that relate latitude to real (2006 US\$) pesticide expenditures were applied to each region's adjusted latitude to predict percent changes in expenditures and yield losses in 2030 relative to 2000.

Pesticide Expenditures and Latitude

Phase 2 Agricultural Resource Management Survey (ARMS) data collected at the field level from barley (2003), corn (1996, 2001, 2005), cotton (1997, 2003, 2007), oat (2005), rice (2000, 2007), sorghum (2003), soybean (1997, 2002, 2006), and wheat (2004) producers were used to estimate a relationship between the natural log of pesticide expenditures per acre, a constant, a time index, regional fixed effects, and a separate latitude coefficient for each crop.¹ The time index was included to account for annual weather variation during 1996-2007. National Agricultural Statistics Service (NASS) production-region fixed effects (not REAP region fixed effects) were included in the regression to account for spatial variation in weather and production possibilities.² The dummy variable for the Corn Belt was excluded.³ Separate coefficient estimates were obtained for each crop by adding independent variables equal to the product of the field's latitude and the crop dummy variables.

¹Pesticide expenditures, which include payments for herbicides, insecticides, and fungicides, were converted to real (2006 US\$) expenditures using the Bureau of Labor Statistics (2010) producer price index for pesticides and other agricultural chemicals.

²In another version of the model, separate annual dummy variables were included for each NASS production region to account for annual weather variation by region and over time more completely. Because the adjusted coefficient of determination was slightly lower in that version of the model and because the estimation results were very similar, we used the results reported in appendix table 6 to specify the pesticide-expenditure and yield-loss impacts used in the analysis.

³Note that the REAP regions are subsets of the major production regions included in the regression.

Appendix table 5

REAP region southward shifts in latitude-temperature space

REAP region	Latitude centroid	Climate Models			
		CNR	CSIRO	ECH	MIROC
			new latitude centroid		
APN	37.13	35.31	35.83	35.65	34.66
APP	35.90	34.12	34.65	34.46	33.55
APS	38.77	36.90	37.42	37.23	36.13
APT	35.89	34.10	34.63	34.44	33.53
CBL	41.18	39.23	39.79	39.55	38.31
CBM	40.66	38.72	39.27	39.04	37.83
CBN	37.93	36.09	36.61	36.42	35.38
CBO	36.63	34.83	35.36	35.17	34.21
CBR	41.08	39.13	39.70	39.45	38.22
DLN	35.61	33.84	34.37	34.18	33.28
DLO	33.56	31.86	32.44	32.21	31.45
DLP	33.47	31.78	32.35	32.12	31.37
DLT	30.25	28.67	29.35	29.03	28.47
LAF	47.58	45.43	46.33	45.70	44.08
LAK	44.87	42.80	43.51	43.10	41.64
LAL	42.98	40.97	41.59	41.28	39.93
LAM	44.19	42.15	42.82	42.45	41.03
MNB	43.32	41.30	41.94	41.61	40.24
MND	39.34	37.44	37.97	37.77	36.64
MNE	45.62	43.53	44.28	43.82	42.31
MNF	48.37	46.21	47.18	46.48	44.81
MNG	42.24	40.25	40.84	40.56	39.26
MNH	36.48	34.68	35.21	35.02	34.07
NPF	46.78	44.65	45.49	44.94	43.36
NPG	43.17	41.16	41.78	41.46	40.10
NPH	39.36	37.47	38.00	37.80	36.67
NPM	41.55	39.58	40.16	39.90	38.64
NTL	42.99	40.98	41.60	41.28	39.93
NTN	40.50	38.56	39.11	38.89	37.69
NTR	42.78	40.77	41.39	41.08	39.75
NTS	40.11	38.19	38.74	38.52	37.34
NTT	38.79	36.91	37.44	37.24	36.15
PAA	45.07	43.00	43.72	43.30	41.82
PAB	46.54	44.42	45.23	44.70	43.14
PAC	36.95	35.13	35.66	35.47	34.49
PAD	39.65	37.74	38.28	38.07	36.92
PAE	48.31	46.14	47.10	46.41	44.74
SPD	31.09	29.48	30.12	29.83	29.22
SPH	34.41	32.69	33.24	33.03	32.21
SPI	28.41	26.88	27.64	27.24	26.81
SPJ	33.70	32.00	32.57	32.35	31.58
SPM	36.22	34.43	34.96	34.77	33.83
SPP	32.71	31.04	31.63	31.39	30.68
SPT	29.16	27.62	28.34	27.97	27.49
STN	34.49	32.76	33.31	33.10	32.28
STP	32.29	30.63	31.23	30.98	30.30
STT	32.92	31.24	31.83	31.59	30.87
STU	27.56	26.07	26.87	26.43	26.05

See table 4 for the sources of the CNR, CSIRO, ECH, and MIROC models.

Note: Figure 4 displays the location of these regions.

This econometric model was used to test the hypothesis that problems with agricultural pests are more severe in southern crop production regions than in northern production regions (Bridges, 1992). Our analysis provided empirical support for the hypothesis for barley, corn, oats, sorghum, soybeans, and wheat; empirical support against this hypothesis for cotton; and empirical support neither for nor against this hypothesis for rice. The maximum likelihood estimates (appendix table 6) indicate a negative and statistically significant relationship between pesticide expenditure and latitude for barley, corn, oats, sorghum, soybeans, and wheat. The estimates indicate a positive and statistically significant relationship for cotton and a positive and insignificant relationship for rice. The estimates suggest that pesticide expenditures in 2030 will increase relative to 2006 for barley, corn, oats, sorghum, soybeans, and wheat; decline for cotton; and remain the same for rice.

Appendix table 6

Pesticide costs per acre versus latitude, by crop, maximum likelihood estimates for the groupwise heteroscedasticity model

dependent variable	Natural log of pesticide expenditures per acre	
observations	19,701	
standard error	1.00	
Adjusted R ²	0.43	
F	840.59***	

Variable	Coefficient	Standard error
intercept	4.1915***	0.0979
year	-0.0386***	0.0015
latitude-barley	-0.0311***	0.0023
latitude-corn	-0.0186***	0.0024
latitude-cotton	0.0066**	0.0026
latitude-oats	-0.0460***	0.0027
latitude-rice	0.0030	0.0027
latitude-sorghum	-0.0250***	0.0026
latitude-soybeans	-0.0267***	0.0025
latitude-wheat	-0.0245***	0.0024
Appalachia	-0.1022***	0.0208
Delta	-0.0815	0.0250
Lake States	-0.0295	0.0181
Mountain	-0.2017***	0.0311
Northern Plains	-0.2066***	0.0159
Northeast	0.0313	0.0312
Pacific	0.1845***	0.0366
Southern Plains	-0.7614***	0.0301
Southeast	-0.0744**	0.0328

*=Statistically significant at the 1-percent level.

**=Statistically significant at the 5-percent level.

***=Statistically significant at the 10-percent level.

Notes: Phase 2 ARMS data for barley (2003), corn (1996, 2001, 2005), cotton (1997, 2003, 2007), oats (2005), rice (2000, 2006), sorghum (2003), soybeans (1997, 2002, 2006), and wheat (2004) were used. Iterated weighted least squares was used assuming groupwise heteroscedasticity, where the groups were the different crops. Ten iterations were required. The null hypothesis of homoscedasticity was rejected ($\chi=1418.12$, $df=7$, $p<0.0001$)