



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

**Fifth Joint Conference on
Agriculture, Food and the Environment**

Proceedings of a Conference Sponsored by
University of Minnesota
Center for International Food and Agricultural Policy

Universita degli Studi di Padova
Dipartimento Territorio e Sistemi Agro-forestali

Agricultural Development Agency – Veneto Region

University of Perugia

University of Bologna – CNR

**SESSION V: COMPUTER SCIENCE AND ENVIRONMENTAL
MANAGEMENT**

**PAPER 3: MANURE APPLICATION PLANNER (MAP):
CONVERSION AND USE IN ITALY**

Antonio Boggia and Wynn Richardson

Center for International Food and Agricultural Policy

University of Minnesota
1994 Buford Avenue, 332 C.O.B.
St. Paul, Minnesota 55108-6040 U.S.A.
Phone: (612) 625-8713
FAX: (612) 625-6245

**Fifth Joint Conference on
Agriculture, Food, and the Environment**

Proceedings of a Conference Sponsored by

University of Minnesota
Center for International Food and Agricultural Policy

Università degli Studi di Padova
Dipartimento Territorio e Sistemi Agro-forestali

Agricultural Development Agency - Veneto Region

University of Perugia

University of Bologna - CNR

Abano Terme - Padova, Italy
June 17-18, 1996

Working Papers are published without a formal review within or the endorsement of the Center for International Food and Agricultural Policy or Department of Applied Economics.

The University of Minnesota is committed to the policy that all persons shall have equal access to its programs, facilities, and employment without regard to race, color, creed, religion, national origin, sex, age, marital status, disability, public assistance status, veteran status, or sexual orientation.

Information on other titles in this series may be obtained from Waite Library, University of Minnesota, Department of Applied Economics, 1994 Buford Avenue, 232 COB, St. Paul, MN 55108-6040, U.S.A.

University of Padua

University of Minnesota

University of Perugia

Agricultural Development Agency

**University of Bologna
CNR**

**5th JOINT CONFERENCE ON AGRICULTURE, FOOD AND THE
ENVIRONMENT**

<p>Manure Application Planner (MAP): Conversion and Use in Italy</p>

by

Antonio Boggia
Istituto di Estimo Rurale e Contabilità
University of Perugia

Wynn Richardson
Center for Farm Financial Management
University of Minnesota

Abano Terme - Padova, 17th - 18th June 1996
ITALY

MANURE APPLICATION PLANNER (MAP): CONVERSION AND USE IN ITALY

Antonio Boggia and Wynn Richardson*

1. Introduction

Many farmers are concerned about the environmental impact of their farming operations. For livestock producers, a primary concern is proper manure utilization and handling. A manure application plan needs to be developed by livestock producers as a means of accountability for their manure management. A manure management plan is a specific set of practices and recommendations developed for a specific farm. These plans would include such items as the application rate, method of application, time of application, etc. By having this type of plan, manure applications could be made that are appropriate and acceptable according to the set of conditions for a given locale.

Producers are equally concerned, however, about the financial status of their farm business. Any manure management plan must also include an economic assessment as well as traditional agronomic considerations.

The Manure Application Planner (MAP), version 3.0 was written and released by the Center for Farm Financial Management at the University of Minnesota to address both the agronomic and economic components of a manure management plan. The software utilizes a mixed-integer form of linear programming, which is a proven optimization routine for manure planning (Schmitt et al., 1994a) and other agricultural applications (Beneke and Winterboer, 1973). This optimization allows an application plan to be easily developed if a farmer does not have an existing application plan. However, if the producer does have an existing plan, the software can also be used to assess the plan and easily make modifications for environmental or economic concerns.

2. History and use of MAP

2.1. History

MAP version 3.0 was developed as a joint project among the Minnesota Extension Service, the Minnesota Natural Resources Conservation Service (NRCS), and the Minnesota Pollution Control Agency (MPCA). Due to this joint support, permitting of farms according to the Minnesota feedlot ordinance and cost-share programs can be

* This paper arises from the common work of the authors in the project for the implementation of an Italian Version of MAP. W. Richardson wrote section 1, 2 and 3; A. Boggia wrote section 4, 5, 6 and 7.

validated by the software's output for the MPCA and the NRCS, respectively.

Besides the use in Minnesota, the Southwestern Illinois Resource Conservation and Development, Inc., and the Illinois Natural Resources Conservation Service also supported the development of version 3.0. These agencies, along with the Illinois Cooperative Extension Service, also use the software to assist farmers with manure application planning.

2.2. Use of Manure Application Planner

As of April, 1996, there are over 340 copies of MAP in use in the United States and Canada. The distribution of these copies by group is shown in Table 1.

Type of User	Number of Copies
Extension	128
NRCS	69
Private business	44
Public agencies	39
Farm management instructors	24
Farmers/ranchers	20
University teaching/research	18

Table 1. MAP Users in the United States/Canada

Since version 3.0 was initially released in September, 1995, the number of copies has more than doubled. This large increase in MAP use may be due in part to recent large manure spills and the resulting media coverage. As people in the United States continue to move to developments or land that border livestock operations, the interaction of farming and nonfarming residents creates an atmosphere in which manure management is a large issue. This close proximity between livestock operations and other residents also exists in Italy, thus the need for a planning tool for Italian conditions that will assist farmers in making manure management decisions.

3. Software Organization

The software is written in Borland Pascal 7.0 and requires an IBM-PC or 100% compatible computer with MS-DOS/PC-DOS 5.0 or higher. The computer also must have an 80386 or later processor, at least 2 megabytes of RAM, one 3.5 inch diskette

drive, and a hard drive with 2 megabytes of free space.

MAP allows for site-specific information to be input, thus making it adaptable for many geographical regions. After initializing the software, the main menu allows the user to go into file creation/management and then into a series of input and output menus. These main menu options will be discussed in the following paragraphs.

Manure Source Input

For whole-farm nutrient planning, the amount of nutrients contained in the farm's manure supply needs to be quantified. For each source/storage facility of manure, the nutrient content from a laboratory analysis and the amount of manure that is available for spreading on an annual basis allows the software to calculate the farm's nutrient supply. If this information is not available for input, the software can estimate manure/nutrient amounts and concentrations using daily manure/nutrient excretion data and average N loss from the storage facility (Midwest Plan Service-18, 1993).

Users of the software should be encouraged to use manure testing regularly due to the wide variation in analyses from seemingly similar livestock operations (Wagar et al., 1994). Manure volume, or quantity, should be calculated from measured storage capacities rather than excretion data because of changes during the handling and storage phases. The manure estimation feature was included primarily for new operations to provide an initial basis for manure management planning.

Field Input

Crop yields are often dependent on the supply of soil nutrients; therefore, the livestock aspects of an enterprise form the nutrient supply and the cropping aspects of the operation form the nutrient demand. Each field is unique in terms of its nutrient demand, or recommendation, because of the cropping rotation, soil test values, and production capacity. Each field's nutrient needs, independent of nutrient source, must be entered individually in the software.

These nutrient needs, or recommendations, form the starting point in calculating the fertilizer replacement value of manure that might be applied to the field. The economic savings from the manure cannot be greater than the costs of the nutrients recommended. The user has the option of using any laboratory's N-P₂O₅-K₂O recommendation. The user also has the option of allowing the software to provide information to calculate a nutrient recommendation based solely on crop removal using crop and yield levels. While this procedure is not encouraged, previous experience by the authors has been that many producers use this method in determining nutrient needs.

To establish environmental guidelines for manure/nutrient applications for each field, the user selects the crop to be grown on a field, which in turn indicates whether the crop is a legume or not, and enters the maximum amount of phosphorus (P_2O_5) that can be applied to the field. The legume information then becomes the initial criterion for N application limits on a field. If the crop is a non-legume, N will be applied to meet the recommendation with no excess available N allowed. If the crop is a legume, the N recommendation is generally zero, but the software then allows manure to be applied to meet the P_2O_5 recommendation under the presumption that the legume will take up the added N with a compensatory decline in N_2 fixation (Peterson and Russelle, 1991; Schmitt et al., 1994b).

The software defaults to a P-based application rate, which allows the maximum P_2O_5 amount being applied to equal the recommended amount of P_2O_5 . However, users can decide if the manure application rate should be N- or P_2O_5 -based, or somewhere in between by entering a maximum P_2O_5 allowable limit. Producers should follow local guidelines in restricting excess P_2O_5 applications when near surface waters, when erosion potential is high, and/or when soil P test values are very high.

An optional entry for each field is selecting the method of manure application. The method of manure application has a direct effect on N loss and N availability, especially in the year of application (Figure 1).

Application Methods Screen 1 of 4

Application Method	Manure Type Beef			% Available	
	-----% N Available-----			P2O5	K2O
	Year 1	Year 2	Year 3		
Bdcast, inc < 12 hrs	60	25	10	80	90
Bdcast, inc < 4 days	45	25	10	80	90
Broadcast, no incorp	25	25	10	80	90
Knife injection	50	25	15	80	90
Sweep injection	60	25	10	80	90
	-	-	-	-	-

PgDn Next Screen
F1 Help

PgUp Prev Screen
<Ctrl> A Exit, Don't Save

<Alt> B Delete
F10 Menu

Figure 1: Example of application methods screen.

While the software has the option of selecting a method of manure application for each field based solely on the N recommendations and economics, we expect that most producers will select specific methods for their fields based on their method of manure handling, the crop, time of application, and/or available equipment.

The distance between the field and the manure source also must be entered for each field. This distance will be important in prioritizing manure applications because of the associated hauling costs.

Economic Input

Economic input is critical in the formulation of the manure management plan created by the software. Fertilizer costs for N-P₂O₅-K₂O are entered as well as a representative application charges for these materials. Combining these costs and the fertilizer recommendations for each field provide the maximum economic value of the manure for each field.

Manure application and hauling costs also should be entered in this section. Manure application costs are a function of application equipment, incorporation operations (if any), and size of equipment, whereas the hauling costs are dependent on the distance to the field and size of tractor. Defaults are provided based on economic datasets for equipment and labor costs (Lazarus, 1996) such that application costs are simply based on method of application/incorporation and hauling costs are dependent on distance to the field. We expect and encourage users to modify these costs to fit the specifics of each individual farm.

Optimization Process

After completion of data entry for the manure supply, crop demand, and economics sections, the software calculates manure application rates for all fields concurrently to optimize the economic value of the manure. This optimization uses a mixed-integer programming algorithm to meet the constraints set forth in the software. These constraints are: 1) the fertilizer recommendations must be met, either from manure or commercial fertilizer sources; 2) for non-legume crops, the amount of N applied cannot exceed the N fertilizer recommendation; 3) the amount of P₂O₅ applied cannot exceed the maximum allowable limit for the field, which may be the P₂O₅ recommendation depending on the maximum allowable limit specified; and 4) as much of the manure as possible must be applied. The objective function of the optimization is to minimize all costs, both commercial fertilizer and manure application and hauling costs.

In meeting these constraints, an initial manure management plan is formulated

for a farm. The majority of users, however, will want to fine-tune parts of this plan to meet some of the personal or logistic constraints unique to that operation. Therefore, each field's rate can be adjusted, with the resulting effects on nutrient excesses or supplemental fertilizer amounts depicted in the output section.

If the producer so chooses, the entire optimization process can be by-passed, with manure application rate and application method being entered along with the field input information. This option can be very useful in evaluating environmental and economic parameters of an existing plan.

Manure Application Output

All of the output can be provided to the user on the screen and/or via the printer. The output information is grouped by field-specific or farm-specific data. The field-specific output includes the manure application rate, supplemental fertilizer rates, if needed, and the quantity of excess nutrients applied, which are defined as the amount of available nutrients applied in excess of what the fertilizer recommendations are for that field. An economic summary also is calculated for each field with two scenarios. In the first, the economic summary is calculated according to the prescribed manure application plan. In the second scenario, the economic summary is calculated assuming no manure is available. The difference between these two scenarios shows the economic implications of using manure as a fertilizer source.

Whole-farm output information is primarily a compilation of the field-by-field information. A main, important exception to this is manure quantity produced/available on a farm compared to its recommended usage. If there is manure "left over" after all of the fields are accounted for, there becomes an environmental concern for this enterprise. Whole-farm economic summaries and supplemental fertilizer quantities needed are also presented.

The printed output also lists some of the parameters used to arrive at the final plans, such as nutrient availability coefficients, field nutrient recommendations, and manure production information. Because all of these parameters can be accessed and changed by the user, the printout allows for verification of these parameters.

4. Manure management issues in Italy

Wherever in Italy there is a discussion regarding the environmental problems connected with livestock, it always comes to the conclusion that a better, proper management of solid and liquid manure is needed. This means basically, that however much one looks at the negative environmental aspects, with livestock everything revolves

around the problem of manure. And this problem is not just connected with the large volumes produced, but also, and especially, with their chemical-physical composition. Besides the odor and ammonia, the greatest worry is with soil and water pollution. This is where the necessity arises for the environmental planning and management of livestock.

Much is said about the nitrogen content of manure and the dangers of it leaching into groundwater. This is undoubtedly a risk to be taken into consideration; however, there is increasing debate as well about the danger of soil and watershed contamination from the growing percentage of heavy metals present, especially for hogs, due to the new trends in feeding. Another important question is that of nutrients. The manure problem is centered around the debate over what is the best form of treatment and disposal from an environmental point of view. Despite new proposals on the subject, at the present time it is impossible to establish what method of disposal best respects the environment. The trend, however, is to give greater attention to feed - it is obvious that the waste is highly dependent on what goes into the animal.

Very strictly connected with manure management is the problem of the odours. The situation is quite serious in Italy, if one considers that in some cases livestock farms are right next to towns. The lack of adequate regulations logically results in the matter being taken to court, where much is left to the discretionary power of the judges.

There are various options for manure management and treatment, but it should be said that wherever it is possible, especially where a careful, balanced diet is provided and water is used prudently, it is still preferable to use liquid and solid manure as fertilizer. This utilization of manure allows animal-breeding to be brought into closer connection with the land and crops. Of course this type of utilization strictly depends on quantities and soil type, and requires careful management. If the manure is not properly stored, processed and applied, the nutrients and fertilizing elements can be lost and sometimes become air, water and soil pollutants. Incorporating manure directly into the soil and applying it when the prevailing winds are not blowing towards populated areas may be other useful precautions. It is always important to treat liquid waste at the farm prior to spreading. Here, too, there are different possible choices, though this is not the appropriate place to list them. Generally, however, it is important to remember that the first step in the proper management of manure for spreading in the field is to have enough storage volume to make sure that spreading is done at the right times during the year, and not being forced to empty the tanks due to a lack of space.

In the past there was in Italy very much attention on an alternative to spreading manure in the field: the treatment of manure in centralized plants, generally cooperative

purifiers. The problem, however, is that these plants do not eliminate the problem of spreading, because in any case the final product is part conditioned sludge and part water to be used for fertilization by irrigation. Other plants seem to be able to produce water which is practically drinkable, but the cost for treatment is very high. Generally, the main problem in all of these centralized plants is the high costs, which have negative repercussions on the producers.

At present, very much attention is given to the application of manure on the fields.

5. Need for MAP in Italy and software conversion

The protecting of physical resources in both urban and rural areas has been for some time the main objective of an environmental policy which aims at restoring and/or improving "environmental quality". This goal can be reached more easily today than in the past, due to the decreasing importance of agriculture devoted solely to greatest possible production of foodstuffs. This is the general framework of the activities of the Italian National Research Council (CNR) in the project RAISA, that is the funding source of the research that includes the Italian conversion of MAP.

The territorial reference unit for the research is the watershed of the Tiber River. The Tiber River watershed covers an area of 17,156 km². In it there are several natural and artificial lakes. Altogether, the area covered by the basin involves 6 regions, 14 provinces and 347 communes, and is comprised of vast areas of mountains or high hills, predominated by woods and pastures, and some middle-low hills and plains areas, where there are intensive farming and livestock activities, especially in the plains along the river and around the lakes. Pig-breeding is one of the most important among livestock activities, with an average of 420,000 head (1990) being raised in the basin. The highest concentration of animals (81%) in the basin are raised in the two Umbrian provinces of Perugia and Terni. The idea of converting MAP to Italian, comes from all the things listed above and in section 4. First of all, the new trend in using manure as a fertilizer. Since there are different rules for manure application on the fields in the territory of almost every different local Environmental Protection Agency, it would be very important to make available a tool for manure management, that can give sound directions and recommendations for manure applications in all the regions. This would avoid that two close livestock enterprises, but one under an Environmental Protection Agency, and another one under another Agency, have to follow different rules for manure management, sometimes with serious economic damages, such as often happen in Italy. So, starting from swine, but in a second time extending to all livestock types,

the Istituto di Estimo Rurale e Contabilita' of the University of Perugia, funded by the above mentioned CNR-RAISA, decided, together with the Center for Farm Financial Management, University of Minnesota, to convert MAP into Italian, to get an Italian version.

The first step in the conversion process involved translating into Italian all the input text, output text, and context-sensitive help messages. Translating the input and output text was accomplished using a software utility program called TRANSPAS. This utility program was developed at the Center for Farm Financial Management, and allows a person familiar with MAP to translate the text seen by the user without having to modify any of the MAP source code. Using this approach only one copy of source code is maintained, so any improvements or corrections made to MAP will be done for all language versions of the software. Translating the help messages simply involved editing the help document in a word processor such as WordPerfect or Microsoft Word.

The second step in the conversion process involved editing the MAP databases (Figure 2).




Figure 2: Map databases

MAP relies on databases to simplify user input, and these were changed for conditions in Italy. An individual user can still customize the databases for a particular region or

farm situation.

6. Case Farm

The farm used for this test of the Italian MAP, is a typical farm of Umbria, with a livestock enterprise, and not very much land, with some different types of crops. In this case, the livestock type is dairy. 70 dairy cows weighing 650kg, and 70 dairy calves, in different stage of weight are present in the farm. The land consists of 47 hectares, and main crops are listed in table 2.

Crops	Hectares
Corn	14.3
Barley	12.5
Sunflowers	2
Wheat	2
Vine	3.3
Olive tree	1.75
Alfalfa	8.6
Set-aside	2.55
TOTAL	47

Table 2: Crops and their area surfaces.

The breed of the cows is Holstein-Friesian. The animals weigh about 50 kg when they are born, and come to 650 kg when they are in full production.

The land is mostly flat, and the soil test remarks that it is clayey, with a good level of organic matter, N, P₂O₅ and K₂O.

The surface area of the farm is divided in 12 fields, and the distance from manure source vary from 20 meters to 3 kilometers. dairy manure is applicated on the fields, apart from the alfalfa field and the set-aside field. So, the available fields for manure application are 10.

The storage system for manure is above ground tank. The manure production has been estimated in 1680 t per year, and the nutrients content per unit of manure has been estimated in 3 kg of N, 1kg of P₂O₅ and 4 kg of K₂O.

After references, is enclosed the copy of the print of the input, that include manure sources, fields and input prices. Per each field it is indicated the name, surface area, crop, nutrient requirements. It is also indicated the result of the optimizer calculations, in terms of application method, manure rate, and distance from the source.

About the nutrient requirements, it must be underlined that it has been indicated only the removal rate of the crop, since the soil test have shown a very good nutrients content.

The print of the output is also enclosed after references. It shows per each field the application method selected by the optimizer, the manure rate, the amount of nutrients per hectare. All the fields need an addition of commercial fertilizer, especially for N and P_2O_5 . The excess nutrient is very common for K_2O , but not for N and P_2O_5 . The manure balance is zero, which means that all manure is used. It is very interesting to look at the source of nutrients: it comes out that most of the nitrogen required by crops is taken from commercial fertilizer, and also a good part of phosphate. About the availability of nutrients, it depends upon the application method, so that there is a very different situation between broadcast, no incorporated and broadcast, incorporated in 12 hours, at least for nitrogen. For phosphate and potash there are no differences.

Coming to the economic output, it comes out that for all the fields apart from no.4 and 5, the total cost per hectare of fertilization with manure is lower than without manure, even if the total for all the 35.85 hectares is a little higher for with manure situation (299,920 lire versus 294,611). This makes even more interesting and effective the use of manure for land fertilization. In fact, using manure is for sure, if well managed, a sustainable environmentally sound farming technique, and can help to solve the problem of manure disposal. If in addition it can be also valuable from the economic point of view, very similar to fertilization only with commercial fertilizer, it becomes extremely interesting for farmers.

MAP is the right tool to manage and monitor all these things, so that manure management can be not only a problem to solve, but also and mainly a way of organizing and optimizing farm resources.

7. Conclusions

As showed in the case farm, MAP can be used for setting up a manure application plan for farms. Thanks to this plan among the others, it is possible to avoid N excess, so that the risk of water and soil pollution can be minimized.

In Italy at present farmers with livestock enterprises are having very hard time, because there are not common regulations for manure management, and the Local Environmental Protection Agencies can close the enterprise at any time, if they find that manure is going to be a problem for the environment and human health. But definitely livestock is still a good source of income for farmers, so it would not make sense to close livestock enterprises to avoid the risk of problems with Environmental Agencies. Since MAP can optimize not only the economic aspect of manure management, but also the environmental side, it can become the guideline for manure application: if for example Local Environmental Protection Agencies would ask for an application plan, MAP could be used. The farmers would have only to respect what is in the MAP output, and they would not take any risk.

The intention of the Istituto di Estimo Rurale e Contabilita' and the Center for Farm Financial Mangement is not only to develop the Italian version for research aims, but they believe that this can be a practical tool, that can be very helpful and play a role for solving the problem of manure management in Italy.

The first step towards this goal will be the presentation of the Italian version of MAP to Extension Agencies, Local Environmental Protection Agencies, and Livestock Producers Associations, which most probably will be held in July. After that, a book on environmental impact of livestock and manure management enclosing a demo disk of MAP, will be diffused in Italy.

MAP is designed to solve a so specific problem, that it can really be the way to allow to farmers to have definite answers about the legitimacy of their activity, and to decision makers to have available a standard model and a control tool objective and equal for different areas.

All these considerations sound like a wish for MAP to meet a good impression by all the people working in the livestock sector, so that it will be able to give a real contribution to the solution of a very important problem for Italian agriculture.

References

Abbozzo P., Mennella V.: 1993, L'impatto delle agrotecnologie nel Bacino del Tevere. Criteri per la valutazione di impatto ambientale, Franco Angeli, Milano.

Abbozzo P., Boggia A., Brunetti M., Environmental quality and hog production, Journal of Environmental Management and Assessment, Kluwer, vol. 41, no. 2, June 1996.

Beneke, R.R., and R. Winterboer. 1973. Linear programming applications to agriculture. Iowa State Univ., Press, Ames, IA.

Lazarus, W. 1996. Minnesota farm machinery economic cost estimates for 1996. Dept. of Appl. Econ., Univ. of Minn., St. Paul.

Midwest Plan Service Committee. 1993. Livestock waste facilities handbook. 3rd ed. Midwest Plan Serv., Iowa State Univ., Ames, IA.

Peterson, T.A., and M.P. Russelle. 1991. Alfalfa and the nitrogen cycle in the Corn Belt. J. Soil Water Conserv. 46:229-235.

Schmitt, M.A., R.A. Levins, and D.W. Richardson. 1994a. A comparison of traditional worksheet and linear programming methods for teaching manure application planning. J. Nat. Resour. Life Sci. Educ. 23:23-26.

Schmitt, M.A., C.C. Sheaffer, and G.W. Randall. 1994b. Manure and fertilizer effects on alfalfa plant nitrogen and soil nitrogen. J. Prod. Agric. 7:104-109.

Schmitt, M.A., D.W. Richardson, et. al. 1995. Manure application planner version 3.0 user's manual. Center for Farm Financial Management, Univ. of Minn., St. Paul.

Wagar, T., M. Schmitt, C. Clanton, and F. Bergsrud. 1994. Livestock manure sampling and testing. 4 pgs. Minn. Ext. Serv. FO-6423, Univ. of Minn., St. Paul.

Manure Application Planner 3.0
 Center For Farm Financial Management
 (C)1995 University Of Minnesota
 User: I.M. Learning
 Wabasha County Extension

Azienda esempio
 Passaggio di Bettona (PG)
 File: PASSAGGI
 Data: 12 Giugno, 1996

*** FONTI DELLE DEIEZIONI ***

Tipo allevamento	Bovini latte
Metodo stoccaggio	Vasca esterna
Descrizione	Frisone
Unita' di misura	t
Deiezioni disp.	1680 t
Kg N per unita'	3
Kg P205 per unita'	1
Kg K20 per unita'	4

Valori elem. nutritivi presi da analisi deiezioni?No

*** APPEZZAMENTI ***

Nome appezz.1 - Vigneto
 Ettari 2,3
 Coltura Vite

Fabbisogno elem. nutr./ha	Vincoli del Modello
Kg N 60	Metodo spandimento Scelte modello
Minimo kg P205 100	
Massimo kg P205 120	
Kg K20 120	

Fonte Deiezioni	Metodo Spandimento	Quantita'/ettaro	Dist. dalla fonte
Bovini latte	Spargim. no interra.	100 t	0,02 km

Nome appezz.2 - Campetta
 Ettari 3
 Coltura Mais

Fabbisogno elem. nutr./ha	Vincoli del Modello
Kg N 280	Metodo spandimento Scelte modello
Minimo kg P205 70	
Massimo kg P205 80	
Kg K20 52	

Fonte Deiezioni	Metodo Spandimento	Quantita'/ettaro	Dist. dalla fonte
Bovini latte	Spargim.int.< 12 ore	100 t	0,05 km

*** APPEZZAMENTI ***

Nome appezz.3 - Cerreto
Ettari 10,5
Coltura Orzo

Fabbisogno elem. nutr./ha	Vincoli del Modello
Kg N 120	Metodo spandimento Scelte modello
Minimo kg P205 30	
Massimo kg P205 40	
Kg K20 30	

Fonte Deiezioni	Metodo Spandimento	Quantita'/ettaro	Dist. dalla fonte
Bovini latte	Spargim.int.< 12 ore	50 t	0,03 km

Nome appezz.4 - Oliveto
Ettari 1,75
Coltura Olivo

Fabbisogno elem. nutr./ha	Vincoli del Modello
Kg N 100	Metodo spandimento Scelte modello
Minimo kg P205 50	
Massimo kg P205 80	
Kg K20 110	

Fonte Deiezioni	Metodo Spandimento	Quantita'/ettaro	Dist. dalla fonte
Bovini latte	Spargim.int.< 12 ore	60,6 t	2 km

Nome appezz.5 - Spaccio
Ettari 5,3
Coltura Mais

Fabbisogno elem. nutr./ha	Vincoli del Modello
Kg N 280	Metodo spandimento Scelte modello
Minimo kg P205 70	
Massimo kg P205 80	
Kg K20 52	

Fonte Deiezioni	Metodo Spandimento	Quantita'/ettaro	Dist. dalla fonte
Bovini latte	Spargim.int.< 12 ore	64,3 t	2,5 km

*** APPEZZAMENTI ***

Nome appezz.6 - Cappuccinelle
Ettari 6
Coltura Mais

Fabbisogno elem. nutr./ha	Vincoli del Modello
Kg N 280	Metodo spandimento Scelte modello
Minimo kg P205 70	
Massimo kg P205 80	
Kg K20 52	

Fonte Deiezioni	Metodo Spandimento	Quantita'/ettaro	Dist. dalla fonte
Bovini latte	Spargim.int.< 12 ore	14,4 t	2,5 km

Nome appezz.7 - Ballitto
Ettari 2
Coltura Frumento T.

Fabbisogno elem. nutr./ha	Vincoli del Modello
Kg N 180	Metodo spandimento Scelte modello
Minimo kg P205 50	
Massimo kg P205 60	
Kg K20 36	

Fonte Deiezioni	Metodo Spandimento	Quantita'/ettaro	Dist. dalla fonte
Bovini latte	Spargim.int.< 12 ore	10 t	2,5 km

Nome appezz.8 - Barca
Ettari 2
Coltura Orzo

Fabbisogno elem. nutr./ha	Vincoli del Modello
Kg N 120	Metodo spandimento Scelte modello
Minimo kg P205 30	
Massimo kg P205 40	
Kg K20 30	

Fonte Deiezioni	Metodo Spandimento	Quantita'/ettaro	Dist. dalla fonte
Bovini latte	Spargim.int.< 12 ore	8,3 t	2,5 km

*** APPEZZAMENTI ***

Nome appezz.9 - Luna
Ettari 1
Coltura Vite

Fabbisogno elem. nutr./ha	Vincoli del Modello
Kg N 60	Metodo spandimento Scelte modello
Minimo kg P205 100	
Massimo kg P205 120	
Kg K20 120	

Fonte Deiezioni	Metodo Spandimento	Quantita'/ettaro	Dist. dalla fonte
Bovini latte	Spargim.int.< 12 ore	36,4 t	2 km

Nome appezz.10 - Sole
Ettari 2
Coltura Girasole

Fabbisogno elem. nutr./ha	Vincoli del Modello
Kg N 100	Metodo spandimento Scelte modello
Minimo kg P205 45	
Massimo kg P205 55	
Kg K20 33	

Fonte Deiezioni	Metodo Spandimento	Quantita'/ettaro	Dist. dalla fonte
Bovini latte	Spargim.int.< 12 ore	9,2 t	3 km

*** PREZZI ***

Fertilizzante Commerciale

Costo per kg N	990
Costo per kg P205	805
Costo per kg K20	725
Costo spandimento	35000 per ettaro

	Includere nel Modello?	Costo Per Tonnellata Spandimento	Costo Per Tonnellata Trasporto per km	Costo Per Metro Cubo Spandimento	Costo Per Metro Cubo Trasporto per km
Fertirrigaz. pioggia	No	-	-	-	-
Fertirrigaz. scorr.	No	-	-	-	-
Iniezione nel terr.	Si	9999999	9999999	2700	600
Spargim. no interra.	Si	2000	600	2000	600
Spargim.int.< 12 ore	Si	2500	600	2500	600
Spargim.int.<4 giorni	Si	2500	600	2500	600

*** RIBPILOGO QUANTITA' DISTRIBUITE ***

Appezzamento	Ettari	Coltura	Metodo spandimento	Quantita' /ha	-- kg per Ettaro ---			kg N Anno 2
					N	P2O5	K2O	
1 - Vigneto	2,3	Vite	Spargim. no interra.	100 t	60	80	360	75
			Fertilizzante commerciale		-	20	-	
2 - Campetta	3	Mais	Spargim.int.< 12 ore	100 t	165	80	360	75
			Fertilizzante commerciale		115	-	-	
3 - Cerreto	10,5	Orzo	Spargim.int.< 12 ore	50 t	83	40	180	38
			Fertilizzante commerciale		38	-	-	
4 - Oliveto	1,75	Olivo	Spargim.int.< 12 ore	60,6 t	100	48	218	45
			Fertilizzante commerciale		-	2	-	
5 - Spaccio	5,3	Mais	Spargim.int.< 12 ore	64,3 t	106	51	232	48
			Fertilizzante commerciale		174	19	-	
6 - Cappuccinelle	6	Mais	Spargim.int.< 12 ore	14,4 t	24	12	52	11
			Fertilizzante commerciale		256	58	-	
7 - Ballitto	2	Frumento T.	Spargim.int.< 12 ore	10 t	16	8	36	8
			Fertilizzante commerciale		164	42	-	
8 - Barca	2	Orzo	Spargim.int.< 12 ore	8,3 t	14	7	30	6
			Fertilizzante commerciale		106	23	-	
9 - Luna	1	Vite	Spargim.int.< 12 ore	36,4 t	60	29	131	27
			Fertilizzante commerciale		-	71	-	
10 - Sole	2	Girasole	Spargim.int.< 12 ore	9,2 t	15	7	33	7
			Fertilizzante commerciale		85	38	-	

*** ELEMENTI NUTRITIVI IN ECCESSO PER ETTARO ***

Appezzamento	Ettari	Coltura	kg N	kg P2O5	kg K2O
1 - Vigneto	2,3	Vite	-	-	240
2 - Campetta	3	Mais	-	10	308
3 - Cerreto	10,5	Orzo	-	10	150
4 - Oliveto	1,75	Olivo	0	-	108
5 - Spaccio	5,3	Mais	-	-	180
6 - Cappuccinelle	6	Mais	-	-	-
7 - Ballitto	2	Frumento T.	-	-	-
8 - Barca	2	Orzo	-	-	0
9 - Luna	1	Vite	0	-	11
10 - Sole	2	Girasole	-	-	-

*** RIEPILOGO COSTI ***

Appezzamento	Ettari	Coltura	-----Con deiezioni-----			---Senza deiezioni---	
			Costo spandimento e trasp.deiez.	Costo fertilizzante commerciale	Costo totale per ha	Costo fertilizzante commerciale	Costo totale per ha
1 - Vigneto	2,3	Vite	462.760	117.530	252.300	602.370	261.900
2 - Campetta	3	Mais	759.000	446.550	401.850	1218750	406.250
3 - Cerreto	10,5	Orzo	1.321.950	757.313	198.025	2096850	199.700
4 - Oliveto	1,75	Olivo	392.424	63.384	260.462	444.500	254.000
5 - Spaccio	5,3	Mais	1.363.636	1176895	479.345	2153125	406.250
6 - Cappuccinelle	6	Mais	346.667	2013917	393.431	2437500	406.250
7 - Ballitto	2	Frumento T.	80.000	461.350	270.675	559.100	279.550
8 - Barca	2	Orzo	66.667	317.942	192.304	399.400	199.700
9 - Luna	1	Vite	134.545	92.082	226.627	261.900	261.900
10 - Sole	2	Girasole	78.833	298.696	188.765	388.300	194.150
Totali	35,85		5.006.483	5745658	299.920	10561795	294.611

*** FONTE DEGLI ELEMENTI NUTRITIVI DISTRIBUITI ***

Fonte	---- kg totali ----		
	N	P2O5	K2O
Bovini latte	2531	1344	6048
Fertilizzante comm.	3907	775	-

*** ANALISI ED UTILIZZAZIONE DELLE DEIEZIONI ***

Fonte	Unita'	--- kg per unita' ---			Deiezioni		Bilancio
		N	P2O5	K2O	dispon.	distrib.	
Bovini latte	t	3	1	4	1680	1680	0

*** DISPONIBILITA' DI ELEMENTI NUTRITIVI ***

Allevamento	Metodo spandimento	% N dispon.	% N dispon.	% N dispon.	% P2O5	% K2O
		Anno 1	Anno 2	Anno 3	disponibile	disponibile
Bovini latte	Spargim. no interra.	20	25	15	80	90
	Spargim.int.< 12 ore	55	25	10	80	90