AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

האוניברסיטה העברית בירושלים
The Hebrew University of Jerusalem

המרכז למחקר בכלכלה חקלאית
The Center for Agricultural
Economic Research

המחלקה לכלכלה חקלאית ומנהל
The Department of Agricultural
Economics and Management

Discussion Paper No. 6.05

The Economic Impact of Global Climate Change on
Mediterranean Rangeland Ecosystems:
A Space-for-Time Approach

by

Aliza Fleischer and Marcelo Sternberg

Papers by members of the Department
can be found in their home sites:

מאמרים של חברי המחלקה נמצאים
גם באתרי הבית שלהם:

<http://departments.agri.huji.ac.il/economics/indexe.html>

P.O. Box 12, Rehovot 76100

ת.ד. 12, רחובות 76100

**The Economic Impact of Global Climate Change on Mediterranean Rangeland
Ecosystems: A Space-for-Time Approach**

Aliza Fleischer^{1*}, Marcelo Sternberg²

¹*Department of Agricultural Economics and Management, The Hebrew University of
Jerusalem, PO Box 12, Rehovot 76100, Israel. E-mail: fleische@agri.huji.ac.il*

* Corresponding author

²*Department of Plant Sciences, Faculty of Life Sciences, Tel Aviv University,
Tel Aviv 69978, Israel. E-mail: MarceloS@tauex.tau.ac.il*

Abstract

Global Climate Change (GCC) can bring about changes in ecosystems and consequently in their services value. Here we show that the urban population in Israel values the green landscape of rangelands in the mesic Mediterranean climate region and is willing to pay for preserving it in light of the expected increasing aridity conditions in this region. Their valuation of the landscape is higher than that of the grazing services these rangelands provide for livestock growers. These results stem from a Time-for-Space approach with which we were able to measure changes in biomass production and rainfall at four experimental sites along an aridity gradient.

Keywords: global climate change, ecosystem, choice modeling, landscape, biomass

1. Introduction

In studies dealing with global climate change (GCC) issues, a division between the life and social sciences is commonly found. Life scientists emphasize the forecasting of different future climate change scenarios or the resulting changes on

ecosystems and their functioning. On the other hand, the economic impact of GCC on the human community is generally dealt with by social scientists. In their analyses, the life science aspects are either assumed or taken as a given from other works. Studies of the economic effects of GCC focus on either market impacts, such as possible changes in farm income, or non-market impacts, such as changes in the value of an ecosystem's services (e.g., life support and aesthetic enjoyment for the human community). Natural rangelands, the ecosystem considered in this study, provide both market services, such as grazing, food supply, and genetic resources, and non-market services, such as landscape, recreation and culture.

In this study we evaluate both types of impact of GCC on natural rangeland ecosystems. We do so by integrating findings from both life science and economic analyses.

Market impacts of GCC have been estimated in several studies mainly by analyzing changes in farm income using the production-function approach (e.g., Decker et al., 1986; Adams, 1989; Adams et al., 1990) or the Ricardian approach (Mendelsohn et al., 1994). These studies have used current market data to evaluate changes that may occur in the future. For example, in studies evaluating non-market impacts of GCC on ecosystems, such as those by Layton and Brown (2000) and Turpie (2003), stated preference techniques were used to elicit the value of the present population's preferences to preserve ecosystems for future generations. The change in ecosystem considered in the former study was forest loss along the Colorado Front Range of the Rocky Mountains. Four alternative levels of forest loss were assumed and analyzed by computerized photographs of a typical mountain range. In Turpie's (2003) study, the researcher used maps of current biomes in South Africa and a simulation of the future distribution of biomes in a scenario that could be brought

about by an increase in the concentration of atmospheric carbon dioxide (CO₂) to 550 ppm. Both studies used only computerized manipulations to illustrate the changes: one of the landscape pictures and the other of the biome maps.

In the present study, an integrated approach was implemented to evaluate the impact of GCC on the value of natural rangeland services along an aridity gradient in Israel. The integration in this study has different dimensions. Firstly, both market and non-market impacts are considered. Secondly, we integrate natural and social science approaches to study natural ecosystems and potential changes in their services. We use real measurements of herbaceous and woody plant biomass at different sites along the aridity gradient as proxies for productivity changes.

The challenge of predicting ecosystem responses to climate changes is based on the multi-dimensional and multi-scale nature of the problem (Osmond et al., 2004). Complex ecological interactions make it difficult to extrapolate from individual species to communities and to predict the ecosystem response when only a few organization levels are targeted. In addition, the lack of realistic climatic scenarios (climate modelling) at the relevant scales adds further complexity to the up-scaling process (Harvey, 2000).

Predictions about ecosystem functioning in relation to GCC along climatic gradients rely on two major research assumptions. The first considers that existing environmental gradients can be used as spatial analogues (climosequences) for future climate change. In this case, environmental and ecological characteristics are described for existing climates in present locations and compared along a gradient. Such predicted climatic scenarios are then imposed on existing conditions, e.g. an increase in rainfall would result in a set of conditions that are similar to current areas in more mesic parts along the climatic gradient. The second assumption is that biotic

responses to climatic changes can be inferred from current species distributions and their correlations with abiotic factors. Such 'climate envelope' approaches use mapped current distributions and predict future distribution solely on changes in abiotic (namely climatic) conditions. This approach is known as the **Space-for-Time approach**.

Another challenge facing researchers analyzing impacts of GCC is the difficulty of experimentally mimicking changes in climatic conditions on larger scales (e.g., large watersheds, whole ecosystems or regions). Most economic papers in this field assume a certain climate scenario for their analysis. Natural scientists, however, perform a more detailed analysis of the impacts on soil characteristics, and changes in the composition and structure of the vegetation and animal community. In the Space-for-Time approach, changes in climatic conditions are simulated by comparing areas that differ naturally in their climatic regimes. Natural climatic gradients, which include environmental factors such as altitude, topography, temperature and rainfall variations, provide a useful framework for studying the effects of climatic changes (Kutiel et al. 1995, Diaz and Cabido 1997, Dunne et al 2003). Moreover, comparisons of ecosystems and biotic communities along gradients are powerful approaches to investigating and understanding the effects of climatic variation on ecosystems (Le Houerou 1990, Koch et al. 1995, Shaw and Harte 2001, Austin 2002). Approaches based on aridity gradients have been frequently used in Mediterranean ecosystems (e.g., Holzapfel et al., 1992; Imeson and Lavee, 1998; Kutiel et al. 2000).

The actual changes that will occur in the future are difficult to predict and even more so to illustrate, due to the complexity of ecosystems. However, the Space-for-Time approach allows one to better illustrate for the general public the different possible scenarios. It also enables linking climate changes to ecosystem processes,

such as measurements of changes in herbaceous and woody biomass (primary productivity) and their economic impact. Since changes in biomass affect both landscape values, i.e., a non-market impact, and feeding costs for sheep and cattle growers, i.e., a market impact, we can evaluate both effects in the same research context. The use of biomass measurements is pivotal to our integrated approach. It is a life science measurement used for the estimation of both types of economic impacts.

The paper is organized as follows: Section 2 describes the sites and data measurements used for the GCC scenarios in the Space-for-Time approach. Sections 3 and 4 depict the evaluation of changes in landscape and feeding costs, respectively. The total welfare evaluation is presented in Section 5. Section 6 lists the conclusions.

2. Description of study sites, plant cover, plant biomass measurements and the GCC scenarios

Four sites were established in 2001 along a climatic gradient in Israel, running from the Galilee in the north to the Negev Desert in the south (gradient length 245 km) (Figure 1). These sites represent, respectively, mesic Mediterranean, Mediterranean, semiarid and arid climatic conditions (Table 1). All sites share the same calcareous bedrock (hard limestone) and are positioned on south-facing slopes. The basic climate is Mediterranean with mild and rainy winters (October-April) and prolonged dry and hot summers. The plants' growing season is closely associated with the temporal distribution of rainfall. The amount of plant biomass at the sites determines the type of landscape, e.g., the higher the biomass, the denser the vegetation. Moreover, the study sites share similar climatic conditions (radiation, temperature, etc.), except for rainfall (Table 1). This links the transformation in landscape and grazing costs to the cardinal issue of the region, water scarcity.

Plant cover and biomass of the herbaceous and woody vegetation were measured at each study site. Five 10 m x 25 m quadrats were randomly selected and marked. Vegetation was monitored in spring (mid-April), during the peak season of primary production. Within each quadrat, plant cover was estimated and perennial species composition inventoried by using two 25-m long transects placed on the edges of each quadrat. On each transect, a point was read every 20 cm, for a total of 125 points per transect, 250 points per quadrat. A point was read using a slender bar positioned vertically to the ground (Müeller-Dombois & Ellenberg 1974). Relative plant cover (in %) was calculated by excluding rock and bare ground cover. Woody vegetation was sampled according to life-form categories: dwarf shrubs (< 0.5 m height), shrubs (> 0.5 m < 2.5 m height) and trees (> 2.5 m height). Herbaceous plant biomass was considered by sampling five 20 x 20 cm quadrats. After harvesting, plants were brought to the laboratory. The samples were then dried in an oven at 80°C for 3 days. After removing from the oven, samples were weighed at room temperature to a resolution of 0.01 g. Woody plant biomass was measured by an indirect procedure. Based on their relative cover, woody biomass estimations were calculated using parameters similar to those presented by Sternberg and Shoshany (2001a, b). In their study they estimated the plant biomass of woody species similar to those found at the study sites. It can be seen in Table 1 that herbaceous and total biomass declines continuously in the transition from mesic Mediterranean to Mediterranean, semiarid and arid. These data enabled us to quantify the landscape changes in Figure 1.

Using the data from the four sites, we were able to simulate four possible scenarios of climate change for the mesic Mediterranean region. In the first, the mesic Mediterranean site maintains the same climate. In each of the other three scenarios, the mesic Mediterranean site evolves into one of the other three climatic zones, i.e., to

Mediterranean, semiarid and arid climates. This allows for four levels of climatic change, ranging from 'no change' to a slight decrease in rainfall in the second scenario (i.e., the mesic Mediterranean site is transformed into a Mediterranean site with a decrease in rainfall) proceeding to a more drastic change (the mesic Mediterranean site transforms to a semiarid site) and finally to a very drastic change (the mesic Mediterranean site is transformed to an arid site). The existence of various site scenarios in the Space-for-Time approach enabled us to contemporaneously measure temperature and precipitation, take current pictures of the landscape and measure the biomass levels at these sites.

3. Evaluating landscape services

Economists have responded to the need to evaluate environmental and natural resources in the absence of markets by developing an array of non-market evaluation methods. Some of the methods depend on markets related to the environmental good, whereas others are based on stated preference techniques. Since the impact of GCC on landscape will occur in the far future, beyond the lifetime of the present population, we had to use a stated preference technique. One of these is the well-known contingent valuation method (CVM). This method is highly controversial and concerns regarding the validity of its results have been expressed as a result of: strategic bias, yea-saying, insensitivity to scope variations, framing and other causes (Bateman et al., 2002; Nunes, 2002).

A more recently developed technique, which seems to better simulate the respondents' choice process, is choice modeling (Bennett and Blamey, 2001). In this technique, the environmental good is described according to its attributes and the levels they take. The different alternatives vary in their attribute levels and

respondents have to choose the alternative they prefer. The attributes in the different alternatives can include environmental damage and abatement costs. By choosing an alternative, the respondents are actually ascribing a value or price to a level of attribute.

The Model

The probability of an individual choosing a specific alternative can be estimated using the standard logit model. However, these models impose three strong restrictions (McFadden, 1973; Train, 1986, 2003): 1) model coefficients are the same for all individuals, i.e., there are no differences in individuals' preferences, 2) the well-known Independence of Irrelevant Alternatives (IIA), and 3) in the case of repeated choices (e.g., where an individual receives a few sets of alternatives to choose from), unobserved factors are assumed to be independent for each decision. Following Train (1998, 1999, 2003), we use the random-parameters logit (RPL) model (also known as mixed logit) for repeated choices. The utility of alternative j for the i^{th} individual is:

$$U_{ij} = X_{ij}\beta_i + \varepsilon_{ij} = X_{ij}\bar{\beta} + X_{ij}\tilde{\beta}_i + \varepsilon_{ij}, \quad (1)$$

where X is a vector of attributes of alternative j , β_i is a random vector with density $f(\beta)$, and ε_{ij} is an iid independent of β_i and X . The coefficient vector for each individual β_i can be expressed as the sum of the mean $\bar{\beta}$ and the individual's deviation from the mean $\tilde{\beta}_i$. The unobserved portion of the utility function by the researcher, $X_{ij}\tilde{\beta}_i + \varepsilon_{ij}$, reflects the individuals' tastes and is thus correlated over alternatives and choices.

Assuming all correlation is due to $\tilde{\beta}_i$, then the probability that an individual i will choose alternative j from a set of alternatives is:

$$P_{ij} = \text{prob}(X_{ij}\beta_i + \varepsilon_{ij} - X_{ik}\beta_i > \varepsilon_{ik}, \forall k \neq j), \quad (2)$$

which implies

$$P_{ij} = \int \left[\frac{e^{x_{ij}\beta_i}}{\sum_{k=1}^m e^{x_{ik}\beta_i}} \right] f(\beta) d\beta. \quad (3)$$

The probability in (3) can be simulated by R draws for β_i from $f(\beta)$ (Train, 2003) as

$$P_{ij} = \frac{1}{R} \sum_{r=1}^R \frac{e^{x_{ij}\beta_{ir}}}{\sum_{k=1}^m e^{x_{ik}\beta_{ir}}}. \quad (4)$$

Following Layton and Brown (2000), the model in (4) is extended to multiple choices, i.e., each respondent receives three sets of alternatives from which he/she has to choose. The attributions of the alternatives vary between sets while preferences of respondents β_i stay the same. The probability is

$$P_{ij} = \frac{1}{R} \sum_{r=1}^R \left[\prod_{t=1}^T \frac{e^{x_{ijt}\beta_{ir}}}{\sum_{k=1}^m e^{x_{ikt}\beta_{ir}}} \right]. \quad (5)$$

Assuming $f(\beta)$ is multivariate normal, then it is possible to simulate the probability of each individual's choice from each set of alternatives and estimate it by maximum likelihood.

Data

The data collection was performed in three stages: focus groups, pre-tests and face-to-face surveys. Focus groups: This stage was based on three focus groups of adults over the age of 18 from different socioeconomic backgrounds. The purpose of this stage was to identify the level of understanding of GCC, landscape, and abatement programs. Another was to identify an acceptable range of bids and the vocabulary used by the participants in describing these issues. This information enabled us to design a first draft of the questionnaire. Pre-test: Extensive pre-testing of the questionnaire was performed with over 50 individuals, and the final version was arrived at. Survey: The survey was administered to a sample of the adult population (above the age of 18) in all 15 cities in Israel having more than 30,000 households. The population in these cities accounts for about half of the 6.8 million Israeli residents. Sample size was set at 500 and the number of respondents from each city was chosen according to the relative weights of the city households. Within each city, respondents were chosen randomly. Each respondent received three different sets of alternatives (see Figure 2 for an example of such a set) and each set contained five possible programs. The use of three sets per respondent allowed for the collection of more information. That is, instead of 500 observations there were 1,500, three per respondent. Questionnaire: The design of the questionnaire relied on the work of Layton and Brown (2000). It starts with a short and simple description of GCC and its possible impact on the eastern Mediterranean region. It ends with questions concerning the demographic and socioeconomic characteristics of the respondents. The main part of the questionnaire contains three sets of alternatives (these sets are denoted a, b, and c in the following discussion) for possible changes in climate in the Galilee, a region with mesic Mediterranean climate. Twelve versions of the questionnaire were administered by alternating the sets. The versions differed in the

order of the sets: one-third of the respondents received the three sets in the order a, b, c, one-third, a, c, b, and one-third c, a, b. Since set c always had higher bids, it was important to mix the sets. Half of the respondents received three sets with a higher level of bids than the other half. Half of the respondents received a scenario in which the time horizon for materialization of the GCC impacts was 100 years and the other half 30 years. The 12 versions and their distribution in the sample appear in Table 2.

Each alternative that the respondents had to choose from had four attributes: landscape, forestation, other abatement measures and bids. The attributes varied as follows: four different landscapes depicted by pictures from the four sites, two levels for forestation (utilizing and not utilizing forestation as a preventive measure), three levels of abatement (none, some, vigorous) to reduce greenhouse gases, and 14 levels of bids ranging from 0 to \$50.

Alternatives set 'a' in Figure 2 depicts five programs that respondents had to choose from. The changes in landscape are demonstrated in the pictures taken at the sites in the spring of 2003 when biomass was at its peak. Pictures in the first row are all taken from the mesic Mediterranean (Galilee) site, while pictures in the second row are from the other three sites: arid, semiarid---appearing twice, and Mediterranean. In program 1, no action is taken, abatement cost is zero and the landscape changes from mesic Mediterranean (Galilee) to arid. In programs 2 and 3, the landscape changes from mesic Mediterranean to semiarid and the abatement cost is \$7.5 per month. The alternatives differ in the abatement method, forestation vs. reduction in greenhouse gases. In program 4, there is a cost of \$15 a month for reduction in greenhouse gases, and forestation changes to a Mediterranean landscape. Program 5 is the most expensive one in the set. For \$20 a month, drastic abatement measures are taken and the mesic Mediterranean landscape is maintained.

Estimates

Equations (3)-(5) are developed under the assumption that the observed part of the utility function is linear in the parameters (Train, 2003). Accordingly, the following variables were entered linearly as the attribute of each program:

1. *cost*---there are 14 values of program costs ranging from \$0 to \$50;
2. *forestation*---dummy variable receives a value of 1 when the method appears in the program and 0 otherwise;
3. *reduction*---dummy variable receives a value of 1 when any level, moderate or vigorous, of greenhouse reduction appears in the program and 0 otherwise;
4. *biomass loss*---tons of biomass are lost in the transition from mesic Mediterranean climate zone to the other climate zones. Measurements of biomass at each of the four sites enabled us to translate the changes in landscape to biomass loss. For example, the transformation from mesic Mediterranean to Mediterranean landscape is caused by a loss of 7.8 tons per hectare (see Table 1).

The parameter of *cost* is expected to receive a negative value for all respondents and thus it is assumed to be constant for all respondents. Chen and Cosslett (1998) and Layton and Brown (2000) used the same assumption to guarantee a negative coefficient for cost and consequently a normal independent distribution for willingness to pay (WTP). All other coefficients are assumed to vary and to have normal distribution. The signs for the variables *reduction* and *forestation* can be either negative or positive, depending on people's preferences for the two methods. In the case of *biomass loss*, although green landscape is held in high esteem in Israel (this was tested in the focus groups), there are still people who would prefer the desert-like landscape. The model parameters were estimated by LIMDEP 8 (2002).

The model estimates can be seen in Table 3. The means and standard deviations of *reduction* and *forestation* reveal the heterogeneity in the population preferences for the two greenhouse-gas-reduction methods. The mean coefficient in the case of *reduction* is 1.66 and the standard deviation is 2.7, that is, 70% of the population likes this method. In the case of *forestation* the coefficient is not significant, that is, the population is indifferent between using or not using this method of abatement.

The ratio of the biomass coefficient to cost coefficient measures the average WTP in order to prevent the loss of 1 ton of biomass per hectare. In this case, the ratio has a normal distribution since *biomass loss* is normally distributed and cost is constant. Thus, the mean WTP is \$2 per ton of biomass loss per hectare with a standard deviation of 7.8. The range of WTP is relatively large, which indicates a wide variation in the population's WTP.

In the Space-for-Time approach, the choice of experimental sites enables us to focus only on changes in rainfall. That is, rainfall is the main factor that varies along the gradient in the production of biomass. As we move south, the amount of rainfall drops, while temperature and other factors remain almost constant. Thus, we can substitute biomass loss with drop in rainfall and estimate, accordingly, the WTP to prevent this change in rainfall. This substitution allows us to link the landscape choice of the respondents to an important factor in climatic change. The alternative variable to *biomass loss*, *drop in rainfall*, measures the drop in rainfall in millimeters in the transition from the mesic Mediterranean climate zone to the other climate zones. The estimated model substituting *biomass loss* with *drop in rainfall* appears in Table 4. The average WTP for the prevention of a drop in rainfall is estimated to be \$0.05 a

year per one millimeter reduction in rainfall, with a standard deviation of 0.13, or \$5 per 100 mm drop in rainfall.

4. Evaluating grazing services

Grazing services of the ecosystem at issue consist of free food for cattle and sheep growers. Sheep and cattle consume mostly herbaceous biomass. Therefore, the more herbaceous biomass there is at the site, the more the grower saves on food costs. Assuming a constant coefficient production function, we can evaluate the change in costs and thus in profits of farmers that depend on these sites. Individual cows and sheep consume 10 and 1.5 kg, respectively, of herbaceous biomass (dry material) per day. Alternatively, growers have to pay \$1 for food per day per cow, \$0.21 per sheep. Based on the total dry herbaceous biomass at each site, Table 5 shows how much growers save in food costs per year. In the mesic Mediterranean area, the savings are naturally the largest and stand at \$83.2 per hectare for cattle and \$116.5 for sheep.

5. Evaluating loss of ecosystem services

Based on the last two sections, we can evaluate the loss in value of the two ecosystem services, landscape and grazing, when climatic conditions change from mesic Mediterranean to any of the other three climates. The landscape is determined by the amount of plant biomass per hectare, and the population values one ton of plant biomass per hectare at \$2. As seen in Table 6, a change in landscape in the northern region from mesic Mediterranean to Mediterranean is valued by the urban population as a \$51.5 million loss in welfare. Alternatively, by looking at changes in rainfall, it is valued at \$39.6 million. Similarly, the transformation to a semiarid climate is valued at, respectively, \$85.5 or \$79.2 million and to an arid one at \$107.6 or \$113.8 million.

In none of the three cases are there any significant differences between alternative values.

If the land is used for cattle growing, then growers will lose \$9.1 per hectare; for sheep raising, the increase in food costs is \$12.7. The total area of grazing land in the mesic region is 630 ha. The loss for cattle growers in the mesic Mediterranean region can range between \$5,733 and \$51,534 a year. In the case of sheep growers, the yearly range of losses in food costs is higher and varies between \$8,001 and \$72,135. Currently, the grazing land in the northern site is used mostly for cattle, thus the loss values for cattle are more relevant.

6. Concluding remarks

The Space-for-Time method provides the population with an illustration of the impact of GCC on landscape in the form of actual photographs of the sites. Furthermore, the use of biomass measurements enables linking these changes in landscape to changes in biomass and, even further, to changes in rainfall. This link lets us assign a value to a climatic variable based on tangible illustrations, rather than on just a narrative describing the changes in climate. This result is made possible by the interdisciplinary nature of this research.

Based on the aforementioned method, we show that the urban population in Israel values green landscape. They are willing to pay for it even though they might not be here when the changes take place. Furthermore, the loss in welfare from the change in landscape is valued much higher than the loss in income for farmers that depend on the land for grazing. The population is willing to pay about \$80 million a year to prevent the mesic Mediterranean landscape from changing to a semiarid one,

whereas cattle and sheep growers will lose \$16 thousand to \$22 thousand, respectively, if this climate transformation occurs.

It should be noted that the result obtained here, whereby the population assigns a higher value to the landscape in rangelands than its additional income to livestock growers, is conditional on the fact that Israel is a high-income country. In the case of low-income countries, we expect the results to be reversed. The ‘free’ feeding services provided by rangelands are significant at low income levels. Moreover, the latter population engages much less in outdoor recreation and thus does not value the landscape as much as high-income countries.

Acknowledgements

This research is funded in the framework of the project GLOWA Jordan River by the German Federal Ministry of Education and Research (BMBF) in collaboration with the Israeli Ministry of Science and Technology (MOST).

- [1] Adams, R. (1989). Global climate change and agriculture: an economic perspective. *American Journal of Agricultural Economics*, 71(5): 1272-9.
- [2] Adams, R., Rosenzweig, C., Pearl, R., Ritchie, J., McCarl, B., Glyer, D., Curry, B., Jones, J., Boote, K., and Allen, H. (1990). Global climate change and U.S. agriculture. *Nature*, 345(6272): 219-24.
- [3] Austin, AT. (2002). Differential effects of precipitation on production and decomposition along a rainfall gradient in Hawaii. *Ecology* 83:328-338.
- [4] Bateman, I.J., Carson, R.T., Day, B., Hanemann, M., Hanley, N., Hett, T., Jones-Lee, M., Loomes, G., Mourato, S., Özdemiroğlu, E., Pearce OBE, D.W., Sugden, R. and Swanson, J. (2002). *Economic Valuation with Stated Preference Techniques*. Cheltenham, UK: Edward Elgar.
- [5] Bennett, J., and Blamey, R. (2001). *The Choice Modeling Approach to Environmental Valuation*. Cheltenham, UK: Edward Elgar.
- [6] Chen, H.z., and Cosslett, S.R. (1998). Environmental Quality Preference and Benefit Estimation in Multinomial Probit Models: A simulation Approach. *American Journal of Agricultural Economics*, 80(3): 512-520.
- [7] Diaz, S. and Cabido, M. (1997). Plant functional types and ecosystem function in relation to global change. *Journal of Vegetation Science* 8: 463-474
- [8] Decker, W., Jones, V., and Achutuni, R. (1986). *The Impacts of Climate Change from increased Atmospheric Carbon Dioxide on American Agriculture*, DOE/NBB-0077. Washington, DC: U.S. Department of Energy.
- [9] Dunne, J.A, Harte, J. and Taylor, K. J. (2003). Subalpine meadow flowering phenology responses to climate change: Integrating experimental and gradient methods. *Ecological Monographs* 73: 69-86.

- [10] Harvey, L.D.D. (2000). Upscaling in climate change research. *Climatic Change* 44: 225-263.
- [11] Holzapfel, C., Schmidt, W., and Shmida, A. (1992). Effects of human-caused disturbances on the flora along a Mediterranean-desert gradient. *Flora*, 186: 261-70.
- [12] Imeson, A., and Lavee, H. (1998). Soil erosion and climate change: the transect approach and the influence of scale. *Geomorphology*, 23: 219-27.
- [13] Koch, G.W., Vitousek, P.M., Steffen, W.L. and Walker, B.H. (1995). Terrestrial transects for global change research. *Vegetatio* 121:53-65.
- [14] Kutiel, P., Lavee, H. and Shoshany, M. (1995) Influence of a climatic gradient upon vegetation dynamics along a Mediterranean-arid transect. *Journal of Biogeography* 22: 1065-1071.
- [15] Kutiel, P., Kutiel, H., and Lavee, H. (2000). Vegetation response to possible scenarios of rainfall variations along a Mediterranean-extreme arid climatic transect. *Journal of Arid Environments*, 44: 277-90.
- [16] Layton, D.F., and Brown, G. (2000). Heterogeneous preferences regarding global climate change. *The Review of Economics and Statistics*, 82(4): 616-24.
- [17] Le Houerou, H.N. (1990). Global change - vegetation, ecosystems, and land use in the southern Mediterranean basin by the mid twenty-first century. *Israel Journal of Botany* 39: 481- 508
- [18] LIMDEP 8 (2002). *Econometric Modeling Guide*. Plainview, NY: Econometric Software, Inc.
- [19] McFadden, D. (1973). Conditional logit analysis of qualitative choice models. In *Frontiers of Econometrics*, ed. P. Zarembka. New York: Academic Press.

- [20] Mendelsohn, R. Nordaus, W.D. and Shaw, D. (1994). The impact of Global Warming on Agriculture: A Ricardian Analysis. *The American Economic Review*, 84(4): 753-771.
- [21] Müller-Dombois, D., and Ellenberg, H. (1974). *Aims and Methods of Vegetation Ecology*. New York: Wiley.
- [22] Nunes, P.A.L.D. (2002). *The Contingent Valuation of Natural Parks*. Cheltenham UK: Edward Elgar.
- [23] Osmond, B., Ananyev, G., Berry, J., Langdon, C., Kolber, Z., Lin, G., Monson, R., Nichol, C., Rascher, U., Schurr, U., Smith, S., and Yakir, D. (2004). Changing the way we think about global climate change research: scaling up in ecosystem experimental science. *Global Change Biology*, 10: 393-407.
- [24] Shaw, M.R. and Harte, J. (2001). Response of nitrogen cycling to simulated climate change: differential responses along a subalpine ecotone. *Global Change Biology* 7:193-210.
- [25] Sternberg, M., and Shoshany, M. (2001a). Influence of slope aspect on Mediterranean woody formations: a comparison of a semiarid and an arid site in Israel. *Ecological Research*, 16: 335-45.
- [26] Sternberg, M., and Shoshany, M. (2001b). Aboveground biomass allocation and water content relationships in Mediterranean trees and shrubs at two climatological regions in Israel. *Plant Ecology* 157: 171-9.
- [27] Train, K.E. (1986). *Qualitative Choice Analysis*. Cambridge: MIT Press.
- [28] Train, K.E. (1998). Recreation demand models with taste differences over people. *Land Economics*, 74(2): 230-9.

- [29] Train, K.E. (1999). Mixed logit models for recreation demand. In *Valuing Recreation and the Environment*, eds.J. Herriges and C. Kling. Northampton, MA: Edwards Elgar.
- [30] Train, K.E. (2003). *Discrete Choice Methods with Simulation*. Cambridge: Cambridge University Press.
- [31] Turpie, J.K. (2003). The existence value of biodiversity in South Africa: how interest, experience, knowledge, income and perceived level of threat influence local willingness to pay. *Ecological Economics*, 46(2): 199-219.

Figure 1: The study sites along the climatic gradient in Israel.

Table 1: Physical and biotic characteristics of the study sites along the aridity gradient. Temperature refers to mean and mean maximum).

Ecosystem type	Rainfall (mm)	Temperature (°C) Min. – Mean - Max.	Elevation (a.s.l)	Soil type	Vegetation formation
Mesic Mediterranean (N 33°0' E 35°14')	780	13.5 - 18.1 - 23.4	500 m	Montmorillonitic Terra Rossa	Closed oak maquis (<i>Quercus calliprinos</i>) and open garigue form dominate shrubs (e.g. <i>Calicotome villosa</i> , <i>Sarcopoterium spinosum</i> , <i>C</i> spp.) and associated herbaceous plants.
Mediterranean (N 31°42' E 35°3')	540	12.8 – 17.7 - 23.6	620 m	Terra Rossa	Dwarf-shrubland dominated by <i>Sarcopoterium spinosum</i> and a high diversity of herbaceous (mostly annual) plant species.
Semiarid (N 31°23' E 34°54')	300	13.2 – 18.4 – 24.8	590 m	Light Brown Rendzina	Dwarf shrubs of <i>Sarcopoterium spinosum</i> and <i>Coridothymus capitata</i> associated with herbaceous (chiefly annual) plant species.
Arid (N 30°52' E 34°46')	90	13.6 - 19.1 - 26.1	470 m	Desert Lithosol	Open vegetation dominated by small shrubs and semi-shrubs such as <i>Zygophyllum dumosum</i> , <i>Artemisia sieberi</i> and <i>Hammada scoparia</i> and sparsely growing desert annuals, geophytes and hemicryptophytes.

Table 2: Distribution of versions by Time Horizon, Bid Levels, and Order of Sets (number of respondents for parentheses).

Time Horizon	30 years (255)		100 years (245)	
Bids	High bids (123)	Low bids (132)	High bids (124)	Low bids (121)
Order (a,b,c) (164)	Version 1 (42)	Version 2 (43)	Version 7 (40)	Version 8 (39)
Order (a,c,b) (172)	Version 3 (40)	Version 4 (48)	Version 9 (42)	Version 10 (42)
Order (c,a,b) (164)	Version 5 (41)	Version 6 (41)	Version 11 (42)	Version 12 (40)

Figure 2: An example of one set of alternatives: respondent had to choose one program from each set.

Program 1 No action	Program 2 Forestation is used to slow down greenhouse effect	Program 3 Reduction in the use of greenhouse gases	Program 4 Forestation and greenhouse-gas	
Landscape in the Galilee ^a will become arid, also loss of plant life will occur	Landscape in the Galilee ^a will become semiarid	Landscape in the Galilee ^a will become semiarid	Landscape in th will have less pl	
\$0 per month	\$7.5 per month	\$7.5 per month	\$15 per month	
Mesic Mediterranean	Mesic Mediterranean	Mesic Mediterranean	Mesic Mediterr:
Arid	Semiarid	Semiarid	Mediterranean

^athe Galilee is the region with mesic Mediterranean climate

Table 3: Estimation of Random Parameter Model with biomass loss.

Variable	Parameter	Value	Std. Error
Cost	Mean of coefficient	-0.033*	0.008
	Std. dev. of coefficient	0	0
Biomass loss	Mean of coefficient	-0.069*	0.018
	Std. dev. of coefficient	0.260*	0.081
Forestation	Mean of coefficient	-0.221	0.194
	Std. dev. of coefficient	2.507*	0.223
Reduction	Mean of coefficient	1.664*	0.794
	Std. dev. of coefficient	2.752*	0.977
R^2 ^(a)		0.18	
Number of obs.		1500= (500 x 3)	

*Denotes significance at 5%.

^a $R^2 = 1 - [(\text{Log-likelihood of the model}) / (\text{Log-likelihood}(\beta = 0))]$.

Cholesky Matrix

	Price	Biomass loss	Forestation	Reduction
Price	0			
Biomass loss	0.025 (0.056)	0.0087 (0.101)		
Forestation	-2.44* (0.297)	0.162* (0.068)	0.562 (0.838)	
Reduction	-1.426 (1.299)	-0.201* (0.056)	0.148 (1.512)	2.34* (1.22)

Note: Standard errors are in parentheses.

*Denotes significance at 5%.

Table 4: Estimation of Random Parameter Model with drop in rainfall

Variable	Parameter	Value	Std. Error
Cost	Mean of coefficient	-0.0498*	0.01
	Std. dev. of coefficient	0	0
Drop in rainfall	Mean of coefficient	-0.0026*	0.0005
	Std. dev. of coefficient	0.0066*	0.0015
Forestation	Mean of coefficient	-0.239	0.189
	Std. dev. of coefficient	2.361*	0.22
Reduction	Mean of coefficient	1.455**	0.845
	Std. dev. of coefficient	2.809*	1.45
R^2 ^(a)		0.19	
Number of obs.		1500= (500 x 3)	

*,**Denotes significance at 5% and 10%, respectively.

^a $R^2 = 1 - [(\text{Log-likelihood of the model}) / (\text{Log-likelihood}(\beta=0))]$.

Cholesky Matrix

	Price	Drop in rainfall	Forestation	Reduction
Price	0			
Drop in rainfall	0.0002 (0.0013)	0.00205 (0.0023)		
Forestation	-2.28* (0.3)	0.0039* (0.001)	0.613 (0.84)	
Reduction	-0.259 (1.4)	0.0049* (0.0014)	-2.75* (1.44)	0.47 (1.58)

Note: Standard errors are in parentheses.

*Denotes significance at 5%.

Table 5: Savings in food costs for cattle and sheep.

	Cattle (\$ per ha)	Sheep (\$ per ha)
Mesic Mediterranean	83.23	116.5
Mediterranean	74.14	103.79
Semiarid	57.66	80.72
Arid	1.45	2.03

Table 6: Yearly loss value of ecosystem services in the transformation from mesic Mediterranean to Mediter

	Drop in rainfall	Total WTP to prevent drop in rainfall ^a (\$ 10 ⁶ ha ⁻¹)	Loss of total biomass (ton ha ⁻¹)	Total WTP to prevent loss of biomass ^b (\$ 10 ⁶ ha ⁻¹)	Loss of herbaceous biomass (ton ha ⁻¹)		
Mesic Med. → Med.	2.4	39.6	7.8	51.5	0.009		
Mesic Med. → Semiarid	4.8	79.2	13.0	85.8	0.256		
Mesic Med. → Arid	6.9	113.85	16.3	107.6	0.818		

^aThe average WTP of \$0.05 is multiplied by 3.3x10⁶ residents of large urban centers and by the drop in rainfall

^bThe average WTP of \$2 is multiplied by 3.3x10⁶ residents of large urban centers and by the loss of total biomass

^cThe difference between mesic Mediterranean region and the other region in saving in food costs is calculated multiplied by 630 ha the total grazing area in the mesic Mediterranean region.

PREVIOUS DISCUSSION PAPERS

- 1.01 Yoav Kislev - Water Markets (Hebrew).
- 2.01 Or Goldfarb and Yoav Kislev - Incorporating Uncertainty in Water Management (Hebrew).
- 3.01 Zvi Lerman, Yoav Kislev, Alon Kriss and David Biton - Agricultural Output and Productivity in the Former Soviet Republics.
- 4.01 Jonathan Lipow & Yakir Plessner - The Identification of Enemy Intentions through Observation of Long Lead-Time Military Preparations.
- 5.01 Csaba Csaki & Zvi Lerman - Land Reform and Farm Restructuring in Moldova: A Real Breakthrough?
- 6.01 Zvi Lerman - Perspectives on Future Research in Central and Eastern European Transition Agriculture.
- 7.01 Zvi Lerman - A Decade of Land Reform and Farm Restructuring: What Russia Can Learn from the World Experience.
- 8.01 Zvi Lerman - Institutions and Technologies for Subsistence Agriculture: How to Increase Commercialization.
- 9.01 Yoav Kislev & Evgeniya Vaksin - The Water Economy of Israel--An Illustrated Review. (Hebrew).
- 10.01 Csaba Csaki & Zvi Lerman - Land and Farm Structure in Poland.
- 11.01 Yoav Kislev - The Water Economy of Israel.
- 12.01 Or Goldfarb and Yoav Kislev - Water Management in Israel: Rules vs. Discretion.
- 1.02 Or Goldfarb and Yoav Kislev - A Sustainable Salt Regime in the Coastal Aquifer (Hebrew).
- 2.02 Aliza Fleischer and Yacov Tsur - Measuring the Recreational Value of Open Spaces.
- 3.02 Yair Mundlak, Donald F. Larson and Rita Butzer - Determinants of Agricultural Growth in Thailand, Indonesia and The Philippines.
- 4.02 Yacov Tsur and Amos Zemel - Growth, Scarcity and R&D.
- 5.02 Ayal Kimhi - Socio-Economic Determinants of Health and Physical Fitness in Southern Ethiopia.
- 6.02 Yoav Kislev - Urban Water in Israel.
- 7.02 Yoav Kislev - A Lecture: Prices of Water in the Time of Desalination. (Hebrew).

- 8.02 Yacov Tsur and Amos Zemel - On Knowledge-Based Economic Growth.
- 9.02 Yacov Tsur and Amos Zemel - Endangered aquifers: Groundwater management under threats of catastrophic events.
- 10.02 Uri Shani, Yacov Tsur and Amos Zemel - Optimal Dynamic Irrigation Schemes.
- 1.03 Yoav Kislev - The Reform in the Prices of Water for Agriculture (Hebrew).
- 2.03 Yair Mundlak - Economic growth: Lessons from two centuries of American Agriculture.
- 3.03 Yoav Kislev - Sub-Optimal Allocation of Fresh Water. (Hebrew).
- 4.03 Dirk J. Bezemer & Zvi Lerman - Rural Livelihoods in Armenia.
- 5.03 Catherine Benjamin and Ayal Kimhi - Farm Work, Off-Farm Work, and Hired Farm Labor: Estimating a Discrete-Choice Model of French Farm Couples' Labor Decisions.
- 6.03 Eli Feinerman, Israel Finkelshtain and Iddo Kan - On a Political Solution to the Nimby Conflict.
- 7.03 Arthur Fishman and Avi Simhon - Can Income Equality Increase Competitiveness?
- 8.03 Zvika Neeman, Daniele Paserman and Avi Simhon - Corruption and Openness.
- 9.03 Eric D. Gould, Omer Moav and Avi Simhon - The Mystery of Monogamy.
- 10.03 Ayal Kimhi - Plot Size and Maize Productivity in Zambia: The Inverse Relationship Re-examined.
- 11.03 Zvi Lerman and Ivan Stanchin - New Contract Arrangements in Turkmen Agriculture: Impacts on Productivity and Rural Incomes.
- 12.03 Yoav Kislev and Evgeniya Vaksin - Statistical Atlas of Agriculture in Israel - 2003-Update (Hebrew).
- 1.04 Sanjaya DeSilva, Robert E. Evenson, Ayal Kimhi - Labor Supervision and Transaction Costs: Evidence from Bicol Rice Farms.
- 2.04 Ayal Kimhi - Economic Well-Being in Rural Communities in Israel.
- 3.04 Ayal Kimhi - The Role of Agriculture in Rural Well-Being in Israel.
- 4.04 Ayal Kimhi - Gender Differences in Health and Nutrition in Southern Ethiopia.
- 5.04 Aliza Fleischer and Yacov Tsur - The Amenity Value of Agricultural Landscape and Rural-Urban Land Allocation.

- 6.04 Yacov Tsur and Amos Zemel – Resource Exploitation, Biodiversity and Ecological Events.
- 7.04 Yacov Tsur and Amos Zemel – Knowledge Spillover, Learning Incentives And Economic Growth.
- 8.04 Ayal Kimhi – Growth, Inequality and Labor Markets in LDCs: A Survey.
- 9.04 Ayal Kimhi – Gender and Intrahousehold Food Allocation in Southern Ethiopia
- 10.04 Yael Kachel, Yoav Kislev & Israel Finkelshtain – Equilibrium Contracts in The Israeli Citrus Industry.
- 11.04 Zvi Lerman, Csaba Csaki & Gershon Feder – Evolving Farm Structures and Land Use Patterns in Former Socialist Countries.
- 12.04 Margarita Grazhdaninova and Zvi Lerman – Allocative and Technical Efficiency of Corporate Farms.
- 13.04 Ruerd Ruben and Zvi Lerman – Why Nicaraguan Peasants Stay in Agricultural Production Cooperatives.
- 14.04 William M. Liefert, Zvi Lerman, Bruce Gardner and Eugenia Serova - Agricultural Labor in Russia: Efficiency and Profitability.
- 1.05 Yacov Tsur and Amos Zemel – Resource Exploitation, Biodiversity Loss and Ecological Events.
- 2.05 Zvi Lerman and Natalya Shagaida – Land Reform and Development of Agricultural Land Markets in Russia.
- 3.05 Ziv Bar-Shira, Israel Finkelshtain and Avi Simhon – Regulating Irrigation via Block-Rate Pricing: An Econometric Analysis.
- 4.05 Yacov Tsur and Amos Zemel – Welfare Measurement under Threats of Environmental Catastrophes.
- 5.05 Avner Ahituv and Ayal Kimhi – The Joint Dynamics of Off-Farm Employment and the Level of Farm Activity.
- 6.05 Aliza Fleischer and Marcelo Sternberg – The Economic Impact of Global Climate Change on Mediterranean Rangeland Ecosystems: A Space-for-Time Approach.