



***The World's Largest Open Access Agricultural & Applied Economics Digital Library***

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search  
<http://ageconsearch.umn.edu>  
[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from AgEcon Search may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

*No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.*

FEEDING AND MANAGEMENT STRATEGIES FOR PRODUCING BEEF FROM HOLSTEIN STEERS  
ON NORTHEAST DAIRY FARMS

Danny G. Fox and Caroline J. Nowak

## INTRODUCTION

There are nearly one million Holstein bull calves born in the Northeast each year. This is a sizeable resource that should be evaluated for fed beef production to determine if it is a potentially viable alternative enterprise for Northeast farms. Previous research at Cornell has presented several beef production systems that would be practical in the Northeast (Knoblauch *et al.*, Woodell). An additional system would be the rearing of the steer calves with the replacement heifers on dairy farms. Many dairy-men have expressed a desire to expand their operation without increasing the milking herd, because of labor, management, facilities limitation or personal desires. Raising the Holstein steer calves produced on dairy farms is an alternative for expanding the farm operation and a means of marketing surplus forages. The purpose of this paper is to outline feeding and management strategies for raising the steers with the replacement heifers and marketing them either as feeder cattle or fed beef. The objectives are to use home-grown forages for as much of the feeding period as possible, maintain efficient growth and produce a quality feeder or slaughter weight steer.

## MANAGEMENT STRATEGY

Successful rearing of these calves for fed beef depends primarily on getting them through the critical starting phase with minimal death loss and chronic health problems, making the transition from milk to lower cost dry diets as quickly as possible, and feeding proper energy and protein supplements for the feedstuffs commonly grown in the Northeast.

One postweaning feeding strategy revolves around feeding the heifers at the recommended plane of nutrition that results in adequate size at breeding (14-15 months of age) and at first calving (24 months), without allowing fat deposits that would cause reproductive problems or a reduction in lifetime milk production (Reid). Table 1 shows rate of gain, dry matter intake and protein content that will result in desired size at breeding and first calving. Dry matter intakes are based on recent studies with Holstein steers (Fox 1981) and on values suggested by the National Research Council (NRC). The most rapid rate of growth for dairy heifers is recommended for the period prior to 8-10 months of age so that they reach puberty by 14-15 months of age. After that age, the rate of gain is reduced to prevent excessive fat deposition.

The potentially appealing feeding strategy for the production of dairy beef is to raise the

steers with the heifers for 9 to 10 months, feeding a ration formulated for the dairy heifers. When the heifers are changed to a lower energy diet, the steers and heifers are separated. The steers are continued on the same or a higher energy level, as the efficiency of growth for fed beef production is highest if the energy level is never lower than that fed at earlier stages of growth (Dexheimer, Meiske and Goodrich). Since actual dry matter intake reaches a constant level at approximately 900 pounds (Fox 1981), as weight increases beyond this weight a higher proportion of the intake will be used for maintenance and the proportion of the gain that is fat increases. When a high energy diet is fed after a period of feeding on a high forage diet, the calves will make some compensatory growth, depending on the length of time on the low plane of energy and on the degree of restriction in preceding growth. Because of these effects, the energy density of the diet must increase to maintain the same rate of gain. Therefore, low energy forages are best utilized for feed during the first part of the feeding period.

The time when heifer ration energy levels are decreased and the steers and heifers are separated, is a logical point to consider the alternative of selling the steers as feeder calves, if good feeder calf markets are available (Milligan, Nowak and Knoblauch).

## RATION FORMULATION

Rations presented in this paper are formulated to feed the steers to slaughter weight utilizing the highest levels of forage that should be fed, considering the principles discussed previously and avoiding discounts for the carcasses that occur when steers are fed a low energy diet continuously (Table 2). Given targeted average daily gain, dry matter intake and crude protein requirement, a microcomputer program was used to evaluate a prespecified ration (Fox, Weiser and Phillips). As the program performed as a ration analyzer, the composition of the ration was adjusted until the balanced ration with the least cost was obtained.

No growth stimulants or feed additives are included when the steers are with the heifers; however, after they are separated it is assumed that the steers are implanted with a growth stimulant every 100 days and that Rumensin is fed at recommended levels (Minish and Fox).

Feeding programs were developed for two feed availability circumstances.<sup>1</sup> The first contains a hay (that is primarily grass), corn silage and purchased corn grain and soybean meal. The second contains hay crop silage (that is mostly legume), corn silage, high moisture

Danny G. Fox is an Associate Professor of Animal Science and Caroline J. Nowak is a Research Support Specialist in Agricultural Economics, Cornell University, Ithaca, New York.

<sup>1</sup> These feeding programs are consistent with the representative farms in Milligan, Nowak and Knoblauch.

Table 1: Expected ad libitum Dry Matter Intake, Average Daily Gain and Crude Protein Requirements; Large Frame Heifers, 2-24 Months of Age.

Age (Months)	Weight Range (lbs.)	Dry Matter Intake lbs./day <sup>a</sup>	Required Average Daily Gain	% Crude Protein in DM <sup>b</sup>
2-4	160-250	6.6	1.4	14
4-6	250-340	8.7	1.6	14
6-8	340-450	10.8	1.9	14
8-10	450-560	13.0	1.8	13
10-12	560-670	15.0	1.7	12
12-14	670-740	16.4	1.2	10
14-16	740-810	18.0	1.2	10
16-18	810-880	19.2	1.2	10
18-20	880-960	20.0	1.2	10
20-22	960-1030	20.5	1.2	10
22-24	1030-1100	21.0	1.2	10

<sup>a</sup> Based on studies of Fox (1981), these values correspond closely to those suggested by the NRC publication, Nutrient Requirements of Dairy Cattle (1978).

<sup>b</sup> Requirements as a percentage of ration dry matter.

Table 2: Dry Matter Intake, Average Daily Gain and Crude Protein Requirements; Steers, 540-1250 lbs.

Weight Range	Average Weight	Dry Matter Intake <sup>a</sup> lbs./day	Minimum Average Daily Gain <sup>b</sup> (lbs./day)	Required % Crude Protein in DM <sup>c</sup>
540-850	690	15.1	2.0	12
850-1250	1050	19.3	2.0	10.5

<sup>a</sup> Steer treated with Rumensin and growth stimulant

<sup>b</sup> Minimum daily gain acceptable as an average for that growth period, considering time on feed, cost of gain and achieving acceptable carcass quality.

<sup>c</sup> Requirement as a percentage of ration dry matter.

shelled corn and purchased soybean meal. Both feeding programs were formulated for three alternative forage compositions: hay only, 50 percent hay crop and 50 percent corn silage on a dry matter basis, and corn silage only. Table 3 contains the average nutrient composition of the feed ingredients used to formulate the rations. See Fox and Nowak for ration compositions.

#### FEEDING PROGRAM: BIRTH TO TWO MONTHS

It is assumed that the calves have a birth weight of 100 pounds and are fed milk replacer and a complete dry calf starter, gaining an average of 60 pounds for the first 60 days (Fox and Warner). On the average, the calves can be expected to consume 40 pounds of commercial milk replacer and 42 pounds of calf starter in this period. The composition of a calf starter developed and used at Cornell is shown in Table 4.

As with the heifers, rearing the steer calves for the first six to eight weeks requires a greater concentration of management than does any other phase of the growth period. It is unlikely that dairymen who do not like starting calves or are having problems rearing the heifers will want to raise steers. Key management factors include adequate colostrum intake within the first few hours after birth, proper sanitation and ventilation, proper feeding of the milk replacer and calf starter, and early detection and treatment of sickness. Procedures for a sound calf-rearing program are outlined by Fox and Warner. It is assumed that practices similar to those prescribed by Fox and Warner are followed, resulting in calves at eight weeks of age that are healthy and therefore capable of maximum performance. Calves are castrated, dehorned, vacci-

nated and wormed between six and eight weeks of age.

#### FEEDING PROGRAM: TWO MONTHS TO SLAUGHTER

Rations were designed for five groupings of weaned calves by size and nutrient requirements. These are: heifers 2-6 months, heifers 6-12 months, heifers 12-24 months, steers 9-14 months, and steers 14-21 months.

The steers are assumed to be with the heifer group until nine months of age. At this age, the steers would be sold as feeders or placed in the two steer groups until slaughter.

Five groups would not be practical on farms with a small number of calves. Under these conditions, an appropriate average of rations should be fed, depending on the average size of the calves being fed together. The ration best suited to the lightest animals is fed to prevent their gains from being suppressed. Also more bunk space should be allowed per head when the variation in size within a pen is large. Since the rations are balanced to be fed *ad libitum*, feed should be available to the cattle at all times.

#### TOTAL FEED REQUIREMENTS

Total feed requirements for rearing the heifers and steers to the different stages of growth are shown in Tables 5-8. The performance simulation program developed for a microcomputer by Fox et al. (1981) was used to determine total feed use. Note that nearly all of the diet for the heifers after the first year of age is forages; supplemental energy and protein are needed, however, for the first year.

Table 3: Nutritional Value of Available Feeds

Feed	Dry Matter %	NE m Mcal/lb	NE q Mcal/lb	Crude Protein %
Mixed mostly grass hay	88	.55	.24	12.5
Mixed mostly legume hay crop silage	47	.58	.27	17
Corn Silage	33	.71	.46	8.5
Dry shelled corn	89	1.02	.67	10
High moisture shelled corn	70	.89	.58	10
Soybean meal (48% CP)	90	.91	.60	53.9
Limestone	99	—	—	—
Dicalcium phosphate	99	—	—	—
Trace mineral salt	99	—	—	—
Urea	89	—	—	281

Table 4: Recommended Complete Calf Starter<sup>a</sup>

Calf Starter	Percent of Ration
Corn and Cob Meal	20
Crushed Oats	25
Beet Pulp	16
Brewer's Grains	10
Soybean Meal (44% CP)	18
Molasses	7
Minerals	4

<sup>a</sup> Fox and Warner

Table 5: Feed Consumed by Heifers on Farms Having Mixed Mostly Grass as the Hay Source

Age, Months	Expected Weight (lbs.)	Hay <sup>a</sup>	Feed (lbs. Consumed as Fed)				
			Corn Silage	Dry Corn <sup>b</sup>	SBM <sup>c</sup>	Dicalcium Phosphate	TM Salt
<u>Hay Based Ration</u>							
2-6	160-340	688	0	337	78	20	5
6-12	340-670	1,458	0	1,200	169	26	13
12-24	670-1100	6,922	0	1,616	0	0	38
Total		9,068	0	3,153	247	46	56
<u>50% Corn Silage/50% Hay Crop Based Ration</u>							
2-6	160-340	432	1,148	135	100	25	5
6-12	340-670	922	2,379	726	253	26	13
12-24	670-1100	4,979	10,195	0	0	40	40
Total		6,333	13,722	861	353	91	58
<u>Corn Silage Based Ration</u>							
2-6	160-340	285	1,785	0	123	25	5
6-12	340-670	0	6,563	0	363	38	13
12-24	670-1100	4,979	10,195	0	0	40	40
Total		5,264	18,543	0	486	103	58

<sup>a</sup> Mixed mostly grass<sup>b</sup> Dry Shelled corn<sup>c</sup> Soybean meal

Table 6: Feed Consumed by Heifers on Farms with Mixed Mostly Legume Hay Crop Silage and High Moisture Shell Corn Based Ration

Age Months	Expected Weight (lbs.)	Hay <sup>a</sup>	Feed (lbs. Consumed as Fed)				
			Corn Silage	HMSC <sup>b</sup>	SBMC <sup>c</sup>	Dicalcium Phosphate	TM Salt
<b>Hay Based Ration</b>							
2-6	160-340	1,168	0	570	22	25	5
6-12	340-670	2,206	0	1,933	113	26	13
12-24	670-1100	13,609	0	1,817	0	0	39
Total		16,983	0	4,320	135	51	57
<b>50% Corn Silage/50% Hay Crop Based Ration</b>							
2-6	160-340	816	1,163	221	55	20	5
6-12	340-670	1,479	2,105	1,299	197	38	13
12-24	670-1100	11,237	7,075	0	0	0	39
Total		13,532	10,343	1,520	252	58	57
<b>Corn Silage Based Ration</b>							
2-6	160-340	550	1,898	0	88	25	5
6-12	340-670	0	6,526	0	334	38	13
12-24	670-1100	11,237	7,075	0	0	0	39
Total		11,787	15,499	0	422	63	57

<sup>a</sup> Hay crop silage<sup>b</sup> High moisture shelled corn<sup>c</sup> Soybean meal

Table 7: Feed Consumed by Steers on Farms Having Mixed Mostly Grass as the Hay Source

Age, Months	Expected Weight (lbs.)	Hay <sup>a</sup>	Feed (lbs. Consumed as Fed)					
			Corn Silage	Dry Corn <sup>b</sup>	SRMC <sup>c</sup>	Urea	Dicalcium Phosphate	Lime- stone
<b>Hay Based Ration</b>								
2-9	160-540	1,320	—	885	152	—	30	—
9-14	540-850	1,287	—	1,053	97	—	11	—
14-21	850-1250	2,244	—	2,026	—	—	—	—
Total		4,851	—	3,964	249	—	41	—
<b>50% Corn Silage/50% Hay Crop Based Ration</b>								
2-9	160-540	832	2,218	475	212	—	35	—
9-14	540-850	842	2,246	637	121	—	11	—
14-21	850-1250	1,367	3,646	1,352	—	20	—	—
Total		3,041	8,110	2,464	333	20	46	—
<b>Corn Silage Based Ration</b>								
2-9	160-540	250	4,817	—	286	—	41	—
9-14	540-850	—	5,816	—	217	—	22	22
14-21	850-1250	—	10,609	400	—	44	20	20
Total		250	21,242	400	503	44	83	42

<sup>a</sup> Mixed mostly grass<sup>b</sup> Dry shelled corn<sup>c</sup> Soybean meal

Table 8: Feed Consumed by Steers on Farms Having Mixed Mostly Legume Hay Crop Silage as the Hay Source

Age, Months	Expected Weight (lbs.)	Feed (lbs. Consumed as Fed)						
		HCS <sup>a</sup>	Corn Silage	HMSC <sup>b</sup>	SBM <sup>c</sup>	Urea	Dicalcium Phosphate	Lime-stone
<b>Hay Based Ration</b>								
2-9	160-540	2,109	—	1,451	75	—	35	—
9-14	540-850	2,048	—	1,719	—	—	11	—
14-21	850-1250	3,227	—	3,196	—	—	19	—
Total		7,384	—	6,366	75	—	65	—
<b>50% Corn Silage/50% Hay Crop Based Ration</b>								
2-9	160-540	1,443	2,055	832	145	—	36	—
9-14	540-850	1,383	1,970	1,145	48	—	11	—
14-21	850-1250	2,309	3,288	2,381	—	—	20	—
Total		5,135	7,313	4,358	193	—	67	—
<b>Corn Silage Based Ration</b>								
2-9	160-540	482	4,876	—	242	—	41	—
9-14	540-850	—	5,862	—	205	—	22	22
14-21	850-1250	—	9,585	822	—	43	39	—
Total		482	20,323	822	447	43	102	22

<sup>a</sup> Hay crop silage<sup>b</sup> High moisture shelled corn<sup>c</sup> Soybean meal

Steer rations require more supplementation of grain than the heifers after a year of age because of the higher rate of gain being supported (Tables 7 and 8). Those rations containing hay crop require less supplemental protein, especially when they contain legumes. However, more corn grain must be fed because of the lower energy content of hay crop compared to corn silage. If the steers are not given growth stimulants and Rumensin, rate of gain and feed efficiency will decrease by approximately 15 percent from 540 pounds to slaughter weight (Minish and Fox). The days required to reach 540 pounds could be reduced by about 30 days from that shown in this paper and feed requirements during this phase would be reduced about 6-8 percent if the steers were implanted with a growth stimulant during this phase. Implanting could be done to the steers but not to the heifers during the time they are fed together.

Fox, D. G. Unpublished data, 1981.

Fox, Danny G. and Caroline Nowak. Feeding and Management Strategies for Producing Beef from Holstein Steers on Northeast Dairy Farms. Dept. Agr. Econ. Staff Paper No. 81-17, Cornell University, 1981.

Fox, D. G. and R. W. Warner. "Guidelines for Producing Beef from Dairy Calves." Cornell Beef Production Reference Manual. Fact Sheet 1250, Dept. An. Sci., Cornell University, 1981.

Fox, D. G., C. J. Sniffen, P. J. VanSoest, D. K. Phillips and P. Robinson. "Grow Diet: A Microcomputer Program for Evaluating Rations of Growing Cattle." Dept. An. Sci., Cornell University, 1981.

Fox, D. G., R. L. Weiser and D. K. Phillips. "Beef Gain: A Microcomputer Program to Predict Performance of Growing Cattle." Dept. An. Sci., Cornell University, 1980.

Knoblauch, Wayne A., Robert A. Milligan, Danny G. Fox and Merri L. Woodell. Economic Utilization of Forages in the Production of Milk and Beef in the Northeast United States. Dept. Agr. Econ. Staff Paper No. 79-29, Cornell University, 1979.

## REFERENCES

Dexheimer, C. E., J. C. Meiske and R. D. Goodrich. "A Comparison of Systems of Feeding Corn Silage." Minnesota Beef Cattle Report. B151, University of Minnesota, 1971.

FEEDING AND MANAGEMENT STRATEGIES FOR PRODUCING BEEF FROM HOLSTEIN STEERS ON NORTHEAST DAIRY FARMS

Milligan, Robert A., Caroline J. Nowak and Wayne A. Knoblauch. Profitability of Feeding Dairy Steers to Feeder and Slaughter Weight on Northeast Dairy Farms. Dept. Agr. Econ. Staff Paper No. 81-16, Cornell University, 1981.

Minish, G. L. and D. G. Fox. Beef Production and Management. Reston, VA: Reston Publishing Co., 1981.

National Research Council. Nutrient Requirements of Dairy Cattle. No. 3, fifth revised edition. Washington, 1978.

Reid, J. T., J. K. Loosli, G. W. Trimberger, K. L. Turk, S. A. Asdell and S. E. Smith. "Effects of Plane of Nutrition During Rearing on Lifetime Performance of Dairy Cattle." Proceedings Cornell Nutrition Conference, pp. 23-26, Cornell University, 1963.

Woodel, Merri Lynn. "An Economic Analysis of Alternative Beef Feedlot Systems in New York State." M.S. thesis, Cornell University, 1981.