



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

EPTD DISCUSSION PAPER NO. 86

**SUSTAINABILITY ANALYSIS FOR IRRIGATION WATER
MANAGEMENT: CONCEPTS, METHODOLOGY, AND
APPLICATION TO THE ARAL SEA REGION**

Ximing Cai, Daene C. McKinney, and Mark W. Rosegrant

Environment and Production Technology Division

**International Food Policy Research Institute
2033 K Street, N.W.
Washington, D.C. 20006 U.S.A.**

December 2001

EPTD Discussion Papers contain preliminary material and research results, and are circulated prior to a full peer review in order to stimulate discussion and critical comment. It is expected that most Discussion Papers will eventually be published in some other form, and that their content may also be revised.

ABSTRACT

Sustainable irrigation water management should simultaneously achieve two objectives: sustaining irrigated agriculture for food security and preserving the associated natural environment. A stable relationship should be maintained between these two objectives now and in the future, while potential conflicts between these objectives should be mitigated through appropriate irrigation practices. Lessons learned from unsustainable water management practices around the world demonstrate the necessity—and growing urgency—of applying sustainability principles to water management in river basins where irrigation is a major factor. This paper presents the operational concepts and analytical framework for sustainability analysis of irrigation water management in irrigation-dominated river basins. These are applied to the Aral Sea region in Central Asia, a region famous for its conflict between sustaining irrigated agriculture and preserving the environment. Maintaining current irrigation practices will lead to worsening environmental and economic consequences. Infrastructure improvements and changes in crop patterns will be necessary to sustain the irrigated agriculture and the associated environment in the region. A penalty tax on salt discharge, as an economic incentive, may help address environmental problems while having only a small effect on irrigation profit.

Keywords: Irrigation Water Management, Sustainability, System Analysis, Aral Sea

ACKNOWLEDGMENTS

The authors are grateful to Claudia Ringler and Siet Meijer for their helpful comments and suggestions on an early draft of this paper.

TABLE OF CONTENTS

1. Introduction.....	1
2. Sustainable Irrigation Water Management: Concepts and Analytical Framework.....	4
3. Case Study: The Aral Sea Problem.....	23
4. Conclusions	43
References	46

Sustainability Analysis for Irrigation Water Management: Concepts, Methodology, and Application to the Aral Sea Region

Ximing Cai¹, Daene C. McKinney², Mark W. Rosegrant¹

1. INTRODUCTION

Water scarcity, pollution, and other water-related environmental and ecological problems have been increasing rapidly in many areas of the world. According to Grigg (1996), “The real crisis in water is a ‘creeping crisis’—it comes on slowly but it demands a response right now.” Sustainable development—that is, development that meets the needs of the present without compromising the ability of future generations to meet their own needs—is a concept that has gained popularity since publication of the 1987 Brundtland Commission Report (WCED 1987). Increasingly, researchers and policymakers are advocating sustainable development as the best approach to today’s and future water problems (Loucks 2000).

For water resources management, sustainability implies a notion of equilibrium that simultaneously satisfies water demands and the preservation of the water resources system. This paper focuses on achieving a sustainable balance between irrigation management—the largest water use worldwide—and environmental preservation. Over the last 30 years irrigated areas have increased rapidly, helping to boost agricultural output and feed a growing population. Irrigation uses the largest fraction of water in almost all countries.

¹ International Food Policy Research Institute.

² Department of Civil Engineering, University of Texas, Austin.

Globally, 70 percent of freshwater diverted for human purposes goes to agriculture, and irrigation water demand is still increasing because the area being irrigated continues to expand. In some countries, the expansion of surface water use appears to be approaching the physical limit, and groundwater abstractions are increasingly exceeding rates of replenishment. Meanwhile, industrial and domestic water demand has been increasing rapidly as a result of increasing economic development and urbanization. In some countries and regions, water is already being transferred out of irrigation and into urban-industrial uses, putting additional stress on the performance of the irrigation sector (Rosegrant and Ringler 2000).

Although the achievements of irrigation in ensuring food security and improving rural welfare have been impressive, past experience also indicates problems and failures of irrigated agriculture. In addition to large water use and low efficiency, environmental concerns are usually considered the most significant problem of the irrigation sector. Environmental problems include excessive water depletion, water quality reduction, waterlogging, and salinization. The marked reduction in annual discharge of some of the world's major rivers—evident in long-term hydrological records (OECD 1998)—has been attributed, in part, to the large water depletion caused by irrigated agriculture. In some basins, excessive diversion of river water for irrigation (and other uses) has brought environmental and ecological disasters to downstream areas, and groundwater pumping at unsustainable rates has contributed to the lowering of groundwater tables and to saltwater intrusion in some coastal areas. Many water quality problems have also been created or aggravated by changes in streamflows associated with agriculture's consumptive uses. Moreover, inappropriate irrigation practices, accompanied by inadequate drainage, have

often damaged soils through over-saturation and salt build-up. The United Nations Food and Agriculture Organization (FAO) estimates that 60 to 80 million hectares are affected to varying degrees by waterlogging and salinity (FAO 1996). Finally, these irrigation-induced environmental problems threaten not only agricultural production systems but also human health and the environment.

Various agencies and researchers have identified broad guidelines for sustainable water resources management. The World Bank (Serageldin 1995) proposes a comprehensive approach, emphasizing economic behavior, the overcoming of market and policy failures, more efficient use of water, greater protection of the environment, and moving to “demand management” from the previously dominant “supply management.” In 1992, the United Nations Conference on Environment and Development (UNCED) in Rio de Janeiro arrived at the conclusion that water should be considered an integral part of the ecosystem, “a natural resource and social and economic good” (UNCED 1992). More recently, the American Society of Civil Engineers (ASCE), associated with the United Nations International Hydrologic Program (UN/IHP) published a monograph on sustainable water resources management (ASCE and UN/IHP 1998). According to that report, “sustainable water resource systems are those designed and managed to fully contribute to the objectives of society, now and in the future, while maintaining their ecological, environmental and hydrological integrity.”

Previous studies have set out several guidelines for sustainable water resources management. They can be briefly outlined as follows:

- successfully accomplish multiple social, economic, and environmental objectives in terms of adequate water quantity and quality;

- maintain stability and flexibility in water supply so as to deal with extreme events such as flooding, drought, excessive waste discharge, and other anticipated stochastic events;
- minimize negative environmental impacts, especially the long-term cumulative negative;
- realize equity to make equitable water rights possible among spatially distributed water demand sites and between current and future generations;
- achieve financial and economic efficiency; and
- adapt to new technology.

No doubt these guidelines could provide some assistance and guidance to those who are actually involved in planning and decision making in specific regions. However, these broad guidelines still need to be translated into operational concepts that can be applied to the planning and management of water resources systems in specific basins. This paper discusses the operational concepts and analytical framework for sustainability analysis in irrigation. These are applied to a case study area, the Aral Sea region in Central Asia, which may have experienced the most serious environmental disaster caused by excessive irrigation (Micklin 1993).

2. SUSTAINABLE IRRIGATION WATER MANAGEMENT: CONCEPTS AND ANALYTICAL FRAMEWORK

2.1. CONCEPTS OF SUSTAINABILITY: DEVELOPMENT VS. PRESERVATION

An integrated system comprises irrigation, crop production, and the environment. Irrigation sustains crop production systems. However, a sole focus on irrigation

development, without taking environmental preservation into consideration, is doomed in many regions of the world. Increased soil and water salinity resulting from extensive irrigation practices have already diminished opportunities to develop the crop production system. Although human society achieved a relatively stable balance between irrigation development and environmental preservation for several thousand years; during the last 30 to 50 years that relationship has been destroyed in some regions by inappropriate irrigation practices.

The purpose of sustainable water resources management is to sustain both the water supply capability and the environment, now and in the future. Water supply capability encompasses both the availability of water and the infrastructure to sustain water supply and use. The environment takes into account the water source and the land and air systems that support human production activities. As water demands in agricultural, municipal, and industrial uses change over time—because of policy and technological changes, among others—the relationship between water use and the environment needs to be continually reviewed and adapted. In river basins where irrigation is the major water use, sustainable water management should ensure a long-term, stable, and flexible water supply to meet crop demands, as well as growing municipal and industrial demands, while at the same time mitigating or preventing negative environmental consequences from irrigation.

Sustainability reflects a systems concept for irrigation water management, that is, applying a set of elements that interact in interdependent fashion. Moreover, sustainability, by its nature, implies a dynamic system whose status is determined by a balance of opposing forces or trends (Svendsen 1987). When an accelerating flow of negative forces reaches a threshold beyond which it is impossible or inordinately costly to

reverse the direction of the change and return to a more favorable equilibrium, the system becomes unsustainable. Another feature of the dynamic system, perhaps the most pervasive one for sustainability issues, is its association with intergroup and intertemporal externalities.

The following section examines how sustainability can be measured and how it can be achieved in the context of irrigation water management.

2.2. MEASURING SUSTAINABILITY: INDICATORS

A set of manageable indicators of sustainability based on broad guidelines and principles are necessary to detect problems as they arise and to provide an early warning system for decision makers. The indicators should be monitored and measured on the basis of the performance of natural systems and anthropogenic interactions, and action should be taken once specified thresholds are passed. In particular, the indicators should be helpful in tracing long-term cumulative environmental changes due to irrigation practices, which can potentially create irreversible problems.

In the context of arid or semi-arid basins where irrigation is the dominant water use, sustainability in irrigation water management can be indicated by

- water supply system reliability, reversibility, and vulnerability,
- environmental system integrity,
- equity in water sharing, and
- economic acceptability.

These indicators, defined at the basin scale, are supposed to be used by basin authorities or related national administrative agencies, instead of individual farmers. An example of indicators designed for farmers can be found in MAF (1997). Moreover, definitions of these indicators should be analytically sound and measurable in a modeling framework, which will be illustrated in the following.

2.2.1. *Reliability, Reversibility, and Vulnerability of the Water Supply System*

Water supply systems are subject to substantial risk because of inherent stochastic variability and a fundamental lack of knowledge. Sustainable water resources management requires a stable water supply system with enough flexibility to deal with various extreme conditions. Risk has been identified as one of the key sustainability issues in water resources management (Simonovic 1997). The traditional measures of system performance (mean value or variance of some variables) are insufficient to capture risk behavior, and additional criteria must be used to quantify recurrence, duration, severity, and other consequences of unsatisfactory system performance. These criteria include reliability, reversibility, and vulnerability (Kundzewicz and Kindler 1995).

Reliability represents the probability of a system's success state and is complementary to risk, which represents the frequency of system failure. Reliability, as used in water resources management, comprises three terms: *occurrence reliability* (the ratio of the number of periods of system success to the number of periods of operation), *temporal reliability* (the ratio of the time the system is in a success state to the total time of operation), and *volumetric reliability* (the ratio of the volume of water supplied to the total volume demanded).

Reversibility or ***resilience*** is the probability the system can recover from failure to some acceptable state within a specified time interval. Fiering (1982) proposed several alternative indices of resilience, including how long the system remains in the satisfactory state and the steady state probability of the system being in the satisfactory state.

Vulnerability represents the severity or magnitude of a system failure. Hashimoto, Stedinger, and Loucks (1982) developed a metric for overall system vulnerability as the expected maximum severity of a sojourn into the set of unsatisfactory states. They emphasized the maximum severity (how bad things are) for each unsatisfactory state and the probability that the failure with the maximum severity would occur.

These risk indicators can be used for various aspects such as water quantity and quality, crop area, yield or production, and flow requirement for environmental and ecological purposes. The selection depends on the specific analytical objectives. The threshold for “system failure” (for example, water supply is 25 percent below water demand), which is a managerial assumption, has to be considered in the quantification of the risk indicators discussed above.

2.2.2. *Environmental System Integrity*

A guiding criterion for sustainable irrigation water management is to minimize the interference of the irrigation system with the associated environmental system, including the effects on the water bodies that receive irrigation water through wind-drift, surface runoff, or drainage to groundwater. In addition, to sustain irrigation profit over the long term, irrigation water management must meet legislative requirements with respect to the environment.

Indicators for environmental system integrity fall into three categories:

(1) Health of aquatic and floodplain ecosystems. Extensive irrigation can affect drinking water health as indicated by bacteria, nutrients, and toxic contaminants, and soil health as indicated by the soil's water-holding capacity, total organic Nitrogen and Carbon, PH value, and the conditions of surface aggregates.

(2) Water quality. Irrigated agriculture affects water quality in several ways, including higher chemical-use rates associated with irrigated crop production, increased field salinity resulting from applied water, accelerated pollutant transport with drainage flows, groundwater degradation due to increased deep percolation to saline formations, and greater instream pollutant concentrations due to flow depletion.

(3) Soil degradation. Irrigation is responsible for soil waterlogging and salinization in many regions where drainage systems are poor; irrigation with traditional furrow systems also causes soil erosion that can be measured by the extent of topsoil losses.

Thus, the adverse environmental effects of irrigation (such as waterlogging and salinization, groundwater pollution, and soil erosion) are often cumulative and may develop to an irreversible state because of long-term poor irrigation management. The measure of these indicators should be connected to both the short-term irrigation practices and performances and the long-term dynamic transmissions through some physical processes.

2.2.3. Equity

Equity is one of the basic concepts of sustainable development. Water is not just an economic resource; it is also a community resource with deep emotional and symbolic

value. Experience has shown that plentiful and clean water flows toward the rich and powerful and away from the poor and powerless (Ingram 1997). Another emerging issue is the relationship between the environment and equity, which needs to better qualify the water needs of ecosystems and the economic, social, and cultural values, function, and services that aquatic ecosystems provide. Thus, equity in water resources management involves complex natural, political, and socioeconomic factors. The ASCE view of equity in sustainable water resources systems would allow people, “now and then” and “here and there” to share the water use right (both benefit and cost) in such a way that no one should be disadvantaged or inadequately compensated (ASCE and UN/IHP1998).

Factors affecting temporal equity and spatial equity in water resources development can be either anthropogenic or natural, or both. Temporal equity is associated with resource depletion and long-term cumulative consequences that may lead to damages or even disasters in the future. Spatial equity often concerns the conflict between upstream and downstream areas in a river basin and the conflict between various water users. As mentioned in the introduction, the continued availability of water for irrigation will be threatened in many regions by rapidly increasing nonagricultural water uses (in industry, households, and the environment), which normally have a higher marginal return than irrigation. The same conflict exists between farmers who plant high-valued crops and those who plant low-valued crops. Continued research is needed to measure the social value, or the public welfare component, of water to ensure that, when water is transferred, the area of origin will retain sufficient water to protect social value and to adequately compensate losses that cannot be prevented (Ingram 1997).

Moreover, water quality is also an equity issue. For example, conflicts arise when upstream users release excessive pollutants into the river and downstream users suffer damage resulting from the poor water quality. The measure of equity can be descriptive and normative. Descriptive measures simply evaluate the dispersion of the benefit using some descriptive statistic. Normative measures are derived from some underlying social welfare function (Tsur and Dinar 1995).

2.2.4. Economic Acceptability

Besides food self-sufficiency, achieving net profit over the long term is the motivating factor that sustains irrigated agriculture. Economically acceptable irrigation systems provide lifestyle and social options for farmers and also contribute to the wider economy and community.

From the perspective of using water more economically, the great challenge in irrigated agriculture is to include the opportunity costs of irrigation water supply, which are often an order of magnitude higher than current charges (Briscoe 1999). Another challenge is to include the long-term economic damage to the environment due to irrigation.

To maintain and improve economic acceptability, some regions will require investments both to enhance water supply capacity and to increase water use efficiency. Given all these considerations, the marginal benefit and marginal cost associated with irrigation development and management can be assessed. When the marginal benefit is less than the marginal cost, the irrigation practice loses its economic acceptability, which implies an unsustainable state.

2.3. ACHIEVING SUSTAINABILITY: AN ANALYTICAL FRAMEWORK

The indicators outlined above provide information for decisionmaking in water resources management. They are not an end in themselves but rather the means to influence the decisionmaking process. How can sustainability be achieved? It can be done through decisions on national or regional agricultural policy, basin/sub-basin water allocation, and water management at the field level. The following describes an analytical framework to support these decisions at various levels.


A river basin forms a natural boundary for water resources planning and management in which water interacts with and, to a large degree, controls the extent of other natural components in the landscape such as soils, vegetation, and wildlife. Human activities can also be organized and coordinated within the river basin unit. Physical processes, such as flow and constituent balances are governed by natural laws and are also affected by human actions, including impoundment, diversion, irrigation, drainage, and discharges from urban areas. Therefore, decisionmaking in water resources management should be based on physical processes and should also take into account artificial “hardware” (infrastructure) and “software” (management policies). A modeling framework can help identify and analyze the issues in the context of the river basin.

Figure 1 presents a diagram of a modeling framework for decisionmaking, showing decision options at the river basin and farm (demand sites) levels.

Water can be used for instream purposes, including hydropower generation, recreation, and waste dilution, and for offstream purposes that are differentiated into agricultural, municipal, and industrial water uses. The rapid increase in municipal and

industrial water demand in many regions of the world will likely not only require increasing transfers of water out of agriculture but also lower the quality of downstream irrigation water if municipal and industrial wastewater discharge is not appropriately controlled.

Figure 1—A decision framework for river basin management modeling



Taking into account instream water requirements for hydropower generation, recreation, and ecological uses involves setting aside sufficient water over space, time, and distribution between instream and offstream water uses.

The social benefit of water uses should be an important component of a basin water management strategy. That means both the positive contribution from the economic value of irrigated agriculture and the environmental damage from salinity and waterlogging, soil erosion, and water quality degradation resulting from irrigation. The social benefit also relates to the benefits and costs of municipal and industrial (M&I) water uses.

The basin water management system should be governed by national or regional agricultural policies, including trade and macroeconomic policies, agricultural input and output prices, government investments in technology and infrastructure, and institutional resource management policies, such as market quotas, water allocation policies, and soil conservation programs. Policy instruments applied directly to water management could include a mix of water prices, penalty taxes on waste discharge and irrigation drainage, and water rights. The impact of these policies on agricultural sector growth and on environmental sustainability in the region should be analyzed along with decisions on water management at the basin/sub-basin and farm levels.

Decisions at the basin/sub-basin and farm levels may be considered separately. Decisions at the basin/sub-basin level should promote an appropriate expansion of infrastructure capacity and optimal operation of existing reservoir, aquifer, and stream systems. They should also take into account water allocation among different sectors and users, especially in transboundary situations, as well as the interaction between those sectors and users. While keeping in mind water rights within a basin, decision makers should consider the efficiency with which water is used at different places in the basin by different users and the degree to which different uses degrade the water quality (Batchelor 1999). Efficiency at the river basin level can be improved by (1) increasing output per unit of evaporated water, (2) reducing losses of usable water to sinks, (3) reducing water pollution, and (4) reallocating water from lower valued to higher valued uses (Seckler 1996).

Batchelor (1999) suggested several ways to improve physical and economic efficiency at the farm level:

- agronomic (for example, improving crop husbandry and cropping strategies);
- technical (for example, installing an advanced irrigation system);
- managerial (for example, adopting demand-based irrigation scheduling systems and better maintaining equipment); and
- institutional (for example, introducing water pricing and improving the legal environment).

Supply and demand management policies may also affect decisionmaking. A portion of the growing demand for water can be met by investing in water supply and utilization systems, and some potential exists for expanding nontraditional sources of water supply (for example, desalinization). However, in many arid or semi-arid areas, water is no longer abundant, and the high economic and environmental costs of developing new water resources limit how much supply can expand. Demand management aims to better utilize existing water resources by curbing inessential or low-value uses through price or nonprice measures. The modeling framework should provide information to analyze trade-offs between the decisions for the two categories. In most cases, new sources and improved demand management are both necessary, and joint decisions for supply and demand should be made for sustainable water resources management.

These water management issues, policies, and decisions are complex and integrated with the social, economic, and environmental development and sustainability of the basin. The process of achieving sustainable water management requires a framework for decisionmaking. This framework should be a dynamic system that includes modeling

components capable of analyzing the effects of the proposed policies and strategies over periods of time long enough to see the cumulative, long-term effects on the system. The components shown in Figure 1 are interrelated, and the interrelationships vary as external inputs (for example, climate variation, population increase, investment in infrastructure) change over time.

2.4 IMPLEMENTATION OF THE ANALYTICAL FRAMEWORK

The modeling framework for sustainability analysis combines traditional water resources management models with indicators that are based on newly developed sustainability criteria. Specifically, the modeling framework needs to simulate the performance of short-term decisions so the sustainability indicator can be quantified by some prescribed mathematical forms. Moreover, the indicators, embedded in the modeling framework, should be used to evaluation rules, and thus form a managerial control to the short-term decisions so as to ensure long-term sustainability. The following section discusses some issues related to implementing this kind of analytical framework.

2.4.1. Integrated Short-Term and Long-Term Objectives

Irrigation water planning and management should balance short-term and long-term objectives. They are neither totally consistent nor totally in conflict with each other. Short-term objectives focus on current benefits, while long-term objectives aim to sustain current and expected benefits into the future. Long-term decisions must account for the long-term consequences of short-term decisions in a way that avoids possible negative future effects of current decisions. Put another way, the long-term situation improves when short-term decisions balance current and future benefits.

2.4.2. Integrated Hydrologic-Agronomic-Economic Modeling Components

The interdisciplinary nature of water resources problems requires the integration of technical, economic, environmental, social, and legal aspects into a coherent analytical framework, so that both economic and environmental consequences of policy choices can be examined. Hydrologic relationships provide a physical basis to evaluate water availability and water quality conditions. Accurate estimates of deep percolation, return flows, and their contaminant concentrations, as well as groundwater levels, are essential to evaluate the environmental effects of irrigation. Assessing the damage resulting from the depletion of water over time is also critical to evaluating the environmental effects of irrigation. Long-term simulations of these processes are necessary to trace the cumulative consequences such as waterlogging, soil salinization, and groundwater quality reduction.

Crop production functions connect water, soil, and other inputs with crop production, which is the fundamental building block for estimating the demand for and value of water in irrigation. An ideal crop-water production model should be flexible enough to address issues at the crop, farm, and basin levels. The production function should allow the assessment of policy-related problems and, in particular, be sufficiently comprehensive to allow the estimation of externality effects. Moreover, the modeling framework should generally include the valuation of nonagricultural uses of water, such as values for domestic demand; commercial, industrial, and mining demand; recreational demand; and environmental demands such as maintenance of instream river flows and flushing of pollutants. (See McKinney et al. (1999) for a comprehensive review of agricultural and nonagricultural water valuation.)


Only by considering all interactive components that benefit from or damage the resource can the optimal use be established from a social standpoint. Thus, with the growing scarcity of water and increasing competition for water across sectors, the economic aspects of water allocation are increasingly important in river basin management. Important economic issues that need to be examined through integrated economic-hydrologic river basin modeling include transaction costs, the effects of allocation mechanisms on agricultural productivity, intersectoral water allocation, the environmental effects of allocations, and property rights for different allocation mechanisms (Rosegrant and Meinzen-Dick 1996). Moreover, institutional relationships present directives aimed at achieving equity in water resources management.

An integrated system should be able to explore the interdependence between economic development and environmental consequences, and between short-term decisions and potential long-term problems. The outcomes of water use can then be examined in terms of efficiency, equity, and environmental impact. Over time, these outcomes change the environment through processes such as salinization and waterlogging, siltation, industrial water pollution, technological change, crop diversification, and legislative and institutional change. These processes are critical for understanding the dynamic changes occurring in the environment associated with water use systems, and for implementing appropriate controls on the actions that drive the system to sustainability thresholds.

2.5. A PROTOTYPE MODELING FRAMEWORK FOR SUSTAINABILITY ANALYSIS

A long-term dynamic modeling framework following the concepts and methodology discussed above was developed and reported in the dissertation of Cai (1999). Figure 2 shows a diagram of the modeling framework.

Figure 2—Components and structure of the modeling framework



The core of the modeling framework is an Inter-Year Control Program (IYCP) and a sequence of Yearly Models (YMs). The yearly model for year y , YM^y , is a short-term (annual) optimization model developed at the river basin scale. The objective function of the model is to maximize total water use net benefit in a river basin for that year. The model includes essential integrated hydrologic, agronomic, and economic components such as (1) flow and pollutant (salt) transport and balance in the river basin network, including the crop root zone; (2) irrigation and drainage processes; (3) crop production functions, including the effects of both water stress and soil salinity; (4) benefit functions for both instream and offstream uses, accounting for economic incentives for salinity

control and water conservation; (5) tax and subsidy systems to induce efficient water allocation, improve irrigation-related capacities, and protect the environment; and (6) institutional rules and policies that govern water allocation.

The model is based on a node-link network with source nodes—such as rivers, reservoirs, and groundwater aquifers—and demand site nodes—such as agricultural, municipal and industrial (M&I), and ecological demand sites, and hydropower stations. Detailed agricultural water demand, instream water uses (including flow release for environmental and ecological use), and hydropower generation are modeled. Details of the yearly model can be found in Cai (1999). In addition, McKinney et al. (1999) provides a comprehensive review of integrated, hydrologic-agronomic-economic models at the basin scale; and Rosegrant et al. (2000) illustrates the application of such a model to the Maipo basin in Chile.

The *IYCP* is a long-term model which uses some prescribed indicators based on sustainability criteria to control relations between short-term irrigation practices and their long-term socioeconomic and environmental consequences (see Figure 2). The long-term decisions include the regulation of inter-year reservoir storage, irrigation and drainage infrastructure improvements, changes in irrigated area and crop patterns, and economic incentives (for example, salt penalty taxes.) The thesis of this framework is that short-term (intra-year) decisions should be controlled by long-term (multi-year) sustainability criteria to reach sustainable planning and management decisions.

The long-term optimization/simulation model of the basin incorporates specific indicators of sustainability, taking into account risk minimization in water supply, environmental conservation, equity in water allocation, and economic efficiency in water

infrastructure development. These indicators are measured by some approximate methods.

Risk is represented by reliability, reversibility, and vulnerability in terms of sustained irrigated area and flow for ecological use, reflecting how much irrigated area/ecological flow can be sustained, how often irrigated area/ecological flow drops below an assumed target, and how serious the deficits are in a long-term time horizon (30 years).

Environmental indicators are identified as worst water and soil salinity conditions in the basin over the long run. Temporal equity (“now and then”) is defined as the standard deviation of the rate of change of total water use benefit in the basin over all years; and spatial equity (“here and there”) is defined as the standard deviation of the average rate of change of water use benefit over all demand sites. Economic indicators are represented by the ratio of the marginal benefits to the marginal costs of additional water infrastructure improvements. Mathematical representations of these indicators are given in Cai, McKinney, and Lasdon (2001).

The results of the sequence of yearly models (YM^y), over a long time horizon under a particular selection of the inter-year decision variables, are used to calculate the indicators, which are then used to evaluate the long-term performance of some policy options represented by the long-term decision variables. To find optimal long-term policies, these modeling processes are implemented in an iterative form as shown in Figure 2. The solution approach to the modeling framework is presented in Cai, McKinney, and Lasdon (2001).

3. CASE STUDY: THE ARAL SEA PROBLEM

The Aral Sea, a land-locked lake, is located in the semi-arid and desert areas of Central Asia (Figure 3). Its level is determined by the inflow of two feeding rivers, the Amu Darya River and the Syr Darya River. The average inflows from the Amu Darya and the Syr Darya rivers once were 72 cubic kilometers (km^3) and 37 km^3 per year, respectively, but have now decreased to a mere trickle because of large-scale irrigation water withdrawals and high water consumption during water distribution, conveyance, and field application.


Figure 3—The Aral Sea basin in Central Asia



Compared with the status in 1960 when the Aral Sea was the world's fourth largest inland lake, by the early 1990s the lake had shrunk to about half its size, was 16 meters below its former level, and three times as salty (Micklin 1993). Figure 4 shows the

evolution of the surface area of the Aral Sea and of the irrigated areas in the basin over the last 80 years.

Figure 4—Irrigated area (million hectares) in the Aral Sea basin and surface area (square kilometers) of the Aral Sea




Source: Based on Micklin 1993.

In fact, the effects of extensive irrigation in the Aral Sea basin extend far beyond the fate of the sea itself. Thirty-five million people have lost access to the lake's water, fish, reed beds, and transport functions. Moreover, far-reaching environmental and ecological problems, such as dust storms, erosion, and poor water quality for drinking and other purposes, have negatively affected human health and economic development in the region. Thus, the Aral Sea disaster is the prime example for unsustainable irrigation development: rapid, large-scale expansion; sole reliance on high-water-use production systems for cotton and rice; poor water distribution and drainage; inefficient irrigation

techniques resulting in enormous losses of irrigation water; and large-scale, non-dose-related uses of fertilizers and pesticides.

Glantz (1999) argues that the Aral Sea region has experienced “creeping” environmental changes, that is, slow-onset, low-grade, and cumulative. These creeping changes can be seen in Figure 5, which shows the upstream and downstream runoff during 1959–98.

Figure 5—Upstream/downstream runoff of the Syr Darya River




Sources: Glantz (1999) for flows during 1959–88; Personal communication, V. Chub, Uzbekistan Main Hydromet, 2000, for flows during 1989–98.

The year 1960 is typically considered the beginning of the deterioration. A so-called “pre-problem” stage started in the mid-1960s, when the first signs of adverse environmental effects began to appear. The downstream flow depletion became more apparent during 1976–81 and full-fledged during 1981–85. The final widespread recognition of the Aral Sea crisis dates to 1986–87, when several threshold levels had been

surpassed. Actions to solve the problem began only in 1990. Unfortunately, by that time, the environmental costs resulting from excessive irrigation surpassed the economic capacity of the newly independent republics in Central Asia (World Bank 1992).

One indicator of the increasing environmental deterioration in the basin was the change in crop yields, as irrigation-induced water quality and soil quality problems led to a reduction in yields. Figure 6 shows the average annual growth rates of seed cotton yield in three Central Asian countries during different periods. All three countries achieved positive growth during 1960–80, but growth turned negative in the 1980s.

Figure 6—Cotton yield annual growth rate in three Central Asian countries



Moreover, the downstream country of Kazakhstan experienced a larger cotton yield reduction, likely because of more serious water and soil quality degradation. Although the changes in the political and economic situation in the region also likely contributed to the

decline in agricultural production in the basin, a significant share was probably caused by growing water and soil quality problems.

For a closer look at sustainability issues in irrigation management, we examine these issues in greater detail in one of the two major basins in the Aral Sea region, the Syr Darya River basin. The river begins at the Pamir and Tien Shan plateaus, crosses the territories of four Central Asia republics (Kyrgyzstan, Tajikistan, Uzbekistan, and Kazakhstan), and terminates in the Aral Sea. The basin's water supply system comprises 9 major tributaries, 11 reservoirs, 23 irrigation distribution systems, and numerous distribution canals.

3.1. CURRENT STATUS OF IRRIGATION MANAGEMENT IN THE SYR DARYA RIVER BASIN

3.1.1. *Risk and Vulnerability of the Water Supply System*

Table 1 shows total renewable water resources, total water withdrawals, and the ratio of withdrawals to water availability in the Syr Darya River basin for selected years from 1961 to 1990 (Glantz 1999). Even in the early 1960s, the basin system was vulnerable in dry years (1961 was a dry year). The high ratios of water withdrawals to renewable water in 1980 and 1990 show that a considerable amount of water in the basin might be reused even in average years.

Table 1—Runoff and irrigation water withdrawals, selected years, Syr Darya basin

	1961	1970	1980	1990
Annual runoff (km ³)	30.0	52.1	39.8	39.3
Irrigation withdrawals (km ³)	27.8	41.0	48.3	43.5
Total withdrawals (km ³)	30.9	47.1	56.8	53.0
Withdrawal/runoff (%)	103	90	143	135

Source: EC (1995).

Raskin et al. (1992) categorized five hydrologic patterns in the basin, with probabilities of occurrence of 3.3 percent (very wet), 16.7 percent (wet), 52.0 percent (normal), 21.3 percent (dry), and 6.7 percent (very dry). The ratio of annual inflow under various hydrologic levels to the normal level is 1.27 (very wet), 1.12 (wet), 1.00 (normal), 0.76 (dry), and 0.59 (very dry). Dry and very dry years occur with a probability of 28 percent. If water withdrawals are at 1990 levels (53.0 km³), then the ratio of withdrawal to runoff needs to be as high as 1.9 in dry years, and 2.4 in very dry years to meet demands. This can only be achieved at high return flow reuse levels. However, because of the high evaporation loss in the region, 70 to 80 percent of water withdrawals are consumed in the basin. Therefore, water withdrawals in dry and very dry years cannot meet the demand at the 1990 level.

Another indicator of water shortage in the basin is per capita water availability, which was estimated at 7,500 cubic meters (m³) in 1950, 2,000 m³ in 1980, and 700 m³ in 2000 (Klotzli 1994). When the per capita water availability is 1,000 m³, the system can suffer high vulnerability (Gleick 1999). This indicator shows that the water supply system in the Syr Darya could be very vulnerable.


3.1.2. Environmental and Ecological System Degradation

Decline of flow to the Aral Sea

Given the high irrigation water withdrawals and the large evaporation losses in the basin, the tremendous decline in flow into the Aral Sea is not difficult to understand. As shown in Figure 5, the decline of downstream flows started in the mid-1960s. Outflows to the lake dropped to a mere trickle in the mid-1970s, with increases only achieved in the 1990s when the Aral Sea problem was first addressed. However, recent inflows to the lake

are not sufficient to reverse the disaster situation. The level, area, and volume of the sea during 1989–98 are shown in Figure 7. Thus, even in the last decade, the drying up of the sea has not slowed and, in fact, the problem has worsened.

Figure 7—Change of level, area, and volume of the Aral Sea in recent years



Source: Personal communication, V. Chub, Uzbekistan Main Hydromet, 2000.

Degradation of soils

Previously fertile, humus-rich, meadow-swamp soils in delta areas in the zone of influence of irrigation have been transformed into low-productivity, sandy-desert soils with much lower fertility. In the early 1990s, 50 percent of the land in the basin was classified as saline. The soil salinity problem varies along the river. In the upper reaches, less than 10 percent of the land has moderate to strong salinity, while in downstream areas more than 50 percent of the irrigated lands are classified as moderately to strongly saline. Salinization is rapidly increasing in the midstream areas that are irrigated with water from the Syr Darya River. For example, the percentage of moderately to strongly saline lands in


the midstream area increased from approximately 26 percent in 1970 to 54 percent in 1995 (EC 1995). Salinization is contributing to the diminishing crop yields and is also creating a health problem because of its effect on potable water supplies. Other irrigation-related soil problems include waterlogging, erosion, and compaction (Klotzli 1994).

Water quality reduction

Although the quality of the natural flows in the Syr Darya meets international water quality standards, it has been adversely affected by anthropogenic activities. Agricultural drainage is the major factor affecting water quality in the middle and lower sections of the basin. Records show that just downstream of the Fergana Valley, a major irrigation district in the upper basin, the average salinity of the river water increases to 1.2 grams per liter (g/l) from a concentration of less than 0.5 g/l entering the valley (Raskin et al. 1992). Salinity conditions vary significantly along the river from upstream to downstream. The mineralization is 0.2–0.7 g/l in the upstream area, 0.7–2.3 g/l in the midstream area, and 9.0–10.0 g/l in the downstream area (EC 1995, Vol. II). Figure 8 plots salinity at selected points in the Syr Darya River from 1950 to 1990.

Some stability and even improvement in the water quality has been achieved over the last 10 years from improved water distribution and irrigation and drainage facilities. Whereas downstream areas suffer most from water and soil salinity problems, some upstream areas, such as the Fergana Valley, also suffer substantial groundwater salinization.

Figure 8—Salinity at selected points in the Syr Darya River from 1950 to 1990



Source: EC 1995.

The highest surface water salinity usually occurs at the midstream reach (inflow to the Kayrakum Reservoir), because of the large drainage load from the Fergana Valley and other upstream and midstream demand sites.

3.1.3. Conflicts in Water Sharing

The Syr Darya River basin is a vital resource for its four Central Asian riparian countries. The transboundary character of the river is a source of tension and conflict among the countries as water allocation directly affects agricultural production and hydroelectric generation. An inequitable allocation of water among the four nations could significantly hurt the economic position of one or more of the republics. Before 1990,

unified water management in the Aral Sea region was carried out under the Soviet law system. Since 1990, the new border situation in Central Asia has created clear-cut upstream-downstream conflicts between the independent countries. All the republics asserted their right to control land, water, and other natural resources within their territories, increasing the complexity of the Aral Sea disaster. Although centralized water management authorities (basin water management agencies) were established for the river basins of the Amu Darya and Syr Darya, for the time being, intergovernment agreements remain the major institution for regulating water allocation among the Central Asian republics.

Typical upstream-downstream water conflicts in the Syr Darya River basin include water depletion, timing problems created by water storage for hydropower production, and water quality deterioration in the lower reaches. Kyrgyzstan controls most of the source flow (70 percent of the runoff) as well as most of the reservoirs regulating the river flow. Uzbekistan, the most populous republic in the region, has only minimal hydraulic control along the course of the river. Kyrgyzstan, which depends on hydropower for most of its energy supply, especially in the winter, wants to save summertime runoff in its reservoirs (Toktogul, the largest in the basin with an active capacity of 14 km³) for hydropower generation in the winter. This causes a major conflict with the downstream republics, which depend on irrigation in the summer growing season. Moreover, Kazakhstan, located downstream of Uzbekistan water uses, complains about low levels of both water quantity and quality, including high concentrations of salt and industrial toxic wastes.

3.1.4. Infrastructure Deterioration

Since independence, the Central Asian republics have lost significant management capacity that has yet to be replaced. Much of the capital that had been accumulated by the former Soviet system has been either consumed or dissipated. As a result, maintenance of important infrastructure such as irrigation and drainage systems and roadways has been deferred; and farm equipment and irrigation machinery are not being adequately serviced and replaced in a timely manner. In all five countries of Central Asia, most irrigation systems are in a state of disrepair. The water control and distribution system is deteriorating, and the aging water supply systems are at risk. Mining and industrial wastes are not monitored, and treatment facilities are not designed to remove industrial toxic wastes. Pollution from industrial mines and uranium enrichment facilities are a major concern. As pollution and water quality problems move downstream to the neighboring republics, they have the potential to cause interstate conflict. Furthermore, the five Central Asian governments, each advancing its own national interest within a regional context, have had to assume operations of systems within their own borders; and large areas of formerly irrigated land have been lost because of the breakdown of the water system.

The water supply and utilization infrastructure in the Aral Sea region requires extensive maintenance and new capital investment. Yet the governments have inadequate financial resources to devote to the energy system; and without major reform, they have little ability to recoup the costs of any expenditure they could make (World Bank 1992).

3.1.5. Potential Approaches to the Aral Sea Problem

The economic and environmental problems in the Aral Sea region will likely worsen. The population could increase by as much as 35 percent over the next 30 years

according to the United Nations (UNESCO 1998). That population growth will increase demands on irrigated agriculture and water resources as the republics promote food self-sufficiency. Additional urban development will further challenge the water supply situation. These problems raise the question of whether such a high level of irrigated agriculture can be sustained while preventing or minimizing adverse environmental and ecological impacts. The answer is at the heart of sustainable water resources management for the basin. Glantz (1999) argues, “It is important that the Aral Sea basin be viewed historically as a ‘meta-ecosystem:’ a system that cannot be separated into its many linked parts. Collective problems must be met with collective solutions.” According to Micklin (1993), the huge hydrogeological changes will take decades to reverse. Reducing the amount of water used for irrigation will have to be part of the solution. However, the economic reliance on a few monoculture crops leaves little room for change (World Bank 1992).

The approaches being suggested to solve the Aral Sea problem include both structural and managerial improvements. Considerable space exists to reduce irrigation withdrawals through improvements in the water distribution and conveyance systems and in the irrigation and drainage systems (Cai 1999). Managerial improvements include structural reforms in agriculture, for example, replacing water-consuming crops, such as cotton and rice, with relatively water-saving crops such as wheat and maize (Micklin 1993), and implementing water prices (as is being done in some form in each republic) (Hutchens and Mann 1998).

Within the region, there are currently very few practical applications of an integrated approach to managing natural resources. Demonstrations of actual applications

of environmental management policies are needed to test the effectiveness of the policies and to introduce new technology. An analytical framework that includes the costs and benefits of these approaches is needed to see the effects of these approaches today and in the future, and to look at the likelihood these approaches can make the region's unsustainable water management sustainable.

3.2. POLICY IMPLICATIONS FOR IRRIGATION PLANNING AND MANAGEMENT: RESULTS FROM THE MODELING FRAMEWORK

The modeling framework described in section 2 has been applied to the Syr Darya River basin in Central Asia over a 30-year period with various hydrologic levels such as very wet (*vw*), wet (*w*), normal (*n*), dry (*d*), and very dry (*vd*). The hydrologic levels were projected by Raskin et al. (1992) and are shown in Figure 8 and other figures. Several policy scenarios have been studied using the model. On the basis of the modeling results, policy implications for planning and management of irrigation water in the basin are discussed in the following.


3.2.1. *Maintaining Current Irrigation Practices*

Maintaining current irrigation practices will put the flow to the Aral Sea at tremendous risk. Assuming the irrigated area will start to decline when crop yields drop below half their maximum value, the first scenario maintains the current irrigated area and crop patterns without improvements to the irrigation and drainage infrastructure. This is defined as the baseline scenario (BAS). This scenario shows that 97 percent of the currently irrigated area can be used during the 30-year study period; there are no years in

which a cutback in planted area of more than 15 percent is required; and the largest annual cutback of irrigated area in the time horizon (30 years) is 11 percent.

Compared to the relatively low risk to irrigation under BAS, the risk to ecological water use is tremendous. Ecological water use is measured as the flow into the Northern Aral Sea from the Syr Darya River. To consider the risk indicators for ecological water use, we define the target flow as the average flow to the Northern Aral Sea from the river during 1965–75. That flow satisfies the inflow requirement to the Aral Sea from the Syr Darya River according to the five-country agreement on flow to the Aral Sea (McKinney and Kenshimov 2000). We then define a failure year as one in which the ratio of water released to the sea is less than 85 percent of the target flow (a 15 percent risk threshold). Under this scenario, only 44 percent of the required flow is released to the sea; in 17 consecutive years in the 30-year time horizon, flow to the sea is less than 85 percent of the target; and in the most serious case, only 1 percent of the target flow can be accessed. (See Figure 9 for the target flow and computed flow by years.)

Figure 9—Flow release to the Aral Sea versus hydrologic year series



Degradation of soil and water quality will probably continue if current irrigation practices are maintained. Soil and water quality in the entire basin will be negatively affected. Modeling results show that the middle stream area will endure the most notable degradation of soil and water quality. Figures 10 and 11 plot the average salt concentration in field drainage and soil salinity in cotton fields under BAS and an alternative scenario (to be described later). From the base year to the last year of the modeling period, the average salt concentration in field drainage will increase from 2.0 g/l to 3.5 g/l, and the average soil salinity in a middle stream area cotton field will increase from 0.7 to 1.3 mmhos/cm (expressed as electric conductivity). A larger increase rate is found in later years of the time horizon for both soil and water salinity.

Figure 10—Salt concentration in field drainage (middle stream) versus hydrologic year series





Figure 11—Soil salinity (middle stream, cotton field) versus hydrologic year series



3.2.2. *An Alternative Scenario of Infrastructure Improvement and Crop Pattern Change*

An alternative scenario (ALT) to the baseline scenario described above includes investment in infrastructure improvements and crop pattern change. Water conveyance and distribution efficiency improves gradually from 0.50–0.67 to 0.70–0.75 in the next 25 years, with larger improvements in downstream areas. Irrigation efficiency gradually improves from 0.55–0.65 to 0.65–0.78, with larger increases in the middle stream area, where a high risk of waterlogging exists, and a moderate level in the downstream area, leaving substantial amounts of water necessary for salt leaching in the crop fields. The drainage system is significantly better for middle stream and downstream areas, with 52–60 percent of the irrigated area drained in the base year and 80 percent of the irrigated area drained by the end of study period. The average annual investment for these improvements is estimated as US\$290 million (Cai, McKinney, and Lasdon 2001).

The cotton in the region is characterized by high water consumption and high salt tolerance, and planting cotton potentially raises soil salinity in the region. The alternative scenario proposes a dramatic crop pattern change to reduce the currently dominant cotton-forage area (from 60 to 40 percent of the total irrigated area) and to increase irrigated area for wheat-maize (from 10 to 32 percent of the total irrigated area). Moreover, it assumes that the total irrigated area will fall by 10 percent within the study period.


Compared to BAS, ALT performs much better in terms of the sustainability indicators. First, risk indicators for ecological flow release to the Aral Sea improve. ALT achieves 84 percent of the target release, versus 44 percent under BAS. In only two consecutive years is the flow to the sea less than 85 percent of the target, versus 17 years

under BAS. And the minimum flow release is 50 percent of the flow target, versus 1 percent under BAS (Figure 9).

Second, as shown in Figures 10 and 11, degradation of water and soil salinity under ALT in future years is not so serious as under BAS, although there is still a slight increase in average salt concentration in field drainage and soil salinity. This implies that additional measures, such as drainage disposal, further reductions in irrigated area, and so on, might be necessary.

Third, ATL results in better economic returns for irrigation than BAS. Figure 12 shows the net irrigation profit under the two scenarios over the time horizon.

Figure 12—Irrigation profit versus hydrologic year series



Under BAS, profits from irrigation decline because of water stress and soil quality (salinity), and year-to-year variations are larger; under ALT, irrigation profits are more stable, and the average in later years is slightly higher than earlier years. Therefore, the modeling results imply that maintaining current irrigation practices (BAS) may hurt the

future water use benefit, while the proposed alternative scenario will avoid the negative effect on the future and sustain irrigated agriculture without significant damage to the environment.


3.2.3. Irrigation versus Hydropower Generation

The long-term model considers hydropower generation by including net hydropower profit in the total benefit of water uses. Because the magnitude of hydropower profit is far less than the irrigation benefit in the whole river basin, hydropower generation will have lower priority than irrigation in the modeling if no additional policy constraint is included. However, as mentioned before, in the real world, the upstream country Kyrgyzstan—which depends on hydropower for most of its power supply, especially in winter—attempts to hold more of the water that comes in during the growing season in the Toktogul Reservoir for hydropower generation in the winter. In one policy proposal Kyrgyzstan does not hold water in the growing season, and in return the downstream countries help Kyrgyzstan with power generation, for example, by trading coal to Kyrgyzstan at a cheap price. The two scenarios discussed above follow this policy, and both result in much lower hydropower generation than in the current reality, especially in dry years.

Another policy assumes that Kyrgyzstan holds enough water in the Toktogul Reservoir to meet the demand for hydropower generation in winter months as much as possible. To see the trade-off between downstream irrigation and upstream hydropower generation, we model this policy scenario (HYP) by putting a higher priority on hydropower generation than irrigation so that the target of hydropower generation will always be met first if possible. Figure 13 shows the irrigation profit gains and energy loss

in nonvegetation months (October–March) of ALT compared to HYP, respectively. Larger gains/losses occur in very dry years (*vd*) and consecutive dry years (*d*), while smaller gains/losses occur in very wet (*vw*) and wet years (*w*).

Figure 13—Irrigation profit gains/energy loss under two scenarios




The largest irrigation profit gain is US\$180 million, associated with energy loss of 3,120 million kilowatt hours, which occurs after three consecutive dry years late in the study period. In very wet years, gains/losses are close to zero. In a normal year, gains/losses are affected by the inter-year flow regulation through the storage system in the basin between that year and its previous and following years.

3.2.4. Effects of Economic Incentives

Modeling results show that taxing salt discharges into the river may help solve environmental problems while having only a small effect on irrigation profit. To illustrate the economic incentive of the salt discharge tax, the tax rate was parametrically varied from US\$0 to US\$200 per ton of salt mass in a short-term yearly model. Figure 14 shows

irrigation profit and salt discharge versus the salt tax rate. Up to US\$50 per ton, a small increase in the tax rate reduces salt discharge significantly while only slightly reducing irrigation profit.

Figure 14—Irrigation profit and salt discharge under various salinity tax rates



Another striking result is that investments to improve water distribution systems and irrigation systems could be financed by taxing the increase in profits resulting from the infrastructure improvements. Annualized investment is estimated at US\$299 million. With a tax rate of US\$10 per ton, without considering the transaction cost of implementing the tax system, the tax collected could be as much as US\$251 million, which is 80 percent of the investment.

4. CONCLUSIONS

Sustainable irrigation water management should reach a two-part objective, simultaneously sustaining both irrigated agriculture (required for food security) and the

associated environment. The way to achieve sustainability is to resolve the conflicts arising from the interactions between water use and the environment, and to balance the benefits between current and future generations. To achieve sustainability, decisions at all the various levels—from crop field management to water allocation at the basin scale and agricultural policy at the regional and national scale—must follow the newly developed sustainability principles.

The complexity involved in irrigation water management necessitates a systems approach to water management analysis. This approach translates broad guidelines for sustainable water resources management into operational concepts that can be applied to water management in irrigation-dominated river basins. Appropriate indicators are defined to measure sustainability, taking into account risk minimization in water supply, environmental conservation, equity in water allocation, and economic efficiency in water infrastructure development. A modeling framework combines intra-year (short-term) decisions with inter-year (long-term) decisions to help find sustainable development patterns in irrigation-dominated river basins. A long-term optimization model of a river basin incorporates specific indicators, so that managerial control of the short-term decisions ensures long-term sustainability.

The concepts and analytical framework are applied to irrigation water management in the Aral Sea region, where current practices have led to an environmental disaster that is likely irreversible in some parts. Lessons from the Aral Sea problem include the need for an appropriate agricultural development strategy, the need for developing and monitoring an early warning system, and the need to take appropriate actions when creeping processes of change are detected. Sustainable resource use in the region must take a long-term view.

Adverse water and soil salinity consequences of irrigation must be fully accounted for, and cumulative changes of water and soil salinity over time must be traced. Trade-offs between short-term irrigation profits and their associated environmental consequences must be carefully dealt with.

Analytical results show that the current irrigated agriculture cannot be sustained because it may put at risk the flow release to the Aral Sea, aggravate the degradation of water and soil salinity, and adversely affect crop yield and harvested area. Improvements in the current infrastructure, such as improving water conveyance/distribution efficiency, upgrading irrigation efficiency and drainage systems, reducing irrigated cotton area, and increasing wheat and maize area, are recommended to sustain agricultural production and the environment in the basin. Annualized investment for the improvements is estimated as US\$299 million.

A penalty tax on salt discharge, as an economic incentive, may help address environmental problems while having only a small effect on irrigation profit. Moreover, investments to improve infrastructure could be financed from taxes on increased profits resulting from the infrastructure improvements. However, the taxation and investments will depend on the development of an interstate agreement on basin infrastructure investments, something that the Central Asian republics still need to develop.

REFERENCES

- ASCE (American Society of Civil Engineers) and UN/IHP (United Nations International Hydrologic Program). Committee on Sustainability Criteria, Water Resources Planning, and Management Division. 1998. *Sustainability criteria for water resources systems*. IV Project M-4.3.
- Batchelor, C. 1999. Improving water use efficiency as part of integrated catchment management. *Agricultural Water Management* 40(2): 249–263.
- Briscoe, J. 1999. The Financing of Hydropower, Irrigation and Water Supply Infrastructure in Developing Countries. *Journal of International Water Resources Development*, Vol.15(4), 459-491.
- Cai, X. 1999. Modeling framework for sustainable water resources management. Unpublished Ph.D. dissertation, University of Texas at Austin.
- Cai, X., D. C. McKinney, and L. Lasdon. 2001. Solving nonlinear water management models using a combined genetic algorithm and linear programming approach. *Advances in Water Resources* 24(6): 667–676.
- EC (European Commission). 1995. *Water resources management & agricultural production in the central Asian republics*, WARMAP Project Report, Vol. 1–6.
- FAO (Food and Agriculture Organization of the United Nations). 1996. *Food production: The critical role of water*. World Food Summit, Rome, Italy.
- Fiering, M. B. 1982. Alternative indices of resilience. *Water Resources Research* 18(1): 33–39.
- Glantz, M. H., ed. 1999. *Creeping environmental problems and sustainable development in the Aral Sea basin*. Cambridge, UK, and New York: Cambridge University Press.
- Gleick, P. H. 2000. “The changing water paradigm: a look at twenty-first century water resources development,” *Water International* 25(1): 127-138.
- Grigg, N. S. 1996. *Water resources management*. New York: McGraw-Hill.
- Hashimoto, T., J. R. Stedinger, and D. P. Loucks. 1982. Reliability, resiliency, and vulnerability criteria for water resources system performance evaluation. *Water Resources Research* 18(1): 14–20.

- Hutchens, A. O., and P. C. Mann. 1998. *Review of water pricing policies, institutions and practices in Central Asia*. Technical Report, U.S. Agency for International Development, Environmental Policy and Technology Project, Central Asia Regional EPT Office, Almaty, Kazakstan.
- Ingram, H. 1997. Water and equity: A direction for research. *The Common Property Resources (CPR) Forum* No. 43.
- Klotzli, S. 1994. *The water and soil crisis in Central Asia—A source for future conflict?* ENCOP Occasional Paper No. 11. Center for Security Policy and Conflict Research Zurich/Swiss Peace Foundation, Berne, Zurich.
- Kundzewicz, Z. W., and J. Kindler. 1995. Multiple criteria for evaluation of reliability aspects of water resource systems. *Proceedings of a Boulder Symposium*. International Association of Hydrologic Sciences (IAHS) Pub. No. 231.
- Loucks, D. P. 2000. Sustainable water resources management. *Water International* 25(1): 3–11.
- MAF (Ministry of Agriculture and Forestry). 1997. *Indicators of sustainable irrigated agriculture*. MAF Technical Paper No. 00/03. Wellington, New Zealand.
- Micklin, P. P. 1993. The shrinking Aral Sea. *Geotimes* 38(4): 14–18.
- McKinney, D.C. and A.K. Kenshimov (eds.) 2000. *Optimization of the Use of Water and Energy Resources in the Syrdarya Basin Under Current Conditions, Technical Report 00-06-W*, Environmental Policies and Institutions for Central Asia (EPIC) Program, US Agency for International Development, Almaty, Kazakhstan.
- McKinney, D. C., X. Cai, M. W. Rosegrant, R. Claudia, and C. A. Scott. 1999. *Modeling water resources management at the basin level: Review and future directions*. SWIM Paper No. 6. Colombo, Sri Lanka: International Water Management Institute.
- OECD (Organization for Economic Co-operation and Development). 1998. *Sustainable management of water in agriculture: Issues and policies*. The Athens Workshop, Athens, Greece.
- Raskin, P., E. Hansen, Z. Zhu, and D. Stavisky. 1992. Simulation of water supply and demand in the Aral Sea Region. *Water International* 17: 55–67.
- Rosegrant, M. W., and C. Ringler. 2000. Impact on food security and rural development of transferring water out of agriculture. *Water Policy* 1(6): 567–586.
- Rosegrant, M.W., C. Ringler, D.C. McKinney, X. Cai, A. Keller, and G. Donoso. 2000. Integrated economic-hydrologic water modeling at the basin scale: the Maipo River basin. *Agricultural Economics*, Vol.24, No.1, pp.33-46.

- Rosegrant, M. W., and R. S. Meinzen-Dick. 1996. *Multi-country research program: Water resource allocation: Productivity and environmental impacts (MP-10). Program statement and methodology*. Washington, D.C.: International Food Policy Research Institute.
- Seckler, D. 1996. *The new era of water resources management: From “dry” to “wet” water savings*. Research Report, No 1. Colombo, Sri Lanka: International Water Management Institute (IWMI).
- Sergageldin, I. 1995. Water resources management: A new policy for a sustainable future. *Water Resources Development* 11(3): 221–231.
- Simonovic, S. P. 1997. Risk in sustainable water resources management. In *Sustainability of water resources under increasing uncertainties*. IAHS Pub. No. 240.
- Svendsen, M. 1987. *Sustainability in irrigated agriculture*. International Irrigation Management Institute (IIMI), Working Paper No. 4. Colombo, Sri Lanka: International Irrigation Management Institute.
- Tsur, Y., and A. Dinar. 1995. *Efficiency and equity considerations in pricing and allocating irrigation water*. Policy Research Working Paper 1460. Washington, D.C.: World Bank.
- UNESCO. 1998. UNESCOSTAT Database. Accessible at <http://unescotat.unesco.org>.
- UNCED (United Nations Conference on Environment and Development). 1992. *Report of the United Nations conference on environment and development*, Chap. 5 and 18, Rio de Janeiro.
- WCED (World Commission on Environment and Development). 1987. *Our common future*. (“The Brundtland Report”) London, UK: Oxford University Press.
- World Bank. 1992. *World development report 1992: Development and the environment*. Washington, D.C.: World Bank.

EPTD DISCUSSION PAPERS

LIST OF EPTD DISCUSSION PAPERS

- 01 *Sustainable Agricultural Development Strategies in Fragile Lands*, by Sara J. Scherr and Peter B.R. Hazell, June 1994.
 - 02 *Confronting the Environmental Consequences of the Green Revolution in Asia*, by Prabhu L. Pingali and Mark W. Rosegrant, August 1994.
 - 03 *Infrastructure and Technology Constraints to Agricultural Development in the Humid and Subhumid Tropics of Africa*, by Dunstan S.C. Spencer, August 1994.
 - 04 *Water Markets in Pakistan: Participation and Productivity*, by Ruth Meinzen-Dick and Martha Sullins, September 1994.
 - 05 *The Impact of Technical Change in Agriculture on Human Fertility: District-level Evidence From India*, by Stephen A. Vosti, Julie Witcover, and Michael Lipton, October 1994.
 - 06 *Reforming Water Allocation Policy Through Markets in Tradable Water Rights: Lessons from Chile, Mexico, and California*, by Mark W. Rosegrant and Renato Gazri S, October 1994.
 - 07 *Total Factor Productivity and Sources of Long-Term Growth in Indian Agriculture*, by Mark W. Rosegrant and Robert E. Evenson, April 1995.
 - 08 *Farm-Nonfarm Growth Linkages in Zambia*, by Peter B.R. Hazell and Behjat Hoijsati, April 1995.
 - 09 *Livestock and Deforestation in Central America in the 1980s and 1990s: A Policy Perspective*, by David Kaimowitz (Interamerican Institute for Cooperation on Agriculture), June 1995.
 - 10 *Effects of the Structural Adjustment Program on Agricultural Production and Resource Use in Egypt*, by Peter B.R. Hazell, Nicostrato Perez, Gamal Siam, and Ibrahim Soliman, August 1995.
 - 11 *Local Organizations for Natural Resource Management: Lessons from Theoretical and Empirical Literature*, by Lise Nordvig Rasmussen and Ruth Meinzen-Dick, August 1995.
-

- 12 *Quality-Equivalent and Cost-Adjusted Measurement of International Competitiveness in Japanese Rice Markets*, by Shoichi Ito, Mark W. Rosegrant, and Mercedita C. Agcaoili-Sombilla, August 1995.
 - 13 *Role of Inputs, Institutions, and Technical Innovations in Stimulating Growth in Chinese Agriculture*, by Shenggen Fan and Philip G. Pardey, September 1995.
 - 14 *Investments in African Agricultural Research*, by Philip G. Pardey, Johannes Roseboom, and Nienke Beintema, October 1995.
 - 15 *Role of Terms of Trade in Indian Agricultural Growth: A National and State Level Analysis*, by Peter B.R. Hazell, V.N. Misra, and Behjat Hoijati, December 1995.
 - 16 *Policies and Markets for Non-Timber Tree Products*, by Peter A. Dewees and Sara J. Scherr, March 1996.
 - 17 *Determinants of Farmers' Indigenous Soil and Water Conservation Investments in India's Semi-Arid Tropics*, by John Pender and John Kerr, August 1996.
 - 18 *Summary of a Productive Partnership: The Benefits from U.S. Participation in the CGIAR*, by Philip G. Pardey, Julian M. Alston, Jason E. Christian, and Shenggen Fan, October 1996.
 - 19 *Crop Genetic Resource Policy: Towards a Research Agenda*, by Brian D. Wright, October 1996.
 - 20 *Sustainable Development of Rainfed Agriculture in India*, by John M. Kerr, November 1996.
 - 21 *Impact of Market and Population Pressure on Production, Incomes and Natural Resources in the Dryland Savannas of West Africa: Bioeconomic Modeling at the Village Level*, by Bruno Barbier, November 1996.
 - 22 *Why Do Projections on China's Future Food Supply and Demand Differ?* by Shenggen Fan and Mercedita Agcaoili-Sombilla, March 1997.
 - 23 *Agroecological Aspects of Evaluating Agricultural R&D*, by Stanley Wood and Philip G. Pardey, March 1997.
-

- 24 *Population Pressure, Land Tenure, and Tree Resource Management in Uganda*, by Frank Place and Keijiro Otsuka, March 1997.
 - 25 *Should India Invest More in Less-favored Areas?* by Shenggen Fan and Peter Hazell, April 1997.
 - 26 *Population Pressure and the Microeconomy of Land Management in Hills and Mountains of Developing Countries*, by Scott R. Templeton and Sara J. Scherr, April 1997.
 - 27 *Population Land Tenure and Natural Resource Management: The Case of Customary Land Area in Malawi*, by Frank Place and Keijiro Otsuka, April 1997.
 - 28 *Water Resources Development in Africa: A Review and Synthesis of Issues, Potentials, and Strategies for the Future*, by Mark W. Rosegrant and Nicostrato D. Perez, September 1997.
 - 29 *Financing Agricultural R&D in Rich Countries: What's Happening and Why?* by Julian M. Alston, Philip G. Pardey, and Vincent H. Smith, September 1997.
 - 30 *How Fast Have China's Agricultural Production and Productivity Really Been Growing?* by Shenggen Fan, September 1997.
 - 31 *Does Land Tenure Insecurity Discourage Tree Planting? Evolution of Customary Land Tenure and Agroforestry management in Sumatra*, by Keijiro Otsuka, S. Suyanto, and Thomas P. Tomich, December 1997.
 - 32 *Natural Resource Management in the Hillsides of Honduras: Bioeconomic Modeling at the Micro-Watershed Level*, by Bruno Barbier and Gilles Bergeron, January 1998.
 - 33 *Government Spending, Growth, and Poverty: An Analysis of Interlinkages in Rural India*, by Shenggen Fan, Peter Hazell, and Sukhadeo Thorat, March 1998. Revised December 1998.
 - 34 *Coalitions and the Organization of Multiple-Stakeholder Action: A Case Study of Agricultural Research and Extension in Rajasthan, India*, by Ruth Alsop, April 1998.
-

- 35 *Dynamics in the Creation and Depreciation of Knowledge and the Returns to Research*, by Julian Alston, Barbara Craig, and Philip Pardey, July, 1998.

 - 36 *Educating Agricultural Researchers: A Review of the Role of African Universities*, by Nienke M. Beintema, Philip G. Pardey, and Johannes Roseboom, August 1998.

 - 37 *The Changing Organizational Basis of African Agricultural Research*, by Johannes Roseboom, Philip G. Pardey, and Nienke M. Beintema, November 1998.

 - 38 *Research Returns Redux: A Meta-Analysis of the Returns to Agricultural R&D*, by Julian M. Alston, Michele C. Marra, Philip G. Pardey, and T.J. Wyatt, November 1998.

 - 39 *Technological Change, Technical and Allocative Efficiency in Chinese Agriculture: The Case of Rice Production in Jiangsu*, by Shenggen Fan, January 1999.

 - 40 *The Substance of Interaction: Design and Policy Implications of NGO-Government Projects in India*, by Ruth Alsop with Ved Arya, January 1999.

 - 41 *Strategies for Sustainable Agricultural Development in the East African Highlands*, by John Pender, Frank Place, and Simeon Ehui, April 1999.

 - 42 *Cost Aspects of African Agricultural Research*, by Philip G. Pardey, Johannes Roseboom, Nienke M. Beintema, and Connie Chan-Kang, April 1999.

 - 43 *Are Returns to Public Investment Lower in Less-favored Rural Areas? An Empirical Analysis of India*, by Shenggen Fan and Peter Hazell, May 1999.

 - 44 *Spatial Aspects of the Design and Targeting of Agricultural Development Strategies*, by Stanley Wood, Kate Sebastian, Freddy Nachtergaele, Daniel Nielsen, and Aiguo Dai, May 1999.

 - 45 *Pathways of Development in the Hillsides of Honduras: Causes and Implications for Agricultural Production, Poverty, and Sustainable Resource Use*, by John Pender, Sara J. Scherr, and Guadalupe Durón, May 1999.

 - 46 *Determinants of Land Use Change: Evidence from a Community Study in Honduras*, by Gilles Bergeron and John Pender, July 1999.
-

- 47 *Impact on Food Security and Rural Development of Reallocating Water from Agriculture*, by Mark W. Rosegrant and Claudia Ringler, August 1999.

 - 48 *Rural Population Growth, Agricultural Change and Natural Resource Management in Developing Countries: A Review of Hypotheses and Some Evidence from Honduras*, by John Pender, August 1999.

 - 49 *Organizational Development and Natural Resource Management: Evidence from Central Honduras*, by John Pender and Sara J. Scherr, November 1999.

 - 50 *Estimating Crop-Specific Production Technologies in Chinese Agriculture: A Generalized Maximum Entropy Approach*, by Xiaobo Zhang and Shenggen Fan, September 1999.

 - 51 *Dynamic Implications of Patenting for Crop Genetic Resources*, by Bonwoo Koo and Brian D. Wright, October 1999.

 - 52 *Costing the Ex Situ Conservation of Genetic Resources: Maize and Wheat at CIMMYT*, by Philip G. Pardey, Bonwoo Koo, Brian D. Wright, M. Eric van Dusen, Bent Skovmand, and Suketoshi Taba, October 1999.

 - 53 *Past and Future Sources of Growth for China*, by Shenggen Fan, Xiaobo Zhang, and Sherman Robinson, October 1999.

 - 54 *The Timing of Evaluation of Genebank Accessions and the Effects of Biotechnology*, by Bonwoo Koo and Brian D. Wright, October 1999.

 - 55 *New Approaches to Crop Yield Insurance in Developing Countries*, by Jerry Skees, Peter Hazell, and Mario Miranda, November 1999.

 - 56 *Impact of Agricultural Research on Poverty Alleviation: Conceptual Framework with Illustrations from the Literature*, by John Kerr and Shashi Kolavalli, December 1999.

 - 57 *Could Futures Markets Help Growers Better Manage Coffee Price Risks in Costa Rica?* by Peter Hazell, January 2000.

 - 58 *Industrialization, Urbanization, and Land Use in China*, by Xiaobo Zhang, Tim Mount, and Richard Boisvert, January 2000.
-

- 59 *Water Rights and Multiple Water Uses: Framework and Application to Kirindi Oya Irrigation System, Sri Lanka*, by Ruth Meinzen-Dick and Margaretha Bakker, March 2000.
- 60 *Community natural Resource Management: The Case of Woodlots in Northern Ethiopia*, by Berhanu Gebremedhin, John Pender and Girmay Tesfaye, April 2000.
- 61 *What Affects Organization and Collective Action for Managing Resources? Evidence from Canal Irrigation Systems in India*, by Ruth Meinzen-Dick, K.V. Raju, and Ashok Gulati, June 2000.
- 62 *The Effects of the U.S. Plant Variety Protection Act on Wheat Genetic Improvement*, by Julian M. Alston and Raymond J. Venner, May 2000.
- 63 *Integrated Economic-Hydrologic Water Modeling at the Basin Scale: The Maipo River Basin*, by M. W. Rosegrant, C. Ringler, D.C. McKinney, X. Cai, A. Keller, and G. Donoso, May 2000.
- 64 *Irrigation and Water Resources in Latin America and the Caribbean: Challenges and Strategies*, by Claudia Ringler, Mark W. Rosegrant, and Michael S. Paisner, June 2000.
- 65 *The Role of Trees for Sustainable Management of Less-favored Lands: The Case of Eucalyptus in Ethiopia*, by Pamela Jagger & John Pender, June 2000.
- 66 *Growth and Poverty in Rural China: The Role of Public Investments*, by Shenggen Fan, Linxiu Zhang, and Xiaobo Zhang, June 2000.
- 67 *Small-Scale Farms in the Western Brazilian Amazon: Can They Benefit from Carbon Trade?* by Chantal Carpentier, Steve Vosti, and Julie Witcover, September 2000.
- 68 *An Evaluation of Dryland Watershed Development Projects in India*, by John Kerr, Ganesh Pangare, Vasudha Lokur Pangare, and P.J. George, October 2000.
- 69 *Consumption Effects of Genetic Modification: What If Consumers Are Right?* by Konstantinos Giannakas and Murray Fulton, November 2000.
-

- 70 *South-North Trade, Intellectual Property Jurisdictions, and Freedom to Operate in Agricultural Research on Staple Crops*, by Eran Binenbaum, Carol Nottenburg, Philip G. Pardey, Brian D. Wright, and Patricia Zambrano, December 2000.
- 71 *Public Investment and Regional Inequality in Rural China*, by Xiaobo Zhang and Shenggen Fan, December 2000.
- 72 *Does Efficient Water Management Matter? Physical and Economic Efficiency of Water Use in the River Basin*, by Ximing Cai, Claudia Ringler, and Mark W. Rosegrant, March 2001.
- 73 *Monitoring Systems for Managing Natural Resources: Economics, Indicators and Environmental Externalities in a Costa Rican Watershed*, by Peter Hazell, Ujjayant Chakravorty, John Dixon, and Rafael Celis, March 2001.
- 74 *Does Quaxi Matter to NonFarm Employment?* by Xiaobo Zhang and Guo Li, June 2001.
- 75 *The Effect of Environmental Variability on Livestock and Land-Use Management: The Borana Plateau, Southern Ethiopia*, by Nancy McCarthy, Abdul Kamara, and Michael Kirk, June 2001.
- 76 *Market Imperfections and Land Productivity in the Ethiopian Highlands*, Stein Holden, Bekele Shiferaw, and John Pender, August 2001.
- 77 *Strategies for Sustainable Agricultural Development in the Ethiopian Highlands*, by John Pender, Berhanu Gebremedhin, Samuel Benin, and Simeon Ehui, August 2001.
- 78 *Managing Droughts in the Low-Rainfall Areas of the Middle East and North Africa: Policy Issues*, by Peter Hazell, Peter Oram, Nabil Chaherli, September 2001.
- 79 *Accessing Other People's Technology: Do Non-Profit Agencies Need It? How To Obtain It*, by Carol Nottenburg, Philip G. Pardey, and Brian D. Wright, September 2001.
- 80 *The Economics of Intellectual Property Rights Under Imperfect Enforcement: Developing Countries, Biotechnology, and the TRIPS Agreement*, by Konstantinos Giannakas, September 2001.
-

- 81 *Land Lease Markets and Agricultural Efficiency: Theory and Evidence from Ethiopia*, by John Pender and Marcel Fafchamps, October 2001.

 - 82 *The Demand for Crop Genetic Resources: International Use of the U.S. National Plant Germplasm System*, by M. Smale, K. Day-Rubenstein, A. Zohrabian, and T. Hodgkin, October 2001.

 - 83 *How Agricultural Research Affects Urban Poverty in Developing Countries: The Case of China*, by Shenggen Fan, Cheng Fang, and Xiaobo Zhang, October 2001.

 - 84 *How Productive is Infrastructure? New Approach and Evidence From Rural India*, by Xiaobo Zhang and Shenggen Fan, October 2001.

 - 85 *Development Pathways and Land Management in Uganda: Causes and Implications*, by John Pender, Pamela Jagger, Ephraim Nkonya, and Dick Sserunkuuma, December 2001.
-