



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

The Use of Food Nutrition Facts Panel Information and Juice Consumption

Hyeyoung Kim, University of Florida, kim1978@ufl.edu

Carlos E. Jáuregui, Florida Department of Citrus, cej@ufl.edu

Jonq-Ying Lee, University of Florida, jonqying@ufl.edu

Selected Paper prepared for presentation at the Agricultural & Applied Economics Association's 2014 AAEA Annual Meeting, Minneapolis, MN, July 27-29, 2014.

Copyright 2014 by Hyeyoung Kim, Carlos E. Jáuregui, and Jonq-Ying Lee. All rights reserved. Readers may make verbatim copies of this document for non-commercial purposes by any means, provided that this copyright notice appears on all such copies.

The Use of Food Nutrition Facts Panel Information and Juice Consumption

Abstract

The Nutrition Labeling and Education Act of 1990 requires that the FDA develop standardized Nutrition Facts Panel (NFP) labels for most packaged food products sold in the United States. The National Health and Nutrition Examination Survey (NHANES) reported more people were using the NFP. Per capita consumption of juices in the United States has decreased steadily, from 8.88 gallons in 2000 to 7.21 gallons in 2010. The purpose of this study is to examine the relationship between the use of the NFP and the consumption of juices in the US using cross-section data provided by the NHANES. We used data from three waves of NHANES: 2005–2006, 2007–08, and 2009–10. Because not all respondents consumed juice during the survey periods, a Tobit model was used in the analysis. Results show that using the NFP label has a negative impact on juice consumption and that there is a negative time trend in juice consumption. These results indicate that the juice industry needs to find a way to combat the negative impact of the NFP label on juice consumption, and to find out what is causing the negative time trend in juice consumption.

Introduction

Poor diet is a significant cause of obesity, heart disease, stroke, cancer, diabetes, osteoarthritis, and other health conditions that impose an economic burden on individuals and society overall (USDA/USDHHS 2011; USDHHS 2010). The prevalence of childhood and adult obesity in the United States more than doubled between the 1976–1980 and 2007–2008 periods (Flegal et al. 2010; Ogden et al. 2010). The high rates of obesity and diet-related health problems have prompted consumers to watch their diets. In 2011, 51% of all adults aged 18 and over reported watching their diet (Mintel Oxygen 2012). Medical costs associated with obesity were estimated as high as \$147 billion, or 10% of all medical costs, in 2008 (Finkelstein et al. 2009; O’Grady and Capretta 2012; Tsai et al. 2011). Growing recognition of the obesity epidemic and the prevalence of diet-related chronic diseases have led to an array of efforts aimed at increasing physical activity and promoting nutritious eating, including changes in the formulation, packaging, labeling, and marketing of food products and beverages (IOM 2010).

While 100% fruit juice has no added sugar, it does contain natural sugar. That is one of the reasons why consumers who try to decrease their sugar intake are drinking less fruit juice. As shown in Table 1, per capita consumption of juices in the United States increased from 5.53 gallons in 1970 to 8.9 gallons in 2000, an increase of 61%, before decreasing steadily to 6.76 gallons in 2011. The decreases of per capita consumptions of orange juice, grapefruit juice, and pineapple juice greatly contributed to the decline in total juice consumption. Buffard (2013) pointed out sugar content as the major culprit for the decrease in per capita consumptions of orange and grapefruit juices. According to the 2012 Internet survey conducted by Mintel Oxygen, one in seven non-consumers of fruit juices and fruit drinks claimed the high caloric content of these products as the reason for their non-consumption of these products. Even though current scientific evidence does not support a relationship between being overweight and fruit juice consumption (O'Neil and Nicklas 2008; Pereira and Fulgoni 2010), there is evidence suggesting that drinking juice is beneficial to one's health (Nicklas et al. 2008, 2010). A recent study by Kim and House (2014) found that beverage consumption was positively correlated to consumer health perception about beverages. This result implies that the publicity about sugar-content warnings may lead to negative perceptions about fruit juice. The downward trend in consumption is a major concern for the citrus juice industry.

The Nutrition Labeling and Education Act of 1990 provides the Food and Drug Administration (FDA) with the authority to require that the nutritional content and health claims on food product nutrition fact panel (NFP) labels are consistent with FDA regulations. The purpose of the NFP label is to provide consumers point-of-purchase nutritional information about products to help them make healthy dietary choices. In addition, the federal government has developed national nutrition guidelines and tools to help consumers obtain useful information about nutrition and

healthy eating from a variety of channels and sources. Despite the text-type information, the percentage of consumers 16 years old and older who do not use or rarely use the NFP information decreased from 43.7% in 2005 to 27.2% in 2007, and decreased again to 22.5% in 2009 (CDC 2013a).

Past studies found that consumers who reported using NFP label information consumed less total food energy and dietary fats than those who reported not using the NFP label information. For example, Ollberding et al. (2011) found that NFP label users had less food-energy and fat intakes than non-users. Temple et al. (2011) studied the food-energy intake of adults consuming lunch in a laboratory setting; they found that the group provided with nutrition labeling information consumed less food energy from low-energy-density and high-energy-density food sources than did the group that received no nutrition labeling information. Other studies also found that NFP information leads consumers to healthier diets (Guthrie et al. 1995; Variyam 2008) and helps improve the overall quality of consumers' diets (Kim et al. 2000, 2001).

The relationship between NFP use and juice consumption has both health implications for consumers and economic implications for the fruit juice industry. The purpose of this study is to examine the relationship between the use of the NFP and the consumption of juices in the United States using cross-section data provided by the National Health and Nutrition Examination Survey (NHANES).

Data

Data from three waves of the NHANES were used: 2005–2006, 2007–08, and 2009–10.

NHANES is a national survey designed to assess the health and nutritional status of adults and children in the United States (CDC 2013b). These surveys collected dietary intake data in two

24-hour recall periods on consumer demographics, self-perceptions of health and nutrition status, and health-related behaviors. We focused on female participants who were 20 years old or older and not pregnant or lactating, who reported trying to lose weight or resisted gaining weight in the past year, and who provided reliable dietary intake information. Among the 31,034 respondents in the survey, 15,095 met our inclusion criteria. Of these 15,095 respondents, only 5,077 (33.6%) had consumed juice in the last year (66.4% of the respondents did not consume any juice).

Covariate Measures

In analyzing the relationship between the usage of food NFP label information and juice consumption, we controlled for consumer demographics, lifestyle, and other factors that may be associated with juice consumption. These factors include income, gender, marital status, race/ethnicity, education (Darmon and Drewnowski 2008; Kim et al. 2000); breakfast consumption (Ruxton and Kirk 1997); dietary supplement consumption (Guenther et al. 2004; Lyle et al. 1998); participation in food assistance programs, such as the Supplemental Nutrition Assistance Program (SNAP; aka food stamp) or the Special Supplemental Nutrition Program for Women, Infants, and Children (aka WIC) (LeBlanc et al. 2006; Wilde et al. 1999); lifestyle behaviors such as alcohol consumption (Breslow et al. 2006; Schuit et al. 2002); and self-rated health status (Watters and Satia 2009; Kim et al. 2012). Table 2 shows the definitions and codes of these variables.

Model and Estimation Procedure

In this study, we assume that the juice consumed by an individual is a function of the above mentioned variables, including how often the individual uses the NFP information. Because not

all respondents reported juice consumption, a Tobit regression was used to estimate (equation 1) consumption. Formally, the relationship can be written as

$$y_i^* = \beta' x_i + \varepsilon_i, \varepsilon_i \sim N(0, \sigma^2)$$

$$y_i = \begin{cases} y_i^* & \text{if } y_i^* > 0 \\ 0 & \text{if } y_i^* \leq 0 \end{cases} \quad (1)$$

where y_i is the quantity of juices consumed by respondent i ; x_i is a vector of determinants of juice consumption as discussed above, including a variable associated with how often the respondent used nutrition label information; β is a vector of the parameters to be estimated; and ε_i is the error term followed using a normal distribution with a zero mean and variance σ^2 . We used STATA (version 12.1; StataCorp, College Station, TX, USA) to generate the descriptive statistics and to perform the regression analyses. Note that since only 35% of the respondents consumed any juice, the Tobit model was estimated using the two-year sample weight. Note that the probability of being uncensored (equation 2) and the impact of changes in the explanatory variable on uncensored juice consumption (equation 3) can be estimated using the following formulas, respectively (Maddala 1983)

$$\frac{\partial \Pr(y_i > 0)}{\partial x_j} = \phi(z) \beta_j \quad (2)$$

$$\frac{\partial E(y_i | y_i^* > 0)}{\partial x_j} = \beta_j \left[1 - z \frac{\phi(z)}{\Phi(z)} - \left(\frac{\phi(z)}{\Phi(z)} \right)^2 \right] \quad (3)$$

where $z = \frac{\beta' x}{\sigma}$, $\phi(z)$ and $\Phi(z)$ are the probability density function and cumulative density

function of z , respectively.

Results

Sample statistics are reported in Table 2. As Table 2 illustrates, 13.8% of the respondents always used the NFP information when shopping. Of the four age groups, the reference age group (those younger than 34 years old) account for about 33% of the sample; and the other three age groups account for about 20% each. The average household size is 3.26 persons and the average annual household income level is \$48,223. Female respondents account for about half of the sample and married respondents account for 47.4% of the sample. Of the respondents, 47.7% are non-Hispanic White, 21.2% are Black, and 27.1% are Hispanic. About 42.5% of the respondents have a college education. Of the respondents, 18.3% are SNAP participants and 9.4% are WIC participants. Most respondents ate breakfast on the two interview days, with 41.8% of the respondents trying to lose weight and 10% of the respondents having tried to resist weight gain during the past year. About 30% of the respondents rated their diet as excellent or very good, and 64.6% rated their diet as good or fair. The reported average juice consumption rate was about six ounces for all participants and close to 16 ounces for juice drinkers.

Regression results are presented in Table 3; the first two columns are the Tobit estimates, which are followed by the probability of being uncensored observations (equation 2), and the marginal impacts on the uncensored observation (equation 3) are listed in the last two columns. As illustrated in Table 3, all estimates are statistically different from zero except the coefficients for Other Race, the dummy for 2007-08, and Losewt. The marginal impact estimates are smaller than the Tobit estimates as expected. As also illustrated in Table 3, the use of the NFP had a negative impact on juice consumption. Results show that if a juice consumer always used the NFP information, his/her juice consumption probability would decrease by 3% and his/her actual

juice consumption would decrease by 15.7 grams, or 3.4% of the average juice consumption based on uncensored observations.

The trend of juice consumption by age, race, and gender is consistent with the findings by Storey et al. (2006). Comparing the age groups, respondents between 35 and 49 years old have the lowest juice consumption, while respondents aged 60 years or older have the highest juice consumption. Black and Hispanic respondents consume more juice than do White respondents. Females consume less juice than do males. Household size is positively related to juice consumption, while household income and being married are negatively related to juice consumption. College education, SNAP and WIC participation, eating breakfast on a regular basis, and taking food supplements are positively related to juice consumption. Kim et al. (2012) also found that 100% fruit juice consumption has a positive relationship with education and a negative relationship with household income. A respondent's BMI and alcoholic beverage consumption are negatively related to juice consumption, and a respondent's diet self-evaluation is positively related to juice consumption. In addition, the number of American adults eating breakfast has declined. According to data from the NHNES, 82% of American adults ate breakfast in 2002, down from 89% in 1971, which may help explain declining fruit juice consumption (IFICF 2008).

The dummies for the second and third waves of NHANES are negative, an indication of a negative trend in juice consumption. It should be noted, however, that the coefficient estimate for the second wave of NHANES is insignificant.

Respondents trying to lose weight consumed less juice than those who did not try to lose weight; however, the coefficient estimate was insignificant. Kim et al. (2012) investigated beverage

consumption habits based on concerns about sugar content and calories. They found that consumers who were concerned about calories drank less 100% fruit juice than did consumers who were concerned about sugar content. This helps explain why consumers concerned about weight control their calorie intake while consumers concerned about health control their nutrition intake.

Among the explanatory variables used in the analysis, race, gender, and eating breakfast, are the most important factors in determining the juice consumption level as measured by the probability of consuming juices and the marginal impact on juice consumption when the respondent had already consumed juices.

Discussion

One hundred percent juices contain no added sugars and current scientific evidence does not support a relationship between being overweight and juice consumption (O'Neil and Nicklas 2008; Pereira and Fulgoni 2011). Furthermore, scientific evidence strongly maintains the nutritional benefits of 100% juices. On the one hand, some studies show that drinking 100% fruit juices is associated with a more nutritious diet overall, including reduced intake of dietary fat, saturated fats, and added sugars (Nicklas et al. 2008, 2010). On the other hand, some studies dispute the nutritional value of juices because they maintain that the sugars in juices are not much different from added sugars (Hamilton 2009; Braun 2014).

The results show that the estimated impact of NFP on juice consumption is relatively small (i.e., 3.6% of average juice consumption). The NFP information is only one of the many sources that consumers use to obtain nutrition information for food purchases. Other nutrition information

sources such as magazine articles and other research results, also have important impact on juice consumption and should be included in research analyses.

Results show that in addition to the NFP label's negative impact on juice consumption, there is also a negative time trend in juice consumption. These results indicate that the juice industry has to find a way to combat the negative NFP label impact on juice consumption, and to find out what is causing the negative time trend in juice consumption.

Identified as the top two “food felons” (Oz and Roizen 2013), refined sugar and sugar syrups damage the human body because they disrupt metabolism and cause cardiovascular and joint inflammation. One of the proposed changes in the NFP labels by the FDA is to add information about added sugar on food product packages. This change may help consumers to differentiate the sucrose, fructose and added refined sugars in juice to reduce the misconception that all sugars are bad and to increase juice consumption.

In an era when consumers are increasingly looking for nutritional information to improve their diet, this study contributes to the literature by exploring the effects of NFP labels on juice consumption. This is an important yet neglected issue in the research. Overall, the results of this study should generate interesting and fruitful discussion about the impact of NFP labels on juice consumption.

References

- Braun, A. 2014. Misunderstanding orange juice as a health drink, *The Atlantic*, February 6
<http://www.theatlantic.com/health/archive/2014/02/misunderstanding-orange-juice-as-a-health-drink/283579> (accessed May 9, 2014).
- Breslow R.A., P.M. Guenther, and B.A. Smothers. 2006. Alcohol drinking patterns and diet quality: The 1999-2000 National Health and Nutrition Examination Survey. *American Journal of Epidemiology* 163:359–366.
- Buffard, K. 2013. New ad could send OJ soaring, *The Ledger*, September 11/12.
<http://www.theledger.com/article/20130911/news/130919794> (accessed May 9, 2014).
- CDC. 2013a. *National Health and Nutrition Examination Survey Data*. National Center for Health Statistics, Centers for Disease Control and, Prevention, United States Department of Health and Human Services, Hyattsville, MD.
http://wwwn.cdc.gov/nchs/nhanes/search/nhanes09_10.aspx (accessed May 9, 2014).
- CDC. 2013b. *NHANES 2005–2006*. National Center for Health Statistics, Centers for Disease Control and, Prevention, United States Department of Health and Human Services, Hyattsville, MD. http://www.cdc.gov/nchs/nhanes/nhanes_questionnaires.htm (accessed May 10, 2014).
- Darmon, N., and A. Drewnowski. 2008. Does social class predict diet quality? *American Journal of Clinical Nutrition* 87:1107–1117.
- Finkelstein, E., J. Trogon, J. Cohen, and W. Dietz. 2009. Annual medical spending attributable to obesity: Payer-and service-specific estimates. *Health Affairs* 28(5):822–831.

- Flegal, K.M., M.D. Carroll, C.L. Ogden, and L.R. Curtin. 2010. Prevalence and trends in obesity among US adults, 1999–2008. *Journal of American Medical Association* 303:242–249.
- Guenther S, R.E. Patterson, A.R. Kristal, K.L. Stratton, and E. White. 2004. Demographic and health-related correlates of herbal and specialty supplement use. *Journal of the American Dietetic Association* 104:27–34.
- Guthrie, J.F., J.J. Fox, L.E. Cleveland, and S. Welch, 1995. Who uses nutrition labeling and what effects does label use have on diet quality? *Journal of Nutrition Education* 27:153–172.
- Hamilton, A. 2009. *Squeezed: What You Don't Know About Orange Juice*. New Haven, CT: Yale University Press.
- IFICF. 2008. *IFIC Review: Breakfast and Health*. Washington, DC: International Food Information Council Foundation.
<http://www.foodinsight.org/Content/76/IFIC%20Brkfast%20Review%20FINAL.pdf>
(accessed May 9, 2014).
- IOM. 2010. *Examination of Front-of-Package Nutrition Rating. Systems and Symbols: Phase I Report of the Institute of Medicine*. Washington, D.C.: The National Academies Press.
- Kim, H. and L.A. House. 2014. Linking consumer health perceptions to consumption of nonalcoholic beverages. *Agricultural and Resource Economics Review* 43(1):1–16.
- Kim, H., L.A. House, G. Rampersaud, and Z. Gao. 2012. Front-of-package nutritional labels and consumer beverage perceptions. *Applied Economic Perspectives and Policy* 34(4):599–614.

- Kim, S-Y, R.M. Nayga Jr, and O. Capps Jr. 2000. The effect of food label use on nutrient intakes: an endogenous switching regression analysis. *Journal of Agricultural and Resource Economics* 25:215–231.
- Kim, S-Y, R.M. Nayga Jr, and O. Capps Jr. 2001. Food label use, self-selectivity, and diet quality. *The Journal of Consumer Affairs* 35(2):346–363.
- LeBlanc, M, B-H Lin, and D. Smallwood. 2006. Food assistance: How strong is the safety net. *Amberwaves* (September).
<http://webarchives.cdlib.org/sw1tx36512/http://www.ers.usda.gov/amberwaves/september06/features/foodassistance.htm> (accessed May 10, 2014).
- Lyle, B.J., J.A. Mares-Perlman, B.E. Klein, R. Klein, and J.L. Greger. 1998. Supplement users differ from nonusers in demographic, lifestyle, dietary and health characteristics. *Journal of Nutrition* 128(12):2355–2362.
- Maddala, G.S. 1983. *Limited-dependent and Qualitative Variables in Econometrics*. Cambridge, UK: Cambridge University Press.
- Mintel Oxygen Website. 2012. Fruit juice and juice drinks—United States, January.
<http://oxygen.mintel.com/display/609344/> (accessed January 10, 2014).
- Nicklas, T.A., C.E. O’Neil, and R. Kleinman. 2008. Association between 100% juice consumption and nutrient intake and weight of children aged 2 to 11 years. *Archives of Pediatrics and Adolescent Medicine* 162(6):557–565.

- Nicklas, T.A., C.E. O’Neil, and R. Kleinman. 2010. The relationship among 100% juice consumption, nutrient intake, and weight of adolescents 12 to 18 years old. *American Journal of Health Promotion* 24(4):231–237.
- Ogden, C.L., M.D. Carroll, L.R. Curtin, M.M. Lamb, and K.M. Flegal. 2010. Prevalence of high body mass index in US children and adolescents 2007–2008. *Journal of American Medical Association* 303:242–249.
- O’Grady, M., and J. Capretta. 2012. Assessing the economics of obesity and obesity interventions. Robert Wood Johnson Foundation, Princeton, N.J.
<http://www.rwjf.org/en/research-publications/find-rwjf-research/2012/03/assessing-the-economics-of-obesity-and-obesity-interventions.html> (accessed May 10, 2014).
- Ollberding, N.J., R.L. Wolf, and I. Contento. 2011. Food label use and its relation to dietary intake among US adults. *Journal of the American Dietetic Association* 111(5):S47–51.
- O’Neil, C.E., and T.A. Nicklas. 2008. A review of the relationship between 100% fruit juice consumption and weight in children and adolescents. *American Journal of Lifestyle Medicine* 2(4):315–354.
- Oz, M., and M.F. Roizen, 2013. The top five food felons. <http://www.success.com/article/drs-oz-roizen-making-you-turns> (accessed May 9, 2014).
- Pereira, M., and V. Fulgoni. 2010. Consumption of 100% fruit juice and risk of obesity and metabolic syndrome: Findings from the National Health and Nutrition Examination Survey 1999–2004. *Journal of American College of Nutrition* 29(6):625–629.

- Ruxton, C.H.S., and T.R. Kirk. 1997. Breakfast: A review of associations with measures of dietary intake, physiology, and biochemistry. *The British Journal of Nutrition* 78(2):199–213.
- Schuit, A.J., A.J.M. van Loon, M. Tijhuis, and M.C. Ocké. 2002. Clustering of lifestyle risk factors in a general adult population. *Preventive Medicine* 35:219–224.
- Storey, M.L., R.A. Forshee, and P.A. Anderson. 2006. Beverage consumption in the US population. *Journal of the American Dietetic Association* 106(12):1992–2000.
- Temple, J.L., K. Johnson, K. Recupero, and H. Suders. 2011. Nutrition labels decrease energy intake in adults consuming lunch in the laboratory. *Journal of the American Dietetic Association* 111(5):S52–55.
- Tsai, A.G., D.F., Williamson, and H.A. Glick. 2011. Direct medical cost of overweight and obesity in the USA: A quantitative systematic review. *Obesity Reviews* 12:50–61.
- USDA. 2012. *Per Capital Data on Products and Food Availability*. United States Department of Agriculture, Washington, D.C. [http://www.ers.usda.gov/data-products/food-availability-\(per-capita\)-data-system.aspx](http://www.ers.usda.gov/data-products/food-availability-(per-capita)-data-system.aspx) (accessed May 9, 2014).
- USDA/USDHHS. 2011. *Dietary Guidelines for Americans, 2010*, seventh edition [United States Department of Agriculture and U.S. Department of Health and Human Services]. Washington, D.C.: United States Government Printing Office.
- USDHHS. 2010. *Nutrition and Weight Status. Healthy People 2020: Topics and Objectives*. United States Department of Health and Human Services, Washington, D.C.

<http://www.healthypeople.gov/2020/topicsobjectives2020/overview.aspx?topicid=29>
(accessed May 9, 2014).

Variyam, J.N. 2008. Do nutrition labels improve dietary outcomes? *Health Economics* 17(6):695–708.

Watters, J.L., and J.A. Satia. 2009. Psychological correlates of dietary fat intakes in African-American adults: A cross-section study. *Nutrition Journal* 8:15.

Wilde, P., P.E. McNamara, and C.K. Ranney. 1999. The effect of income and food programs on dietary quality: a seemingly unrelated regression analysis with error components. *American Journal of Agricultural Economics* 81:959–971.

Table 1. Per capita juice consumption, 1970–2011

Year	Orange	Grapefruit	Apple	Grape	Pineapple	Other ^a	Total
1970	3.72	0.57	0.54	0.22	0.27	0.22	5.53
1975	4.66	0.69	0.58	0.23	0.21	0.33	6.70
1980	4.95	0.58	1.10	0.25	0.31	0.23	7.41
1985	4.81	0.61	1.55	0.23	0.34	0.23	7.76
1990	3.25	0.91	1.74	0.28	0.50	0.34	7.02
1995	4.73	0.59	1.59	0.45	0.38	0.34	8.09
2000	5.54	0.53	1.80	0.34	0.30	0.39	8.90
2001	5.15	0.54	1.79	0.33	0.31	0.42	8.54
2002	5.02	0.47	1.80	0.37	0.32	0.38	8.37
2003	4.90	0.40	1.95	0.40	0.34	0.47	8.46
2004	4.94	0.38	2.13	0.38	0.27	0.41	8.51
2005	4.76	0.23	1.87	0.51	0.26	0.45	8.08
2006	4.39	0.20	2.22	0.44	0.27	0.45	7.98
2007	4.13	0.29	2.28	0.56	0.22	0.44	7.92
2008	3.79	0.30	2.11	0.45	0.27	0.46	7.38
2009	3.92	0.26	2.09	0.38	0.27	0.47	7.40
2010	3.74	0.22	2.21	0.37	0.21	0.45	7.21
2011	3.59	0.25	1.75	0.42	0.23	0.53	6.76

^aOther juices include lemon, lime, cranberry, and prune juices.

Source: USDA (2012) Available at [http://www.ers.usda.gov/data-products/food-availability-\(per-capita\)-data-system.aspx](http://www.ers.usda.gov/data-products/food-availability-(per-capita)-data-system.aspx).

Table 2. Sample statistics and variable definitions, NHANES, 2005-06 thru 2009-10 (N=15,095)

Variable	Definition	Mean	S.D.
Nfp5	Always use the NFP =1; otherwise = 0	0.138	0.345
≤34 years	≤34 years (reference group)	0.328	0.470
35–50 years	35–50 years = 1; otherwise = 0	0.245	0.430
50–64 years	50–64 years = 1, otherwise = 0	0.205	0.404
65+ years	65+ years = 1; otherwise = 0	0.221	0.415
Household Size	household size (persons)	3.260	1.695
Income	household income (\$000)	48.223	29.457
Female	Female = 1; otherwise = 0	0.508	0.500
Married	married =1; otherwise = 0	0.474	0.499
white	Non-Hispanic white (reference group)	0.477	0.499
OTHER	other races = 1; otherwise = 0	0.040	0.196
Black	Non-Hispanic black = 1; otherwise = 0	0.212	0.409
Hispanic	Hispanic = 1; otherwise = 0	0.271	0.444
College	Had college or more education = 1; otherwise = 0	0.425	0.494
FSP	food stamp program participants = 1; otherwise = 0	0.183	0.387
WIC	WIC participants = 1; otherwise = 0	0.094	0.292
BRFT	Days ate breakfast in the two recall days (0–2)	1.692	0.585
DSDCOUNT	# of food supplements taken	1.025	1.748
BMI	body mass index	28.619	6.783
ALQ	# alcoholic drink/day	1.551	2.518
Hdiet1	Self-rated diet quality is excellent or very good	0.299	0.458
Hdiet2	Self-rated diet quality is good or fair =1; otherwise = 0	0.646	0.478
T2	NHANES 2007-08 = 1; otherwise = 0	0.333	0.471
T3	NHANES 2009-10 = 1; otherwise = 0	0.361	0.480
Losewt	Tried to lose weight in the past year = 1; otherwise = 0	0.418	0.493
Ngainwt	Tried not to gain weight in the past year = 1; otherwise = 0	0.096	0.294
Juice	Juice consumption full sample (g)	155.676	310.953
Juice (> 0)	Juice consumption for juice drinkers only (subsample, g)	462.828	381.205

Table 3. Tobit regression results^a

Juice Consumption	Tobit Estimates		$\partial \Pr(y_i > 0)/\partial x_j$		$\partial E(y_i y^* > 0)/\partial x_j$	
	Coef.	Std. Err.	dy/dx	Std. Err.	dy/dx	Std. Err.
Nfp5	-59.32*	26.10	-0.030*	0.013	-15.67*	6.89
35–50 years	-176.60*	25.61	-0.089*	0.013	-46.64*	6.71
50–64 years	-89.86*	28.99	-0.045*	0.015	-23.73*	7.65
65+ years	37.13**	28.03	0.019**	0.014	9.81**	7.39
Household size	9.75**	6.76	0.005**	0.003	2.58**	1.78
Income	-0.54**	0.34	0.0003**	0.0002	-0.14**	0.09
Female	-112.02*	18.54	-0.056*	0.009	-29.59*	4.87
Married	-36.27*	19.89	-0.018*	0.010	-9.58*	5.25
Other Race	14.39	41.80	0.007	0.021	3.80	11.04
Black	264.35*	22.18	0.133*	0.011	69.82*	5.66
Hispanic	167.49*	23.04	0.084*	0.012	44.24*	5.99
College	95.89*	19.18	0.048*	0.010	25.32*	5.06
FSP	40.56*	25.28	0.020*	0.013	10.71*	6.67
WIC	42.74**	32.36	0.022**	0.016	11.29**	8.54
BRFT	182.68*	18.04	0.092*	0.009	48.25*	4.71
DSDCOUNT	16.86*	4.99	0.008*	0.002	4.45*	1.31
BMI	-5.12*	1.51	-0.003*	0.001	-1.35*	0.40
ALQ	-11.51*	4.38	-0.006*	0.002	-3.04*	1.16
Hdiet1	147.96*	45.46	0.075*	0.023	39.08*	11.99
Hdiet2	71.61*	42.95	0.036*	0.022	18.91*	11.34
T2	-25.94	22.38	-0.013	0.011	-6.85	5.91
T3	-34.58*	21.67	-0.017*	0.011	-9.13*	5.72
Losewt	-25.56	20.50	-0.013	0.010	-6.75	5.41
Ngainwt	50.97*	30.45	0.026*	0.015	13.46*	8.03
Constant	-553.03*	72.28				
Sigma	689.433*	17.740				

^aWeighted Tobit estimates.*Statistically different from zero at $\alpha = 0.05$ level.**Statistically different from zero at $\alpha = 0.10$ level.