



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

**An assessment of variable importance when predicting greenhouse gas emissions, beef
output and land use of German dairy farms**

Monika Zehetmeier

Technische Universität München, 85350 Freising
Email: monika.zehetmeier@tum.de

Helmut Hoffmann

Technische Universität München, 85350 Freising
hoffmann@wzw.tum.de

Johannes Sauer

Technische Universität München, 85350 Freising
Email: jo.sauer@tum.de

Donal O'Brien

Teagasc, Moorepark, Ireland
Email: Donal.O'Brien@teagasc.ie

***Selected Poster prepared for presentation at the Agricultural & Applied Economics Association's
2014 AAEA Annual Meeting, Minneapolis, Minnesota, 27-29 June 2014***

Copyright 2014 by Zehetmeier Monika, Hoffmann Helmut, Sauer Johannes, O'Brien Donal. All rights reserved. Readers may make verbatim copies of this document for non-commercial purposes by any means, provided that this copyright notice appears on all such copies.



Conclusions



The dominance analysis approach enabled a ranking of a set of regressors with different units in terms of relative importance.

The implementation of the indicator „beef output per kg FPCM³“ helped to identify those variables that have a high potential to reduce GHG emissions and land use per kg FPCM BUT could result in a shift of GHG emissions between production systems.

Objectives

This study investigates GHG emissions, beef output and land use per kg of FPCM of commercial dairy farms from two regions in Germany as affected by breed.

Those variables will be identified that have a high potential to mitigate GHG emissions i.e. (a) are highly variable between farms, (b) have a high level of contribution on GHG emissions (c) do not show a trade off with beef output per kg FPCM and land use per kg FPCM.

Material and Methods

Dairy farms from Bavaria with dual purpose Fleckvieh cows (South-Dual purpose) and from Nordrhein-Westfalia with milk breed Holstein-Friesian dairy cows (West- Milk breed) were investigated. GHG emissions and land use were calculated using a LCA approach.

Table 1: Value of farm characteristics for the investigated dairy farms (2010) (FPCM=fat and protein corrected milk yield)

Item	Unit	South-Dual purpose 27		West-Milk breed 26	
		Mean	(max-min)	Mean	(max-min)
Dairy cows	#	86	(145-49)	149	(457-67)
Milk yield	kg FPCM/cow per year	8559	(9840-7507)	9596	(10680-8186)
Adjusted replacement rate	%	29	(55-14)	27	(51-15)
Calving interval	days	380	(416-359)	410	(461-380)
Feed intake dairy cow	kg DM*/cow/year	7081	(8816-6153)	7686	(8700-7033)

*Dorfner et al. (2010), *DM=dry matter*

Variables that are high contributors to e.g. GHG emissions, beef output or land use and show a high degree of variability are defined as ‘important parameters/variables’.

Most important variables were identified using multiple linear regression and dominance analysis, “relaimpo” package in R (Azen and Budescu, 2003; Groemping, 2006).

$$LMG(x_k) = \frac{1}{p} * \sum_{i=0}^{p-1} \left(\sum_{S \subseteq \{x_1, \dots, x_p\}, |S|=i} \frac{seqR^2(\{x_k\} | S)}{\binom{p-1}{i}} \right)$$

where LMG(xk) equals the average over model sizes i of average improvements in R² when adding regressor xk to a model of size i without xk, seqR²({xk}|S) equals = additional R² when adding xk to a model with the regressors in set S (Groemping, 2006).

Results

- Lower GHG emissions (p<0.05) within West- Milk breed dairy farms but also considerable lower potential beef output
- No statistical significant difference between land use
- Wide range in GHG emissions and land use within the Investigated regions and production systems




Fig 1: GHG emissions, potential beef output and land use per kg of FPCM of investigated dairy farms

Milk yield and replacement rate had the highest impact on variation of GHG emissions of both dairy farm groups.

A trade off between GHG emissions per kg of FPCM and potential beef output per kg of FPCM was shown in the case of milk yield and replacement rate.




Fig 2: Linear regression with variance decomposition indicating the percent of variance in GHG emissions and potential beef output per kg of FPCM accounted for by predictor variables

An increase in milk yield per day of life (joint indicator of milk yield/cow, replacement rate and age of first calving) (Roemer, 2011) captures a high percentage of variance in GHG emission outcomes.




Fig 3: GHG emissions per FPCM as a function of milk yield per day of life (milk yield per cow per life divided by days of life from birth to culled)

References & Contact

References:

Azen, R., Budescu, D.V., 2003; Dorfner, G. und Hofmann, G., 2012; Groemping, U., 2006; Roemer, A., 2011.

¹ Technische Universität München, Germany

² Teagasc, Moorepark, Ireland

³ fat and protein corrected milk

Contacts :



Monika Zehetmeier
Technische Universität München, Agricultural Production and Resource Economics, Akademie 14, 85350 Freising Weihenstephan, Germany
&
Bavarian Research Institute
Menzinger Straße 54
80638 München, Germany

Tel. 089 17800 228

E-Mail: monika.zehetmeier@tum.de