AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

ELSEVIER

Agricultural Economics 22 (2000) 91–103

AGRICULTURAL
ECONOMICS

www.elsevier.com/locate/agecon

The external costs of pasture weed spread: an economic assessment of serrated tussock control

R.E. Jones *, D.T. Vere, M.H. Campbell

Co-operative Research Centre for Weed Management Systems and NSW Agriculture, Orange Agricultural Institute, Forest Road, Orange, New South Wales 2800, Australia

Received 1 February 1999; received in revised form 16 August 1999; accepted 8 September 1999

Abstract

The external cost associated with the spread of pasture weeds such as serrated tussock (*Nassella trichotoma*) is an important economic problem. This problem is complicated in many parts of south-eastern Australia where low rainfall and low soil fertility prevent the economic viability of control of this weed through pasture improvement. A consequence of serrated tussock spread in this region has been calls for increased public intervention in its control. However, because there have been no attempts to measure the external costs of serrated tussock spread, one of the major economic grounds on which this activity might be justified has not been quantified. The purpose of this paper is to provide this information. A stochastic simulation model is developed to determine the size of the external cost associated with the spread of serrated tussock and to evaluate the economic benefits of a range of control scenarios. It is concluded that on low rainfall-low soil fertility country the socially optimal control option for serrated tussock is to retire land from agriculture and re-vegetate it with trees. ©2000 Elsevier Science B.V. All rights reserved.

Keywords: External costs; Weed spread; Stochastic simulation; Australia

1. Introduction

One of the least studied areas of weed research has been to evaluate the external costs of weed spread. External costs result when the actions of individuals impose uncompensated costs on others. Spreading weeds impose external costs where they adversely affect the economic welfare of other landholders, both private and public. These costs indicate a divergence between the private and socially desirable optimal level of weed control. They are thus a component of the overall social costs attributable to weeds and vary according to

the rate of spread, the nature of the production systems affected and the prevailing environment. Where the rate of spread is rapid, the greater is the private–social divergence in control requirements and the stronger is the rationale for public intervention in facilitating weed control (Auld et al., 1987).

External costs are important in relation to weed spread because they are a primary indication of the concept of market failure that is a central issue in proposals for policies of public intervention in weed control. Pannell (1994) considered that market failure pervaded most of the weed management activities of governments. Of the 11 types of market failure that were categorised as being relevant to all weeds, nine were characterised by external costs. The first two categories concerning the spread of weeds between farms

* Corresponding author. Tel.: +612-63913960;

fax: +612-63913975

E-mail address: randall.jones@agric.nsw.gov.au (R.E. Jones)

and from farms to the environment are particularly relevant to pasture weeds. Pannell considered that the appropriate public response to these external cost problems was through direct control and legislation where farms were the source of weed spread in the expectation that such activity would generate greater social benefits than the operations of the free market.

Serrated tussock (*Nassella trichotoma*) is a major weed of pastures in south-eastern Australia given that it has no grazing value because of its high fibre and low protein content and has a propensity to invade and colonise desirable pastures. The main components of the economic problem caused by serrated tussock are the reduced livestock production caused by infestations in pastures, the costs of control with herbicides and improved pastures and the external costs of spread. While the first two components have been well researched (Vere et al., 1980; Edwards and Freebairn, 1982; Vere et al., 1993; Jones and Vere, 1998), the third has not. The external cost aspect of the serrated tussock problem is now assuming greater prominence as the weed becomes more concentrated in the more marginal agricultural areas. While there has been a long history of successful serrated tussock control by landholders and local government, it is apparent that such direct control methods have been ineffective under unfavourable environmental and economic conditions (Vere et al., 1993). The problem of serrated tussock in these areas persists because many landholders cannot undertake profitable control and this results in the infestation of other areas.

Serrated tussock is now well established in large areas of south-eastern Australia in which the soil fertility and rainfall environments prevent economic control under the preferred method of replacement by improved pastures. Heavy infestations of the weed in such areas are a source of infestation to both neighbouring and distant lands as serrated tussock seeds can disperse over long distances. Recent surveys in 1994 (Gorham, unpublished) and in 1997 (Jones and Vere, 1998) found that the total area of serrated tussock in New South Wales was 9 and 30% higher than an earlier estimate by Campbell (1987). Of a total area of approximately 887,000 ha infested by serrated tussock (Jones and Vere, 1998), 14.1, 24.7 and 61.2% was classified as being heavily, moderately and lightly infested, respectively. The Monaro region had 18.1% of the total area of serrated tussock and importantly, 28%

of the heavy infestations which was double the state proportion of heavy to total infestation. In 10 years, heavy serrated tussock areas in the Monaro increased from 2400 to 35000 ha. This temporal change evidence indicates that serrated tussock is a prominent example of the external cost problems caused by weed spread.

The external cost aspects of the spread of serrated tussock are now attracting considerable public concern. Recognition that the problem of this weed in low potential agricultural country has moved beyond the private control decision context has given rise to proposals for increased levels of public intervention (Anon., 1998). As Pannell (1994) has indicated, the presence of market failure and net social benefits comprises the economic rationale for such activity. As the existence of external costs is a foremost condition in the demonstration of market failure, there is a need to evaluate these costs in a logical and consistent manner.

The purpose of this paper is to evaluate options for reducing the external costs associated with the spread of serrated tussock. A stochastic simulation model is developed which incorporates two options for controlling serrated tussock by landholders and one option for public control. While the paper does not attempt to develop a case for public intervention in serrated tussock control, such policies are closely linked with any analysis of the external cost problem.

2. Modelling weed spread

Auld and Coote (1980, 1981) described the rate of weed spread in terms of; (i) population growth rate at a primary infection site, (ii) the proportion of annual population increase which is dispersed beyond the boundaries of the infection site; (iii) the area over which the fraction (ii) is dispersed, and (iv) the susceptibility of invaded areas to colonisation. It has been suggested that invading weed species have a constant exponential rate of population growth (Harper, 1977). Auld and Coote (1980, 1981) used an exponential population growth model for a weed-affected farm represented by Eq. (1).

$$P_n = \frac{P_0}{100} \left[1 + \frac{c}{100} \right]^n, \quad n = 1, \dots, T \quad (1)$$

where P_n is the population in terms of the percentage of area infested in year n (taking values between

Fig. 1. Exponential weed spread.

0 and 100), P_0 is the initial population, c is a constant rate factor, and T is the time to full infestation. In this model, population growth continues until $P_n = 100$, when complete infestation occurs and P_n remains constant. For the spread of serrated tussock, values of 15–20 years to reach full infestation have been observed in naturalised pasture systems. To illustrate the exponential growth model a hypothetical weed spread function is presented in Fig. 1 using parameter values of $P_0 = 2$ and $T = 15$. To derive the function in Fig. 1 the first step is to determine the constant, $(1 + c/100)$, by substituting the appropriate values for P_n and T into Eq. (1).

$$P_{15} = \left(1 + \frac{c}{100}\right)^{15} = 100$$

$$15 \left[\log_{10} \left(1 + \frac{c}{100}\right) \right] = \log_{10}(100)$$

$$\log_{10} \left(1 + \frac{c}{100}\right) = 0.1333$$

$$1 + \frac{c}{100} = 1.359$$

The annual levels of weed infestation illustrated in Fig. 1 were thus estimated from the following equation:

$$P_n = \frac{P_0}{100} [1.359]^n \quad \text{with} \quad P_0 = 2$$

For a weed such as serrated tussock, where wind is the main seed dispersal agent, a small proportion of seeds are distributed significant distances from the parent plant. Approximately half of the seeds fall near

the parent prior to the inflorescences being dispersed by wind, the remainder are transported in the inflorescence to spread on and beyond the farm. In relation to the annual increment in population growth, it is reasonable to assume that the dispersal fraction from one farm to others is a constant from year to year. If a fixed proportion, s , of the annual population increase occurs beyond the farm then Auld and Coote (1980, 1981) propose annual population growth (on the farm) as;

$$P_n = \frac{P_0}{100} \left[\left(1 + \frac{c}{100}\right) \left(1 - \frac{s}{100}\right) \right]^n, \quad n = 1, \dots, T \quad (2)$$

In this study the annual population growth equation as proposed by Auld and Coote (1980, 1981) is reformulated to Eq. (3). This equation has two developments. First, notation (i) is introduced to represent the specific farm infested and, second, it is now a Markov equation as infestation in year n is a function of infestation in year $n-1$ and not the initial infestation in year 0. This latter feature facilitates the modelling of a dynamic system.

$$P_{i,n} = \frac{P_{i,n-1}}{100} \left[\left(1 + \frac{c_i}{100}\right) \left(1 - \frac{s_i}{100}\right) \right], \quad n = 1, \dots, T \quad (3)$$

Eq. (3) implies that the weed population growth on farm i is reduced to the extent that some of the population growth is dispersed beyond the farm boundary. The first term within the brackets of Eq. (3), $(1 + c_i/100)$, represents the proportional increase in weed infestation while the second term, $(1 - s_i/100)$, represents the proportion of the new infestation dispersed beyond the farm. Consequently, $P_{i,n}$ represents the level of infestation only on farm i in year n , not the total level of infestation derived from the previous farm infestation, $P_{i,n-1}$.

Now consider a neighbour, farm j . Weed population growth on this farm will be a function of increasing density from previous infestations on farm j , plus additional infestations from the dispersal fraction on farm i . This is represented as follows;

$$P_{j,n} = \frac{P_{j,n-1}}{100} \left[\left(1 + \frac{c_j}{100}\right) \left(1 - \frac{s_j}{100}\right) \right] + \frac{P_{i,n-1}}{100} \left[\frac{s_i}{100} \right], \quad n = 1, \dots, T \quad (4)$$

This process of weed spread may continue with farm j (and possibly farm i) dispersing a fraction of infestation to farm k and so on. Additionally, there is the potential for infestation dispersal from farm j to farm i . If farm j has no initial weed presence, i.e. $P_{j,0} = 0$, then the only source of weed infestation is from colonisation by weed dispersal from farm i . However, once this takes place then $P_{j,n} > 0$ and population growth can occur from infestations within the farm.

As the parameter s is a proportional increase in infestation from farm i to farm j , it is independent of the size of farm j . Consequently, the area of infestation occurring on farm j in any given year from farm i is calculated as a function of the infested area on farm i by the dispersal fraction. Auld and Coote (1980, 1981) restricted the parameter s to occur only following population saturation at the farm level. Given the nature of seed spread from serrated tussock there appears little evidence to impose such a constraint and the approach adopted here was to allow spread at all population levels.

3. The serrated tussock model

A stochastic simulation model was developed to evaluate management options for reducing the external cost of serrated tussock. Rainfall is the major stochastic variable as it affects pasture productivity, and therefore potential livestock stocking rate, and the establishment success of perennial pastures and trees. The principles of weed spread are explicitly incorporated in the model as is the impact of serrated tussock on pasture and livestock performance.

The model used a Latin hypercube sampling procedure and was solved for 5000 iterations over 25 years at a discount rate of 5%. The solution included estimated means, standard deviations and percentile values for net present value (NPV) for each specific area. The model estimated the benefits of a range of control scenarios and ranked the results according to stochastic dominance (Anderson et al., 1977).

3.1. Land types

The effect of serrated tussock spread on- and off-farm was estimated from Eqs. (3) and (4). The private cost from weed spread was determined by ap-

plying Eq. (3) to estimate weed density, while Eq. (4) was used to determine the associated external costs.

The external costs of serrated tussock and the benefits of its control were estimated for two representative land types in a low rainfall environment. Area A comprised of a naturalised pasture on non-arable land with low soil fertility and a heavy serrated tussock infestation. Area B comprised of a fertilised mix of naturalised pasture in arable high fertility country and with no initial serrated tussock. The distribution of spread of serrated tussock over area B , through the dispersal proportion s , was assumed to be uniform. The alternative land types are illustrated in Fig. 2 and are represented by the two concentric rings, the inner circle being area A while the outer ring is area B . The external boundary of area B represents the outer limit to which new infestations can occur. Areas A and B can be thought of as representing farms i and j , but here area B represents an aggregation of three to four farms affected by seed dispersal from farm i (i.e. area A). By treating area B as one large farm, infestation of neighbouring farms in area B is endogenously incorporated through the first term of Eq. (4).

Although serrated tussock seeds have been observed to travel up to 20 km, the large proportion of seeds which lead to new infestations occur on neighbouring farms and consequently travel much smaller distances. Accordingly, the dispersal proportion was confined to a radius of 5 km from the source. It is assumed that weeds are distributed uniformly in area A and the source of the infestation is taken to be the midpoint of the radius of area A , r_A .

Given the relationship between the dispersal proportion (s) and the distance seeds can travel (d) there is a direct relationship between the size of area A and the circumference of area B and accordingly its area. Setting area A to be 1000 ha, the radius (r_A) is calculated at 1.784 km ($r = 0.1\sqrt{A/\pi}$)¹. If $d = 5$ the radius of area B (r_B) is 5.892 km [i.e. $5 + 0.5(1.784)$] and the size of area B is calculated as;

$$B = \pi 100r_B^2 - A$$

¹ The value 0.1 is simply a constant for converting from hectares to kilometres. Likewise the value 100 in the following equation for calculating the size of area B is a constant for converting kilometres to hectares.

Fig. 2. Land types for measuring weed spread.

where B and A are the sizes of the respective areas. From this process area B was determined to be 9906 ha.

3.2. Control scenarios and annual decisions

Three separate serrated tussock control scenarios were proposed for area A . Control options in area B were not included as the objective was to evaluate the external cost of weed spread imposed on this region. A 'no control' scenario was used to estimate the impact of each of the control scenarios upon the external cost due to weed spread. A 'no control' scenario is not a serious management option as serrated tussock is a declared noxious weed and by legislation must be controlled. The three control scenarios are described as follows.

Scenario 1: Naturalised pasture with periodic herbicide control. This scenario involves the spraying of a herbicide on a naturalised pasture supporting sheep when serrated tussock exceeds 50% of the area. This is the least cost scenario and given that no additional con-

trols are imposed (such as competitive pasture species) it can be expected that serrated tussock will need to be controlled again within 7–10 years. The main species in this pasture are winter growing *Danthonia* spp., *Trifolium subterraneum*, annual grasses (*Vulpia* spp., *Hordeum leporinum*), and summer-growing *Bothriochloa macra*, *Microlaena stipoides*, *Themeda triandra* and *Stipa* spp. Given the nature of summer and winter production from these pastures, annual rainfall distribution is considered an important determinant of pasture production. As the herbicides used to control serrated tussock can have a negative effect upon some of these species, no livestock can be supported in the year of control. After a herbicide treatment it can take up to 3 years until a pasture fully recovers and, consequently, stocking rates of 50 and 80% of full carrying capacity in the years 2 and 3 after control are assumed.

Scenario 2: Introduced perennial pasture. A successful method of serrated tussock control is to sow a competitive introduced perennial pasture, such as *Trifolium subterranean* with *Phalaris aquatica* and/or *Dactylis glomerata*, after spraying. If successfully es-

established, these species can replace serrated tussock on infested areas and provide long-term control. On non-arable low fertility country, the probability of successful establishment is low due to the difficulty in establishing pastures by aerial seeding in dry years. The two critical periods for rainfall which determine whether establishment will be successful are June to August and September to December. If rainfall is deficient in either of these two periods, a perennial pasture will fail to establish and re-sowing will be necessary when seasonal conditions permit. Rainfall from May to December determines production once the pasture is established from the air. The treated area should be rested from grazing for 1 year following sowing to ensure maximum weed control from sown pastures. As it takes 3–4 years for a newly established perennial pasture to reach full production, pasture production in years 2 and 3 following the establishment were set to be 50 and 80% of full production, respectively.

Scenario 3: Tree establishment. Vere et al. (1993) demonstrated that it is unprofitable to control serrated tussock by pasture improvement in non-arable areas with low rainfall and low soil fertility. Campbell and Nicol (1996) proposed that the establishment of trees to control weeds would be a more sustainable option in such situations. Tree varieties that achieve this goal are *Pinus radiata* and various species of *Eucalyptus*. The objective of tree planting is solely for weed control and no financial returns, either from agroforestry or firewood, are to be expected. This means that the land is withdrawn from agricultural production. Rainfall between May and December determines the success or failure of tree establishment. If drought conditions prevail during this period then tree establishment has failed and would need to be repeated the following year. Once trees have established, moderate variations in climate do not have a significant effect upon the efficacy of serrated tussock control by trees.

The capital and annual variable costs for the scenarios are given in Table 1. The details of the annual management decisions for the three control scenarios are given in Table 2.

3.3. Control effects

The dispersal proportion s_A (i.e. dispersal from area A) was set at 5 which, due to a lack of more objective

Table 1

Capital and variable costs associated with control strategies

	Capital cost (\$/ha)	Variable cost (\$/ha)
Naturalised pasture — herbicide control	100	—
Maintain naturalised pasture	—	15
Establish perennial pasture	286	—
Maintain perennial pasture	—	90
Establish trees	725	—

Table 2

Annual decisions associated with control strategies

Decision	Description	Scenario
1	Control naturalised pasture with herbicide	1
2	Graze naturalised pasture — year 2	1
3	Graze naturalised pasture — year 3	1
4	Graze naturalised pasture — year 4 and onwards	1
5	Establish an introduced perennial pasture (phalaris)	2
6	Graze perennial pasture — year 2	2
7	Graze perennial pasture — year 3	2
8	Graze perennial pasture — year 4 and onwards	2
9	Establish trees	3
10	Grow trees	3

information, was obtained from anecdotal evidence. For Scenario 3 it was assumed that trees reduced seed dispersal by 95% from a combination of the tree cover and increased competition from other plant species that establish once serrated tussock and livestock are removed from the system. The number of years until full infestation, T , was set at 15, 30 and 100 for scenarios 1, 2 and 3, respectively. The effect of the different control methods was to reduce the level of infestation in the year of treatment. The new level of infestation following control was 3% (i.e. $P_{A,n} = 3$, a 97% weed kill) for each control decision. The initial weed infestations were set in the model at $P_{A,0} = 80$ and $P_{B,0} = 0$.

3.4. Climatic effects

Rainfall is a critical determinant of pasture production, and therefore carrying capacity, and the efficacy of the control decisions. Four climatic scenarios were developed to represent the normal range of rainfall

Table 3
Rainfall probability distributions

Rainfall period	Distribution type	Mean	Standard deviation
Annual	Lognormal	480.86	142.33
June to August	Lognormal	83.42	48.44
September to November	Lognormal	125.90	67.75
May to December	Normal	289.26	90.85

conditions in the region — drought, dry, median and favourable. Monthly rainfall data for the township of Dalgety between 1896 and 1994 was used to construct each climatic scenario and determine the associated probabilities of occurrence.

Four rainfall periods were derived; annual, June to August, September to November, and May to December. Annual rainfall was used to estimate production of native pastures. The two rainfall periods June to August and September to November were used to estimate the success or failure of establishing a perennial pasture. The period May to December was used to estimate the production of an established perennial pasture and the success or failure of tree establishment.

The appropriate probability distribution for rainfall is expected to be either a normal distribution or a lognormal distribution to account for any possible skewness in the data. For the four rainfall periods various probability distributions were tested for goodness of fit and the distributions and parameters chosen are given in Table 3.

3.5. Pasture production

The production of naturalised and perennial pastures was based on the daily pasture growth curves for poor, average and good growing conditions, represented by the 25th, 50th and 75th percentiles (McDonald, 1995). These descriptions corresponded with the dry, median and favourable descriptions used in this analysis and McDonald's seasonal trends were used for both naturalised and perennial pastures, with a drought category included and calculated at the 10th percentile of pasture dry matter production (Table 4). An 'expert' panel approach was used to estimate the production differences in the monthly growth curves for different soil fertility and rainfall conditions.

The livestock enterprise considered was a Merino wether system producing 21 μ m wool. The potential

Table 4
Climatic effects on pasture production by seasonal growing condition (kg DM/ha)

	Growing conditions			
	Drought	Dry	Median	Favourable
<i>High fertility</i>				
Perennial pasture	2421	4841	9630	16416
Good native pasture	1159	2318	4725	9486
Poor native pasture	587	1174	2490	4847
<i>Medium fertility</i>				
Perennial pasture	1936	3873	7701	13133
Good native pasture	927	1854	3780	7589
Poor native pasture	470	939	1992	3878
<i>Low fertility</i>				
Perennial pasture	1452	2905	5778	9850
Good native pasture	695	1391	2835	5691
Poor native pasture	352	704	1494	2908

Table 5
Climatic effects on stocking rate by seasonal growing condition (DSE/ha)

	Growing conditions			
	Drought	Dry	Median	Favourable
<i>High fertility</i>				
Perennial pasture	0.5888	1.1761	4.7201	16.0406
Good native pasture	0.5058	1.0117	3.9781	6.1422
Poor native pasture	0.0854	0.1708	0.5754	0.9399
<i>Medium fertility</i>				
Perennial pasture	0.4710	0.9409	3.7761	12.8325
Good native pasture	0.4046	0.8094	3.1825	4.9138
Poor native pasture	0.0683	0.1366	0.4603	0.7519
<i>Low fertility</i>				
Perennial pasture	0.3533	0.7057	2.8321	9.6244
Good native pasture	0.3035	0.6070	2.3869	3.6853
Poor native pasture	0.0410	0.1025	0.3452	0.5639

livestock carrying capacity² for each soil fertility, pasture type and climatic scenario were determined from a linear programming model of southern New South Wales (Table 5) which incorporated the seasonality of pasture supply and livestock feed demands (Jones and Vere, 1998).

² Stocking rates are reported on the basis of dry sheep equivalents (DSE) per hectare. A DSE is a rating based on the amount of energy required to maintain a 50 kg wether per annum.

3.6. Weed effects

Following the approach of Denne (1988) and Jones and Vere (1998) it was assumed that there is a linear relationship between weed density and pasture yield loss. Thus, if a serrated tussock infestation is 40% of an area, there is expected to be a 40% reduction in pasture production.

Cousens (1985) has argued that the appropriate yield loss function for annual crops is a rectangular hyperbola because at low densities weeds are most competitive to crops and hence cause a maximum marginal reduction in yield. The effect of an increase in weed numbers at low densities is additive. However, when the density is high increased intra-specific weed competition tends to reduce the marginal yield loss. The function that Cousens derived was;

$$Y_L = \frac{ID}{1 + (ID/A)} \quad (5)$$

where Y_L is percentage yield lost because of weed competition, D is weed density (plants per square metre), I is the percentage yield loss per unit weed density as weed density approaches zero, and A is an estimate of the maximum yield loss of a weedy crop relative to the yield of a weed free crop.

This hyperbolic yield-loss function has been validated (Cousens et al., 1984; Martin et al., 1987) for annual crops where weeds are measured in terms of some land unit, such as plants per square metre. There have been few attempts in pasture systems to either validate the hyperbolic function or estimate the most appropriate functional form for yield loss when weed infestations are reported on a percentage infestation. As a result, in the absence of any better information the simpler linear yield loss relationship has been adopted here.

3.7. Economic effects

The economic performance measure used in this analysis is the NPV of annual profit over a 25 year simulation period. NPVs were calculated separately for area A , area B and the combined total of the two areas. The annual profit function for both area A and area B was calculated as follows;

$$\pi_{i,n} = H_i \left[\left(1 - \frac{P_{i,n}}{100} \right) SR_{s,k,n} GM - AC_n - K_n \right] \quad (6)$$

where $\pi_{i,n}$ is the profit of the i th region in year n , H is the area (hectares) of the i th region, $P_{i,n}$ is the level of infestation on the i th area in year n , $SR_{s,k,n}$ is the stocking rate of the k th pasture type in the s th season in year n (Table 5), GM is the annual sheep enterprise gross margin (\$15.40 per DSE), AC are annual operating costs and K are capital costs (Table 1). The parameter i can take values of A for area A , B for area B and T for the total area where $\pi_{T,n} = \pi_{A,n} + \pi_{B,n}$. For scenario 3, SR was set to zero so $\pi_{A,0} = -725,000$ (i.e. -725×1000 ha) and zero thereafter. The discount rate used was 5%.

4. Results

4.1. Level of serrated tussock infestation

The expected level of serrated tussock infestation of both area A and area B for each control scenario (Fig. 3) was derived from a deterministic version of the model. Application of control scenario 3 resulted in infestation levels close to zero in both areas and was not plotted. Implementation of scenarios 1 and 2 both reduced the level of infestation on area A , however, within a 7–10-year period significant re-infestation had occurred and control was again necessary. Reliance upon either scenario 1 or 2 to control serrated tussock infestations in area A resulted in a significant level of infestation in the previously unaffected area B within 20 years.

4.2. Benefits of the control options

The ‘no control’ scenario for area A was run in addition to the three control scenarios to compare the economic benefits of each scenario for area A , area B and the total area. Reported in Table 6 are the minimum, maximum, mean and standard deviation of NPV for each scenario.

Area A: The ‘no control’ scenario resulted in a mean NPV for area A of $-\$211,409$. Scenario 2 was the only control option that gave a positive mean NPV for area A ($\$314,642$). Relying upon scenario 1 caused economic losses ($-\$358,047$) from the combination

Fig. 3. Infestation of serrated tussock on areas A and B for scenarios 1 and 2.

Table 6
Simulation model NPV results (\$)

	Minimum	Maximum	Mean	Standard deviation
<i>No control</i>				
Area A	-211409	-211409	-211409	0
Area B	164901	2454092	1377136	334035
Total	-46508	2242683	1165727	334035
<i>Scenario 1</i>				
Area A	-368935	-347031	-358047	3298
Area B	4439285	7562788	6051184	482760
Total	4070786	7215757	5693137	485597
<i>Scenario 2</i>				
Area A	-186775	776226	314642	154602
Area B	4675246	8305960	6323640	519017
Total	5016550	8276932	6638282	505698
<i>Scenario 3</i>				
Area A	-1348073	-690476	-693326	43194
Area B	5648109	9450052	7724302	523733
Total	4957632	8759576	7030976	524939

of high control costs and low stocking rates. Tree establishment has high capital costs and no annual income, consequently the mean NPV of scenario 3 (−\$693,326) was the lowest of all the scenarios.

Area B: The 'no control' scenario resulted in the lowest mean NPV (\$1,377,136). Scenario 3 resulted in the highest mean NPV (\$7,724,302) followed by sce-

nario 2 (\$6,323,640) and scenario 1 (\$6,051,184). To calculate the external cost associated with the spread of serrated tussock, a 'no weed spread' scenario was run (mean NPV \$7,741,601) with the external cost for each scenario being the difference in the mean NPVs. This resulted in calculated external costs of \$1,690,417, \$1,417,961 and \$17,299 in terms of mean

NPV for scenarios 1, 2 and 3, respectively. On the basis of these results, adoption of scenario 3 to control serrated tussock in area A would be preferred by landholders in area B as it resulted in the minimum external cost being imposed upon them.

The total area: An economic trade-off is involved in protecting production in area B by establishing trees in area A. The opportunity cost to area A (\$1,007,968 mean NPV) is represented by the difference in mean NPV between scenarios 2 and 3. Therefore, the combined result from areas A and B are required to determine the socially preferable scenario. Scenario 3 resulted in the greatest mean NPV for the total area (\$7,030,976), which was significantly greater than scenario 1 (\$5,693,137 mean NPV) and scenario 2 (\$6,638,282 mean NPV).

4.3. Risk aversion and stochastic dominance

Ranking the results on the basis of mean NPV implies risk neutrality by an individual decision maker. There is evidence that most Australian farmers are risk averse (Bardsley and Harris, 1987) and consequently if risk attitudes are taken into account the rankings of the scenarios may change. For example, for the total area the standard deviation of NPV for scenario 3 was greater than the standard deviation of scenarios 1 and 2, implying a greater degree of income variability and risk.

Testing strategies for stochastic dominance is a means of ranking alternative strategies when risk preferences are unknown (Anderson et al., 1977). Stochastic efficiency rules are applied by undertaking pairwise comparisons of the cumulative distribution functions (CDFs) of the scenarios (Fig. 4). On the basis of this analysis scenario 3 was preferred. As scenario 3 exhibited first-degree stochastic dominance no further testing, such as for stochastic dominance with respect to a function (Meyer, 1977a, b) was required.

4.4. Sensitivity analyses

Sensitivity analyses were applied to a number of parameters that were considered variable. These were a higher level of soil fertility for area A, different sizes of infestation, a greater dispersal distance, the population dispersal proportion and a lower initial infestation

Table 7

Sensitivity analysis on soil fertility, size of area A, dispersal distance and initial infestation (\$ NPV)

	Mean	Standard deviation
<i>Medium fertility</i>		
No control	1623510	283455
Scenario 1	4246150	348457
Scenario 2	4783324	395678
Scenario 3	4656215	355860
<i>Area A of 500 ha</i>		
No control	1991501	290245
Scenario 1	4360789	347024
Scenario 2	4594381	364026
Scenario 3	4618485	353106
<i>Dispersal distance 10 km</i>		
No control	9168800	1179900
Scenario 1	17476060	1335309
Scenario 2	17942930	1369164
Scenario 3	17992120	1345908
<i>Dispersal proportion 10%</i>		
No control	787503	311811
Scenario 1	5765205	471238
Scenario 2	6716380	492717
Scenario 3	7035977	511164
<i>Initial infestation 20%</i>		
No control	2585173	335323
Scenario 1	3763278	367922
Scenario 2	4652386	415935
Scenario 3	4658397	391635

on area A. Reported in Table 7 are the results from the sensitivity analysis in terms of mean and standard deviation of NPV for the total area.

Effect of alternative soil fertility: The analysis was repeated for the case where the soil fertility of area A was medium instead of low. Higher soil fertilities have greater potential pasture production and stocking rates and thus, greater opportunity costs from their removal from agriculture. Consequently, the optimal solution for controlling weed spread may differ under alternative soil fertility conditions. Adopting a higher soil fertility for area A resulted in scenario 2 being the preferred control option as it had the (marginally) highest mean NPV.

Alternative size in infested area: The size of many motherload areas in the Monaro region may be smaller than the assumed 1000 ha. Reducing the size of area A to 500 ha tested the effect of the size of the infested area upon the robustness of the results. There was no

Fig. 4. Cumulative distribution functions for the alternative scenarios.

change in the optimal solution with scenario 3 remaining the preferred control option, although it was only marginally superior to scenario 2.

Seed dispersal distance: The analysis purposely used a conservative estimate of the dispersal distance, d . Given the uncertainty of the distance with which new infestations can occur from the original source, parameter d was increased from 5 to 10 km. The effect of this change was to increase the size of area B from 9906 to 36,271 ha. Despite the size of the external cost increasing with the increase in dispersal distance, there was no change in the relative rankings of the scenarios.

Population dispersal proportion: The population dispersal proportion parameter, s , was obtained from anecdotal evidence. The effect of variations in the parameter was tested arbitrarily by increasing the parameter value from 5 to 10%. This only had a marginal effect on the mean NPV for the three scenarios and did not change the preferred ranking.

Initial level of infestation: The analysis was repeated for a significantly lower level of initial infestation of serrated tussock. At an initial infestation of 20% there was little difference between scenarios 2 and 3. The larger standard deviation associated with scenario 2 suggests that scenario 3 would remain the preferred

control option even for lower levels of initial infestation on area A .

5. Summary and discussion

An important cost associated with weeds is the external cost due to weed spread beyond farm boundaries. A stochastic simulation model was developed to determine the size of the external cost from the pasture weed serrated tussock in south-eastern Australia and to evaluate the benefits of alternative control scenarios. The model measured the rate of spread both within an area already infested with serrated tussock and an adjoining area, not previously infested, which was susceptible to weed spread from wind-borne seeds. Both areas were assumed to be in a low rainfall region and the infested area was of low soil fertility.

Three separate control scenarios for the infested area were assessed, (i) periodic use of a herbicide in a naturalised pasture system, (ii) herbicide application followed by establishment of an introduced perennial pasture and (iii), herbicide control followed by tree establishment. The scenario that resulted in maximum returns to society was the retirement of the infested area from agriculture and re-vegetation with

trees, a result that held for risk attitudes ranging from risk-neutral to risk-averse. This scenario minimised the external cost from serrated tussock spreading to the adjoining area. The scenario that maximised returns to the private landholder within the infested area was control with a herbicide followed by establishment of an introduced perennial pasture. The commonly adopted control option of periodic herbicide use in a naturalised pasture system gave negative returns. The reasons for the adoption of this scenario are compulsion by legislation, and severe budget constraints combined with the high capital costs of establishing an introduced perennial pasture. Sensitivity analyses on the results indicated that they were robust for a range in variation in key parameters. The ranking of the options changed when the infested area was assumed to be of medium instead of low fertility, in which case the perennial pasture control scenario became socially optimal.

This analysis indicates that there is a clear divergence between the socially optimal form of serrated tussock control, re-vegetation with trees, and that which is privately optimal to an individual decision-maker on the infested area, introduced perennial pasture. Pursuing the privately optimal form of control will result in continued and significant external costs to neighbouring landholders. This represents an example of market failure, a necessary condition before any form of government intervention can be justified.

The policy implications of these results mainly relate to the issues of justifying some form of intervention in weed control and to the more difficult policy of acquiring marginal agricultural and environmentally sensitive land from private landholders. The first issue is more readily addressed where the external costs and market failure aspects of the problem have been established. The second issue is more difficult because of the often politically sensitive nature of land retirement. The community's acceptance of a land acquisition and rehabilitation policy is expected to be mixed. Some individuals, particularly those affected by the external costs, will support this policy while others will resist moves to retire their land from agriculture. There are few Australian precedents for such a policy and the political acceptance is unknown. An alternative and perhaps more efficient option to direct government intervention may be for landholders affected by weed

spread to instigate some form of collective action at a regional or catchment level, such as purchasing or rehabilitating these marginal lands. This represents market driven policy response to the problem rather than reliance upon government intervention.

The estimated benefits for scenario 3 are considered conservative as they only consider the agricultural benefits from a reduction in the external costs of weed spread. There are a range of environmental amenity values attached to native ecosystems which are likely to be improved by the re-vegetation of agricultural lands, particularly if it involves native species. In addition, there may be benefits also associated with tree planting in terms of reduced soil loss and stream turbidity, a reduction in hydraulic loading with consequent improved dryland salinity effects, and the encouragement of a greater natural bio-diversity. Environmental benefits may also be derived from retiring marginal agricultural land and revegetation with trees due to the cessation of periodic chemical use to control serrated tussock.

The results indicate that care needs to be taken when designing an economic analysis of weed spread. Proper consideration must be given to adjoining land that is negatively impacted upon by spread from infested areas. If an analysis focuses on the benefits and costs of control solely within the infested area, then the derived preferred strategies are unlikely to closely correlate with those that are socially optimal.

References

- Anderson, J.R., Dillon, J.L., Hardaker, J.B., 1977. *Agricultural Decision Analysis*. Iowa State University Press, Ames.
- Anon., 1998. Proc. of Workshop on Recovering the Monaro. Cooma.
- Auld, B.A., Coote, B.G., 1980. A model of a spreading plant population. *Oikos* 34, 287–292.
- Auld, B.A., Coote, B.G., 1981. Prediction of pasture invasion by *Nassella trichotoma* (gramineae) in south east Australia. *Prot. Ecol.* 3, 271–277.
- Auld, B.A., Menz, K., Tisdell, C., 1987. *Weed Control Economics*. Academic Press, London.
- Bardsley, P., Harris, M., 1987. An approach to the econometric estimation of attitudes to risk in agriculture. *Aust. J. Agric. Econ.* 31 (2), 112–126.
- Campbell, M.H., 1987. Area and distribution of serrated tussock (*Nassella trichotoma* (Nees) Arech.) in New South Wales, 1975 to 1985. *Plant Prot. Quart.* 2 (4), 161–164.

- Campbell, M.H., Nicol, H.I., 1996. Establishing trees on non-arable land to control weeds. In: Proc. 11th Australian Weeds Conference, Melbourne, pp. 493–496.
- Cousens, R., 1985. A simple model relating yield loss to weed density. *Ann. App. Biol.* 107, 239–252.
- Cousens, R., Peters, N.C.B., Marshall, C.J., 1984. Models of yield loss — weed density relationships. In: Proc. 7th International Symposium on Weed Biology, Ecology and Systematics, Paris, pp. 367–374.
- Denne, T., 1988. Economics of nassella tussock (*Nassella trichotoma*) control in New Zealand. *Agric. Ecosystems and Environ.* 20, 259–278.
- Edwards, G.W., Freebairn, J.W., 1982. The social benefits from an increase in productivity in a part of an industry. *Rev. Mktg. Agric. Econ.* 50 (2), 193–210.
- Harper, J.L., 1977. *Population Biology of Plants*. Academic Press, London.
- Jones, R.E., Vere, D.T., 1998. The economics of serrated tussock in New South Wales. *Plant Prot. Quart.* 13 (2), 70–76.
- McDonald, W., 1995. Matching pasture production to livestock enterprise requirements for Tableland and Slopes districts of NSW. NSW Agriculture, unpublished mimeo.
- Martin, R.J., Cullis, B.R., McNamara, D.W., 1987. Prediction of wheat yield loss due to competition by wild oats (*Avena* spp.). *Aust. J. Agric. Res.* 38, 487–499.
- Meyer, J., 1977a. Choice among distributions. *J. Econ. Theory* 14 (2), 326–336.
- Meyer, J., 1977b. Second-degree stochastic dominance with respect to a function. *Int. Econ. Rev.* 18 (2), 477–487.
- Pannell, D., 1994. Economic justifications for government involvement in weed management: a catalogue of market failures. *Plant Prot. Quart.* 9, 131–137.
- Vere, D.T., Sinden, J.A., Campbell, M.H., 1980. Social benefits of serrated tussock control in New South Wales. *Rev. Mktg. Agric. Econ.* 48 (3), 123–138.
- Vere, D.T., Auld, B.A., Campbell, M.H., 1993. Economic assessments of serrated tussock (*Nassella trichotoma*) as a pasture weed. *Weed Tech.* 7, 776–782.

Guide for Authors

Agricultural Economics publishes articles covering the range of work done on agricultural economics, divided into three categories. (1) Disciplinary work: improvement of theories, techniques and descriptive knowledge of economics and its contributing disciplines such as statistics, mathematics and philosophy. (2) Multi-disciplinary subject matter areas: energy, technical change, institutional change, natural resources, farm management, rural communities, marketing, human development and the environment – areas which are important to fairly well-defined groups of public and private decision-makers facing well-defined sets of problems. (3) Problem solving: the definition, solution and management of specific practical problems. Work in each of these three categories may deal with teaching, extension and out-reach, consulting, advising, entrepreneurship and administration, as well as research. All of these may require knowledge of values, non-monetary as well as monetary. The Editor and Editorial Board, under the general direction of the IAAE's President, Executive Committee and Council, are charged with implementing Journal policy to serve members of IAAE around the world.

Submission of manuscripts

Submission of an article is understood to imply that the article is original and is not being considered for publication elsewhere. Upon acceptance of the article by the journal, the author(s) will be asked to transfer the copyright of the article to the publisher. This transfer will ensure the widest possible dissemination of information.

Papers for consideration should be submitted to:

Dr S.R. Johnson
Editor-in-Chief *Agricultural Economics*
218 Beardshear Hall
Iowa State University
Ames, IA 50011-2020
USA
Tel. (+1-515) 294 6635
Fax (+1-515) 294 9781
E-mail: ecbalm@iastate.edu

Book reviews will be included in the journal on a range of relevant books which are not more than 2 years old. Book reviews will be solicited by the Book Review Editor. Unsolicited reviews will not usually be accepted, but suggestions for appropriate books for review may be sent to the Book Review Editor:

C.L. Delgado
International Food Policy Research Institute
2033 K Street, NW
Washington, DC 20006
USA

Electronic manuscripts

Electronic manuscripts have the advantage that there is no need for the rekeying of text, thereby avoiding the possibility of introducing errors and resulting in reliable and fast delivery of proofs.

For the initial submission of manuscripts for consideration, hardcopies are sufficient. For the processing of *accepted papers*, electronic versions are preferred. After *final acceptance*, your disk plus two, final and

exactly matching printed versions should be submitted together. Double density (DD) or high density (HD) diskettes (3.5 or 5.25 inch) are acceptable. It is important that the file saved is in the native format of the wordprocessor program used. Label the disk with the name of the computer and wordprocessing package used, your name, and the name of the file on the disk. Further information may be obtained from the Publisher.

Preparation of manuscripts

1. Manuscripts should be written in English. Authors whose native language is not English are strongly advised to have their manuscripts checked by an English speaking colleague prior to submitting. Authors in Japan please note: Upon request, Elsevier Science Japan will provide authors with a list of people who can check and improve the English of their paper (*before submission*). Please contact our Tokyo office: Elsevier Science Japan, 1-9-15 Higashi-Azabu, Minato-ku, Tokyo 106; Tel. (03)-5561-5032; Fax (03)-5561-5045.
2. Submit the original of your manuscript (*plus two copies omitting author names for double-blind refereeing process*). Enclose the original illustrations and two sets of photo-copies (three prints of any photographs).
3. Manuscripts should be typewritten, typed on one side of the paper, with wide margins and double spacing throughout, i.e. also for abstracts, footnotes and references. Every page of the manuscript, including the title page, references, tables, etc. should be numbered in the upper right-hand corner. However, in the text no reference should be made to page numbers; if necessary, one may refer to sections. Underline words that should be in italics, and do not underline any other words. Avoid excessive usage of italics to emphasize part of the text.
4. Manuscripts in general should be organized in the following order:

Title (should be clear, descriptive and not too long)

Name(s) of author(s)

Affiliation(s)

Present address(es) of author(s)

Complete correspondence address to which the proofs should be sent

Any (short) additional information concerning research grants, etc., may be included on the title page under the address(es). If this information is long, please include it in the text, either at the end of the introduction or in a separate acknowledgment section preceding the references.

Abstract

JEL Classification code(s)

Keywords (indexing terms), normally 3–6 items

Introduction

Material studied, area descriptions, methods, techniques

Results

Discussion

Conclusion

Acknowledgments

References

Tables

Figure captions

5. In typing the manuscript, titles and subtitles should not be run within the text. They should be typed on a separate line, without indentation. Use lower-case lettertype.
6. SI units should be used.
7. If a special instruction to the copy editor or typesetter is written on the copy it should be encircled. The typesetter will then know that the enclosed matter is not to be set in type. When a typewritten character may have more than one meaning (e.g., the lower case letter l may be confused with the numeral 1), a note should be inserted in a circle in the margin to make the meaning clear to the typesetter. If Greek letters or uncommon symbols are used in the manuscript, they should be written very clearly, and if necessary a note such as "Greek lowercase chi" should be put in the margin and encircled.
8. Elsevier reserves the privilege of returning to the author for revision accepted manuscripts and illustrations which are not in the proper form given in this guide.

Abstracts

The abstract should be clear, descriptive and not longer than 400 words.

JEL Classification

At least one classification code according to the classification system for Journal Articles as used by the Journal of Economic Literature should be supplied.

Tables

1. Authors should take notice of the limitations set by the size and lay-out of the journal. Large tables should be avoided. Reversing columns and rows will often reduce the dimensions of a table.
2. If many data are to be presented, an attempt should be made to divide them over two or more tables.
3. Drawn tables, from which blocks need to be made, should not be folded.
4. Tables should be numbered according to their sequence in the text. The text should include references to all tables.
5. Each table should be typewritten on a separate page of the manuscript. Tables should never be included in the text.
6. Each table should have a brief and self-explanatory title.
7. Column headings should be brief, but sufficiently explanatory. Standard abbreviations of units of measurement should be added between parentheses.
8. Vertical lines should not be used to separate columns. Leave some extra space between the columns instead.
9. Any explanation essential to the understanding of the table should be given as a footnote at the bottom of the table.

Illustrations

1. All illustrations (line drawings and photographs) should be submitted separately, unmounted and not folded.
2. Illustrations should be numbered according to their sequence in the text. References should be made in the text to each illustration.

3. Each illustration should be identified on the reverse side (or – in the case of line drawings – on the lower front side) by its number and the name of the author. An indication of the top of the illustrations is required in photographs of profiles, thin sections, and other cases where doubt can arise.
4. Illustrations should be designed with the format of the page of the journal in mind. Illustrations should be of such a size as to allow a reduction of 50%.
5. Lettering should be in Indian ink or by printed labels. Make sure that the size of the lettering is big enough to allow a reduction of 50% without becoming illegible. The lettering should be in English. Use the same kind of lettering throughout and follow the style of the journal.
6. If a scale should be given, use bar scales on all illustrations instead of numerical scales that must be changed with reduction.
7. Each illustration should have a caption. The captions to all illustrations should be typed on a separate sheet of the manuscript.
8. Explanations should be given in the typewritten legend. Drawn text in the illustrations should be kept to a minimum.
9. Photographs are only acceptable if they have good contrast and intensity. Sharp and glossy copies are required. Reproductions and photographs already printed cannot be accepted.
10. Colour illustrations cannot usually be included, unless the cost of their reproduction is paid for by the author.

References

1. All publications cited in the text should be presented in a list of references following the text of the manuscript. The manuscript should be carefully checked to ensure that the spelling of author's names and dates are exactly the same in the text as in the reference list.
2. In the text refer to the author's name (without initial) and year of publication, followed – if necessary – by a short reference to appropriate pages. Examples: "Since Peterson (1993) has

shown that..." "This is in agreement with results obtained later (Peterson and Kramer, 1994, pp. 12–16)".

3. If reference is made in the text to a publication written by more than two authors the name of the first author should be used followed by "et al.". This indication, however, should never be used in the list of references. In this list names of first author and co-authors should be mentioned.
4. References cited together in the text should be arranged chronologically. The list of references should be arranged alphabetically on authors' names, and chronologically per author. If an author's name in the list is also mentioned with co-authors the following order should be used: publications of the single author, arranged according to publication dates – publications of the same author with one co-author – publications of the author with more than one co-author. Publications by the same author(s) in the same year should be listed as 1974a, 1974b, etc.
5. Use the following system for arranging your references:

a. *For periodicals*

Arnade, C.A., 1994. Testing two trade models in Latin American agriculture. *Agric. Econ.* 10, 49–159.

b. *For books*

Polopolus, L.C., Alvarez, J., 1991. Marketing sugar and other sweeteners. *Developments in Agricultural Economics*, 9. Elsevier, Amsterdam, 361 pp.

c. *For edited symposia*

Edwards, F., Spawton, T., 1991. Pricing in the Australian wine industry: a marketing perspective. In: Botos, E.P. (Ed.), *Vine and Wine Economy. Proceedings of an International Symposium*, 25–29 June 1990, Kecskemét, Hungary. *Developments in Agricultural Economics*, 8. Elsevier, Amsterdam, pp. 203–212.

d. *For multi-author books*

Grinnell, G.E., 1992. Economics of energy in Agriculture. In: Fluck, R.C. (Ed.), *Energy in Farm Production. Energy in World Agriculture*, 6. Elsevier, Amsterdam, pp. 33–46.

e. *For unpublished reports, departmental notes, etc.*

Setzinger, A.H., Paarlberg, P.L., 1990. The export enhancement program: How has it affected wheat exports? *Agric. Inf. Bull.* 575, U.S. Department of Agriculture, Washington, DC.

6. Do not abbreviate the titles of periodicals mentioned in the list of references; alternatively use the International List of Periodical Title Word Abbreviations.
7. In the case of publications in any language other than English, the original title is to be retained. However, the titles of publications in non-Latin alphabets should be transliterated, and a notation such as "(in Russian)" or "(in Greek with English abstract)" should be added.
8. In referring to a personal communication the two words are followed by the year, e.g., "(J. McNary, personal communication, 1984)".

Formulae

1. Formulae should be typewritten, if possible. Leave ample space around the formulae.
2. Subscripts and superscripts should be clear.
3. Greek letters and other non-Latin or handwritten symbols should be explained in the margin where they are first used. Take special care to show clearly the difference between zero (0) and the letter O, and between one (1) and the letter I.
4. Give the meaning of all symbols immediately after the equation in which they are first used.
5. For simple fractions use the solidus (/) instead of a horizontal line, e.g. $I_p/2_m$ rather than $\frac{I_p}{2_m}$.
6. Equations should be numbered serially at the right-hand side in parentheses. In general only equations explicitly referred to in the text need be numbered.
7. The use of fractional powers instead of root signs is recommended. Also powers of e are often more conveniently denoted by exp.
8. Levels of statistical significance which can be mentioned without further explanation are $*P < 0.05$, $**P < 0.01$ and $***P < 0.001$.
9. In chemical formulae, valence of ions should be given as, e.g., Ca^{2+} and CO_3^{2-} , not as Ca^{++} or CO_3^{--} .

10. Isotope numbers should precede the symbols, e.g., ^{18}O .
11. The repeated writing of chemical formulae in the text is to be avoided where reasonably possible; instead, the name of the compound should be given in full. Exceptions may be made in the case of a very long name occurring very frequently or in the case of a compound being described as the end product of a gravimetric determination (e.g., phosphate as P_2O_5).

Footnotes

1. Footnotes should only be used if absolutely essential. In most cases it will be possible to incorporate the information in normal text.
2. If used, they should be numbered in the text, indicated by superscript numbers, and kept as short as possible.

Nomenclature

1. Authors and editors are, by general agreement, obliged to accept the rules governing biological nomenclature, as laid down in the *International Code of Botanical Nomenclature*, the *International Code of Nomenclature of Bacteria*, and the *International Code of Zoological Nomenclature*.
2. All biotica (crops, plants, insects, birds, mammals, etc.) should be identified by their scientific names when the English term is first used, with the exception of common domestic animals.
3. All biocides and other organic compounds must be identified by their Geneva names when first used in the text. Active ingredients of all formulations should be likewise identified.
4. For chemical nomenclature, the conventions of the *International Union of Pure and Applied Chemistry* and the official recommendations of the *IUPAC-IUB Combined Commission on Biochemical Nomenclature* should be followed.

Copyright

1. An author, when quoting from someone else's work or when considering reproducing an illus-

tration or table from a book or journal article, should make sure that he is not infringing a copyright.

2. Although in general an author may quote from other published works, he should obtain permission from the holder of the copyright if he wishes to make substantial extracts or to reproduce tables, plates, or other illustrations. If the copyright-holder is not the author of the quoted or reproduced material, it is recommended that the permission of the author should also be sought.
3. Material in unpublished letters and manuscripts is also protected and must not be published unless permission has been obtained.
4. A suitable acknowledgement of any borrowed material must always be made.

Proofs

1. Copy editing of manuscripts is performed by the staff of Elsevier. The author is asked to check the galley proofs for typographical errors and to answer queries from the copy editor.
2. Elsevier, at its discretion, is entitled to recover from the author of any paper or report published in the journal, any cost occasioned by alterations made by the author in the printer's proofs other than correction of typesetting errors and essential additions which update information in the paper; the latter preferably as sentences at the end of existent paragraphs or as new paragraphs.

Offprints

1. Fifty offprints will be supplied free of charge.
2. Additional offprints can be ordered on an offprint order form, which is included with the proofs.
3. UNESCO coupons are acceptable in payment of extra offprints.

AGRICULTURAL ECONOMICS HAS NO PAGE CHARGES
--