



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

**An Econometric Decomposition of Aggregate Data:
Application to the EU-15 Average Cattle Slaughter Weight**

Jacinto F. Fabiosa

Center for Agricultural and Rural Development
Iowa State University
Ames, Iowa

V. Premakumar

WEFA
Eddystone, PA

**CARD Working Paper 98-WP 205
November 1998.**



ABSTRACT

The level of aggregation of reported data has limited the proper accounting of live animals in most, if not all, livestock models, making the consistency of their underlying biologics suspect. We develop an econometric methodology to decompose a wide range of aggregate data. This method is applied to estimate a disaggregated average cattle slaughter weight for calves, cows, and other cattle from total beef production data in the European Union. The estimated model gave acceptable statistical properties and estimated slaughter weights that closely correspond to reported sample-based slaughter weights across all cattle categories. The model has a good fit, no serial correlation, and highly significant coefficients with consistent signs. Standard validation statistics suggest that the model is able to track well both the mean and variability of the actual average cattle slaughter weight. Furthermore, other decompositions using the same methodology were successful in other countries—New Zealand, and Mexico; other sectors—sheep; and other aggregate data—dairy and beef calving rates.

AN ECONOMETRIC DECOMPOSITION OF AGGREGATE DATA: APPLICATION TO THE EU-15 AVERAGE CATTLE SLAUGHTER WEIGHT

Most, if not all, models used in generating long-term projections of the world livestock sector do not have a complete and proper accounting of the stock and flow of live animals that ensure consistency in the underlying biologics of the model. Ideally, only the flow of live animals from either a technical relationship (e.g., death and calf crop) or decision of producers (e.g., slaughter number, slaughter weight, and breeding herd addition) should have a behavioral functional specification in a model, and the live animal stock should only be derived through an accounting identity.

The EU-15 CPPA model (USDA) includes only the meat side with no specification of any live animal inventory number. The EU-15 AgLink model (OECD), on the other hand, includes a specification of live animal inventory number as a behavioral equation, but accounting of live animal flow such as calf-crop, slaughter, and death is absent. The EU-15 FAPRI model (FAPRI, 1997) comes close to including specifications for live animal inventory and live animal number flow.

The primary reason for the absence of a clear accounting of live animal stocks and live animal flows in many models is dictated by what data are available. In the case of livestock data, both USDA's PS&D and OECD's database report aggregated variables that do not easily detail the number of live animals. This is glaringly true for cattle. For example, cattle death is reported as a total number with no breakdown of death by animal category. Also, although the cow inventory is divided into dairy and beef cows, the calves born is not similarly disaggregated. In some cases, the number of cattle slaughtered is not divided into categories such as, cow, calf, or other cattle. Moreover, it is not easy to derive a meaningful average cattle slaughter weight from aggregate production data.

There are a number of compelling reasons why proper accounting of live animal stock and live animal flow is necessary in modeling the livestock sector. First and, this is

the only way that the underlying biologics in the model can be consistent. Also there are technical and policy impacts specific to particular animal categories that can only be captured when the specification allows a disaggregated accounting of live animal stocks and flows. Good examples are the recent foot-and-mouth disease (FMD) outbreak in Taiwan and the classical swine fever (CSF) outbreak in the European Union (EU). These outbreaks may have differential impacts on piglets, sows, and other swine. If such detail of differential impact is reported, as in the CSF in the EU-15, then it is easy to incorporate that impact in a model where live animal stocks and live animal flows are properly accounted.

Although our econometric decomposition is general enough to be useful in other types of aggregate data decomposition, the focus of our application is the derivation of average cattle slaughter weight by animal category (i.e., calf, cow, and other cattle) from aggregate production data. Derivation of average cattle slaughter weight by simply dividing total beef production with total number of cattle slaughtered is not very meaningful for behavioral specification. When beef production comes from three different animal categories with different average weights, then an aggregate slaughter weight is not only a function of beef-feed price ratio and technology but also the relative proportions of the categories of animals slaughtered. That is, holding all else constant, average slaughter weight will decline if more calves are slaughtered relative to cows and other cattle. If there are technical and policy impacts that change this mix of categories of animals slaughtered, then a simple average slaughter weight equation is suspect. In this case, it is necessary to derive a representation of the average slaughter weight by category of animal. Although not available from reported data, the methodology developed here derives them from the reported aggregate production and slaughter number data.

This method is applied in the EU-15 because there are policy measures in its beef-cattle sector that give a classic example of a policy-induced need for a derivation of average cattle slaughter weight by animal category for a detailed accounting of live animal stock and flows. In response to the 1996 Bovine Spongiform Encephalopathy (BSE),

policies were introduced to reduce beef supply in the market as a strategy to support prices. These policies have animal category specific intended impacts.¹ For example, two specific policies were targeted to reduce availability of calves for beef production as well as reduce slaughter weight. The first is the calf-processing scheme that provides a premium of 120 European currency units (ECUs) for 20-day old male dairy calves and 150 ECUs for beef calves. The second is the early marketing premium scheme, which awards a basic premium of 50 ECUs for veal calves slaughtered at a weight 15 percent below the 1995 national average. These schemes are expected to reduce beef and veal production by 200,000 metric tons per year. With these schemes, the number of calves slaughtered may increase, causing the average slaughter weight to decline. Without a slaughter weight specification that is disaggregated into calves, cows, and other cattle, the impacts of these policies cannot be captured accurately in the model.

Model

The proposed econometric decomposition exploits the fact that even if the data are highly aggregated and if the variability in the final explanatory variables still allows identification of the parameters uniquely associated with each of its underlying components, then an econometric decomposition is possible. Although the same approach can be applied to similar aggregate data such as cattle death and calf-crop, the application developed here is on the average cattle slaughter weight by category of cattle.

If total beef production is given and there are three categories of cattle slaughtered – calf, cows, and other cattle – deriving an average slaughter weight for each animal type is difficult, but not impossible. The following econometric decomposition can be applied. Consider the identity, where total beef production is equal to the sum of its underlying components including production from calves, cows, and other cattle, i.e.,

$$BVSPR_t = CCKWV_t.CCKCV_t + CCKWB_t.CCKBR_t + CCKWC_t.CCKCO_t, \quad (1)$$

¹ An Over 30 Months Scheme (OTMS) is a compensation scheme that removes cattle aged more than 30

where $BVSPR$ is total beef production, $CCKWV$, $CCKWB$, and $CCKWC$ is the average slaughter weight for calves, cows (i.e., breeding herd), and other cattle, respectively. The number slaughtered for each animal category is $CCKCV$, $CCKBR$, and $CCKCO$. Although slaughter weight data are not available, we represent the unreported average slaughter weight for calves slaughtered as,

$$[2] \quad CCKWV_t = \alpha_0 + \alpha_1 \left(\frac{PB_t}{PF_t} \right) + \alpha_2 T + \varepsilon_{1t},$$

where PB is the price of beef, PF is the price of feed, T is a trend to capture technology, ε is an independently, and identically distributed (iid) error process, and α is a vector of parameters.² A similar specification can be given for the unreported slaughter weight for cows and other cattle slaughtered. Substituting the respective slaughter weight equations in [1] we get

$$[3] \quad BVSPR_t = (\alpha_0 + \alpha_1 \left(\frac{PB_t}{PF_t} \right) + \alpha_2 T + \varepsilon_{1t}).CCKCV_t + \\ (\beta_0 + \beta_1 \left(\frac{PB_t}{PF_t} \right) + \beta_2 T + \varepsilon_{2t}).CCKBR_t + (\delta_0 + \delta_1 \left(\frac{PB_t}{PF_t} \right) + \delta_2 T + \varepsilon_{3t}).CCKCO_t$$

Although the parameters of the unobserved slaughter weight equations (i.e., the vector of parameters α , β , and δ) can be estimated using only reported data (i.e., production and slaughter number) from equation [3], the properties of the parameter estimates are suspect because of the error structure in [3].

Diving both sides by the total number of cattle slaughtered
($CCKTN=CCKV+CCKBR+CCKO$),

months of age from the human and animal food chain in the UK.

² A vector of other regressors can be included in [2].

$$[4] \quad CCKW_t = (\alpha_0 + \alpha_1 \left(\frac{PB_t}{PF_t} \right) + \alpha_2 T + \varepsilon_{1t}) \cdot KCVS_t +$$

$$(\beta_0 + \beta_1 \left(\frac{PB_t}{PF_t} \right) + \beta_2 T + \varepsilon_{2t}) \cdot KBRS_t + (\delta_0 + \delta_1 \left(\frac{PB_t}{PF_t} \right) + \delta_2 T + \varepsilon_{3t}) \cdot KCOS_t + \varepsilon_t,$$

where $CCKW$ is the average slaughter weight, $KCVS$, $KBRS$, and $KCOS$, are the share of calf, cow, and other cattle slaughtered in the total slaughter. Equation [4] expresses average slaughter weight as a function of beef-feed price ratio, technology, and the mix of cattle slaughtered with the parameters of the cattle slaughter weight by cattle category.

Equation [4] has no intercept and has an error structure of

$$[5] \quad \varepsilon_t = \varepsilon_{1t} \cdot KCVS_t + \varepsilon_{2t} \cdot KBRS_t + \varepsilon_{3t} \cdot KCOS_t .$$

With an additional assumption that the error processes across the average weight equation for each category of animal are identical, equation [4] is further reformulated to derive a good estimating equation. Imposing this assumption and re-arranging terms,

$$[6] \quad CCKW_t = \alpha_0 + (\beta_0 - \alpha_0) \cdot KBRS_t + (\delta_0 - \alpha_0) \cdot KCOS_t +$$

$$\alpha_1 \cdot \frac{PB_t}{PF_t} + (\beta_1 - \alpha_1) \cdot \frac{PB_t}{PF_t} \cdot KBRS_t + (\delta_1 - \alpha_1) \cdot \frac{PB_t}{PF_t} \cdot KCOS_t +$$

$$\alpha_2 \cdot T + (\beta_2 - \alpha_2) \cdot T \cdot KBRS_t + (\delta_2 - \alpha_2) \cdot T \cdot KCOS_t + \varepsilon_t' .$$

Equation [6] has several acceptable properties. First, it can be estimated as a linear equation. The multiplicative explanatory variables can be defined as a new variable (i.e.,

$$\frac{PB_t}{PF_t} = \frac{PB_t}{PF_t} \cdot KBRS_t)$$

and the coefficients can be estimated as an aggregate with the

individual coefficients derived after estimation (i.e., $\gamma_1 = (\beta_1 - \alpha_1)$ and then

$\beta_1 = (\gamma_1 + \alpha_1)$.³ Of more importance, however, the error term is well-behaved, ensuring

unbiased and consistent estimates. That is, if $\varepsilon_{1t} = \varepsilon_{2t} = \varepsilon_{3t}$, then

$$\varepsilon_t = \varepsilon_{1t} \cdot KCVS_t + \varepsilon_{2t} \cdot KBRS_t + \varepsilon_{3t} \cdot KCOS_t = \varepsilon_t' = \varepsilon_{it} \text{ for } i=1,2,3.$$

³ The coefficients can also be estimated directly using a nonlinear estimator.

The approach outlined in this section can be applied to decompose other highly aggregated data.

Data and Results

Meat production and live animal inventory were taken from USDA's PS&D database, while prices were taken from OECD's database. In the case of the EU-15, although a longer series was available, estimation used only the period from 1980 to 1994. In 1995, Austria, Finland, and Sweden joined the EU. Their data cannot be included easily because the EU-12 cattle sector was heavily influenced by the milk quota policy, where producers were penalized with a super-levy for producing above the quota, which was not binding to these new members.

Mean values of some key variables are given in Table 1. The average slaughter weight was 265.80 kilograms per head carcass weight. The slaughter rates (number slaughtered relative to animal inventory) for each animal category are very close, 23.57 percent for calves, 25.63 percent for cows, and 28.51 percent for other cattle. Slaughter of other cattle account for close to half of total slaughter at 47.69 percent, followed by cow slaughter share at 28.51 percent, and calf slaughter share at 23.80 percent.

The average slaughter weight model has very acceptable statistical properties. Table 2 shows that the model has a good fit, with an R^2 at 0.974 and adjusted R^2 at 0.964. Although the DW-statistic of 1.593 is in the inconclusive range, with a critical value at 1 percent significance level at $d_l=0.488$ and $d_u=1.704$, the DW-statistic is more towards the upper limit that would accept the hypothesis of no serial correlation. Except for the coefficient associated with calf weight, all the parameters are highly significant. Of more importance, the signs of the coefficients are also consistent with expectations; that is, beef price has a positive impact on the slaughter weight (elasticity of 0.0653), while otherwise for feed price.

Moreover, the methodology successfully derived and estimated slaughter weights for calves, cows, and other cattle from total beef production data. To validate the

methodology, Table 3 compares the estimated average slaughter weight by animal category with some reported weights from sample information. Several countries report maximum slaughter weight for calves in the EU's early marketing premium scheme. The average for France, Germany, and Italy of 109 kilograms per head is very close to the estimated average slaughter weight for calves of 108.⁴ The reported slaughter weight for cows and other cattle is taken from the Cattle Grading Scheme of some EU members. For cow slaughter weight, the average for Belgium, Denmark, Germany, and Italy is 228 kilograms per head, which is very close to the estimated average cow slaughter weight of 233 kilograms per head. There is more variability of slaughter weight in the other cattle category. The range for Germany is 319 to 432, or an average of 376 kilograms per head, which is close to the estimated average of 370 kilograms per head.

Standard validation statistics of the average cattle slaughter weight model are given in Table 3. Both the mean and standard deviation of the actual and predicted series are very close. This good fit is similarly shown in the low percent mean error, 0.004 percent. The correlation of the actual and predicted series is very high at 0.987. The decomposition of the MSE suggests that almost all of the prediction error is due to random process and none of it is due to bias or other structural error. Moreover, the model is able to track well not only the mean of cattle slaughter weight but also its variability. Theil statistics approach zero, suggesting a very good fit.

Further validations were performed with the decomposition of other aggregate data. A disaggregated average cattle slaughter weight was estimated for New Zealand and Mexico. Beef production in both countries are grass-fed in their cattle production system. The estimated average weights shown in Table 1 are very reasonable. For example, an attaché report gives the export beef slaughter weight New Zealand in 1996 at 253 kilograms per head, which is very close to the estimated average slaughter weight for other cattle in New Zealand at 255 kilograms per head (USDA, AGR #NZ7006).

⁴ France, Germany, and Italy represent 52 percent of total EU cattle inventory.

Moreover, the same approach was applied in other animals—in New Zealand’s sheep sector, and in the EU’s dairy and beef calf-crop—with similar success.⁵

Conclusion

The general issue addressed in this paper concerns the lack of proper and complete accounting of live animal stock and flow in most, if not all, livestock models used in generating long-term projections for the world. The common problem is the high level of aggregation in available data that do not lend easily to detailed accounting.

An econometric methodology was developed to decompose aggregate data and was applied to derive and estimate a behavioral specification of average cattle slaughter weight by cattle category in the EU cattle sector. The model gave acceptable statistical properties: good fit, no serial correlation, highly significant coefficients, consistent signs of coefficients that give estimated average slaughter weight close to reported slaughter weight across all cattle categories. Standard validation statistics suggest that the model is able to track well both the mean and variability of actual slaughter weight series. Furthermore, other validations were successfully applied to New Zealand and Mexico, to New Zealand’s sheep sector, and to other aggregate data for EU’s dairy and beef calving rates.

⁵ Full results of additional validations can be obtained from the authors.

Table 1. Means of variables and elasticity

Variables	Mean
Mean of Variables	
Slaughter Weight (kg/head carcass)	265.80
Calf Slaughter Rate (%)	23.57
Cow Slaughter Rate (%)	25.63
Other Cattle Slaughter Rate (%)	28.51
Proportion of Calves Slaughtered (%)	23.80
Proportion of Cows Slaughtered (%)	28.51
Proportion of Other Cattle Slaughtered (%)	47.69
Elasticity of Slaughter Weight to Price	0.0653

Table 2. Average slaughter weight for cattle in the EU-15, New Zealand, and Mexico

Parameter	Independent	EU-15		N. Zealand		Mexico	
		Coef	S. Dev	Coef	S. Dev	Coef	S. Dev
α_0	Calf Weight	0.108	0.088	0.0764	0.0301	0.0750	
β_0	Cow Weight	0.233	0.048	0.1679	0.0808	0.1479	0.0568
δ_0	Intercept	0.245	0.021	0.1718	0.0869	0.1219	0.0718
δ_1	Trend	0.004	0.001	0.0024	0.0017	0.0039	0.0014
δ_2	Price Ratio	0.018	0.008	0.0001	0.0001	0.0001	0.0001
Diagnostic							
	R^2	0.974		0.832		0.794	
	Adj R^2	0.964		0.756		0.701	
	DW	1.593		1.591		1.115	

Table 3. Estimated and reported slaughter weights (from sample information)

	Model Estimate	Reports
Average Calf Slaughter Weight	108	109
France (52%)		108
Germany		103
Italy		117
Average Cow Slaughter Weight	233	228
Belgium		255
Denmark		200
Germany		240
Italy		216
Average Other Cattle Slaughter Weight	370	376
Germany		319 - 432

Table 4. Cattle average slaughter weight model validation

	Actual		Predicted	
	Mean	Std Dev	Mean	Std Dev
Slaughter Weight	0.2556	0.0114	0.2556	0.0113
Statistics of Fit				
Mean Error			0.0000	
Percent Mean Error			0.0044	
Mean Absolute Error			0.0015	
Mean Absolute Percent Error			0.5682	
Root Mean Square Error			0.0018	
Root Mean Square Percent Error			0.6900	
R-square			0.9742	
Theil Forecast Error Statistics				
Mean Square Error			3.1E-06	
Correlation			0.9870	
Bias			0.0000	
Regression			0.0000	
Disturbance			1.0000	
Variance			0.0070	
Co-variance			0.9930	
Theil U1			0.0069	
Theil U			0.0035	

REFERENCES

Food and Agricultural Policy Research Institute (FAPRI). FAPRI Integrated Livestock and Poultry Model. 1997.

Meat and Livestock Council. *European Handbook: The Common Agricultural Policy*. Volume I. Paris, France, 1997.

OECD. Agriculture Database, (electronic-disk), 1997, Paris, France.

SAS Institute Inc., SAS/ETS User's Guide, Version 6, Second Edition, Cary, NC: SAS Institute Inc., 1993.

U.S. Department of Agriculture, Economic Research Service. PS&D Database, (electronic), June 23, 1997.
<<http://mann77.mannlib.cornell.edu/data-sets/international/93002/>>

U.S. Department of Agriculture. Foreign Agriculture Service (FAS). New Zealand Livestock Semi-Annual Report, AGR Number NZ7006. 1997.

Filename: WP_JAY2.doc
Directory: C:\My documents\Downloaded Papers
Template: C:\Program Files\Microsoft Office\Templates\Normal.dot
Title: An Econometric Decomposition of Aggregate Data:
Subject:
Author: Valued Gateway Customer
Keywords:
Comments:
Creation Date: 10/20/98 10:36 AM
Change Number: 3
Last Saved On: 12/09/98 11:46 AM
Last Saved By: Valued Gateway Customer
Total Editing Time: 31 Minutes
Last Printed On: 03/18/99 2:10 PM
As of Last Complete Printing
Number of Pages: 13
Number of Words: 2,684 (approx.)
Number of Characters: 15,304 (approx.)