



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

**Achieving Efficiency and Equity in Irrigation Management:
An Optimization Model of the El Angel Watershed, Carchi, Ecuador©**

Elizabeth M. Evans, David R. Lee, Richard N. Boisvert

Department of Applied Economics and Management,
Cornell University, Ithaca, NY 14853, USA

Blanca Arce

Department of Animal Science
Cornell University, Ithaca, NY 14853, USA

&

Tammo S. Steenhuis

Department of Biological and Environmental Engineering
Cornell University, Ithaca, NY 14853, USA

ABSTRACT

The objective of this paper is to address the problems of inefficiency and inequity in water allocation in the El Angel watershed, located in Ecuador's Sierra region. Water is captured in a high-altitude region of the watershed and distributed downstream to producers in four elevation-defined zones via a system of canals. Upstream and downstream producers face radically different conditions with respect to climate and terrain. A mathematical programming model was created to study the consequences of addressing chronic water scarcity problems in the watershed by shifting water resources between the four zones. The model captures the nature of water use by humans, crops and dual purpose cattle. Its objective function maximizes producer welfare as measured by aggregate gross margin, subject to limited supplies of land, labor and water. Five water allocation scenarios are evaluated with respect to efficiency in land and water use and equity in income distribution. Results reveal that although water is the primary constrained resource downstream, in the upstream zones, land is far more scarce. The current distribution of water rights does not consider these differences and therefore is neither efficient nor equitable. Improvements in efficiency (resource use) and equity (income distribution) are associated with (1) a shift of water to the lower zone, and (2) the use of lower levels of irrigation intensity upstream. Furthermore, the scenarios that result in the most efficient use of resources also bring the greatest degree of equity in income distribution, indicating that these may be complementary, not conflicting, goals.

KEYWORDS: Mathematical programming, water allocation, efficiency, equity.

©Copyright 2002 by Nelson Villoria and David R. Lee. All rights reserved. Corresponding author: Prof. David R. Lee, DRL5@cornell.edu

INTRODUCTION

Agriculture demands more water than any other single activity, requiring 69 percent of the world's water supply (Holden and Thobani, 1996). In many countries, efforts to raise levels of agricultural production through increases in cultivated land, cropping intensity and yields have led to a greater dependence on irrigation. This pressure has been most severe in developing nations, where water resources are often scarce and many irrigation systems are primitive.

Although Ecuador has estimated water resources of 38,372 cubic meters per capita, nearly four times the worldwide average of 10,800 cubic meters per capita, the distribution of water is uneven and is heavily concentrated in the country's Amazon and Coastal regions (Sotomayor, 1996). It is estimated that 95 percent of the country's irrigation structures utilize conventional canal technologies (Sotomayor, 1996). The upstream-downstream asymmetry in water availability and low levels of irrigation technology lead to an inequitable and inefficient allocation of water among users. Absent a strong institutional system of water management, there are few incentives for upstream producers to use water efficiently, which leads to uncertainty for downstream producers, and in many cases, conflict. This study examines the effects of alternative management scenarios aimed at improving efficiency and equity in water allocation. Although focused specifically on Ecuador's Sierra region, the study is illustrative of water management problems faced widely in the developing world.

The canals of the El Angel watershed in the highlands of Carchi, Ecuador were constructed over one hundred years ago to provide water to the several large haciendas which then dominated the 100,000-hectare region. Today, these canals transport water a distance of over 60 kilometers, from altitudes of more than 4,000 meters above sea level (masl), to hotter, drier regions as low as 1,500 masl. The primary source of water is the *páramo*, an area of humid alpine grasslands in the upper reaches of the watershed whose sponge-like terrain is well-suited to the capture, storage and transfer of water. The five canals that

are the focus of this study are largely non-reinforced, open structures, highly susceptible to losses from evaporation, spillage, seepage and theft.¹

The five canals are located in the eastern portion, or left margin of the watershed. As demonstrated by Prato and Wu (1996), the “target-level” of a resource management policy plays a significant role in determining its environmental and economic costs and benefits. Brooks et al. (1994) argue for policymaking at the watershed level, noting that most natural processes occur within watershed boundaries. Beaulieu et al. (1998), however, caution that most farmers make decisions at the field level, and do not consider the watershed-level effects of their decisions. Furthermore, policymaking often comes down to a tradeoff between economic and environmental objectives due to the fact that watershed boundaries rarely coincide with political boundaries (Brooks et al., 1994; Okumu et al., 1999).

The study region is divided according to elevation into three primary zones: upper, middle and lower; the middle zone is further divided to account for the diversity in production systems found in the region.² As one travels downstream, temperatures, levels of solar radiation and rates of evapotranspiration rise, and soils are increasingly sandy and susceptible to erosion. While the upper zone is oriented primarily toward pasture (cattle) and potato production, a more diverse group of crops can be produced in the middle and lower zones. However, the degree of dependence on irrigation in these downstream zones is much greater. Table 1 summarizes key characteristics of each zone.

In the midst of Ecuador’s agrarian reform, water rights or “concessions” were divided among new landowners along the canals. Although a number of regional water agencies were established to enforce the nation’s 1972 Water Law, in reality, minimal control is exercised by the state, and in most cases, management occurs at the local level (Proaño, 2000). Community members form groups called *juntas* to enforce concessions and carry out maintenance tasks for a particular canal. The effectiveness of the *juntas* varies widely. Most failures can be attributed to a lack of resources, poor communication and farmers’ unwillingness to participate.

¹ The five canals studied are: 1) San Vicente de Pusir de los Baños, 2) San Vicente de Pusir del Voladero, 3) Cúnquer, 4) Chulunguaza, and 5) El Tambo.

² The four zones are referred to as “upper,” “middle 1,” “middle 2,” and “lower” in this paper.

Without effective enforcement of water rights, many producers base water use decisions on availability rather than actual crop water requirements, and as a result, regularly use canal water beyond their legal allocations. Downstream farmers are at a natural disadvantage, their access being limited to that portion of canal water that has not been used upstream. These circumstances have led to a number of water-related conflicts between rural and urban communities, between producers at different altitudes and between small- and large-scale producers (Poats et al., 1998).

The objective of this paper is to address the problems of inefficiency and inequity in water allocation in the El Angel watershed. A comprehensive, crop-livestock mathematical programming model is described which allows the simulation of various water allocation scenarios. The current system of water management is then evaluated with respect to the achievement of efficiency and equity objectives. Various alternative water allocation strategies are examined. These alternative strategies are compared, revealing the consequences of shifting water resources between watershed zones. Finally, the relationships between efficiency and equity objectives are discussed.

AN OPTIMIZATION MODEL OF CROP AND LIVESTOCK PRODUCTION

The mathematical programming model is designed to maximize aggregate gross margin from agricultural production in the El Angel watershed. The model's activities and constraints characterize the nature of water use for humans, crops and livestock over a 12-month period. The technical coefficients that quantify resource requirements were estimated for a representative farm in each zone.³

Mathematical programming models seek to determine the optimum allocation of constrained resources among competing activities (Hazell and Norton, 1986), and are widely utilized in studies of resource allocation in agriculture, including water allocation. Combining mathematical programming and simulation modeling techniques, Bernardo et al. (1988) find opportunities for improved water

³ Characterization of the representative farms is based on data collected in a joint survey, data obtained from the International Potato Center (CIP) and a number of secondary sources. The survey, conducted from January – September, 2000, is a collaboration between the MANRECUR Project, Cornell University and Universidad Central de Ecuador. The CIP data is from the project “Agricultural Chemical Use and the Sustainability of Potato Production in the Andean Zone,” collected between April 1990 and December 1992 in a watershed directly adjacent to the El Angel watershed.

management in the Columbia River Basin, given that decreasing the water supply by 40 percent at the farm level results in only a 10 percent decrease in economic returns. Garg and Ali (1998) use a two-level mathematical programming model for the Lower Indus Basin in Pakistan to determine the optimal cropping pattern and sowing dates to efficiently use scarce water resources. Unlike these two studies, the focus of this model is not limited to water use for crops, nor to the analysis of resource use for livestock, as in Nicholson (1990) and Berger (1999). Rather, this study simultaneously incorporates both crop and livestock activities in evaluating water allocation.

Model objective function. The model's objective function maximizes aggregate gross margin (total revenue less variable costs) from crop and livestock production:

$$\begin{aligned} \text{MAX} \quad & \sum_{i=1}^4 \sum_{j=1}^J \sum_{k=1}^2 [(GMCROP_{ijk})(CROP_{ijk})] + \sum_{i=1}^2 \sum_{k=1}^2 \sum_{n=1}^{16} [(GMA_{ikn})(A_{ikn})] \\ & + \sum_{i=1}^2 \sum_{k=1}^2 [(RMILK_{ik})(MILK_{ik})] - \sum_{i=1}^2 \sum_{k=1}^2 [(SUPCOST_{ik})(SUP_{ik})] - \sum_{i=1}^4 \sum_{k=1}^2 [(WCOST_{ik})(WUSE_{ik})] \end{aligned} \quad (1)$$

where activities are defined for the i^{th} zone, the j^{th} crop, the n^{th} (of 16) herd category, and the k^{th} season (wet or dry) and where GMCROP = per-hectare gross margin; CROP = number of hectares of each crop activity; GMA = per-animal gross margin, calculated using the revenue from animal sales only; A = number of cattle in each herd category; RMILK = per-liter price of milk; MILK = number of liters of milk sold; SUPCOST = per-kilogram cost of animal feed supplements; SUP = number of kilograms fed; WCOST = per-cubic meter tax on water use; and WUSE = total use of canal water.

Model activities. Four categories of activities are included in the model: human, crop, animal and water (Table 2). Human activities are included to quantify non-agricultural water requirements in each zone, based on population estimates and water consumption data in Montenegro (1998).

Composite crop activities are defined over a 12-month period, and include all of the processes that are involved with the production of a crop, including soil preparation, fertilization, irrigation and harvest. Activities are defined for each season to account for variations in irrigation requirements due to lower levels of precipitation in the dry season. Certain crop activities are defined at multiple levels of irrigation intensity to account for the fact that farmers often exhibit profit-maximizing rather than yield-

maximizing behavior when allocating irrigation water among crops (Doyle, 1990; Santos, 1998). The exceptions are potatoes and improved pasture, for which reductions in irrigation showed little or no impact on yields.

The yield response to water is estimated for each crop using the relationship between relative yield (Y_a/Y_m) and relative evapotranspiration (ET_a/ET_p), established by Doorenbos and Kassam (1979):⁴

$$\frac{Y_a}{Y_m} = \prod_{i=1}^4 \left[1 - k_{y_i} \left(1 - \frac{ET_{a_i}}{ET_{p_i}} \right) \right] \quad (2)$$

where “i” denotes each period (of four) in the plant’s growth cycle; Y_a = actual yield; Y_m = maximum yield; k_{y_i} is the yield response factor for each stage i of the crop’s growth cycle; ET_{a_i} = total amount of actual evapotranspiration during period i; and ET_{p_i} = total amount of potential evapotranspiration during period i.

The objective of using this method was not to precisely calculate the relationship between water use and yields. This was not possible due to the high data requirements of econometric and simulation methods. Rather, as intended by Doorenbos and Kassam, the equation is used “to quantify the effect of water stress” on the studied crops in the watershed. The right-hand side of Equation 2 is a number between zero and one, often referred to as the crop’s water stress index (WSI). A WSI closer to one indicates a lower level of water stress.

ET_p data are taken from the comprehensive study by Montenegro (1998) of the use, management and distribution of irrigation and potable water in the El Angel watershed. These data were calculated using the FAO software CROPWAT.⁵ Based on average levels of precipitation in each zone as well as soil moisture observations, ET_a is calculated using the method described in Thornthwaite and Mather (1957) and later in Dunne and Leopold (1978). This method was chosen primarily due to the instrumentation and data limitations associated with working in the remote study area.

⁴ Previous applications of this method include Bernardo et al. (1988) and Paudyal and Das Gupta (1990).

⁵ CROPWAT version 5.7, issued in 1992 is a decision support system developed by the Land and Water Development Division of FAO. It is commonly used to calculate reference evapotranspiration, as well as

Doorenbos and Kassam define Y_m as “the harvested yield of a high-producing variety, well-adapted to the given growing environment, including the time available to reach maturity, under conditions where water, nutrients and pests and diseases do not limit yields.” Y_m values are most accurately estimated with the use of experimental data that account for the effects of a number of variables, including crop variety, temperature and length of the growing season. These experimental values must then be adjusted to according to actual farming conditions (Doorenbos and Kassam, 1979). Due to a lack of experimental field data, Y_m values used in this study were obtained from interviews with local farmers and from secondary sources such as Proaño and Paladines (1998) and Espinosa et al. (1996) that are specific to the region or to a comparable region. These maximum yield values compare conservatively with those presented by Doorenbos and Kassam for tropical climatic regions. To solve for yield levels for the various irrigation-defined crop activities, the difference between ET_{a_i} and ET_{p_i} during each period of a crop’s growth cycle is decreased by a percentage that represents the extent to which irrigation is used to fill any evapotranspiration deficit. Y_a is then calculated according to Equation 2.

In the case of dual purpose cattle production, activities are included for the upper zone and middle zone 1 only. Herd category and nutrient requirement activities are subdivided to account for differences in animal age, sex, weight, weight gain and activity. The model solves for the number of animals in each of the seven cattle age groups, which are further subdivided by sex and activity. Activities are also included to quantify herd size in animal units (AUs), used later in stocking rate restrictions. Based on the total herd size, the model calculates aggregate nutrient requirements,⁶ the amount of *canal* water required to supplement animal feeds to meet water requirements, the kilograms of feed supplements purchased to meet nutritional requirements, and levels of milk production.

Finally, the model solves for the levels of water use in each zone. Like all other activities, water use activities are included for the dry and wet seasons separately.

crop water and irrigation requirements.

⁶ Required quantities of dry matter, crude protein, metabolizable energy and water.

Model constraints. The model's constraints can be divided into five categories: human, crop, animal, land and labor, and water (Table 2). Human constraints are included to define the human population in each zone, in each season, which is fixed throughout the year.

Although a shift of resources to the production of high-valued crops such as potatoes and anise could result in significant income growth, such a dramatic change is unlikely in the short run, given the potential problems associated with resource bottlenecks, nutritional deficits and price variability. To be consistent with the short-run nature of the model and historical production patterns in the region, constraints are included for each zone to ensure minimum levels of crop diversity.

Similar to animal activities, animal constraints are driven by the nutritional requirements of the various herd categories. The first of six types of animal-related constraints divide the herd into age-, sex- and activity-based categories. These constraints establish relationships between the sizes of age groups, the numbers of males and females, the numbers of sale and replacement heifers and the numbers of lactating and dry cows. Herd size is converted to AUs in the second group of animal constraints. The third and fourth types of animal constraints respectively define aggregate herd nutrient requirements and ensure that these nutrient requirements are met. These constraints determine the levels of canal water and nutritional supplements consumed by the herd. The fifth set of constraints limits the herd size according to maximum stocking rates typical to the region. Finally, a set of milk constraints defines the liters of milk that are produced, consumed and sold.

Land and labor constraints limit the total supply of these resources in each zone in each season. The land supply consists of actual or potentially canal irrigable sectors, as identified by Montenegro (1998). Labor supply is limited based on estimates of the total population active in agriculture and the number of days worked per month, and is assumed to be non-transferable across zones.

The last category of constraints defines water constraints. Although efforts are currently underway to systematically measure water volume at multiple points along several of the canals, for the purposes of this analysis, it was impossible to identify how much water is taken from the *individual* canals in each zone. Thus, in the model, canal water is considered to be a global resource among the five

canals. Of the five canals, four begin in the *páramo*, above the upper zone, and one initiates at an altitude of 3,200 masl and is used only by residents of middle zone 2 and the lower zone. To allow for multiple intakes of water along the canals, the first set of constraints defines water use in each zone, with water use from each source defined separately. The second set limits total water use from each source to be less than or equal to the estimated supply.⁷ In addition, for certain allocation scenarios, constraints are included that limit water use from both sources at the zone-level. Water constraints are explained in greater detail in Evans (2001).

DEVELOPMENT OF MODEL TECHNICAL COEFFICIENTS

The study of resource allocation on a farm requires empirical specification of relationships between underlying biological and economic processes (Nicholson, 1990). This section describes the calculation of technical coefficients to quantify the resource requirements for humans,⁸ crops and animals. To account for differential resource availability by season, total requirements of a resource or input are disaggregated by wet and dry season when necessary.

Although farmers in the watershed often utilize a single hectare for the production of multiple crops, for ease of measurement, technical coefficients for crops are calculated on a per-hectare basis. Crop irrigation requirements, which account for the majority of demand for canal water, are calculated using the FAO Penman-Monteith method (Allen et al., 1998). The crop's total water requirement during month t (REQ_t) is calculated using the formula:

$$REQ_t = ET_{o_t} \times k_{c_t} \quad (3)$$

where ET_{o_t} is the total amount of reference evapotranspiration during month t and k_{c_t} is the crop coefficient corresponding to the appropriate month of crop growth. The formula for a crop's total irrigation requirement (IREQ) during its growth cycle is:

⁷ The estimated supply from each source is calculated by varying average canal volume (Proyecto DRI, 1991) according to assumed monthly fluctuations, and summing the monthly values over the dry and wet seasons. These fluctuations are calculated from weekly volume data collected by the local MANRECUR Project from November, 1999 to October, 2000 in the canal San Vicente de Pusir de los Baños.

⁸ Based on data reported in Montenegro (1998), a fixed quantity of 3.75 cubic meters of water per month is allocated per human for such uses as drinking, washing, cooking and watering small gardens. The net

$$IREQ = \sum_{t=1}^T \frac{REQ_t - EP_t}{IE} \quad (4)$$

where t is the total T months in the plant's growth cycle; REQ_t is the crop's water requirement during month t ; EP_t is the amount of effective precipitation during month t ; and IE is indicative of the level of inefficiency in the system of water distribution.

A number of sources of data were relied upon for the calculation of irrigation requirements. Precipitation data were obtained from Ecuador's National Center for Hydrologic Resources (CNRH)⁹ and from the International Potato Center.¹⁰ Reference evapotranspiration data were from Montenegro (1998). Effective precipitation was calculated using the method described in Brouwer and Heibloem (1986). Monthly k_c values specific to the region were obtained for most crops from Apollin and Eberhart (1998). Finally, factors reported in Montenegro (1998), reflecting the degree of efficiency in the capture, conduction and application of irrigation water in the watershed's system of canals, were used to determine gross irrigation requirements.

Data used in the derivation of crop enterprise budgets were for the most part collected by B. Arce during the course of fieldwork in the year 2000. Costs were broken down into six major categories: labor, equipment, seed, fertilizer, pest control and transportation. Levels of investment vary widely between the crops that are produced for commercial sale (potatoes, corn, anise), and those which are largely produced for home consumption (wheat, sweet potato, white carrot). Costs and yields are adjusted for crop activities where lower irrigation intensities are assumed (Table 3).¹¹

Enterprise budgets for dual purpose cattle production include the costs of labor, vaccines, vitamins and minerals that are directly attributed to each animal. Costs of producing improved pasture

amounts of human water use in each zone are then adjusted for the effects of conveyance inefficiencies.

⁹ As reported by Montenegro (1998) and collected from the meteorological station at El Angel.

¹⁰ Data are monthly measures of rainfall (mm) at four representative meteorological stations: Chalpatán (3,360 masl); El Angel (3,055 masl); Bolívar (2,640 masl); and San Vicente de Pusir (1,870 masl).

¹¹ Although not identical, this formulation is similar to a separable programming formulation to approximate diminishing marginal productivity of an input (Hazell and Norton, 1986).

are counted as a cost of crop production in the objective function. Costs are identified separately for lactating and non-lactating animals, primarily due to the differences in labor requirements.

The herd's energy, protein and water requirements determine the amount of pasture, concentrate and canal water needed for dual purpose cattle production. Utilizing assumptions regarding the quantity and quality of daily milk production per cow, as well as average weights and weight gains, the daily feed requirements of each type of cattle were determined.¹²

Both native and improved pasture are common in the upper zone; in middle zone 1, however, native pasture dominates. Although the use of improved pasture has been shown to allow gains in herd size and production per cow (Brockington et al., 1992), due to the costs of labor, seed and fertilizer, cultivation of improved pasture involves a greater investment for the producer. Because the nutritional content of both native and improved pasture varies between seasons and between years, average levels of nutritional content are calculated and used as technical coefficients in the model.¹³

Grain concentrates fed in the watershed generally consist of wheat bran and corn grain. In the model both are assumed to be available for purchase in unlimited quantities, and because they are manufactured, content is assumed to not vary significantly between seasons or years.

MODEL VALIDATION AND WATER ALLOCATION SCENARIOS

The process of model validation includes a comparison of the results of an Initial solution to historical production patterns. Generally, the similarity between the model's solution and historical land use is greatest in the upper zone, and decreases as one moves downstream. Based on historical data, the average percentage of cultivated land utilized for the representative crops in the left margin is greatest in the upper zone (83 percent), decreases in middle zone 1 (64 percent) and middle zone 2 (49 percent), and

¹² Equations for calculating the requirements of heifers and cows are taken primarily from Berger (1999); they are based largely on the published requirements of the National Research Council (1989). Requirements for steers are based on the National Research Council's (1996) requirements for beef cattle.

¹³ Average nutritional contents of native and improved pasture for each season over a period of three years are determined from data provided by Instituto Nacional Autónomo de Investigaciones Agropecuarias (INIAP).

is at its minimum in the lower zone (30 percent).¹⁴ This result is expected, given the greater diversity in production made possible by the higher temperatures and levels of solar radiation characteristic of downstream areas. Historical land use patterns for the entire watershed show that the percentage of cultivable land left fallow is also greatest in the lower zone (33 percent), which might partially account for the relatively low percentage of land sown in the representative crops (Proaño and Paladines, 1998).

The Initial solution, in which herd size is endogenous, includes no cattle activities. This result supports the contention by Nicholson et al. (1994), that in most Latin American countries, “highland areas typically provide greater net returns when dedicated to agricultural enterprises other than livestock” (p. 312). For the small and medium-sized producers who dominate the watershed, it is likely that cattle are raised for reasons other than the monetary income they generate. The dual purpose production system is considered to be especially suited to tropical regions, given its sustainability with respect to resource use and consistency with the demand for milk and meat (Restrepo et al., 1991). Furthermore, in comparison to crop production, dual purpose cattle production provides a steady income, which mitigates the risk inherent in variable crop prices and the effects of between-harvest periods when savings are low.

For these reasons, constraints are added to ensure minimum acceptable herd sizes for the upper zone and middle zone 1, where minimum herd size is based on cattle population estimates by Arce et al. (1993) and Proaño and Paladines (1998). The validated model is used in all subsequent scenarios for the study of water allocation.

To address the concerns of policymakers, five scenarios are modeled to reveal the effects of water allocation on the use of resources and the distribution of income among producers. First, in the Enforced Rights Allocation scenario, water is allocated from the canals strictly according to the pattern of ownership of water concessions between the four zones. This provides a starting point for policymakers, as it represents the possible consequences of enforcing the set of laws that currently govern water use.

Although this analysis reveals flaws in the current system, in reality, concessions are not enforced. Because much of the responsibility of water management is left to local residents, actual water

¹⁴ Average percentages of cultivated area in the representative crops in each zone from 1996-1999, based

use is driven by the laws of nature, rather than by the system of concessions. Thus, relative allocation is largely a function of location; upper zone farmers require less than their legal allocation, and the surplus water flows downstream. Middle zone producers are therefore in a position to use water beyond their legal allocation. The Actual Allocation scenario, designed to represent reality and used as a baseline of comparison for further analysis, portrays the distribution of water that results when illegal water use is allowed by middle zone producers.

Assuming the goal of policymakers goes beyond the identification of current system failures, it is useful to study alternatives to the Enforced Rights and Actual Allocation scenarios. Three additional scenarios are analyzed that specifically address the concepts of efficiency and equity. In the Efficient Allocation scenario, water is allocated by the basic economic criterion that the value of its marginal product is equal among all uses. This scenario is an efficiency “ceiling” for policymakers, representing the highest achievable level of aggregate income, given current resource availability and relative prices.

If policymakers instead seek to give some priority to equity considerations through resource (re)allocation, they must first clearly define their equity goal. In the Land-based Allocation scenario, equity is defined in terms of water allocation – an equal amount of water is allocated per hectare of land throughout the watershed. In contrast, when the goal is defined in terms of an equitable income distribution, a preferable standard for water allocation is that given by the Irrigation-based Allocation scenario, in which water is allocated in proportions that reflect the differences in average per-hectare irrigation requirements.¹⁵

MODEL RESULTS

Table 4 reports the activity levels for the five modeled scenarios. Water use in each zone is reported separately for the dry and wet seasons. Crop production activities are differentiated according to the levels of irrigation intensity. Herd size in animal units and the volume of milk produced in the upper

on survey data collected in 2000.

¹⁵ In both the Land-based and Irrigation-based Allocation scenarios, human water use is subtracted from the total water supply prior to establishing the proportions of water allocated for agricultural use in each zone.

zone and middle zone 1 are also presented. Table 5 displays total income and income distribution, represented by various measures, between the four zones in each case.

Key Results. The results of the Enforced Rights Allocation scenario demonstrate that if water were allocated strictly according to the legal concessions in the watershed, resource use and income distribution would be neither efficient nor equitable. Although irrigation requirements are relatively low in the upper zone, producers there own more than 71 percent of the total volume of assigned water.¹⁶ Water is used in a manner that maximizes agricultural income in the upper zone, until land restrictions become binding and significant surplus water results. If downstream farmers are restricted from using this excess water, more than 50 percent of the area's total water supply, and 40 percent of total land supply, go unused. Middle zone 1 producers respond by shifting resources to the production of dual purpose cattle fed by native pasture, activities that require much less water. The restrictions in resource use and resulting shifts to less profitable activities have a severe impact on income generated. As shown in Table 5, total income in the Enforced Rights Allocation scenario is limited, and its distribution is highly inequitable, heavily favoring upper zone producers.

From these results it appears that the illegal use of water by middle zone farmers is in fact consistent with the goals of efficiency and equity. The Actual Allocation scenario, which relaxes the restrictions on middle zone water use, clarifies the importance of location along the canals to water use and income distribution. Although lower zone farmers receive their full legal allocation (approximately 25 percent of total supply in both seasons), they earn only 11 percent of aggregate income (Tables 4 and 5). Despite the fact that the majority of water is used in middle zone 2 and the lower zone, it is not enough to offset the limitations imposed on downstream producers by differences in per hectare irrigation requirements. Both income per capita and income per hectare decrease as one travels downstream.

One alternative to the current system of water management is to allocate water according to a standard of economic efficiency. Water is allocated to the uses that bring the greatest return; hence, total income in the Efficient Allocation scenario is the greatest of the five studied scenarios. Not surprisingly,

¹⁶ Water concession information was obtained from Ecuador's National Center for Hydrologic Resources

maximizing the efficiency of water use also brings the most efficient use of land resources; the percentage of land supply that is cultivated in the lower zone is at its maximum, 56 percent. Similar to the results of Bernardo et al. (1988), the scenario demonstrates the potential for efficiency gains due to water reallocation. In comparison to the Actual Allocation scenario, although water use in the upstream zones decreases by an average of 49 percent in the dry season, the average decrease in income in these three zones is less than 11 percent due to more efficient water use.

Results from the model reveal that policies designed to achieve equity as defined by water use versus income distribution criteria will have quite different effects on both resource use and income distribution, and are not likely to be supported by the same groups in the watershed. The Land-based Allocation scenario, which incorporates the standard of equity in water use, makes available an equal amount of water, at the water's source, to each hectare of land in the watershed, ignoring the differences in conveyance inefficiencies and irrigation requirements between zones. Due to these technological and climatological factors, the value of this water is much greater to upstream producers than downstream producers, and income distribution favors the upper zone and middle zone 1. An examination of Table 4 reveals a similarity between the results of the Land-based Allocation scenario and the Actual Allocation scenario. This suggests that, in reality, per-hectare agricultural water allocation is relatively equitable among zones. This does not imply that per hectare water *use* is equal between zones, as more water is required per hectare downstream, resulting in more land being left fallow.

In the Irrigation-based Allocation scenario, which attempts to achieve equity in income distribution, the average seasonal irrigation requirements in each zone are summed and the proportion of the total average requirement attributed to each zone in each season is used as a basis for limiting agricultural water use. These proportions naturally increase as one travels downstream, and the scenario allocates more water and more income to the lower zone, in comparison to the Land-based Allocation scenario. Lower zone farmers respond by increasing production of all crops, with greatest increases in high-value crops such as corn and anise. As is the case with the Efficient Allocation scenario, the shift of

water to the lower zone is partly achieved by reducing the levels of irrigation intensity in the upper zone and middle zone 1. Of the five scenarios, the Irrigation-based Allocation scenario does in fact result in the most equitable distribution of income.¹⁷

The combined results of the five water allocation scenarios clarify important relationships between water allocation and the achievement of efficiency and equity objectives. The Actual and Land-based Allocation scenarios demonstrate that an equitable distribution of water will not maximize efficiency in resource use nor equity in income distribution. In contrast, the Efficient and Irrigation-based Allocation scenarios result in the most efficient use of water and land resources, as well as the most equitable distribution of income, indicating that these two goals can be consistent with one another. Summarizing the results, improvements in efficiency (in resource use) and equity (in income distribution) are generally associated with (1) a shift of water to the lower zone, and (2) the use of lower levels of irrigation intensity upstream.

It is important to note, however, that even if policymakers were to take action to change water use according to these standards, it is unlikely that all residents of the watershed would support such an initiative. Although a water management policy may potentially improve the efficiency of resource use and increase total income in the watershed, it may not be implemented because a majority of farmers are better off without the policy (Ray, 1997). This is true of upper zone farmers in the El Angel watershed, who are currently favored with respect to the ownership of water rights as well as position along the canals, and who have no incentive to support such a change.

Resource valuation. A comparison of shadow prices (reported in Evans, 2001) reveals a relationship between the allocation of water and the value of land.¹⁸ In the water allocation scenarios which favor the upper zone (the Enforced Rights Allocation

¹⁷ As a percentage of per capita income in the highest-income zone, the difference in per capita income between the highest-income zone and the lowest-income zone is 23 percent in this scenario, compared to 95 percent, 75 percent, 50 percent and 76 percent in the Enforced Rights, Actual, Efficient and Land-based Allocation scenarios, respectively.

¹⁸ Labor is never a limiting resource in any scenario, and therefore its shadow price is always zero. The same can be said for the shadow price of land in the lower zone. In all modeled scenarios, water

scenario and the Land-based Allocation scenario), the shadow price of land is greatest in the upper zone, and decreases as one goes downstream. Conversely, with respect to water allocation, the Irrigation-based Allocation scenario favors downstream producers. In this scenario, the value of land is greater in middle zone 1 and middle zone 2 than in the upper zone. Thus, changes in water allocation would not only influence farmers' annual incomes earned from crop and cattle production, but could also simultaneously influence producers' wealth by either increasing or decreasing the value of their land assets.

Taxation strategies. The shadow price of water in the Efficient Allocation scenario, \$0.053/m³ in the dry season and \$0.036/m³ in the wet season, is an estimate of the value generated per cubic meter of water when the marginal value of water is equal among uses. These dry and wet season values are used as a basis for the study of water use taxation strategies. Water use is taxed at these levels; the tax is then gradually increased in the upper zone and middle zone 1 and gradually decreased in middle zone 2 and the lower zone.

The series of tax strategies studied using the model demonstrate that taxation can be used as a tool for shifting water resources and altering levels of efficiency and equity. As the tax on upstream water use is increased and the tax on downstream water use is decreased, the marginal value of water to downstream producers increases relative to upstream producers, and water and income therefore flow to the lower zone. At the same time, aggregate income increases and aggregate tax revenue decreases, due to the simultaneous increase in water use and decrease in tax rates downstream. Levels of production, and therefore income, in the upper zone, middle zone 1 and middle zone 2 prove to be quite insensitive to

limitations restrict land use in this zone to less than 60 percent of the total available land supply.

the imposition of taxes on water use, with primary changes occurring in production of the same crops at lower irrigation levels and the shift of land from improved to native pasture.

CONCLUSIONS AND POLICY IMPLICATIONS

Results from this study indicate that the actual allocation of water from the five studied canals, characterized by middle zone water use beyond the legal allotment, is more consistent with the goals of economic efficiency and income equity than the allocation of water that would result if current water rights were strictly enforced. Water rights are not strictly enforced, as local water user groups are hindered by a lack of government involvement, limited access to monetary and technological resources and poor communication. Under these circumstances, a producer's ability to obtain and utilize water resources to earn income is largely dependent upon his or her proximity to the water's source. Upstream farmers have no incentive to use water efficiently, and in the face of uncertainty, downstream farmers have an incentive to use water illegally.

By examining alternative standards for water allocation, this study provides guidelines for policymakers who seek to change these incentives. On one end of the policy spectrum is the strict enforcement of current concessions, which results in an inordinate waste of both water and land. At the other end is an economically efficient allocation of water that maximizes the value of agricultural output. Although this standard results in the greatest level of total income, it also requires the most dramatic shift of water resources to the lower zone.

Water can be used more efficiently in the lower zone for a number of reasons. First, the sensitivity of income to changes in water use is relatively small in the three upstream zones, due to the fact that crops in these zones may be produced at lower levels of irrigation intensity without severely reducing yields. Second, yield losses associated with reduced irrigation intensities in the lower zone offset the gains that could be made from using this water elsewhere, given relative price levels. This is largely the result of the third factor driving water downstream, the limited upstream land supply. Much of the water flows to the lower zone because the land constraints become binding in the three upstream zones.

Given the differences in the relative values of land and water resources, allocating an equal amount of water to each hectare of land will not provide the necessary incentives for improvement. If agricultural water could be allocated in proportions that reflect these differences, upstream farmers would have an incentive to utilize water more efficiently, and downstream farmers would have a sufficient supply to ensure a more equitable distribution of income.

The existence of a market for water rights might provide the necessary incentives to alter water use patterns. Unfortunately, due to a lack of communication between water user groups and government organizations, current water rights are not reliably recorded and local water organizations have neither the tools nor the authority to make and enforce decisions with respect to water use. Furthermore, water ownership is tied to land, which seriously inhibits the transferability of water in accordance with differences in its marginal value product among uses.

Results from this analysis also show that, if effectively implemented and enforced, a water use tax could be used to redistribute water as well as raise funds for canal improvements. Given the minimal impact of the tax on cropping patterns, tax revenue could also be used to equalize income distribution without significantly changing production levels through direct payments to targeted taxpayers.

In conclusion, achieving efficiency in resource use and equity in income distribution requires a significant transfer of water resources to the lower zone, largely accomplished through a shift to lower irrigation intensity crop activities upstream. However, at this time, the design and implementation of a policy to stimulate such a shift would be a difficult task, given the vastly different conditions faced by upstream and downstream users and the resulting incentives for water use. An alternative would be to make a one-time investment in technology to modernize the canals and improve levels of efficiency, thereby reducing the locational discrepancies between zones.

The implications and conclusions of this study are of relevance not only to the Andean Sierra, but could be applied to any region or watershed facing the challenges of increasing agricultural production, food security and achieving greater efficiency and/or equity in the allocation of irrigation resources. Using the terminology of Omezzine et al. (1998), for those watersheds where the “horizontal” expansion

of agricultural areas is not possible due to limitations in land and water supply, methods must be determined to “vertically” expand yields by increasing land and water efficiency and productivity. The mathematical programming model could be modified to reflect the meteorological and agricultural characteristics of other watersheds to determine the relative value of constrained resources and to provide guidelines for policymakers with respect to resource allocation.

REFERENCES

- Allen, Richard G, Luis S. Pereira, Dirk Raes, and Martin Smith. 1998. *Crop evapotranspiration: Guidelines for computing crop water requirements*. FAO Irrigation and Drainage Paper No. 56, Rome, Italy.
- Apollin, Frederic and Christophe Eberhart. 1998. *Metodologías de Análisis y Diagnóstico de Sistemas de Riego Campesino*. Sistema de Capacitación para el Manejo de los Recursos Naturales Renovables (CAMAREN), Quito, Ecuador.
- Arce, Blanca, Victor Barrera and Jovanny Suquillo. 1993. *Caracterización del Sistema de Producción del Pequeño Productor del Cantón Espejo, Provincia del Carchi*. Instituto Nacional Autónomo de Investigaciones Agropecuarias (INIAP), Fundación para el Desarrollo Agropecuario (FUNDAGRO), Quito, Ecuador.
- Beaulieu, Jeffrey, Dave Bennett, Steven Kraft and Raja Sengupta. 1998. “Ecological-Economic Modeling on a Watershed Basis: A Case Study of the Cache River of Southern Illinois.” Annual Meetings of the American Agricultural Economics Association (AAEA). Salt Lake City, UT.
- Berger, Ariadna Maria. 1999. *Optimal Resource Allocation in Argentine Dairy Farms: A Study of Management Strategies, Risk Considerations and Herd Size*. M.S. thesis, Cornell University, Ithaca, NY.
- Bernardo, D.J., N.K. Whittlesey, K.E. Saxon and D.L. Bassett. 1988. “Irrigation Optimization Under Limited Water Supply.” *Transactions of the ASAE* 31 (3), 712-719.
- Brockington, N.R., J.M. Veil and R. Zoccal. 1992. “Herd Dynamics and Management Strategies for Small-Scale Milk Production Systems in Southeast Brazil.” *Agricultural Systems* 39 (1992), 201-225.
- Brooks, Kenneth N., Peter F. Ffolliott, Hans M. Gregersen and K. William Easter. 1994. *Policies for Sustainable Development: the Role of Watershed Management*. Environmental and Natural Resources Policy and Training Project (EPAT)/Midwest Universities Consortium for International Activities, Inc. (MUCIA). Policy Brief No. 6.
- Brouwer, C. and M. Heibloem. 1986. *Irrigation Water Management*. Training Manual No. 3. FAO, Rome, Italy.
- Doorenbos, J. and A.H. Kassam. 1979. *Yield Response to Water*. FAO Irrigation and Drainage Paper No. 33, Rome, Italy.

- Doyle, C.J. 1990. "Application of Systems Theory to Farm Planning and Control: Modelling Resource Allocation." in *Systems Theory Applied to Agriculture and the Food Chain*. J.G.W. Jones and P.R. Street, co-editors. Elsevier Applied Science, London.
- Dunne, Thomas and Luna B. Leopold. 1978. *Water in Environmental Planning*. W.H. Freeman and Company, San Francisco, CA.
- Espinosa, Patricio, Rocio Vaca, Jorge Abad and Charles C. Crissman. 1996. *Raíces y Tubérculos Andinos: Cultivos Marginados en el Ecuador*. Centro Internacional de la Papa (CIP). Ediciones Abya-Yala, Quito, Ecuador.
- Evans, Elizabeth. 2001. *Efficiency and Equity in Water Allocation: An Optimization Model of the El Angel Watershed, Carchi, Ecuador*. M.S. thesis, Cornell University, Ithaca, NY.
- Garg, N.K. and Abbas Ali. 1998. "Two-level optimization model for Lower Indus Basin." *Agricultural Water Management* 36, 1-21.
- Hazell, P.B.R. and R.D. Norton. 1986. *Mathematical Programming for Economic Analysis in Agriculture*. Macmillan Publishing Co., New York, NY.
- Holden, Paul and Mateen Thobani. 1996. *Tradable Water Rights: A Property Rights Approach to Resolving Water Shortages and Promoting Investment*. Policy Research Working Paper, The World Bank, Washington, D.C.
- Montenegro, Marybel M. 1998. *Estudio Sobre Uso, Manejo y Distribución de Agua Para Riego y Consumo Humano*. Consejo Nacional de Recursos Hídricos, Quito, Ecuador.
- National Research Council (NRC). 1989. *Nutrient Requirements of Dairy Cattle*. Sixth Revised Edition Update 1989. National Academy Press, Washington, D.C.
- National Research Council (NRC). 1996. *Nutrient Requirements of Beef Cattle*. Seventh Revised Edition 1996. National Academy Press, Washington, D.C.
- Nicholson, Charles F., David R. Lee and Richard N. Boisvert. 1994. "An Optimization Model of the Dual-Purpose Cattle Production System in the Humid Lowlands of Venezuela." *Agricultural Systems* 46, 311-334.
- Nicholson, Charles Frederick. 1990. *An Optimization Model of Dual Purpose Cattle Production in the Humid Lowlands of Venezuela*. M.S. thesis, Cornell University, Ithaca, NY.
- Okumu, B.N., M.A. Jabbar, D. Colman and N. Russell. 1999. "Bio-economic Modeling of Watershed Resources in Ethiopia." Annual Meetings of the American Agricultural Economics Association (AAEA). Nashville, TN.
- Omezzine, Abdallah and Lokman Zaibet. 1998. "Management of modern irrigation systems in Oman: allocative vs. irrigation efficiency." *Agricultural Water Management* 37, 99-107.
- Paudyal, G.N. and A. Das Gupta. 1990. "A nonlinear chance constrained model for irrigation planning." *Agricultural Water Management* 18, 87-100.
- Poats, Susan V., Osvaldo Paladines, Jenny Valencia, Hernán Valencia and Galo Medina. 1998. *Manejo Colaborativo y Uso Apropiado de Recursos Naturales en la Ecoregión de la Cuenca del Río El Ángel, Carchi, Ecuador*. Proyecto MANRECUR II, Quito, Ecuador.

- Prato, Tony and Shunxiang Wu. 1996. "Alternative Spatial Criteria for Targeting Soil and Water Quality Improvements in an Agricultural Watershed." *Review of Agricultural Economics* 18, 293-301.
- Proaño, Mauricio. 2000. "¿Abundancia o Escasez? Concesiones, Conflictos, Poderes y Políticas en el Manejo del Agua en la Cuenca del Río El Angel." Presentation to the Consorcio Carchi, March 3, 2000, Carchi, Ecuador.
- Proaño, Mauricio and Osvaldo Paladines. 1998. *Análisis de los Sistemas Agropecuarios de los Pequeños Productores de la Cuenca del Río El Angel, Provincia del Carchi, Ecuador*. MANRECUR Project, Quito, Ecuador.
- Proyecto DRI. 1991. *Estudio de Rehabilitación de Acequia*. Mira, Ecuador.
- Ray, Isha. 1997. *A Farming Systems Analysis of Irrigation Institutions*. Ph.D. thesis, Stanford University. UMI Dissertation Services, Ann Arbor, MI.
- Restrepo, J.I., E. Murgueitio and T.R. Preston. 1991. "Milk production systems in tropical Latin America." in *Feeding Dairy Cows in the Tropics*. FAO Animal Production and Health Paper 86, Rome, Italy, 189-198.
- Santos, Francisco L. 1998. "Evaluation of Alternative Irrigation Technologies based upon Applied Water and Simulated Yields." *Journal of Agricultural Engineering Resources* 69, 73-83.
- Sotomayor V., Jorge. 1996. *Perfil de Riego de la República del Ecuador*. Instituto Internacional del Manejo de la Irrigación (IIMI), Quito, Ecuador.
- Thornthwaite, C.W. and J.R. Mather. 1957. *Instructions and tables for computing potential evapotranspiration and the water balance*. Laboratory of Climatology, Publication No. 10, Centerton, NJ.

Table 1. Breakdown of the Left Margin of the El Angel Watershed by Zone

Zone	Elevation range (masl)	Land (hectares)	Human population	Average farm size (hectares)	Average precipitation (mm/year)	Average evapotrans. (mm/year)	Production system
Upper	4,000-3,100	552	6,356	3.88	1,046	722	Dual purpose cattle, potatoes barley
Middle 1	3,099-3,000	910	4,022 ^a	3.37 ^a	955	903	Dual purpose cattle, potatoes barley, wheat
Middle 2	2,999-2,400	732			807	1,100	Corn, black beans, peas
Lower	2,399-1,500	747	2,244	3.15	416	1,421	Anise, corn sweet potato, white carrot

Sources: Montenegro (1998), Proaño and Paladines (1998), MANRECUR Project, CIP.

^a

Table 2. Model Activities and Constraints^a

Activities	Constraints
Human Population in each zone Crop Hectares of crop/pasture production - For each zone - For each level of irrigation intensity Animal Primary age groups Activity- and sex-based subgroups - Male/female calves - Replacement/sold heifers - Steers - Lactating/dry cows - Culled cows Animal units Aggregate nutrient intake - Dry matter - Crude protein - Metabolizable energy - Water Canal water requirement Purchased supplemental feeds Milk activities - Production - Consumption - Sale Water Levels of water use - From each source of water supply ^b - For each zone	Human Define populations in each zone Crop Equate seasonal production - For each zone - For each level of irrigation intensity Animal Divide age groups by sex and activity Relate sizes of age groups Convert herd to animal units (AUs) Define aggregate nutrient reqts. Ensure nutrient reqts. are met Determine cattle canal water reqt. Restrict stocking rate Define milk use - Production - Consumption - Sale Land and Labor Restrict land use based on supply Restrict labor use based on supply Water Define aggregate water supplies ^b Define total water use in each zone Restrict water use \leq water supply Restrict water use in each zone ^c

^a Most activities and constraints are defined for both the dry and wet seasons. The complete algebraic formulation of the model, including over 200 individual activities and constraints, is reported in Evans (2001).

^b Water supplies enter at the top of the watershed and at the end of middle zone 1.

^c For certain water allocation scenarios.

Table 3. Summary of Assumed Crop Input Costs, Yields and Prices

		Price ^a (\$/mt)	100% Irrigation Cost ^a (\$/ha)	Yield (mt/ha)	67% Irrigation Cost ^a (\$/ha)	Yield (mt/ha)	33% Irrigation Cost ^a (\$/ha)	Yield (mt/ha)
Upper	Potatoes ^b	120.00	824.84	15.91				
	Imp. Pasture ^{b,c}	n/a	175.31	19,247				
	Barley	195.00	248.45	2.27	242.25	2.19	236.05	2.10
Middle 1	Potatoes ^b	120.00	824.84	15.91				
	Barley	195.00	248.45	2.27	227.17	1.83	207.91	1.43
	Wheat	211.00	263.73	2.27	246.36	1.94	230.01	1.64
Middle 2	Corn	243.18	345.10	3.86	334.55	3.65	324.10	3.44
	Beans	440.00	222.41	1.00	217.04	0.92	211.72	0.84
	Peas	580.00	214.89	1.11	200.23	0.78	187.43	0.49
Lower	Corn	243.18	345.10	3.86	282.39	2.46	237.05	1.46
	Anise	1,585.58	250.34	0.90	231.24	0.54	217.97	0.30
	Sweet Potato	102.94	318.01	9.00	272.77	5.10	242.48	2.53
	White Carrot	100.00	318.01	9.00	273.46	5.16	243.33	2.60

Data sources: Arce, fieldwork (2000), Proaño and Paladines (1998), CIP (1990-1992).

^a Prices and costs effective as of June, 2000.

^b Potatoes and improved pasture are assumed fully irrigated.

^c Yield data for improved pasture are annual values of kgDM/ha. Cost data for improved pasture are totals for a 3-year period.

Table 4. Zone-level Water Use, Crop Production and Cattle Production for the Modeled Scenarios

			Enforced Rights		Actual		Efficient		Land-based		Irrigation-based		
Zone	Water Use (m³)		Dry	Wet	Dry	Wet	Dry	Wet	Dry	Wet	Dry	Wet	
	Upper		1,143,833	773,807	1,143,828	773,806	474,824	630,065	1,143,833	773,807	406,295	538,982	
	Middle 1		240,770	241,118	1,416,278	867,522	1,184,950	788,766	1,416,278	867,522	697,307	622,748	
	Middle 2		240,768	254,072	2,212,597	2,578,976	625,422	1,278,029	1,514,288	1,699,156	1,162,979	1,479,559	
	Lower		1,585,635	1,776,103	1,585,635	1,776,103	4,073,128	4,404,215	1,572,107	1,760,767	3,286,868	4,459,792	
Zone	Crop Production		Irrig. %	Hectares	Irrig. %	Hectares	Irrig. %	Hectares	Irrig. %	Hectares	Irrig. %	Hectares	
	Upper	Potatoes	100%	353.49	100%	353.49	100%	318.24	100%	353.49	100%	203.56	
		Barley	100%	35.70	100%	35.70	100%	32.14	100%	35.70	33%	109.65	
		I. Pasture	100%	162.81	100%	162.81	100%	15.36	100%	162.81	100%	0.00	
		N. Pasture	0%	0.00	0%	0.00	0%	186.25	0%	0.00	0%	238.80	
	Middle 1	Potatoes	100%	35.22	100%	240.68	100%	240.68	100%	240.68	100%	240.68	
		Barley	100%	0.00	100%	336.25	100%	336.25	100%	336.25	100%	67.66	
		Barley	33%	49.20	33%	0.00	33%	0.00	33%	0.00	33%	268.59	
		Wheat	33%	18.65	100%	127.42	33%	127.42	100%	127.42	33%	127.42	
		N. Pasture	0%	806.93	0%	205.66	0%	205.66	0%	205.66	0%	205.66	
		Middle 2	Corn	100%	0.00	100%	181.78	100%	0.00	100%	159.41	100%	80.84
			Corn	33%	24.85	33%	0.00	33%	262.74	33%	57.99	33%	154.48
	Beans		33%	38.45	100%	281.26	33%	406.54	33%	336.38	33%	364.11	
	Peas		100%	12.73	100%	268.96	100%	62.72	100%	178.22	100%	132.57	
	Lower	Corn	100%	68.09	100%	68.09	100%	173.55	100%	67.44	100%	205.81	
		Anise	100%	66.93	100%	66.93	100%	185.88	100%	66.29	100%	143.22	
		S. Potato	100%	14.74	100%	14.74	100%	40.93	100%	14.60	100%	31.54	
		W. Carrot	100%	7.64	100%	7.64	100%	21.21	100%	7.56	100%	16.34	
Zone	Cattle Production		AUs	Milk (l)	AUs	Milk (l)	AUs	Milk (l)	AUs	Milk (l)	AUs	Milk (l)	
	Upper		391	139,080	391	139,080	391	139,080	391	139,080	391	139,080	
	Middle 1		1320	469,980	391	139,080	391	139,080	391	139,080	391	139,080	

Table 5. Breakdown of Income Distribution by Scenario and Zone

		Enforced Rights	Actual	Efficient	Land- based	Irrigation- based
Total Income		\$631,631	\$1,168,852	\$1,294,589	\$1,137,726	\$1,127,115
Upper	Zone income	\$392,159	\$392,159	\$361,869	\$392,159	\$251,235
	Percent of total	62.09%	33.55%	27.95%	34.47%	22.29%
	Income per hectare ^a	\$710	\$710	\$656	\$710	\$455
	Income per capita ^b	\$493	\$493	\$455	\$493	\$316
Middle 1	Zone income	\$82,533	\$359,656	\$347,041	\$359,656	\$314,181
	Percent of total	13.07%	30.77%	26.81%	31.61%	27.87%
	Income per hectare ^a	\$91	\$395	\$381	\$395	\$345
	Income per capita ^b	\$69	\$303	\$292	\$303	\$264
Middle 2	Zone income	\$24,358	\$284,456	\$226,694	\$254,599	\$242,316
	Percent of total	3.86%	24.34%	17.51%	22.38%	21.50%
	Income per hectare ^a	\$33	\$389	\$310	\$348	\$331
	Income per capita ^b	\$24	\$285	\$227	\$255	\$243
Lower	Zone income	\$132,581	\$132,581	\$358,985	\$131,312	\$319,383
	Percent of total	20.99%	11.34%	27.73%	11.54%	28.34%
	Income per hectare ^a	\$177	\$177	\$481	\$176	\$428
	Income per capita ^b	\$121	\$121	\$329	\$120	\$292

^a Calculated based on total land supply.^b Based on total population dependent upon agriculture.