



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

JAAKKO KANGASNIEMI AND THOMAS REARDON*

Demographic Pressure and the Sustainability of Land Use in Rwanda¹

Abstract: Increasing land scarcity forces Rwandan farmers to expand the area under food crops at the expense of pasture, fallow, and forest. Since the non-cropping uses of land provide more vegetative cover against erosion than most food crops, land scarcity appears to be associated with unsustainable land uses. However, demographic pressure also pushes farmers to grow crops in dense associations, which increases vegetative cover on cultivated fields. The estimated relationship between farm size and protective crop cover depends crucially on how the measure of vegetative cover is adjusted to account for high cropping densities. Without any adjustment, the association between land scarcity and erosive land use is strong; with the adjustment used here, it disappears, except for high altitude areas, where bananas, the only major food crop that protects land well against erosion, do not grow well.

INTRODUCTION

Does demographic pressure make agriculture less sustainable in developing countries? Literature presents two conflicting hypotheses (National Research Council, 1993). The pessimistic, 'Ricardian,' hypothesis is that when population grows, farmers tend to mine their soils and expand cultivation to marginal, easily erodible land. The higher the rate of population growth and the more fragile the environment, the greater is the rate of erosion.

The optimistic hypothesis is that because land-scarce farmers depend on land productivity for food security, they attempt to prevent degradation. Above all, they use their relatively abundant labour to adopt more sustainable labour-intensive land use practices and to invest in soil conservation. In Rwanda, the pessimistic hypothesis has much credibility. The most densely populated country on the African continent, Rwanda has an average farm size below one hectare (DSA, 1992). Farming has already expanded to cover the easily erodible hillsides. Almost one-third of cultivable fields are now located on slopes of 20 degrees or more. In a 1991 survey by the Ministry of Agriculture, 40 percent of field blocks were reported by their cultivators as degraded.

Yet economic theory could support either hypothesis. On the one hand, it suggests that land-scarce poor farmers may have a high rate of time preference and may sacrifice long-term sustainability for immediate food security. On the other hand, theory suggests that as population growth changes the relative endowments of land and labour, it also may change relative prices so that using labour to maintain the productivity of land becomes more attractive. In the end, the question is empirical and depends significantly on the characteristics of the land uses (crops) farmers can choose in a specific environment.

To understand fully what determines how sustainable are the land uses farmers choose, ideally one would first model the determinants of land use, and then model their impacts on the farming environment. The space here is too brief for both, so we focus on the second, and ask directly whether increasing land scarcity, reflected in miniaturization of farms, is associated with Rwandan farmers having unsustainable land uses. This

* Visiting Specialist, Michigan State University, USA, Outposted in the Division of Agricultural Statistics (DSA), Ministry of Agriculture (MINAGRI), Kigali, Rwanda, Michigan State University, USA, respectively.

addresses an important unresolved debate in a context that should tell much about the future sustainability of agriculture in Africa.

We examine this longitudinal question with cross-sectional data by comparing how well the land uses chosen by more and less land-scarce farmers protect the land against erosion. Land use is here defined to include the allocation of land to different uses (crops, pasture, fallow, and woodlot) and cultural practices such as intercropping and crop planting density. This definition excludes treatment of soil conservation investments, which are studied with the same data base by Clay and Reardon (1994).

DATA AND METHODS

This study uses data for 821 sample households that derive from the detailed rural household survey conducted in 1991 by the Division of Agricultural Statistics (DSA), Ministry of Agriculture and Livestock of Rwanda.

Our approach has four steps. First, we categorize farm households into landholding quartiles according to their land scarcity using the criterion of cultivable land per adult equivalent.² All pastures, fallows, and woodlots are considered cultivable. Second, we compare over quartiles the average index of the protective crop cover on farm fields, without controlling for other factors. Third, we repeat the comparison with stratifications for agroclimatic zone and for altitude.

Fourth, to control better for agroclimatic factors we use regression analysis with observations aggregated over fields to farm-household observations. The control variables include rainfall and altitude. In addition, a multiplicative dummy variable is introduced to allow the coefficient for farm size to differ for high altitudes where bananas do not grow well (MINAGRI, 1978). The hypothesis is that since bananas are highly caloric and also protect the land well against erosion, the areas where few bananas are grown would show a stronger link between erosive land uses and demographic pressure.

The dependent variable in the analysis is an index of the protective vegetative cover provided by crops or other land use such as pasture, known as the C-value (erosion index measuring crop protective cover). Soil scientists use C-values together with data on slope, rainfall, and soil type to predict erosion (Wischmeier and Smith, 1978). The C-value itself is not a measure of erosion but, other things being equal, a lower C-value indicates less erosion. Depending on the varieties and cultural practices, the protective cover provided by a given crop varies widely and its C-value needs to be estimated empirically for different environments. The crop-specific C-values used in this paper are estimated in Lewis (1986) and Lewis, Clay and Dejaegher (1988) based on Rwandan field data.

Although C-values are specific to cultural practices such as weeding and mulching, the C-value estimates used here are the same for all farm-size quartiles. However, it is known that all major crops show 50–100 percent higher yields on the fields of the most land-scarce quartile compared to those of the least land-scarce quartile (Uwamariya, Kangasniemi, and Reardon, 1993). This suggests that small farmers (in terms of cultivable land per adult equivalent) use much more labour per unit of land than large farmers. The impact of this on the protective crop cover is unclear, since while some ways of using additional labour (weeding) are likely to expose land to erosion, others (mulching) clearly increase soil protection and crop cover.

Since more than one-half of cultivated fields in Rwanda are intercropped, a key question is how to estimate C-values for crop associations. Because few empirically-

based estimates were available for crop associations, we used C-values estimated for purely cropped fields to compute C-values for the associations. Our unadjusted C-value is an average of crop-specific C-values weighted by the estimated land shares of each crop in a given association. However, DSA's data show that planting densities on intercropped fields are higher than on purely cropped fields, which suggests that there is more vegetation, and thus more crop cover against erosion. Also DSA's field experiments indicate that the average C-value of the crops grown in the association overstates the C-value (exposure to erosion) of the association (Lewis 1986). In the literature, reduced erosion is mentioned as one of the reasons farmers continue to rely on intercropping (Fussel and Serafini, 1985).

To adjust C-values for high densities we divide the unadjusted C-value by the sum of the densities of the crops grown in the association (for technical details, see Kangasniemi, 1993). On average, the adjustment reduces C-values (increases soil protection, *ceteris paribus*) of the intercropped fields from 0.20 to 0.13.

Rwanda's main export crop, coffee, is usually mulched and has a very low C-value (0.02). Also forest ($C = 0.06$), fallow ($C = 0.10$), and pasture ($C = 0.10$) cover the land well. Of the main food crops, only bananas have a low C-value (0.04), whereas beans (0.19), tubers (sweet potato: 0.23, potato: 0.22, and cassava: 0.26), and cereals in particular (maize: 0.35 and sorghum: 0.40) provide much less protective cover against erosion.

Table 1 shows the differences in land use by farm-size quartiles. The smallest farms, compared to the other quartiles, dedicate more of their cultivable land to bananas, tubers, cereals, and coffee. They cultivate a larger share of their land at the expense of pasture, fallow and forest.

Table 1 Land Allocation by Farm Size Quartile

	Farm Size Quartile (Ares/AE)				C-value
	0-9 ares	9-15 ares	15-26 ares	26- ares	
Cereals	12%	12%	10%	9%	0.35-0.40
Tubers	26%	24%	22%	13%	0.22-0.26
Beans	16%	15%	12%	10%	0.19
Fallow/Pasture	9%	13%	19%	30%	0.10
Forest	5%	5%	8%	13%	0.06
Banana	18%	19%	18%	15%	0.04
Coffee	6%	5%	5%	4%	0.02
Other	7%	7%	7%	6%	0.02-0.35

Source: DSA/MINAGRI Survey Data, Agricultural Year 1991.

Note: Means shares for seasons A and B, weighted by farm size.

RESULTS

Results of the C-value calculations for the cultivated, cultivable, and intercropped fields of each farm size quartile at the national level are shown in Table 2. All differences between the quartiles that show up in the table are significant at the 0.1 percent level.

Table 2 *Crop Cover Indices by Farm Size Quartile*

	Cultivable Fields		Cultivated Fields		Intercropped Fields		Total Density
	Density- adjusted C-value	Un- adjusted C-value	Density- adjusted C-value	Un- adjusted C-value	Density- adjusted C-value	Un- adjusted C-value	
Farm Size per Adult Equiv.							
0– 9 ares	0.13	0.17	0.14	0.19	0.11	0.19	1.889
9– 15 ares	0.13	0.17	0.14	0.18	0.13	0.20	1.758
15–26 ares	0.13	0.16	0.14	0.18	0.13	0.20	1.698
26– ares	0.12	0.14	0.14	0.17	0.14	0.20	1.585

Source: DSA/MINAGRI Survey Data, Agricultural Year 1991.

Notes: Means weighted by field size. All visible differences significant at 1 percent level.

When land use on *all fields* is examined, the unadjusted C-values show that land-scarce households allocate their land to uses that provide less protective cover against erosion than the uses chosen by those with more land. However, the density-adjusted C-values show no clear relationship, suggesting that small farmers (in terms of cultivable land per adult equivalent) make up much of the difference by growing crops in higher densities.

This key result is compatible with the finding that there are no significant differences between farm size quartiles in the share of fields reported as degraded (Clay, 1993).

Table 3 *Erosivity (C-value, or Crop Cover) Indices by Farm Size, Zone and Altitude*

	Agroclimatic zone					Altitude (m)	
	North West	South West	North Central	South Central	East	–1900	1900–
Farm Size per Adult Equiv.							
0–9 ares	0.17	0.11	0.12	0.11	0.11	0.12	0.17
9–5 ares	0.18	0.12	0.13	0.11	0.11	0.12	0.16
15–26 ares	0.17	0.12	0.12	0.11	0.11	0.12	0.14
26– ares	0.15	0.10	0.11	0.11	0.10	0.12	0.12

Source: DSA /MINAGRI Survey Data, Agricultural Year 1991.

Notes: Mean Adjusted C-values of cultivable fields weighted by field size. All visible differences between quartiles significant at 1 percent level.

When only *cultivated fields* are compared, neither measure suggests that demographic pressure pushes farmers to grow more erosive crops. Due to the higher densities on their intercropped fields, land-scarce farmers may actually generate less exposure to erosion on their cultivated fields. Thus, the reason they do worse or at least not better than larger farmers when all fields are compared is that they need to cultivate a larger share of their land at the expense of pasture, fallow, and forest.

The examination of the subset of *intercropped fields* shows that land-scarce farmers have substantially higher cropping densities on their intercropped fields. This covers their intercropped fields better against erosion than do their less land-scarce neighbours.

The result that the degree of protective crop cover is almost the same on small and on large farms holds well in three of the five agroclimatic zones (Table 3). In the North West,

agricultural land uses are more erosive than elsewhere, particularly on small farms. The reason presumably is that much of the North West is of too high an altitude for bananas, which are the only major food crop that covers the land well against erosion. Differences in banana production may also explain the results for the North Central zone. To explore this further, comparisons were also made separately for altitudes below and above 1900 metres. At the high altitudes, which cover roughly one-quarter of Rwanda's cultivable land, land scarcity was strongly associated with erosive land use practices.

Table 4 shows that using regression analysis to control for rainfall and altitude with does not change the basic results presented above. The first equation confirms the finding that when the estimates of crop cover are adjusted downward for densely intercropped fields, farm size is not significantly associated with erosive land use, except for the high altitude areas. The most important determinant of crop cover is altitude, largely because few bananas are grown at high altitudes.

Table 4 Regression Results for Crop Cover Index (Cultivable Fields)

Independent Variable	Dependent Variable	
	Adjusted C-value	Unadjusted C-value
Intercept	0.00641	0.09889
(<i>t</i>)	(0.70)	(2.28)
Rainfall (cm)	-0.00025	-0.00085
(<i>t</i>)	(-5.88)*	(-4.25)*
Altitude (100m)	0.00871	0.00592
(<i>t</i>)	(14.52)*	(2.08)
Land (ares/ad.eq.)	-0.00003	0.00340
(<i>t</i>)	(0.54)	(12.88)*
Land*dummy for alt.> 1900 metres	-0.00048	0.00008
(<i>t</i>)	(-4.09)*	(-0.15)
Adj. R Square	0.24	0.21

Source: DSA/MINAGRI Survey Data, 1991.

Note: * significant at the 1 percent level.

The second equation (Table 4, right column) illustrates that without the adjustment for density, the result is entirely different. If small farmers are not 'given credit' for their higher cropping densities in the calculation of the C-values, land scarcity appears to make land use much more erosive. In contrast, the association of crop cover with altitude becomes insignificant if the higher densities at low altitudes are not accounted for.

CONCLUSIONS

Although increasing land-scarcity forces Rwandan farmers to cultivate a larger share of their land at the expense of forest, pasture, and fallow, it also encourages them to grow more perennials and to grow crops in dense associations. While the expansion of cultivation contributes to erosion, especially when it occurs on easily erodible steep slopes, perennials and dense associations are a form of intensification that makes land use more sustainable. The estimated net impact of these changes depends crucially on how one adjusts the estimates of vegetative crop cover for high cropping densities. With the

adjustment used here, small farmers do not appear to have substantially more erosive land uses than large farmers, except in the high-altitude areas.

Our adjustment for high cropping densities reduces the estimated C-values on the intercropped fields of the smallest farm-size quartile almost by one-half, which may be too generous. On the other hand, no adjustment was made for purely cropped fields, which probably also have higher cropping densities on small farms than on large farms. Without field experiments, it is impossible to say whether our adjustment is too optimistic.

With or without adjustment for high cropping densities, land uses appear erosive at high altitudes, where few bananas are grown. Crop cover on these areas is poor and is becoming less protective with increasing land scarcity. Moreover, fields in high altitude areas are much steeper than elsewhere, which makes them even more vulnerable to erosion. At high altitudes, finding ways to channel the additional labour provided by population growth to soil conservation through investments such as anti-erosion ditches and terraces and land use practices such as mulching and agroforestry is a major challenge. Since much of the environmentally sustainable labour-based intensification at lower altitudes involves banana production, agricultural research on banana varieties that grow well on high altitudes might have a high payoff in terms of environmental sustainability and long-term food security.

In sum, our cross-sectional comparisons give only limited support for the hypothesis that demographic pressure makes land use less sustainable. Although this gives some optimism regarding the environmental impacts of population growth in the nearby future, it does not change the fact that during the recent past population growth has already forced Rwandan farmers to cultivate marginal lands. On the steep slopes, the current levels of crop cover are insufficient to control erosion. The current rate of land degradation is alarming, and measures to reduce it are badly needed. To design policies that promote soil conservation investments and land use practices that protect the soil against erosion, especially on easily erodible areas, decision makers need information on the determinants of investments and land use. Research that provides such information deserves priority in Rwanda and other countries facing similar problems.

NOTES

¹ We are grateful to Dan Clay for his suggestions. We are also grateful for USAID/Kigali's Rwanda Add-on and to USAID/ARTS/FARA/FSP's Productivity Add-on for support for this research as part of the Food Security II Cooperative Agreement at Michigan State University, and to the Division des Statistiques Agricoles (DSA), Ministère d'Agriculture et de l'Élevage, Rwanda for use of data. We are also grateful for comments on an earlier version by Akin Adesina. The ideas and interpretations expressed herein are those of the authors and are not necessarily shared by the sponsoring agencies.

² Conversion into adult equivalents is based on Miniplan (1988).

REFERENCES

- Clay, D. C., 1993, 'Fighting an Uphill Battle: Demographic Pressure, the Structure of Landholding, and Land Degradation in Rwanda', unpublished draft manuscript, Michigan State University, Department of Sociology, East Lansing.
- Clay, D. and Reardon, T., 1994, 'Determinants of Farm-Level Conservation Investments by Rwandan Farm Households', Contributed Paper to the International Association of Agricultural Economists Twenty-Second Conference, Harare, Zimbabwe, pp. 210–221 of this volume.

- DSA, 1992, *Enquête Nationale Agricole 1990: Production, Superficie, Rendement, Elevage et Leur Evolution 1984-1990*, Publication DSA No. 26, Division Des Statistiques Agricoles, Ministère de l'Agriculture et de l'Elevage, Republique Rwandaise, Kigali, Rwanda.
- Fussel, L.K. and Serafini, P.G., 1985, 'Crop Associations in the Semi-Arid Tropics of West Africa: Research Strategies Past and Future', in Ohm, H.W. and Nagy, J.G. (eds.), *Appropriate Technologies for Farmers in Semi-Arid West Africa*, Purdue University.
- Kangasniemi, J., 1993, 'Estimating Vegetative Cover (C-values) for Crop Association', Division of Agricultural Statistics, unpublished research note, Ministry of Agriculture and Livestock, Republic of Rwanda.
- MINAGRI, 1978, *Guide Pratique de l'Agronome: Première Partie: Les Principales Cultures du Rwanda: Les Conditions de Production*, Ministère de l'Agriculture et de l'Elevage, Kigali, Rwanda.
- MINPLAN, 1988, *Enquête Nationale sur le Budget et la Consommation des Ménages (Milieu Rural), Volume 4: Consommation Alimentaire en Milieu Rural*, Ministère du Plan, République Rwandaise, Kigali, Rwanda.
- National Research Council (US), 1993, Committee on Sustainable Agriculture and the Environment in the Humid Tropics, *Sustainable Agriculture and the Environment in the Humid Tropics*, National Academy Press, Washington, D.C.
- Lewis, L.A., 1986, 'Predicting Soil Loss in Rwanda', unpublished research note, Graduate School of Geography, Clark University, Worcester, Maryland, and Division of Agricultural Statistics, Ministry of Agriculture and Livestock, Republic of Rwanda, Kigali, Rwanda.
- Lewis, L.A., Clay, D.C. and Dejaegher, Y.M.J., 1988, 'Soil Loss, Agriculture, and Conservation in Rwanda: Toward Sound Strategies for Soil Management', *Journal of Soil and Water Conservation*, Vol. 43, No. 5, pp.418-421.
- Uwamariya, L., Kangasniemi, J. and Reardon, T., 1993, 'La Productivité Agricole au Rwanda, 1989-1990: La Productivité Moyenne de la Terre, du Travail, et la Rentabilité de la Terre, et les Fonctions de Production', draft Working Paper, Division Des Statistiques Agricoles, Ministère de l'Agriculture et de l'Elevage, Republique Rwandaise, Kigali, Rwanda.
- Wischmeier, W.H. and Smith, D.D., 1978, *Predicting Rainfall Erosion Losses, a Guide to Conservation Planning*, Agricultural Handbook No. 537, US Department of Agriculture, Washington, D.C.

DISCUSSION OPENING — K.J. Thomson (*University Of Aberdeen, UK*)

The paper addressed directly a major issue in a country currently suffering a human disaster depicted nightly on our TV screens.

In the introduction, two potentially interesting theoretical issues are raised, though not (understandably) resolved. First, it is suggested that rate of time preference depends on poverty and the land resources of farm families. Second, it is suggested that the land/labour ratio affects the sustainability of land management practices (in passing, it may be noted that in developed countries, non-sustainable land management occurs, often in labour scarce regions. Either of these issues might reward further more general analysis.

The empirical analysis in the paper is of course crucially dependent on the nature and reliability of the C-values and the adjustment process undertaken by the authors for intercropping. Since the C-value itself, and the adjustment arithmetic, are technical measures directly related to actual erosion, the result may be sensitive to the nature of the dependent variable rather than the erosion damage ultimately in focus. Since land management apparently makes a great difference to yields (+50 to 100 per cent), and presumably to erosion too, more sophistication may be justified here. Turning to the results, the differences in the tables are rather small (one or two points) although asserted to be statistically significant at the 1 per cent level, and some information on variability in the data would be welcome.

The paper thus, represents an interesting but partial attack on a crucial problem in African agriculture, as elsewhere. Its narrow data base and analytic focus inevitably suggest the need for further information on such matters as the land management practices of the farming communities involved, including their responses to increased erosion itself. Meanwhile, one can only hope that such analyses are not overwhelmed by events such as the current emergency in Rwanda.

GENERAL DISCUSSION — Michelle Veeman, Rapporteur (*University of Alberta, Canada*)

Discussion of the paper by Rekha Mehra included comments on elements of the social structure that limit women's access to land and a query about the impact of matrilineal, as versus patrilineal descent on women's land rights. Dr Mehra argued that the impact of matrilineal descent in providing women with access to land is reduced due to the prevalence of patriarchal societies and the tendency for women's location to change after marriage. Other points of discussion included possibilities and limitations of women's access to employment and the question of whether increased access to land, as versus other employment opportunities, would better increase family food security. The need for targeting of programs, such as credit, specifically toward women was advocated by one discussion participant and disputed by another. A challenge to this author, repeated by several speakers, queried the demonstration of a clear linkage between women's access to land and adverse environmental effects.

Hugo De Groote's paper on women's income versus family income as a determinant of food security for Mali was, in the absence of the presenter, briefly summarized by the discussant, Benon Gazinski, who noted that it is based on a survey of male household heads. (Dr De Groote was not able to present his paper until a later session). This issue was raised by a participant who noted that surveys of male household heads tend to elicit different responses and policy prescriptions than when women are interviewed.

Numbers of comments on the paper by Jaako Kangasniemi and Thomas Reardon were directed to the focus on cropping patterns, specifically inter-cropping, relative to a broader range of resource management practices to combat land degradation, and a lack of information on the erodibility index and their adjustments to this index. The apparent small difference between size classes shown in the tables were the subject of several comments. The suggestion, in the verbal presentation, that commodity tax and price policy might be useful to encourage less erosive cropping practices was also challenged by several speakers. The authors responded that although differences between size classes appeared to be small, on average, there are sizeable differences in erodibility at high altitudes. While acknowledging that the scope of the paper is narrower than the broad problems of erosion and population pressure, Kangasniemi and Reardon noted that farmer's investment and labour-intensive practices to control erosion, such as ditching, by small farmers are expressly excluded from this paper but are considered in another paper; presented at this conference, based on the same data set (Clay and Reardon, pp.210–219 of this volume).

Participants in the discussion included D.G.R. Belshaw, Steven Franzel, Theodora S. Hyuha, Rosebud Kurwijib, Bongwe Njobe, Mark Odhiambo, P.C. Sarkar, Nana M. Tanko and Joachim von Braun.