



*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

**Give to AgEcon Search**

AgEcon Search

<http://ageconsearch.umn.edu>

[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

*No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.*

## Effect of citric acid and acetic acid on the performance of broilers

M.Z. Islam, Z.H. Khandaker<sup>1</sup>, S.D. Chowdhury and K.M.S. Islam

Department of Poultry Science and <sup>1</sup>Department of Animal Nutrition, Bangladesh Agricultural University, Mymensing-2202, Bangladesh

### Abstract

An experiment was conducted with commercial broilers to investigate the effects of feeding citric acid, acetic acid and their combination on their performance and to determine the economic competence of using citric acid and acetic acid in broiler rations. A total number of 108 one day old straight run broiler chicks were distributed to four dietary treatments i.e. 0 % citric or acetic acid (A) , 0.5% citric acid (B), 0.5% acetic acid (C) and their combinations 0.5% citric acid and 0.5% acetic acid (D). The birds were reared in cages. Body weight gains, feed conversion ratio (FCR), mortality, dressing yield and carcass characteristics were recorded. The performance showed significant increase in body weight gain ( $P<0.05$ ) when compared with the control during 0-5 weeks of age on 0.5% citric acid (B). Feed consumption increased ( $P<0.05$ ) at 2<sup>nd</sup> and 3<sup>rd</sup> weeks of age on 0.5% citric acid (B). Feed conversion was found higher ( $P<0.05$ ) during 0-5 weeks of age in treatment B when compared to those of the other treatments. Carcass characteristics were not affected by dietary treatments. The highest mortality was found in treatment A and treatment C due to excessive hot. It may be concluded that use of 0.5% citric acid in the diet of broilers may have better performance in respect of live weight gain and feed conversion.

**Keyword:** Acetic acid, Citric acid, Broiler performance

### Introduction

The feed additives have a number of beneficial effects like control of pathogenic microorganisms and enhance the growth of beneficial microorganisms (Shane, 1999). Antibiotics possess these beneficial effects but their use in the poultry industry has been intensively controversial because of the development of bacterial resistance and potential consequences on the human health. So, the alternatives to antibiotics are researched. Among these compounds, organic acids are promising alternatives (Hyden, 2000). Health of the gut is one of the major factors governing the performance of birds and thus, the economics of poultry production (Samik *et al.*, 2007) and the profile of intestinal microflora play an important role in gut health. Dietary organic acids and their salts are able to inhibit microbial growth in the food and consequently to preserve the microbial balance in the gastrointestinal tract. In addition, by modifying intestinal pH, organic acids also improve the solubility of the feed ingredients, digestion and absorption of the nutrients (Patten and Waldroup, 1988; Owings *et al.*, 1990; Skinner *et al.*, 1991; Adams, 1999).

Both, the feed industry and the poultry production sector, still suffer from huge losses due to the contamination of food with pathogenic bacteria and their related impacts in the animal, such as lower weight gains or even increased mortality. Poultry performance and feed efficiency are closely interrelated with the qualitative and quantitative microbial load of the host animal, including the load in the alimentary tract and in the environment. Organic acids like citric acid and acetic acid have been used in diets due to their positive effect on health and growth of bird. More recently, the ban on antibiotics as a growth promoter in the European Union and the resulting pressures on meat exporters around the world, have increased interest in organic acids to attain performance improvements in growing swine and poultry. As the uses of organic acids are becoming more acceptable to feed manufacturers,

poultry producers and consumers, there is a growing interest in substituting them for antibiotic as growth promoters (Calsen, 1999). Citric acid and acetic acid are used as the substitute of antibiotic growth promoters in many countries of the world (Estieve *et al.*, 1997). But use of citric acid and acetic acid as substitute of antibiotic growth promoter in Bangladesh is a new phenomenon. The effects of citric acid and acetic acid as substitute of antibiotic have not yet been evaluated much under Bangladesh condition. Therefore, the present study was undertaken to evaluate the effect of feeding citric acid and acetic acid and their combination on the performance of broiler and to determine its economic impact in broiler production.

## Materials and Methods

The experiment was conducted with 108 one day old straight run broiler chicks (Hubbard Classic) for a period of 35 days. The chicks were randomly divided into 4 equal treatment groups (A, B, C and D) each having 27 chicks. Each treatment was subjected to 3 equal replications of 9 chicks each. The diets were formulated with commonly available feed ingredients is shown in Table 1. The dietary treatments were A (control diet) and B, C and D were supplemented with 0.5% citric acid, 0.5% acetic acid and 0.5% citric acid + 0.5% acetic acid respectively with drinking water. Dry mash feed was supplied on *adlibitum* basis. Fresh clean drinking water was made at all the times. Adequate sanitary measures were taken during the experimental period. The birds were housed in cages of 120cm × 76cm.

**Table 1. The ingredients and chemical composition of control diet**

Ingredients	Amount in the diet (%)
Maize	51.75
Soybean meal	42.00
Soybean oil	4.00
Salt	0.25
Di- Calcium Phosphate	0.50
Calcium premix	1.00
Vitamin-Mineral premix	0.75
DL-Methionine	0.15
Choline Chloride 60%	0.05
<b>Chemical composition</b>	<b>Amount (%)</b>
Dry matter	85.00
Crude protein	22.21
Crude fibre	5.88
Ether extract	1.76
Nitrogen free extract	48.41
Ash	6.96
ME(kcal/kg DM)*	3241.22

Calculated according to Wiseman (1987)

At the age of day 4 and 14, birds were vaccinated against Infectious Bursal Disease (IBD) using Bursine-2. Chicks were also vaccinated with B.C.R.D.V on 8<sup>th</sup> day. To evaluate the treatment effect, weight gain, feed conversion ratio, mortality, dressing percentage, economy of broiler production were recorded and calculated. At the end of experiment, two birds from each treatment were selected randomly to record the dressing yield, organs weight and cut up parts. Feed samples were analyzed for dry matter (DM), crude protein (CP), ether extract

(EE), crude fiber (CF), nitrogen free extract (NFE), and total ash by following the method of AOAC (1990). Duplicate samples were analyzed and the average value was taken. Collected and calculated data were analyzed for analysis of variance (ANOVA). The significant differences were identified by LSD among the treatments.

## Results and Discussion

### Body weight gain

Effect of organic acids inclusion in broiler ration on live weight gain is presented in Table 2. Significant ( $P<0.05$ ) difference in body weight of birds among the groups were observed at all ages. Birds on treatment C showed lower ( $P<0.05$ ) weight gain than control group (A) and treatment B showed the highest ( $P<0.05$ ) weight gain. Treatment D showed improved growth when administration of both citric acid in diets and acetic acid in water was done together. The growth retardation in treatment C seemed to be a consequence of a depressed water intake induced by application of acetic acid in water. The result is in agreement with Schuhmacher *et al.* (2006), who found lower weight gain. Highest weight gain on 0.5% citric acid agreed with previous findings of Shen-HuiFang *et al.* (2005); Denil *et al.* (2003) and Stipkovits *et al.* (1992) where improved weight gain was observed with administration of citric acid in diets at 0.3, 0.5 and 0.7%, respectively. The results contradict with the findings of previous researchers Pinchasov *et al.* (2000) where depressed weight gain was observed with application of acetic acids in diets.

**Table 2. Live weight gains at different weeks in different treatments**

Age (weeks)	Dietary treatments				SEM	Level of Sig.
	A	B	C	D		
Initial weight	46.2 ± 0.59	46.1 ± 0.56	46.4 ± 0.28	46.1 ± 0.33	0.37	
1	84.5 ± 4.37 <sup>a</sup>	88.7 ± 1.64 <sup>a</sup>	73.1 ± 3.32 <sup>b</sup>	70.3 ± 2.03 <sup>b</sup>	2.61	*
2	210.3 ± 12.19 <sup>a</sup>	219.8 ± 11.48 <sup>a</sup>	175.0 ± 4.37 <sup>b</sup>	176.6 ± 7.84 <sup>b</sup>	7.72	*
3	286.4 ± 5.89 <sup>ab</sup>	299.2 ± 15.19 <sup>a</sup>	247.6 ± 3.71 <sup>c</sup>	281.0 ± 3.82 <sup>b</sup>	6.10	*
4	396.8 ± 13.63 <sup>b</sup>	447.2 ± 24.06 <sup>a</sup>	368.5 ± 28.51 <sup>b</sup>	372.2 ± 9.62 <sup>b</sup>	16.78	*
5	403.5 ± 23.97 <sup>c</sup>	451.5 ± 16.72 <sup>b</sup>	424.7 ± 26.21 <sup>bc</sup>	507.9 ± 2.90 <sup>a</sup>	16.79	*
0-4	977.9 ± 22.27 <sup>b</sup>	1054.0 ± 20.86 <sup>a</sup>	864.2 ± 24.77 <sup>c</sup>	900.1 ± 13.85 <sup>c</sup>	17.11	*
0-5	1381.4 ± 9.88 <sup>b</sup>	1506.3 ± 4.16 <sup>a</sup>	1289.0 ± 19.86 <sup>c</sup>	1408.0 ± 19.21 <sup>b</sup>	12.25	*
Final body weight (g)	1427.6 ± 10.15 <sup>b</sup>	1552.4 ± 4.12 <sup>a</sup>	1335.4 ± 20.09 <sup>c</sup>	1454.1 ± 19.09 <sup>b</sup>	12.09	*

A= Control diet; B= Control diet + 0.5% citric acid; C= Control diet + 0.5% acetic acid; D= Control diet + 0.5% citric acid + 0.5% acetic acid; ±= Standard deviation; SEM= Standard Error Mean; Figure having different superscript in the same row differ significantly ( $P<0.05$ ); \* = 5% level of significance; NS= non significant.

### Feed intake

The average feed intake of birds fed on different diets is shown in Table 3. It is evident that average feed intake was lower in treatment A and higher in treatment D and differed statistically ( $P<0.05$ ) only at 2<sup>nd</sup> and 3<sup>rd</sup> week of age. These results contradict with the finding of previous researchers (Darko *et al.*, 1991; Frigg *et al.*, 1983 and Stipkovits *et al.*, 1992) where depressed feed intake was observed. During (0-4 weeks) of age feed intake was the highest in treatment B (1787.00g) and the lowest in treatment C (1681.00g). During (0-5 weeks) of age feed intake was the highest in treatment B (3118.56g) and the lowest in treatment A (2913.16g) but difference was non-significant ( $P>0.05$ ). The lower feed intake in treatment C was accompanied by retarded growth to be the consequence of depressed water intake by the application of acetic acid in water.

**Table 3. Feed intake (g) at different weeks of experimental birds in different treatments**

Age (weeks)	Dietary treatments				SEM	Level of Sig.
	A	B	C	D		
1	122.3 ± 1.42	124.1±1.0	120.0 ± 6.19	117.0 ± 0.64	2.88	NS
2	320.9 ± 15.94 <sup>a</sup>	325.9±5.70 <sup>a</sup>	292.2 ± 3.02 <sup>b</sup>	301.1±2.94 <sup>b</sup>	7.05	*
3	514.5 ± 14.61 <sup>bc</sup>	532.9 ± 7.86 <sup>ab</sup>	499.1 ± 22.62 <sup>c</sup>	561.1 ± 14.70 <sup>a</sup>	13.02	*
4	738.2 ± 39.24	804.4 ± 39.49	769.9 ± 58.60	801.9 ± 13.26	33.56	NS
5	1217.2 ± 105.67	1331.2± 84.39	1348.8±166.65	1320.8± 12.64	87.58	NS
0-4	1696.0 ± 70.46	1787.0± 42.69	1681.0± 72.30	1781.0± 30.81	46.62	NS
0-5	2913.2 ± 142.90	3118.6±126.99	3029.9±237.88	3101.9±106.8	125.20	NS

A= Control diet; B= Control diet + 0.5% citric acid; C= Control diet + 0.5% acetic acid; D= Control diet + 0.5% citric acid + 0.5% acetic acid; ±= Standard deviation; SEM= Standard Error Mean; Figure having different superscript in the same row differ significantly (P<0.05); \*= 5% level of significance; NS= non significant.

### Feed conversion

The effect of organic acid supplementation on feed conversion is presented in Table 4. It is evident that FCR differ significantly (P<0.05) among treatments at all ages. Better feed conversion was found in treatment B and lower in treatment C during 0-5 weeks of age. The highest feed conversion on the administration of citric acid was in agreement with the findings of Afsharmanesh *et al.* (2005) who found higher feed conversion with the administration of citric acid in poultry.

**Table 4. Feed conversion ratios of birds in different treatments**

Age (weeks)	Dietary treatments				SEM	Level of significance
	A	B	C	D		
0-2	1.5 ± 0.02 <sup>b</sup>	1.5 ± 0.06 <sup>b</sup>	1.7 ± 0.09 <sup>a</sup>	1.7 ± 0.08 <sup>a</sup>	0.281	*
0-3	1.7 ± 0.03 <sup>b</sup>	1.6 ± 0.01 <sup>b</sup>	1.8 ± 0.04 <sup>a</sup>	1.9 ± 0.05 <sup>a</sup>	0.630	*
0-4	1.7 ± 0.03 <sup>b</sup>	1.7 ± 0.01 <sup>b</sup>	1.9 ± 0.04 <sup>a</sup>	1.9 ± 0.04 <sup>a</sup>	0.630	*
0-5	2.1 ± 0.12 <sup>b</sup>	2.1 ± 0.08 <sup>b</sup>	2.4 ± 0.16 <sup>a</sup>	2.2 ± 0.01 <sup>ab</sup>	0.089	*

A= Control diet; B= Control diet + 0.5% citric acid; C= Control diet + 0.5% acetic acid; D= Control diet + 0.5% citric acid + 0.5% acetic acid; ±= Standard deviation; SEM= Standard Error Mean; Figure having different superscript in the same row differ significantly (P<0.05); \*= 5% level of significance

### Carcass characteristics

#### Organs weight

It is evident from the Table 5 that dressing percentage for treatment A, B, C and D were 54.98, 55.40, 51.90 and 56.80 % respectively which did not differ significantly (P>0.05). The results are in well agreement with the previous findings (Kahraman *et al.*, 1997) where no significant effect was observed. The highest (56.7%) value for carcass yield was found in treatment D and the lowest (51.8%) value was found in treatment C.

**Table 5. Carcass characteristics of broilers in different treatments**

Parameters	Dietary treatments				SEM	Level of Sig.
	A	B	C	D		
Live weight (g)	1146.7±152.0	1300.0±0.00	1150.0±132.0	1150.0 ± 150.0	102.74	NS
<b>Organ weights (% live weight)</b>						
Killed weight	89.6 ± 3.56	89.9 ± 3.67	89.2 ± 2.63	89.9 ± 1.42	2.43	NS
Shank weight	4.6 ± 0.66	4.9 ± 0.44	4.9 ± 0.43	4.9 ± 0.13	0.373	NS
Head weight	3.9 ± 0.45	3.9 ± 0.00	3.5 ± 0.38	3.2 ± 0.31	0.851	NS
Giblet weight	7.3 ± 0.60	7.2 ± 0.80	7.3 ± 0.81	7.2 ± 0.59	0.578	NS
Skin weight	16.2 ± 1.12	15.1 ± 1.24	15.7 ± 1.32	14.8 ± 0.19	0.873	NS
Visceral weight	7.8 ± 0.95	7.7 ± 0.39	7.3 ± 0.35	7.7 ± 0.36	0.466	NS
Carcass yield	54.98 ± 2.24	55.4 ± 2.66	51.9 ± 4.63	56.8± 3.08	2.67	NS

A= Control diet; B= Control diet + 0.5% citric acid; C= Control diet + 0.5% acetic acid; D= Control diet + 0.5% citric acid + 0.5% acetic acid; SEM= Standard Error Mean; NS= non significant.

In dietary treatment D the dressing yield was improved by about 3.46 % when compared with the control group. This result did not agree with previous findings of Garcia *et al.* (2000) who found decrease carcass yield. The increased dressing yield on dietary treatment D might be due to increasing live weight on 0.5% citric and acetic acid. The result partially agreed with Sapra and Mehta (1990), who found increased edible meat yield with increasing body weight. Per cent giblet weight was not affected by dietary treatments. Weight of shank in different treatments did not differ significantly ( $P>0.05$ ) among different dietary treatments. Head weight was similar in all treatments. Skin and feather weight per cent did not differ significantly ( $P>0.05$ ) among treatments.

### Economy of broiler production

The production cost of broiler in different dietary treatments is shown in Table 6. The feed cost was highest (Tk.1559.49) in treatment D and the lowest (Tk. 1464.48) in treatment A (control group). The addition of 0.5% citric acid in diets and 0.5% acetic acid in water resulted in an increased feed cost in treatment B, C and D against treatment A (Control group). Cost per kg live weight of broiler was the highest in treatment C (Tk. 69.21) followed by treatment A (60.35), B (60.70) and D (63.77) respectively. Net profit per kg live broiler was the highest in treatment A (Tk. 19.70) followed by treatment B (Tk 19.30), C (Tk.10.70) and D (Tk.16.20) respectively. The highest total net profit was observed in dietary treatment B and the lowest total net profit was observed in treatment C as compare to treatment A (Control group).

**Table 6. Production cost of broiler in different dietary treatments**

Parameters	Dietary treatments			
	A	B	C	D
Total feed cost	1464.48	1568.04	1523.30	1559.49
Total chick cost	378	378	378	378
Management cost <sup>1</sup>	484.59	537.35	502.68	566.20
Total acid cost	-	60.31	90.77	156.27
Total production cost (Tk.)	2327.07	2543.73	2494.75	2503.69
Total production cost (Tk.)/kg live weight	60.35	60.70	69.21	63.77
Total sale priced (Tk.)	3084.48	3352.32	2883.60	3140.64
Total net profit (Tk.)	757.48	808.59	388.25	636.95
Net profit (Tk.)/kg live weight	19.70	19.30	10.70	16.20

A= Control diet; B= Control diet + 0.5% citric acid; C= Control diet + 0.5% acetic acid; D= Control diet + 0.5% citric acid + 0.5% acetic acid.

NB-cost of feed calculated based on ingredients and test substances added.

<sup>1</sup> Management cost assuming approximately 33% of total cost (except chick cost).

It may be concluded that supplementation of 0.5% citric acid (B) in the diet showed positive effect on live weight, feed intake and feed conversion efficiency with no detrimental effect on carcass characteristics.

## References

- Adams, C. 1999. Poultry and dietary acids. *Feed Int.*, 20 (19): 1370-1372.
- Afsharmanesh, M. and Pourreza, J. 2005. Effects of calcium, citric acid, ascorbic acid, vitamin D<sub>3</sub> on the efficacy of microbial phytase in broiler starters fed wheat based diets. *Internat J. Poult Sci.*, 4(6): 418-424.
- Callesen, J. 1999. Commercial feed products and feed mixes for piglets. *Lansuvalget for Svin, Report*. 409, January: 20-21.
- Darko, S., Jelena, P.G. Zdravka, B. and Kanizaj, V. 1991. Effect of therapeutic level of tiamutin on higher toxicity of monensin in weaned pigs. *Veterinarski arhiv.*, 61(2): 67-82.
- Denli, M., Okan, F. and Celik, K. 2003. Effect of dietary probiotic, organic acid and antibiotic supplementation to diets on broiler performance and carcass yield. *Pak. J. Nutr.*, 2(2): 89-91.
- Estieve, G.E., Brufau, J. and Perez, A.K. 1997. Bioefficacy of enzyme preparations containing beta-glucanase and xylanase activities in broiler diets based on barley or heat in combination with flavomycin. *Poult Sci.*, 76: 1726-1737.
- Frigg, M., Broz, J. and Weber, G. 1983. Compatibility studies of ionophore antococcidials with various antibiotics and chemotherapeutics in broiler chicks. *Arch. Geflugelk.*, 47(5): 213-220.
- Garcia, R.G., Ariki, J., Moraes, V.M.B., Kronka, S.N., Borges, S.A., Murata, L.S. and Campos, V.A. 2000. Isolated or combined action of organic acids and growth promoter in broiler rations. *Revista Brasileira de Ciencia Avicola.*, 2(2): 149-154.
- Hyden, M. 2000. Protected acid additives. *Feed Internat.*, 7: 14-16.
- Kahraman, R., Abas, I., Bostan, K., Tanor, A., Kocabagli, N. and Alp, M. 1997. Effects of organic acids and yeast culture on performance, ileum pH and Enterobacteriaceae population of broilers. *Pendik Veteriner Mikrobiyoloji Dergisi*. 28(2): 171-180.
- Owings, W.J., Reynolds, D.L., Hasiak, R.J. and Ferket, P.R. 1990. Influence of dietary supplementation with *Streptococcus faecium* M-74 on broiler body weight feed conversion; carcass characteristics and intestinal microbial colonization. *Poult. Sci.*, 69: 1257-1264.
- Patten, J.D. and Waldroup, P.W. 1988. The use of organic acids in broiler diets. *Poult. Sci.*, 67: 1182- 1187.
- Pinchasov, Y. and Elmalich, S. 2000. Broiler chick responses to anorectic agent: dietary acetic and propionic acids and the digestive system. Faculty of Agriculture, Hervev University of Jerusalem, Rehovot, Israel. (<http://www.ncbi.nlm.nih.gov/sites/entrez>).
- Samik, K.P., Gobinda, H., Manas, K.M. and Gautam, S. 2007. Effect of organic acid salt on the performance and gut health of broiler chicken. *J. Poult. Sci.*, 44: 389-395.
- Sapra, K.I. and Mehta, R.K. 1990. A comparative study on addition Livol (Hervol growth promoter) and some chemical growth promoters in the diet of broiler chicks. *Indian J. Anim. Prod. and Manag.*, 6: (2), 115-118.
- Schuhmacher, K., Bafundo, W., Islam, K.M.S., Aupperle, H., Glaser, R., Schoon, H.A. and Groppe, J.M. 2006. Tiamutin and semduramicin: Effects of Simultaneous Administration on Performance and Health of Growing Broiler Chickens. *Poult. Sci.*, 85: 441-446
- Shane, S. 1999. The antibiotics issue. *Poult. Internat.*, 38: 46- 50.
- Shen-HuiFang, Han-ChuiWang and Du-BingWang. 2005. Effects of citric acid on production performance of Three Yellow chicken. *China Poult.*, 27(16): 14-15.
- Skinner, J.J., Izat, A.L. and Waldroup, P.W. 1991. Research Note: Fumaric acid enhances performance of broiler chicken. *Poult. Sci.*, 70: 1444- 1447
- Stipkovits, L., Csiba, E., Laber, G. and Bruch, D.G.S. 1992. Simultaneous treatment of chickens with salinomycin and tiamutin in feed. *Avian Dis.*, 36: 11-16.
- Wiseman J. 1987. Feeding of Non-ruminant Livestock. Butterworths and Co.Ltd., London. p. 10.