

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search http://ageconsearch.umn.edu aesearch@umn.edu

Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.

Genotype-environment interaction and stability analysis of four fine rice varieties

P. L. Biswas¹, U. K. Nath², S. Ghosal¹ and A. K. Patwary²

¹Plant Breeding Division, BRRI, Gazipur and ²Department of Genetics and Plant Breeding, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh. Email: ujjalnath@gmail.com

Abstract

Genotype-environment interactions through different stability parameters and performance traits of four fine rice genotypes were studied. The traits were; days to 50% flowering, days to maturity, plant height (cm), number of effective tillers per hill, panicle length (cm), number of fertile grains per panicle, number of sterile grains per panicle and yield (t/ha) in four fine rice genotypes across nine environments along with experimental farm of Genetics and Plant Breeding department, Bangladesh Agricultural University (BAU) Mymensingh. Significant differences were observed for genotypes, environments and genotypes–environment interaction. Stability analysis after Eberhat and Russell's model suggested that the genotypes used in study were more or less responsive to environmental changes. Most of the genotypes performed better in Comilla. BAU125 was found stable for effective tillers per hill and comparatively less sensitive to other genotypes in panicle length and number of sterile grains per panicle. BR5 was stable for days to maturity and plant height whereas Kalizira performed better than other genotypes for fertile grains per panicle. In general, only the genotype BAU125 was found stable for effective tillers per hill, panicle length and lowest number of sterile grains per panicle.

Keywords: Fine rice, Genotype-environment interaction, Stability

Introduction

Rice feeds more than half of world population. The demand of fine rice is always high although it possesses lower yield potentiality than HYV (High Yielding Variety). Yield is a complex character which is dependent on a number of other characters and is highly influenced by many genetic factors as well as environmental fluctuation. Genotypes which can adjust its phenotypic state in response to environmental fluctuations in such a way that it gives maximum stable economic return can be termed as well "buffered" or stable (Allard and Bardshaw, 1964). In plant breeding programme, potential genotypes are usually evaluated in different environments before selecting desirable ones. For stabilizing yield it is necessary to identify the stable genotypes suitable for wide range of environments. To identify such genotypes, G-E interactions are of major concern for a breeder, because such interactions confound the selection of the superior cultivars by altering their relative productiveness in different environments (Eagles and Frey, 1977). Varietals study in yield with respect to wide range of environments is one of the most desired properties of genotypes to fit the crop under available cropping pattern. So, wider adaptability and stability are prime consideration in formulating efficient breeding programme. Stability analysis is a good technique for measuring the adaptability of different crop varieties to varying environments (Morales et al., 1991). Therefore, present study was undertaken to estimate the G-E interactions through stability parameters and performance of some traits of four fine rice genotypes across environments and to identify the suitable genotypes for future breeding programme.

Materials and Methods

The experiment was conducted at nine different environments i.e. nine districts viz. Dhaka (Env.-01), Comilla (Env.-02), Sylhet (Env.-03), Barisal (Env.-04), Bogra (Env.-05), Dinajpur (Env.-06), Rajshahi (Env.-07), Jessore (Env.-08) along with experimental farm, Department of Genetics and Plant Breeding, Bangladesh Agricultural University (BAU), Mymensingh (Env.-09), Bangladesh. Four fine rice genotypes viz.BAU125, BR5, Kalizira and Kataribhog were used in the study. The experiment was laid out in a randomized complete block design with three replications in each environment. The unit plot size in a replication was 6m in length and 5 m in width, 25 rows of 1000 hills per genotypes keeping row to row distance 20 cm and hill to hill distance 15 cm. In all the experimental locations, same operational procedures were followed. Seeds were sown on 7th July, 2000 and 32 days old seedlings were

Genotype-environment interaction and stability analysis of rice varieties

transplanted on 9th August, 2000. Fertilizer was applied @100: 90: 70 kg ha⁻¹ of urea, TSP and MOP. One-third of urea, whole amount of TSP and MOP were applied as basal dose during final land preparation. Remaining two-third of the urea was applied in two splits, first at 21 days after transplanting and the other before panicle initiation. Normal cultural practices were followed as and whenever necessary. Data on eight characters were considered and presented in this paper. The genotypes– environment interactions and stability analysis were done following the method suggested by Eberhart and Russell (1966) also quoted by Singh and Chaudhury (1985) and Dabholkar (1992).

Results and Discussion

The results of the combined analysis of variance after Eberhart and Rusell's model are presented in Table 1. The mean squares for genotypes and environments for all the traits under study were highly significant ($p \le 0.01$); suggesting the existence of considerable variation among genotypes as well as environments. The genotype-environment interaction when tested against pooled error was found significant for all the characters, indicating that all the traits were highly influenced by the change in environments leading to extension of analysis for estimating stability parameters. The linear portion of G-E interaction was highly significant ($p \le 0.01$) for all the characters except three such as plant height, number of effective tillers per hill and number of fertile grains per panicle. Thus the prediction of the genotypes in the environments appeared to be feasible for five characters under study.

Table 1. Combined analysis of variance (MS) for eight characters in a genotype-environmen interaction study in fine rice after Eberhart and Russell model											
Source of variation		Days to	Days to	Plant	No.	Panicle	No. of fertile	No. sterile	Yield		

Source of variation	d.f.	Days to 50% flowering	Days to maturity	Plant height (cm)	No. effective tillers/hill	Panicle length (cm)	No. of fertile grains / panicle	No. sterile grains / panicle	Yield (t/ha)
Genotypes	3	24.705**	52.889**	300.005*	4.434**	8.298**	2873.045**	367.959**	0.434**
Environments	8	10.881**	87.815**	528.609**	23.328**	68.008**	3586.151**	531.017**	2.927**
Gen x Env.	24	3.026**	4.417**	265.325*	2.545**	7.679**	895.631**	196.806**	0.257**
Env. +Gen.x Env.	32	1.663**	8.422**	110.379**	2.580**	7.587**	522.757**	93.452**	0.308**
Env.(Linear)	1	29.012**	234.168**	1409.800**	62.192**	181.368**	9563.168**	1415.972**	7.804**
Gen. x Env.(Linear)	3	1.979**	5.052**	46.965	0.983	9.567**	178.219	300.188**	0.230**
Pooled deviation	28	0.653**	0.721**	70.766**	0.622	1.168**	236.8**	24.068**	0.049**
Pooled error	72	0.245	0.163	2.622	0.492	0.195	40.826	5.032	0.002

*p≤0.05 and ** p≤ 0.01

The mean performance of the individual genotype along with their stability parameters ($p_{i,b}$, and s^2d_i) for days to 50% flowering is presented in Table 2. From the environmental mean it was observed that Env.-02 was the most favorable environment. Analysis of the stability parameters of the individual genotypes indicated that all the genotypes showed individual linear response. However, two of the genotypes BR5 and Kataribhog showed combined b_i and s^2d_i sensitivity which suggested that both linear and non linear components were responsible for significant genotype–environment interaction. Genotype BAU125, Kalizira and Kataribhog had the negative phenotypic indices, therefore, they were desirable genotypes for this character. Genotypes BAU125 and Kalizira showed b_i value higher than unity with non-significant s^2d_i values, indicating its suitability only for favorable environment condition. Estimation of all of three stability parameters, it appeared that there was no stable genotype against this character.

Days to maturity

The stability parameters along with the average days to maturity of different genotypes and environment are presented in the Table 2. From the environmental mean it was observed that Env.-5 was the favorable and Env.-3 was the unfavorable one. Analysis of stability parameters of individual genotypes indicated that three genotypes (BAU125, Kalizira and Kataribhog) out of four showed combine b_i and s^2d_i sensitivity suggesting either linear or non-linear component alone or their cumulative effects were responsible for significant genotype-environment interaction. Genotype BR5 showed b_i value higher than unity with non-significant s^2d_i indicating their suitability for in better environmental condition. All the genotypes except BR5 had the negative phenotypic indices, therefore, they were desirable genotypes for this character. Estimating all the stability parameters it appeared that genotype BR5 was relatively stable in changing environment.

2

Table 2. Different important agronomical traits including average days to 50% flowering, days to maturity, panicle length, number of sterile grains per panicle and yield: phenotypic index (Pi), regression (bi) and deviation from regression (S²di) for four rice genotypes evaluated under nine different environments using Eberhart and Russell model

	Characters																		
		Da	ays to 50%	flowering			Days to maturity							Panicle length					
Env.	BAU125	BR5	Kalizira	Katari	Mean	Env.	BAU125	BR5	Kalizira	Katari	Mean	Env.	BAU125	BR5	Kalizira	Katari	Mean	Env.	
						Index						Index						Index	
Env.1	102.8	108.5	104.5	101.3	104.3	1.85	142	148	144	141	143.8	2.46	22.9	22.2	23.9	24.3	23.3	0.36	
Env.2	100.7	104.2	101.5	100	101.6	-0.86	139	142	137	134	138	-3.37	23.2	21.4	24.7	24.3	23.4	0.4	
Env.3	101.2	106	103.5	103.7	103.6	1.14	144	149	146	144	145.8	4.38	26.2	21.5	22.7	24.7	23.8	0.78	
Env.4	101.2	103.3	102.2	103.2	102.5	0.02	141	143	139	139	140.5	-0.87	22.4	16.9	13.1	16.7	17.3	-5.7	
Env.5	99.2	104.8	101.2	101.7	101.8	-0.73	137	141	136	137	137.7	-3.62	24.7	22.5	24.3	22.3	23.5	0.48	
Env.6	100.2	104	108.8	102	102	-0.44	141	146	143	139	142.3	0.88	25.1	22.4	22.3	22.3	23.1	0.06	
Env.7	99.3	103.5	101.2	102	101.5	-0.94	139	144	137	139	139.6	-1.62	26.4	25.7	25.6	27.7	26.4	3.38	
Env.8	100.5	102.8	102.5	102.7	102.1	-0.32	141	144	139	140	141	-0.37	23.2	22.2	23.7	21.7	22.7	-0.29	
Env.9	101.5	105.2	101.8	102.3	102.7	0.27	142	147	144	141	143.5	2.13	23.3	21.9	24.1	24.7	23.5	0.51	
Mean	100.72	104.7	102.24	102.09	102.4		140.7	144.9	140.6	139	141.4		24.16	21.86	22.7	23.18	22.98		
Pi	-1.72	2.26	-0.19	-0.35			-0.7	3.52	-0.78	-2.04			1.18	-1.12	-0.26	0.2			
bi	1.039**	1.532**	1.122**	0.363*			0.728**	1.004**	1.353**	0.943**			0.405*	0.910**	1.478**	1.207**			
S ² di	0.006	0.714*	-0.157	0.901*			0.107*	-0.054	0.457**	1.100**			1.24	0.290*	1.647**	0.764**			

*p≤0.05 and ** p≤ 0.01

Table 2. Continued

		Characters												
Env.		Numb	per of sterile g	grains per pan	icle		Yield							
	BAU125	BR5	Kalizira	Katari	Mean	Env. Index	BAU125	BR5	Kalizira	Katari	Mean	Env. Index		
Env.1	12.2	30.2	25.5	28.2	24	-0.02	3.4	3	2.8	3.1	3.1	0.29		
Env.2	12.1	12.9	31.6	26.1	20.7	-3.36	3.5	4.1	3.5	3.1	3.5	0.77		
Env.3	12	28.2	18.7	26.9	20.5	-3.57	2.9	2.5	2.4	2.7	2.6	-0.16		
Env.4	25.8	20.3	20.6	21	21.9	-2.11	3.7	3.5	3.4	3.1	3.4	0.64		
Env.5	14.6	25.4	24.4	17.6	20.5	-3.57	2.5	2.4	1.7	2.2	2.2	-0.64		
Env.6	11.5	30.6	31.1	34.2	26.9	2.82	3.4	2.4	2.3	2.8	2.7	-0.55		
Env.7	12.8	15.4	16.2	17.2	15.4	-8.65	3.2	2.3	2.4	2.8	2.7	-0.7		
Env.8	17.3	33.7	24.5	35.7	27.8	3.76	2.3	1.5	2.2	2.2	2.1	-0.03		
Env.9	17.3	34	34.8	68.8	38.7	14.69	2.7	2.4	2.4	2.9	2.6	-0.08		
Mean	15.07	25.64	25.27	30.21	24.04		3.07	2.66	2.56	2.74	2.8			
Pi	-8.98	1.59	1.22	6.16			0.31	-0.1	-0.2	-0.02				
bi	0.145	0.853**	0.698**	2.304**			0.888**	1.433**	1.064**	0.616**				
S ² di	18.389**	27.270**	15.238**	15.658**			0.038**	0.076**	0.047**	0.022**				

*p≤0.05 and ** p≤0.01

Genotype-environment interaction and stability analysis of rice varieties

Plant height

The average plant height (cm) of individual genotype over nine environments with their regression coefficient and deviation from regression are presented in Fig. 1a. From the environmental mean it was observed that Env.-2 was the most favorable than others for producing lowest plant height. The lowest plant height produced by the genotype BR5. The phenotypic indices of Kalizira, BR5 and BAU125 were positive, therefore, they are desirable genotypes for this character. The two genotypes Kalizira and Kataribhog showed combined bi and s^2d_i sensitivity suggesting both linear and non-linear component were responsible for genotype-environment interaction. The genotype BAU125 and BR5 showed significant s^2d_i values which indicate that they were more affected by the environmental fluctuations i.e. performance of these genotypes over environments were unpredictable. In all the three parameters, genotype BR5 appeared to be stable than other genotypes that having bi value near to unity with positive phenotypic indices.

Effective tillers

The average number of effective tillers per hill of the individual genotype along with p_{i} , bi and s^2d_i are presented in Fig. 1b. From the environmental mean it was observed that Env.-7 had the highest 13.5 tillers per hill and Env.2 had the lowest 9.3 tillers per hill. Analysis of stability parameters of individual genotype indicated that none of the genotypes showed combine b_i and s^2d_i sensitivity and as a result there is no cumulative effect of linear and non-linear components for significant genotype-environment interaction. BAU125 and BR5 had the positive phenotypic indices, therefore, they were desirable for this character. BR5, Kalizira and Kataribhog had the higher b_i value with non-significant s^2d_i indicating their suitability for growing better environmental conditions. Taking all the three stability parameters, genotype BAU125 was considered stable and desirable genotype for number of effective tillers per hill.

Panicle length

The stability parameters along with the average length of panicle (cm) in different genotypes and environment are presented in Table 2. From the environmental mean it was observed that Env.-7 was the favorable and Env.-4 was the most unfavorable one. Analysis of stability parameters of individual genotypes indicated that all the four except one (BAU125) showed combined b_i and s^2d_i sensitivity suggesting either linear or non-linear component alone or their cumulative effects were responsible for significant genotype-environment interaction. Genotypes BAU125 and Kataribhog had the positive phenotypic indices, therefore, they were desirable for this character. The significant b_i value with non-significant s^2d_i of BAU125 genotype indicating its suitability for growing in better environmental condition. Considering all the three parameters (p_i b_i and s^2d_i) the genotype BAU125 was found relatively stable among all the genotypes.

Fertile grains per panicle

The average number of fertile grains per panicle of individual genotype over nine environments with their regression co-efficient and deviation from regression are presented in Fig. 1c. From the environmental mean it was observed that Env.-7 was the most favorable and Env.-9 was the unfavorable one. The highest number fertile grains per panicle (143.18) were produced by BR5 and the lowest number of fertile grains per panicle (104.45) was produced by BAU125. The phenotypic indices of BR5 and Kalizira were positive, therefore, they were desirable genotypes for this character. Analysis of the stability parameters of the individual genotypes indicates that the genotypes BAU125, BR5 and Kalizira showed combined bi and s²d_i sensitivity suggesting both linear and non-linear components were responsible for genotype-environment interaction. Kalizira had higher bi value with non –significant s²d_i indicating its suitability for growing in better environmental condition. In all the stability parameters genotype Kalizira was found comparatively stable among others for this character.

4





Fig. 1a. Average plant height and phenotypic index (Pi), regression (bi) and deviation from regression (S²di) for four rice genotypes evaluated under nine different environments using Eberhart and Russell model



Fig. 1b. Average number of effective tillers per hill and phenotypic index (Pi), regression (bi) and deviation from regression (S²di) for four rice genotypes evaluated under nine different environments using Eberhart and Russell model



Fig. 1c. Average number of fertile grains per panicle and phenotypic index (Pi), regression (bi) and deviation from regression (S²di) for four rice genotypes evaluated under nine different environments using Eberhart and Russell model

Sterile grains per panicle

The mean performance of the individual genotype along with their stability parameters for the number of sterile grain per panicle are presented in Table 2. From the environmental mean it was observed that Env.-7 was the most favorable for the production less number of sterile grains per panicle (15.4). BAU125 genotype produced the lowest average number of sterile grains per panicle (15.07) where Kataribhog showed the highest number of sterile grains per panicle (30.21). Three genotypes BR5, Kalizira and Kataribhog showed combined b_i and s^2d_i sensitivity which suggested that both linear and non-linear components were responsible for significant genotype-environment interaction. The phenotypic indices of BR5, Kalizira and Kataribhog were positive, therefore, they were undesirable genotypes for this character. Genotype BAU125 showed non-significant b_i and significant s^2d_i value indicating it produced higher number of sterile grains per panicle only for highly favorable environment otherwise not. Evaluating all the three parameters (p_i b_i and s^2d_i) the genotype BAU125 was found relatively stable for less number of sterile grains per panicle produced among all the genotypes.

Yield

The average yield performance of the individual genotype along with $p_i b_i$ and $s^2 d_i$ are presented in Table 2. From the environmental mean it was observed that Env.-2 had the highest mean yield (3.5 t/ha) and the lowest one was obtained in Env.-8 (2.1 t/ha). This indicated that Env.-2 was the most favorable one and the majority of the genotypes had the capacity to exploit that environment to confer the highest yield. Over all the environments, BAU125 showed the highest yield (3.7 t/ha) and Kalizira showed the lowest (2.56 t/ha). Analysis of the stability parameters of the individual genotype indicated that all the four genotype showed combined bi and $s^2 d_i$ sensitivity suggesting both linear and non-linear component were responsible for significant genotype-environment interaction. Genotypes BR5, Kalizira and Kataribhog showed negative phenotypic indices while BAU125 was positive. BAU 125, BR5 and Kalizira and Kataribhog showed significant $s^2 d_i$ values, thus prediction of their performance over environment would not be authentic. In all the three parameters ($p_i b_i$ and $s^2 d_i$) it was observed that all the genotypes were sensitive to environments i.e. none of the genotypes were found stable against this character.

Biswas et al.

The study suggests that genotype BR5 may be selected for stability in days to maturity and plant height. Genotype BAU125 may be selected for effective tillers per hill, panicle length and lowest number of sterile grains per panicle .These materials can be used in fine rice breeding program as a source of genes for stability.

References

- Allard, R.W. and Bradshaw, A.D. 1964. Implication of Genotype-Environment interactions in applied plant breeding. Crop Sci. 4: 503-508.
- Dabholkar, A.R., 1992. Elements of Biometrical Genetics. Concept Publishing Company. New Delhi, India. pp: 338-359.
- Eagles, H.A. and Frey, K.J. 1977. Repeatability of the stability variance parameter in oats. Crop Sci. 17:253-256.
- Eberhart, S.A. and Russel, W.L. 1966. Stability parameters for comparing varieties. Crop. Sci. 6: 36-40.
- Morales, A.C., Paragas, A.A. and Carangal, V.R. 1991. Phenotypic stability for grain yield in mungbean (*Vigna radiata* (L) Wilczek). FLCG Newsletter. 16: 12-15.
- Singh, R.K. and Chaudhary, B.D. 1985. Biometrical Methods in Quantitative Genetic Analysis (rev. Ed.). Kalyani Publishers, New Delhi, India. pp: 252-269.