



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

Starch and flour extraction and nutrient composition of tuber in seven cassava accessions

M. S. A. Fakir, M. Jannat, M.G. Mostafa and H. Seal¹

Department of Crop Botany, ¹Department of Agricultural Chemistry, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh, Email: fakirmsa@gmail.com

Abstract

Cassava (*Manihot esculenta* Crantz) roots (tubers) are used as staple food. Starch extracted from tubers is widely utilized as raw materials in industries. Dry matter (DM) content, starch and flour extraction and proximate composition were investigated in seven cassava accessions (Coc-A₁, Kh-A₂, Cow-A₃, Sa-A₄, Me-A₅, Va-A₆ and Sy-A₈) in 2010-2011. Leaf DM varied from 20.51% in Me-A₅ to 29.01% in Sy-A₈; that of stem from 27.24% in Va-A₆ to 32.10% (average of Sy-A₈, Me-A₅ and Sa-A₄); and that of tuber from 37.30% in Kh-A₂ to 45.26% in Sy-A₈. Starch was extracted by blending chopped tuber followed by decantation. Tubers were sliced, sun dried and milled into flour. Tuber starch content (fresh wt. basis) varied between 15.04% in Sy-A₈ and 24.97% (average of Coc-A₁ and Me-A₅); that of peel from 4.54% in Va-A₆ to 5.85% in Coc-A₁. Crude protein varied from 1.80% (average of Kh-A₂, Cow-A₃ and Sy-A₈) to 4.53% in Va-A₆. Crude fiber content varied from 1.95% (average of Sa-A₄ and Coc-A₁) to 4.27% in Cow-A₃. Cyanogens present in cassava plant escape as hydrogen cyanide (HCN) during harvesting and processing. Variation for HCN existed and it was 140.95 mg/kg fresh tuber (average of Sy-A₈ and Coc-A₁) to 546.0 mg/kg fresh tuber in Va-A₆. There was no detectable HCN in the extracted flour and starch. It may be concluded that genetic variation for DM, starch, protein and HCN existed in seven cassava accessions, and Coc-A₁ may be a better one due to its lower HCN, higher DM and starch content.

Keywords: Cassava roots, DM, HCN, Proximate composition

Introduction

Cassava (*Manihot esculenta* Crantz) is a perennial, subtropical, woody shrub, grown as annual and is valued for its underground starchy tubers (roots) (Grace, 1977; Purseglove, 1988; Islam *et al.*, 2008). Cassava is the important source of energy as staple food for more than 500 millions people in Africa, Latin America and Asia (Hillocks, 2002). Tuber of cassava is also used as raw materials in the garment, bakery, food and Pharmaceutical industries (Bokanga *et al.*, 1994; IITA, 2011; Fakir *et al.*, 2012). Cassava root can be consumed raw as a snack or just after boiling like sweet potato. Cassava is a mesophyte and can be grown in fallow high land, hill slope, rice field dyke ('ail') and other unutilized high lands. It can be grown in poor soils and produce fairly good yield where other crops almost fails and provide food security (Siritunga *et al.*, 2004). In Bangladesh, tribal people of Madhupur, Garo Hill, Natrokona and Chittagong Hill Tract region locally grows and eat cassava. There is a good potentiality of cultivating and introducing this crop to other areas in Bangladesh. Private entrepreneurs like 'Rahman and Chemical Co.' and 'Bharsha group' have also been producing cassava in the country through contact farmers in Comilla, Tangail and Chittagong Hill Tracts and utilising tuber for starch, glucose and glucose syrup production. There is no reliable statistics of areas and production in Bangladesh.

Dry matter (DM) determination in different genotypes of cassava is important since nutrition and energy calculation is based on magnitude and nature of DM content. Higher DM content naturally would provide greater yield once partition of the same is increased to the economic part. Boerboom (1978) and Islam *et al.* (2008) investigated dry weight of plant parts and observed that leaf, stem and tuber contained 30-60% DM. Cassava tuber contains about 70% moisture, 20-30% carbohydrate, 1.0-1.8% crude protein, 1.5-3.5% crude fiber, 0.35-0.45% fat and 8-28 mg HCN/kg of dry mass (Purseglove, 1988; Charles *et al.*, 2005). Cassava plant parts are rich source of vitamins and minerals. Though tuber is the main product of cassava plant, its young branch and leaf is also edible both for human and animal (Fakir *et al.*, 2010). Tuber is the main source of starch and minerals; leaf is the rich source of protein, vitamins and minerals. Cassava flour (10-30%) in combination with wheat flour is used in bread industry to reduce pressure on wheat (Grace, 1977). Therefore, cassava flour along with mixture with wheat flour can be used to make nutritious food and food products. Its tuberous root contains (fresh weight basis) 30-40% dry matter (DM) and 25-30% starch. Nutritionally, cassava contains potassium, iron, calcium, vitamin A, folic acid, sodium, vitamin C, vitamin B-6, and protein (Montagnac *et al.*, 2009). Nutritional quality especially protein can be added in composite flours (tapioca) in cassava-soya, cassava-peanut bread. The protein content of 'tapioca' is about 10% and the macaroni is nearly twice as nutritious as rice (Grace, 1977). Supplemental use of cassava flour and starch in combination with wheat flour may reduce pressure and thus would enhance food security in Bangladesh.

Cassava flour is used for making bread and other products like Chips, jam, jelly and chutneys. Flour extraction from cassava tuber depends on reduction of moisture. Cassava drying aims at reducing its water content to less than 15%. IITA (2005) cites four factors that influence drying of cassava (chips flour and starch). These are temperature, airflow, humidity and tumbling frequency. Starch from cassava tuber can be extracted in two simple, wet and sun dry, methods. In the wet method, fresh tubers are blended with adequate water and the filtrate is decanted to starch. The starch obtained is then dried in the sun and is ready for use. In wet method fresh tubers are certainly be used for flour production immediately after harvest, within 72 hrs, since fresh roots deteriorates quickly. In sun dry methods, fresh tubers are chopped into small pieces and are sun dried until desirable moisture content is achieved. The dried cubes are ground and milled into flour. The methods of flour extraction, therefore, also influence the degree of detoxification.

All parts of the cassava plants contain cyanogens that are hydrolyzed to hydrocyanic acid (HCN) that escapes into the air during harvesting and processing (Bokanga *et al.*, 1994; Fakir *et al.*, 2009). At harvest of cassava tubers, the amount of HCN acids varies from harmless to lethal. The lethal dose of HCN is 50mg/kg fresh weight. Hydrolysis of glycoside by the enzyme can be accelerated by soaking the roots in water, by crushing or cutting them or by heating. HCN content of cassava roots could thus be reduced to a great extent by treating in such ways (Fakir *et al.*, 2010). Although literatures on botany, physiology, nutrient content and HCN is available at home and abroad (Grace, 1977; Alves, 2002; Anonymous, 2004; Charles *et al.*, 2005; Islam *et al.*, 2008; Fakir *et al.*, 2009, 2010, 2011), there is scanty information in Bangladesh on methods of flour and starch production and proximate composition of cassava tuber (Jannat, 2011; Fakir *et al.*, 2012). So it was necessary (i) to estimate the dry matter content of plant parts; (ii) to investigate the methods of flour/starch extraction; and (iii) to determine the proximate composition of tuber of seven cassava accessions.


Materials and Methods

Plant establishment: The experiment was conducted at the field laboratory of the Department of Crop Botany, Bangladesh Agricultural University, Mymensingh during November, 2010 to October, 2011. Healthy and uniform size (about 15cm with 5 to 6 nodes) 8 months old stem cuttings of seven cassava accessions (Coc-A₁; Kh-A₂; Cow-A₃; Sa-A₄; Me-A₅; Va-A₆ and Sy-A₈) were planted at 1m x 1m spacing in 16 m² plots. Stem cuttings were, collected from previous experiment, planted horizontally at 10 to 12 cm soil depth with one cutting per hole. The cuttings were treated for ten minutes with the fungicides 'Bavistin' @ 5 g/10 L before planting. Seven treatments (seven accessions) were planted in a randomized complete block design (RCBD) with four replications.


Crop sampling and harvesting: After 10 months, four plants excluding border ones were randomly selected from each plot and harvested, cleaned and brought to the laboratory for further observations.

Peeling and dry matter content: After harvest, tubers were peeled and cut into small cubes (2 cm long and 0.5 cm wide) for dry matter content. The dry matter contents of the different plant parts of cassava were determined by Benesi *et al.* (2004). The samples were sun dried following oven drying (65°C for 72 h) until constant weight and cooled in desiccators and weighed immediately. The drying and weighing steps were repeated until consecutive constant weights were achieved.

Starch extraction: Starch was extracted using the wet method described by Benesi *et al.* (2004). Fresh tubers were washed, peeled, chopped into approximately 1 cm cubes and then pulverized in a high-speed blender (Model KING, Osaka, Japan) for 5 min. The pulp was suspended in ten times its volume of water, stirred for 5 minutes and filtered using double fold cotton cloth. The filtrate was allowed to stand for 2 h for the starch to settle and the top liquid was decanted and discarded. Water was added to the sediment and the mixture was stirred again for 5 minutes. Filtration was repeated as before and the starch from the filtrate was allowed to settle. After decanting the top liquid, the sediment (starch) was sun dried for 24 h and stored. The starch of peel was also extracted by the same procedure. Following is flow diagram of starch extraction.



Dry matter (DM) content of starch, pulp and flour: DM content of starch was determined by the same procedure described earlier. The pulp (broken tissue plus fibers in moistened condition) after filtration was also sun dried and kept at 10-12% moisture for making flour. Flour was obtained after grinding dried pulp followed by sieving. Following is the flow diagram for flour production:



Tubers proximate composition: Crude protein (CP), crude fiber (CF), crude fat (EE), total carbohydrate (NFE) and ash were determined by AOAC (1990) and hydrogen cyanide (HCN) by Grace (1977).

Data collection and analysis: The data were compiled and analyzed to find out the statistical significance. The means for all the collected data were calculated and the mean differences were evaluated by Duncan's New Multiple Range Test (Gomez and Gomez, 1984).

Results and Discussion

Dry matter (DM) content of plant parts

Peel: The dry matter content of peel varied significantly ($P \leq 0.05$) among the seven cassava accessions and it was greater in Coc-A₁, Kh-A₂ and Me-A₅ (average of 24.58%) than in the Sa-A₄ and Cow-A₃ (average of 21.72%), Va-A₆ (18.99%) and Sy-A₈ (16.62%) (Table 1). **Tuber:** The tuber DM content varied significantly ($P \leq 0.05$) from 37.30% in Kh-A₂ to 45.26% in Sy-A₈ (Table 1). It was greater in Sy-A₈ (45.26%) than Va-A₆ (43.79%), Cow-A₃, Coc-A₁, Sa-A₄ and Me-A₅ (average of 41.08%) and Rh-A₂ (37.30%). On an average, tuber contained 41.53% DM i.e. 58.47% water in the current research. **Stem:** Stem DM content was greater in Sa-A₄, Sy-A₈ and Me-A₅ (average of 32.10%) than Coc-A₁ (29.97%) Cow-A₃ (28.72%) and Va-A₆ (27.24%) (Table 1). **Petiole:** The DM content of petiole varied significantly ($P \leq 0.05$) with greater magnitude in Cow-A₃ (31.91%) than in Va-A₆ (29.32%), Kh-A₂ (28.02%), Me-A₅ and Coc-A₁ (average of 26.42%) and Sa-A₄ (21.66%) (Table 1). **Leaf lobe:** Leaf lobe DM varied from 20.51% in Me-A₅ to 29.01% in Sy-A₈. The dry matter content of Leaf lobe was significantly greater in Sy-A₈ (29.01%) than Coc-A₁ and Cow-A₃ (average of 26.97%), Va-A₆ (24.36%), Kh-A₂ and Sa-A₄ (average of 22.50%) and Me-A₅ (20.51%) (Table 1).

Table 1. Dry matter (DM) content of plant parts of seven cassava accessions

Accessions	Dry Matter content (%)				
	Peel	Tuber	Stem	Petiole	Leaf lobe
Coc-A ₁	24.79 a	41.38 c	29.97 b	26.81d	27.28 b
Kh-A ₂	24.20 a	37.30 d	29.10 bc	28.02 c	22.79 d
Cow-A ₃	21.85 b	40.88 c	28.72 c	31.91 a	26.67 b
Sa-A ₄	21.59 b	40.58 c	32.64 a	21.66 e	22.22 d
Me-A ₅	24.75 a	41.49 c	31.80 a	26.04 d	20.51 e
Va-A ₆	18.99 c	43.79 b	27.24 d	29.32 b	24.36 c
Sy-A ₈	16.62 d	45.26 a	31.88 a	29.12 bc	29.01 a
Mean	21.83	41.53	30.19	27.55	24.69

In a column, numbers followed by different letters differ significantly at $P \leq 0.05$ by DMRT

Genetic variations for DM content of plant parts existed in the current research. Howeler and Cadavid (1983), Boerboom (1978) and Islam *et al.* (2008) also noted genetic variation for DM in different cassava accessions. Mean DM content was greater in tuber (41.53%) than stem (30.19%), petiole (27.55%), leaf lobe (24.69%) and peel (21.83%) (Table 1). Boerboom (1978) and Islam *et al.* (2008) also observed similar results in fewer accessions. Howeler and Cadavid (1983), however, observed higher DM of tuber (50-60%). This difference could be due to variation in genotype and locations between the experiments.

Dry matter (DM) content of starch, pulp and flour

Starch: The DM content of starch of cassava varied significantly ($P \leq 0.05$) among the seven accessions and it was greater in Sa-A₄ (88.11%) than in Kh-A₂ and Me-A₅ (average of 86.90%), Cow-A₃ (84.97%), Coc-A₁ (83.12%) (Table 2). **Pulp:** The DM content of pulp varied significantly ($P \leq 0.05$) among the seven cassava accessions and it was greater in Coc-A₁, Cow-A₃ and Kh-A₂ (average of 90.76%) than in Sa-A₄ and Me-A₅ (average of 86.2%), Va-A₆ (85.16%) and Sy-A₈ (81.72%) (Table 2). **Flour:** The DM of flour varied from 80.38% in Va-A₆ to 89.35% Kh-A₂ (Table 2). It was greater in Kh-A₂ (89.35%) than in Coc-A₁ (88.08%), Cow-A₃ and Me-A₅ (average of 85.88%) and Sa-A₄ (84.77%) (Table 2).

Table 2. Dry matter content of tuber, starch, pulp and flour in seven cassava accessions

Accessions	Dry matter content (%)		
	Starch	Pulp	Flour
Coc-A ₁	83.12 d	92.41 a	88.08 b
Kh-A ₂	86.73 b	90.95 a	89.35 a
Cow-A ₃	84.97 c	90.93 a	85.97 c
Sa-A ₄	88.11 a	85.35 bc	84.77 d
Me-A ₅	87.08 ab	87.05 b	85.80 c
Va-A ₆	82.35 de	85.16 c	80.38 e
Sy-A ₈	81.92 e	81.72 d	80.48 e
Mean	84.90	87.65	84.98

In a column, numbers followed by different letters differ significantly at $P \leq 0.05$ by DMRT

Preparation of cassava products such as flour, pulp, starch, chips, pellets etc. depends on DM content. Therefore, DM determination of starch, pulp and flour is important. Generally, 10-12% moisture i.e. 88-90% DM is good for flour production (IITA, 2005). However, in the current research flour moisture content varied from 10-20%. It could be due to improper drying in the sun. For short term storage flour must be dried within the stipulated range.

Peeling ratio, starch and flour yield, and HCN content of tuber

Peeling ratio: Weight of the peel to tuber i.e. peeling ratio of cassava tuber varied significantly ($P \leq 0.05$) among the seven cassava accessions and it was greater in Coc-A₁ and Sa-A₄ (average of 19.62%) than in Me-A₅ (17.79%), Kh-A₂ (16.42%), Sy-A₈ (14.40%) and Va-A₆ (12.91%) (Table 3). **Starch content:** Tuber starch content varied from 15.04% in Sy-A₈ to 25.05% in Coc-A₁. It was significantly greater in Coc-A₁ and Me-A₅ (average of 24.97%) than Kh-A₂ and Cow-A₃ (average of 21.73%), Sa-A₄ and Va-A₆ (average of 17.19%) and Sy-A₈ (15.04%) (Table 3). **Peel starch content:** The Starch content of peel of cassava and it was greater in Coc-A₁ (5.85%) than in Cow-A₃, Sa-A₄ and Me-A₅ (average of 5.32%), Kh-A₂ (5.19%), Sy-A₈ (4.85%) and Va-A₆ (4.54%) (Table 3). **Flour Yield:** Flour content varied from 101.89 g/kg (average of Kh-A₂, Cow-A₃ and Va-A₆) to 201.86 g/kg (in Sy-A₈). It was significantly greater in Sy-A₈ (201.86 g/kg) than Coc-A₁, Sa-A₄ and Me-A₅ (average of 162.44 g/kg) and Kh-A₂, Cow-A₃ and Va-A₆ (average of 101.89 g/kg) (Table 3).

Tuber HCN content: The HCN content of fresh tuber varied significantly ($P \leq 0.05$) among the seven cassava accessions (Table 3). It was greater in Va-A₆ (546.0 mg/kg fresh weight) than in Sa-A₄ (324.0 mg/kg fresh weight), Cow-A₃ (244.4 mg/kg fresh weight), Kh-A₂ and Me-A₅ (average of 215.6 mg/kg fresh weight) and Sy-A₈ and Coc-A₁ (average of 140.95 mg/kg fresh weight) (Table 3).

Table 3. Peeling ratio, starch and flour yield, and hydrogen cyanide (HCN) contents (fresh wt. basis) in seven cassava accessions

Accessions	Peeling ratio (peel to tuber)	Starch % (fresh wt.)		Flour yield (g/kg)	Tuber HCN (mg/kg)
		Tuber	Peel		
Coc-A ₁	19.89 a	25.05 a	5.85 a	162.03 b (16.20) ⁺	137.20 e
Kh-A ₂	16.42 c	21.56 b	5.19 c	107.59 c (10.76)	217.01 d
Cow-A ₃	17.35 bc	21.91 b	5.30 b	098.14 c (09.81)	244.42 c
Sa-A ₄	19.35 a	17.61 c	5.27 b	166.35 b (16.63)	324.06 b
Me-A ₅	17.79 b	24.90 a	5.54 b	158.94 b (15.89)	214.28 d
Va-A ₆	12.91 e	16.78 c	4.54 e	99.94 c (09.99)	546.01 a
Sy-A ₈	14.40 d	15.04 d	4.85 d	201.86 a (20.18)	144.70 e
Mean	16.87	20.41	5.22	142.12 (14.20)	261.04

In a column, numbers followed by different letters differ significantly at $P \leq 0.05$ by DMRT, +: Figures within parenthesis indicate percentage flour yield

Genetic variation for tuber mean starch content (15.04-25.05 %) was observed in the current study (Table 3). This supports the results of Sajeev *et al.* (2003) who also observed variations in starch (20-25 %) in fresh tuber. Of the 100 g fresh tuber, there were 20.41 g starch, 14.20 g flour and the rest, 65.39 g water on mean weight basis (Table 3). On fresh weight basis, current cassava starch yield was greater (7.25 t/ha, mean of seven accessions, data not shown) than potato and sweet potato (6 t/ha), rice (3.2 t/ha), corn (2.5 t/ha) and wheat (2.4 t/ha) (Jannat, 2011). Variation also existed for HCN content in tuber. Lethal dose of HCN is 50 mg/kg fresh weight. But HCN could almost be removed by proper processing. HCN produced during harvesting, chopping, boiling, drying, blending and other postharvest operations are almost removed (Bokanga *et al.*, 1994; Fakir *et al.*, 2009, 2010). The flour and starch thus produced was free of toxicity since HCN was not detected there.

Tuber proximate composition

Crude protein: The crude protein was greater in Va-A₆ (4.53%) than in Coc-A₁ and Sa-A₄ (average of 2.92%), Kh-A₂, Cow-A₃ and Sy-A₈ (average of 1.80%) (Table 4). **Crude fiber:** Crude fiber content significantly varied from 1.95% in Coc-A₁ and Sa-A₄ to 4.27% in Cow-A₃ (Table 4). It was significantly greater in Cow-A₃ (4.27%) than Coc-A₁ and Sa-A₄ (average of 1.95%). **Crude fat (Etheral extract, EE):** EE content of cassava varied significantly ($P \leq 0.05$) and it was greater in Va-A₆ (1.87%) than Sa-A₄ (0.80%) (Table 4). **Ash:** The ash or mineral content of cassava varied significantly ($P \leq 0.05$) among the seven accessions and it was greater in Me-A₅ and Va-A₆ (average of 5.53%) than Cow-A₃ and Kh-A₂ (average of 3.65%) (Table 4). **Total Carbohydrate (Nitrogen free extract, NFE):** The NFE content of cassava tuber varied significantly ($P \leq 0.05$) among the seven accessions (Table 4). It was greater in Kh-A₂ (91.10%) than in Me-A₅ (87.92%) and Va-A₆ (84.35%) (Table 4).

Cassava tuber is used as staple food, feed and vegetables (Grace, 1977; Charles *et al.*, 2005; Fakir *et al.*, 2010). Proximate composition such as crude protein, crude fiber, crude fat, ash and NFE of tuber is important. (Hang *et al.*, 2007). Genetic variation for proximate composition in seven accessions was observed and therefore, selection for genotype with good nutrients is possible. High protein (4.53%) in Va-A₆ may not be useful since it also contained high HCN (546 mg/kg fresh wt.). However, nature and duration of processing like chopping, threshing, boiling etc. reduces as much as 95% HCN (Montagnac *et al.*, 2009; Fakir *et al.*, 2009, 2010). Therefore, processing is important in this case. Genetic variations for DM content of plant parts; and flour, starch and HCN content of tuber existed. There is, therefore, opportunity for selection of improved genotype(s) for higher DM, flour and starch yield. Further, selected genotypes could be used as parent(s) to develop high yielding varieties. It appears that the Coc-A₁ would be a better accession based on DM and nutrients contents.

Table 4. Proximate composition of tuber of seven cassava accessions

Accessions	Proximate composition of tuber (%)				
	Crude protein	Crude fiber	Crude fat	Ash	NFE [†]
Coc-A ₁	2.83b	1.66c	1.52ab	4.55ab	89.42bc
Kh-A ₂	1.92c	2.45bc	1.18abc	3.84b	91.18a
Cow-A ₃	1.68c	4.27a	1.44abc	3.47b	89.12bc
Sa-A ₄	3.01b	2.24c	0.80d	4.04ab	89.90ab
Me-A ₅	2.56bc	2.88bc	1.33abcd	5.47a	87.92c
Va-A ₆	4.53a	3.60ab	1.87a	5.60a	84.35d
Sy-A ₈	1.81c	2.65bc	0.86cd	4.41ab	90.26ab
Mean	2.62	2.82	1.29	4.48	88.88

In a column, numbers followed by different letters differ significantly at $P \leq 0.05$ by DMRT, [†]: Nitrogen free extract

References

- Alves, A.A.C. 2002. Cassava biology and physiology, *In*: Cassava biology, production and utilisation. eds. Hillocks, R.J., Thresh, J.M., Beltotti, A.C., p. 67-90, CAB Intl. Oxford
- Anonymous, H. 2004. Hydrogen cyanide and cyanides: human health aspects, Concise Int. Chem. Assessment Doc. 61, World Health Org.
- AOAC (Association of Official Agricultural Chemists) 1990. 15th ed. Association of official analytical chemistry, Washington DC.
- Benesi, I.R.M., Labuschagne, M.T., Dixon, A.G.O. and Mahungu, N.M. 2004. Stability of native starch quality parameters, starch extraction and root dry matter of cassava genotypes in different environments, *J. Sci. Food Agric.*, 84: 1381-1388.
- Boerboom, B.W.J. 1978. A model of dry matter distribution in cassava (*Manihot esculenta* Crantz). *Netherlands J. Agric. Sci.*, 26: 267-277.
- Bokanga, M., Ekanayake, I.J., Dixon, A.G.O. and Proto, M.C.M. 1994. Genotype-environment interaction for cyanogenic potential in cassava. *Acta. Hort.*, 375:131-139.
- Charles, A.L., Sriroth, K. and Tozou-chi, H. 2005. Proximate composition, mineral contents, hydrogen cyanide and phytic acid of 5 cassava genotypes. *Food Chem.*, 92(4): 615-620.
- Fakir, M.S.A., Mostafa, M.G. and Seal, H.P. 2009. Food security in Bangladesh: Evaluation of cassava (*Manihot esculenta*) morphotypes based on hydrogen cyanide acid toxicity and protein content of tuber. Poster presented *In* Natl. Symp. Climate change, plant protection & food security interface, Assoc. Advan. Plant protection, BCKV, Kalyani, India, 17-19 Dec., 2009, p. 59
- Fakir, M.S.A., Mostafa, M.G. and Seal, H.P. 2010. Food security in Bangladesh: Selection, nutritional status evaluation of processing technique of cassava strains for use as a potential human and animal food. Poster presented *In* Intl. Conf. "Food security during challenging times" Univ. Putra Malaysia, Selangor, Malaysia, 5-7 July, 2010. p. 218-220.
- Fakir, M.S.A., Talukder, M.H.R., Mostafa, M.G. and Rahman, M.S. 2011. Debranching effect on growth and yield in cassava. *J. Agrofor. Environ.*, 5(1): 1-5.
- Fakir, M.S.A., Mostafa, M.G., Jannat, M., Islam, F. and Seal, H.P. 2012. Dry mass content of plant parts, flour extraction and nutrient contents of tuber of cassava accessions. Abst. *In* Souvenir, 3rd Intl. Seed Conf., 'Quality seed and food security under changing climate', Seed Sci Soc. Bangladesh, Bangladesh Agric. Univ., Mymensingh, Bangladesh, 8-10 Feb, 2012. p. 41.
- Gomez, K.A. and Gomez, A.A. 1984. Statistical procedure for agricultural research. 2nd ed. John Wiley and Sons, New York. p. 680.
- Grace, M.R. 1977. Cassava processing. *Plant production and protection series*, No. 3. FAO, Rome.
- Hang, P., Preston, K.A. and Nassar, J.H. 2007. Proximate composition, hydrogen cyanide of two cassava genotypes. *Phyto Chem.*, 56 (5): 215-217.
- Hillocks, R.J. 2002. Origin, distribution and economic importance: Cassava in Africa. *In*: Cassava biology, production and utilisation. eds. Hillocks, R.J., Thresh, J.M., Beltotti, A.C., CAB Intl. Oxford, p. 40-45,
- Howeler, R.H. and Cadavid, L. F. 1983. Accumulation and distribution of dry matter and nutrients during a 12 month growth cycle of cassava. *Field Crops Res.*, 7: 123-139.
- IITA (International Institute of Tropical Agriculture), 2011. Research highlights, P.M.B. 5320, Oyo state, Ibadan, Nigeria. p. 12.
- Islam, A., Islam, A.T.M.T., Mostafa, M.G. and Fakir, M.S.A. 2008. Effect of branch number on growth and yield in two cassava morphotypes. *Bangladesh J. Agric.*, 1(1): 1-6.
- Jannat, M. 2011. Dry mass content of plant parts, flour extraction and nutrient contents of tuber of cassava (*Manihot esculenta*) accessions, MS Thesis, Dep. Crop Botany, Bangladesh Agric. Univ. Mymensingh.
- Montagnac, J.A., Christopher, R.D. and Tanumi, S.A. 2009. Processing technique to reduce toxicity and antinutrients of cassava for use as staple food. *Comprehensive Rev. Food Sci. Food Safety*, 8:17-27.
- Purseglove, J.W. 1988. *Tropical Crops dicotyledons*, ELBS, Longman, U.K.
- Sajeev, M.S., Moorthy, S.N., Kailappan, R. and Rani, V.S. 2003. Gelatinisation characteristics of cassava starch settled in the presence of different chemicals. *Starch.*, 55: 213-221.
- Siritunga, D., Garzon, D.A., White, W. and Seyre, R.T. 2004. Over-expression of hydroxynitrile lyase in transgenic cassava root accelerates cyanogenesis and food detoxification. *J. Plant Biotechnol.*, 2: 37-43.