



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

Prevalence of helminth parasites in sheep (*Ovis aries*) in Tangail district, Bangladesh

A. Sangma, N. Begum, B. C. Roy and M. O. Gani¹

Department of Parasitology, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh and ¹Bangladesh Livestock Research Institute, Savar, Dhaka, E-mail: bcroy.dvm51@gmail.com

Abstract

Prevalence of helminths of sheep in relation to age, sex, nutritional status, management system and flock size was studied at Tangail district, Bangladesh from July to December 2010 by fecal examination. A total of 190 sheep were examined of which 154 (81.1%) were positive for one or more species of helminth parasites. Seven species of helminths were identified, of them three species were trematodes, namely, *Fasciola gigantica* (8.4%), *Paramphistomum* spp. (44.2%) and *Schistosoma indicum* (3.7%); four species were nematodes, namely, *Bunostomum* sp (19.0%), *Trichuris* spp. (2.1%), Strongyles (62.6%) and *Strongyloides* spp. (9.5%). No cestodes were identified. Prevalence of helminths was significantly ($p < 0.01$) higher in young sheep aged >1-2 year (92.7%) than adult aged > 2 years (83.3%) and lamb aged ≤ 1 year (63.6%). Higher prevalence was recorded in female than in male sheep. In relation to nutritional status and flock size, prevalence of helminths were significantly ($p < 0.01$) higher in poor health and large flock sized animals. It is suggested that helminth infection is widely prevalent in sheep in Tangail district of Bangladesh.

Keyword: Prevalence, Helminths, Sheep, Tangail District

Introduction

Parasitism is of supreme importance in many agro-ecological zones and still a serious threat to the livestock economy worldwide (Vercruysse and Claerebout, 2001). Sheep are known to suffer from various endoparasites of which helminth infection are of great importance. Helminth infections remain one of the major constraints to small ruminant production in tropics (FAO, 1992; Diaz *et al.*, 2000). Infection with gastrointestinal nematodes is regarded as one of the important factor causing production losses of livestock. Tangail district situated in central region of Bangladesh is very prospective for sheep rearing due to its geo- climatic condition. But the sheep rearing is hindered by various problems of which parasitic diseases might be one of the major problems because the mild winter and the long summer including the rainy season create a favorable environmental condition for the survival of various parasites with their intermediate host such as snail (Islam, 1969 and Haq and Sheikh, 1968). There is no precise report available on parasitic diseases of sheep in Bangladesh. Therefore, the present study was conducted to find out the prevalence of helminths of sheep at Tangail district and to determine the effect of age, sex, nutritional condition, management system and flock size of sheep, which would provide a basis for the understanding of different helminth parasites of sheep in formulating the control measures of the parasitic diseases.

Materials and Methods

The study was conducted for a period of six months from July to December 2010 at Tangail district of Bangladesh. Fecal samples of sheep were collected from the different areas of Tangail district and carried to the Laboratory for morphological examination of the parasites with their developmental stages. One hundred and ninety sheep were selected randomly irrespective of age, sex, nutritional status, management system and flock size. The age of the sheep were 5 months and above. Age of the sheep was determined by examining teeth as described by Rahman and Hossain (1997). According to age, sheep were divided into three groups, namely, lamb (≤ 1 year), young (> 1-2 years) and adult (> 2 years). The nutritional condition of sheep was categorized into two groups, namely, poor health and healthy according to eye inspection and body condition (Rahman and Hossain, 1997). The selected sheep were reared either in semi intensive or free range grazing system. The flock size was divided into three groups as small flock consisting of sheep 4-8, Medium flock had 9-12 sheep and the large flock were >12 sheep. After collection of all relevant information, sheep were restrained properly and all possible hygienic measures were maintained and feces were collected directly from the rectum. Some fresh fecal samples were collected from the ground immediately after voiding. A total of 190 samples were collected and

about 5-10 gm of feces were collected from each sheep and kept in separate polythene bag, tied carefully, numbered properly and preserved in 10% formalin. The fecal samples were examined by Stoll's Ova counting technique and identification of egg of different helminths was performed by their characteristic morphological features as described by Soulsby (1982) and Rahman *et al.* (1996). Statistical analyses were carried out by using Statistical Package for Social Sciences (SPSS-11.5) technique using F test to detect the significant differences of mean values of eggs per gram of feces of identified helminths. Odds ratio of different parameters were calculated according to the formula given by Schlesselman (1982).

Results and Discussion

Overall prevalence of helminths in sheep

A total of 190 sheep were examined through fecal sample examination, of which 154 were infected with one or more species of helminths indicating 81.1% overall prevalence. Seven species of helminths were identified; of them three were trematodes, namely, *Fasciola gigantica* (8.4%; Fig. 1), *Paramphistomum* spp. (44.2%; Fig. 2), and *Schistosoma indicum* (3.7%; Fig. 3); four were nematodes, namely, strongyles (62.6%; Fig. 4), hookworm (19%; Fig. 5), *Trichuris* spp. (2.1%; Fig. 6), and *Strongyloides* spp. (9.5%; Fig. 7). No cestodes were detected. It was observed that prevalence of Strongyles (62.6%) was the highest, whereas *Trichuris* spp. (2.1%) was the lowest (Table 1). Egg per Gram of Feces (EPG) was determined. EPG count was the highest in case of strongyles infection (100-1500) followed by *Paramphistomum* spp. (100-1100), hookworm (100-400), *Fasciola gigantica* (100-300), *Strongyloides* spp. (100-200), *Schistosoma indicum* (100-100) and *Trichuris* spp. (100-100). Mean EPG count was also higher in case of strongyles infection (430.3±99.9) followed by *Paramphistomum* spp. (339.3±73.6), hookworm (233.3±23.9), *Fasciola gigantica* (162.5±20.2) and *Strongyloides* spp. (116.7±9.6). A low parasitic burden was in case of *Schistosoma indicum* and *Trichuris* spp. (100±0.0).

Table 1. Overall prevalence of helminths in sheep in Tangail District

Name of parasites	No. of positive cases (N=190)	Prevalence (%)	Egg per gram of feces (EPG)	
			Range	Mean ± SE
<i>Fasciola gigantica</i>	16	8.4	100-300	162.5±20.2 ^d
<i>Paramphistomum</i> spp.	84	44.2	100-1100	339.3±73.6 ^b
<i>Schistosoma indicum</i>	7	3.7	100-100	100±0.0 ^f
Hookworm	36	19.0	100-400	233.3±23.9 ^c
<i>Trichuris</i> spp.	4	2.1	100-100	100±0.0 ^f
Strongyles	119	62.6	100-1500	430.3±99.9 ^a
<i>Strongyloides</i> spp.	18	9.5	100-200	116.7±9.6 ^e
Total	154*	81.1	100-1500	211.7±32.5
P value	0.0055**			

* = Total number of animals infected is less than the summation of individual infection because same animal was infected by more than one type of helminths

N = Number of sheep examined

**= P<0.01, figures in the 5th column having different superscript varies significantly (p<0.01)

Age related prevalence of helminths in sheep

Age of the host had an effect on the prevalence of helminths in sheep. Prevalence of helminths in sheep was significantly higher (p<0.01) in young sheep (87.0%) than in adult (83.3%) and in lamb (70.9%). Young sheep were 3.8 and 1.3 times more susceptible than lambs and adults, whereas adults were 2.9 times more susceptible to helminth infection than lambs. Young sheep were infected by 7 different species of parasites. The adults and lambs were infected by 6 and 4 species, respectively. Prevalence of helminths in lambs was the highest in case strongyles (52.7%) followed by that of *Paramphistomum* spp. (31.0%), hookworm (20%) and *Fasciola gigantica* (7.3%). Prevalence of helminths in young sheep was

the highest in case of strongyles (72.5%) followed by that of *Paramphistomum* spp. (56.5%), hookworm (17.4%), *S. indicum* (7.3%), *Fasciola gigantica* (5.8%), *Strongyloides* spp. (4.4%), *Trichuris* spp. (2.9%). Prevalence of helminths in adults was the highest in case of strongyles (60.6%) followed by that *Paramphistomum* spp. (42.4%), *Strongyloides* spp. (22.7%), hookworm (19.7%), *F. gigantica* (12.1%) and *S. indicum* (3.0%; Table 2).

Table 2. Age related prevalence of helminths in sheep in Tangail District

Age	Name of parasites	No. of positive cases	Prevalence (%)	Egg per gram of feces (EPG)		Odds ratio
				Range	Mean \pm SE	
Lamb (\leq 1year) n=55	<i>Fasciola gigantica</i>	4	7.3	100-200	200.0 \pm 0.0	Young vs Lamb = 3.8
	<i>Paramphistomum</i> spp.	17	31.0	100-800	270.6 \pm 62.3	
	Hookworm	11	20.0	100-200	200.0 \pm 0.0	
	Strongyles	29	52.7	100-1000	455.2 \pm 55.4	
	Sub total	39*	70.9	100-1000	281.4\pm29.4^a	
Young ($>$ 1-2year) n=69	<i>Fasciola gigantica</i>	4	5.8	100-100	100.0 \pm 0.0	Young vs Adult = 1.3
	<i>Paramphistomum</i> spp.	39	56.5	100-1100	389.7 \pm 50.6	
	<i>Schistosoma indicum</i>	5	7.3	100-100	100.0 \pm 0.0	
	Hookworm	12	17.4	100-300	250.0 \pm 33.7	
	<i>Trichuris</i> spp.	2	2.9	100-100	100.0 \pm 0.0	
	Strongyles	50	72.5	100-1500	360.0 \pm 56.5	
	<i>Strongyloides</i> spp.	3	4.4	100-100	100.0 \pm 0.0	
	Sub total	60*	87.0	100-1500	200\pm20.1^c	
Adult ($>$ 2year) n=66	<i>Fasciola gigantica</i>	8	12.1	100-300	175.0 \pm 36.6	Adult vs Lamb =2.9
	<i>Paramphistomum</i> spp.	28	42.4	100-1000	310.7 \pm 53.3	
	<i>Schistosoma indicum</i>	2	3.0	100-100	100.0 \pm 0.0	
	Hookworm	13	19.7	100-400	246.2 \pm 31.3	
	Strongyles	40	60.6	100-1200	500.0 \pm 72.0	
	<i>Strongyloides</i> spp.	15	22.7	100-200	120.0 \pm 10.7	
	Sub total	55*	83.3	100-1200	242.0\pm34.0^b	
P value		0.0006**				

n = No. of sheep examined

* = Total number of animals infected is less than the summation of individual infection because same animal was infected by more than one type of helminthes

**= P<0.01

Figures in the 6th column having different superscript varies significantly (p<0.01).

Sex related prevalence of helminths in sheep

The prevalence of helminth infection was higher in female (83.3%) than in the male (79.3%) sheep. Female sheep were 1.2 times more susceptible than male. In male, prevalence was the highest in case of strongyles (55.7%) followed by that of *Paramphistomum* spp. (47.2%), *Strongyloides* spp. (8.5%), *F. gigantica* (6.6%), *S. indicum* (6.6%), hookworm (5.7%) and *Trichuris* spp. (3.8%). In female, prevalence was the highest in case of strongyles (71.4%) followed by that *Paramphistomum* spp. (40.5%), hookworm (35.7%), *F. gigantica* (10.7%) and *Strongyloides* spp. (10.7%; Table 3).

Nutritional status related prevalence of helminths in sheep

The prevalence of helminth infection was significantly (p<0.01) higher in poor health sheep (84.4%) than that in healthy sheep (76.5%). Poor health sheep were 1.7 times more susceptible than the healthy sheep. In poor health sheep, prevalence was the highest in case of strongyles (82.6%) followed by that of *Paramphistomum* spp. (52.3%), hookworm (27.5%), *F. gigantica* (12.8%), *Strongyloides* spp. (12.8%), *S. indicum* (6.4%) and *Trichuris* spp. (2.8%). On the other hand, in healthy sheep, prevalence was the highest in case of strongyles (35.8%) followed by that of *Paramphistomum* spp. (33.3%), hookworm (7.4%), *Strongyloides* spp. (4.9%), *F. gigantica* (2.5%) and *Trichuris* spp. (1.2%; Table 4).

Table 3. Sex related prevalence of helminths in sheep in Tangail District

Sex	Name of parasites	No. of positive cases	Prevalence (%)	Egg per gram of feces (EPG)		Odds ratio
				Range	Mean \pm SE	
male (n=106)	<i>Fasciola gigantica</i>	7	6.6	100-100	114.3 \pm 14.3	Female vs Male = 1.2
	<i>Paraphistomum</i> spp.	50	47.2	100-1100	346.0 \pm 47.3	
	<i>Schistosoma indicum</i>	7	6.6	100-100	100.0 \pm 0.0	
	Hookworm	6	5.7	100-400	300.0 \pm 44.7	
	<i>Trichuris</i> spp.	4	3.8	100-100	100.0 \pm 0.0	
	Strongyles	59	55.7	100-1500	540.7 \pm 60.7	
	<i>Strongyloides</i> spp	9	8.5	100-200	133.3 \pm 16.7	
	Sub total	84*	79.3	100-1500	233.5\pm26.2^b	
Female (n=84)	<i>Fasciola gigantica</i>	9	10.7	100-300	200.0 \pm 28.8	
	<i>Paraphistomum</i> spp.	34	40.5	100-600	329.4 \pm 39.1	
	Hookworm	30	35.7	100-400	220.0 \pm 16.2	
	Strongyles	60	71.4	100-1000	321.7 \pm 36.9	
	<i>Strongyloides</i> spp.	9	10.7	100-100	100.0 \pm 0.0	
	Sub total	70*	83.3	100-1000	234.2\pm24.2^a	
P value		0.062 ^{NS}				

N = No. of sheep examined

* = Total number of animals infected is less than the summation of individual infection because same animal was infected by more than one type of helminths

NS= Not significant

Table 4. Nutritional status related prevalence of helminths in sheep in Tangail District

Nutritional status	Name of parasites	No. of positive cases	Prevalence (%)	Egg per gram of faeces (EPG)		Odds ratio
				Range	Mean \pm SE	
Poor health (n=109)	<i>Fasciola gigantica</i>	14	12.8	100-300	110.0 \pm 10.0	Poor health vs Healthy = 1.7
	<i>Paraphistomum</i> spp.	57	52.3	100-1100	415.8 \pm 59.5	
	<i>Schistosoma indicum</i>	7	6.4	100-100	100.0 \pm 0.0	
	Hookworm	30	27.5	100-400	228.6 \pm 28.6	
	<i>Trichuris</i> sp.	3	2.8	100-100	100.0 \pm 0.0	
	Strongyles	90	82.6	100-1500	560.0 \pm 82.1	
	<i>Strongyloides</i> spp	14	12.8	100-200	100.0 \pm 0.0	
	Sub total	92*	84.4	100-1500	230.6\pm30.0^a	
Healthy (n=81)	<i>Fasciola gigantica</i>	2	2.5	100-100	250.0 \pm 22.4	
	<i>Paraphistomum</i> spp.	27	33.3	100-500	276.1 \pm 29.7	
	Hookworm	6	7.4	100-200	100.0 \pm 0.0	
	<i>Trichuris</i> spp.	1	1.2	100-100	234.5 \pm 18.8	
	Strongyles	29	35.8	100-600	100.0 \pm 0.0	
	<i>Strongyloides</i> spp.	4	4.9	100-100	376.2 \pm 37.9	
	Sub total	62*	76.5	100-600	125.0\pm13.1^b	
P value		0.0037**				

n = No. of sheep examined

* = Total number of animals infected is less than the summation of individual infection because same animal was infected by more than one type of helminths

** = P < 0.01

Management system related prevalence of helminths in sheep

Management system had a profound effect on the prevalence of helminth infection in sheep. The prevalence of helminth infection was higher in sheep reared in free range grazing system (82.8%) than that in sheep reared in semi-intensive grazing system (79.1%). Free range grazing sheep were 1.3 times more susceptible than semi-intensive grazing sheep. In free range grazing sheep, prevalence was the highest in case of *Paramphistomum* spp. (63.6%) followed by that of strongyles (31.3%), *F. gigantica* (13.1%), hookworm (12.1%), *Strongyloides* spp. (3.0%) and *S. indicum* (6.0%). On the other hand, in semi-intensive grazing sheep, prevalence was the highest in case of strongyles (96.7%) followed by that of hookworm (26.4%), *Paramphistomum* spp. (23.1%), *Strongyloides* spp. (16.5%), *Trichuris* spp. (4.3%), *F. gigantica* (3.3%) and *S. indicum* (1.1%; Table 5).

Table 5. Management system related prevalence of helminths in sheep in Tangail District

Management system	Name of parasites	No. of positive cases	Prevalence (%)	Egg per gram of feces (EPG)		Odds ratio
				Range	Mean \pm SE	
Free Ranging (n=99)	<i>Fasciola gigantica</i>	13	13.1	100-300	114.3 \pm 14.3	Free Ranging Vs Semi-intensive = 1.3
	<i>Paraphistomum</i> spp.	63	63.6	100-1100	253.8 \pm 30.7	
	<i>Schistosoma indicum</i>	6	6.1	100-100	100.0 \pm 0.0	
	Hookworm	12	12.1	100-200	237.5 \pm 20.7	
	Strongyles	31	31.3	100-900	500.0 \pm 48.1	
	<i>Strongyloides</i> spp.	3	3.0	100-100	120.0 \pm 10.7	
	Sub total	82*	82.8	100-1100	220.9\pm24.8^a	
Semi-intensive (n=91)	<i>Fasciola gigantica</i>	3	3.3	100-200	200.0 \pm 28.8	
	<i>Paraphistomum</i> spp.	21	23.1	100-1000	413.3 \pm 51.6	
	<i>Schistosoma indicum</i>	1	1.1	100-100	100.0 \pm 0.0	
	Hookworm	24	26.4	100-400	225.0 \pm 25.0	
	<i>Trichuris</i> spp.	4	4.3	100-100	100.0 \pm 0.0	
	Strongyles	88	96.7	100-1500	319.6 \pm 52.8	
	<i>Strongyloides</i> spp.	15	16.5	100-200	100.0 \pm 0.0	
	Sub total	72*	79.1	100-1500	208.3\pm22.6^b	
P value		0.065 ^{NS}				

n = No. of sheep examined

* = Total number of animals infected is less than the summation of individual infection because same animal was infected by more than one type of helminths

NS= Not significant

Flock size related prevalence of helminths in sheep

The prevalence of helminth infection was varied significantly ($p < 0.01$) among different flock size of sheep. The highest infection of helminth was in sheep of large flock sheep (89.1%) followed by the sheep in small (81.8) and medium flock (73.9%). Sheep of large flock were 2.9 and 1.8 times more susceptible than medium and small flocked sheep, respectively whereas, small flocked sheep were 1.6 times more susceptible in helminth infection than medium flocked sheep. In small flocked sheep, prevalence of helminths was the highest in case of strongyles (72.7%) followed by hookworm (24.2%), *Paramphistomum* spp. (22.7%), *Strongyloides* spp. (12.1%), *F. gigantica* (7.6%), *S. indicum* (3.0%) and *Trichuris* spp. (3.0%). Prevalence of helminths in medium flocked sheep was the highest in case of *Paramphistomum* spp. (52.2%) followed by strongyles (46.4%), hookworm (14.5%), *Strongyloides* spp. (7.3%), *F. gigantica* (5.8%) *Trichuris* spp. (4.4) and *S. indicum* (1.5%). On the other hand, in large flocked sheep, prevalence was the highest in case of strongyles (70.9%) followed by *Paramphistomum* spp. (60.0%), hookworm (18.2%), *F. gigantica* (12.7%), *Strongyloides* spp. (9.1%), *S. indicum* (7.3%) and *Trichuris* spp. (3.6%; Table 6).

Table 6. Flock Size related prevalence of helminths in sheep in Tangail District

Flock size	Name of parasites	No. of positive cases	Prevalence (%)	Egg per gram of feces (EPG)		Odds ratio
				Range	Mean \pm SE	
Small (4-8 no.) n=66	<i>Fasciola gigantica</i>	5	7.6	100-200	100.0 \pm 0.0	Small vs Medium = 1.6
	<i>Paramphistomum</i> spp.	15	22.7	100-900	257.1 \pm 47.7	
	<i>Schistosoma indicum</i>	2	3.0	100-100	100.0 \pm 0.0	
	Hookworm	16	24.2	100-300	255.0 \pm 24.5	
	<i>Trichuris</i> spp.	2	3.0	100-100	100.0 \pm 0.0	
	Strongyles	48	72.7	100-1300	511.6 \pm 56.0	
	<i>Strongyloides</i> spp.	8	12.1	100-200	150.0 \pm 22.4	
	Sub total	54	81.8	100-1300	228.9\pm25.1^a	
Medium (9-12 no.) n=69	<i>Fasciola gigantica</i>	4	5.8	100-300	100.0 \pm 0.0	Large vs Medium=2.9
	<i>Paramphistomum</i> spp.	36	52.2	100-1100	243.2 \pm 30.6	
	<i>Schistosoma indicum</i>	1	1.5	100-100	100.0 \pm 0.0	
	Hookworm	10	14.5	100-400	207.7 \pm 21.1	
	<i>Trichuris</i> spp.	32	4.4	100-1000	100.0 \pm 0.0	
	Strongyles	32	46.4	100-1000	352.4 \pm 64.2	
	<i>Strongyloides</i> spp.	5	7.3	100-100	100.0 \pm 0.0	
	Sub total	51	73.9	100-1100	171.9\pm19.3^c	
Large (>12 no.) n=55	<i>Fasciola gigantica</i>	7	12.7	100-200	176.9 \pm 23.1	Large vs small =1.8
	<i>Paramphistomum</i> spp.	33	60	100-1100	481.8 \pm 64.7	
	<i>Schistosoma indicum</i>	4	7.3	100-100	100.0 \pm 0.0	
	Hookworm	10	18.2	100-200	200.0 \pm 0.0	
	<i>Trichuris</i> spp.	2	3.6	100-100	100.0 \pm 0.0	
	Strongyles	39	70.9	100-1500	423.5 \pm 70.3	
	<i>Strongyloides</i> spp.	5	9.1	100-200	100.0 \pm 0.0	
	Sub total	49	89.1	100-1500	226.0\pm26.4^b	
P value		0.0001**				

n = No. of sheep examined

* = Total number of animals infected is less than the summation of individual infection because same animal was infected by more than one type of helminths.

**= P<0.01

Figures in the 6th column having different superscript varies significantly (p<0.01).



Fig. 1. Egg of *Fasciola gigantica*



Fig. 2. Egg of *Paramphistomum* spp.



Fig. 3. Egg of *Schistosoma indicum*

Fig. 4. Egg of Strongyles



Fig. 5. Egg of Hookworm

Fig. 6. Egg of *Trichuris* spp.Fig. 7. Egg of *Strongyloides* spp.

Parasitism is of supreme importance in many agro-ecological zones and still a serious threat to the livestock economy worldwide (Vercruysse and Claerebout, 2001). Sheep are known to suffer from various endoparasites of which helminth infections are of great importance and remain one of the major constraints to small ruminant production in tropics (FAO, 1992; Diaz *et al.*, 2000).

The present study indicates that about 81.1% sheep were found infected with one or more helminths, namely, *Fasciola gigantica* (8.4%), *Paramphistomum* spp. (44.2%), *Schistosoma indicum* (3.7%), hookworm (19.0%), *Trichuris* spp. (2.1%), strongyles (62.6%) and *Strongyloides* spp. (9.5%). Similar results were reported by Azad *et al.* (1997), Mbap and Chiroma (1998) and Oncel (2000). On the other hand, Mazid *et al.* (2006) in Mymensingh district, Bangladesh and Bezubik *et al.* (1969) in Poland reported 94.7% and 95.0% helminths infection in sheep, respectively. The present finding is higher than the earlier findings of Ijaz *et al.* (2009), Gadahi *et al.* (2009), Asif *et al.* (2007) in Pakistan, Siddiqua (2010) in India, Khan *et al.* (2010) in Pakistan and Krishnappa *et al.* (1992) in India who reported 70.7%, 63.5%, 62%, 53.3%, 44.2% and 37.2% prevalence of helminths in sheep, respectively. The present findings is in agreement with the earlier findings of Siddiqua (2010) in India, Bhuyan (1970) in Bangladesh, Mir *et al.* (2008) in India and Mazyad and Hi (2002) in Egypt who recorded 5.4%, 8.3%, 10.0% and 11.8% fascioliasis in sheep, respectively. However, these results differ from the findings of Paz-silva *et al.* (2003) in Spain, Ratnaparkhi *et al.* (1993) and Bhatia *et al.* (1989) in India who reported higher (83.3%, 81.4% and 25.8%, respectively) rate of *Fasciola* infection in sheep. The findings of the present study also differ from previous findings of Celep *et al.* (1995) in Turkey, Azad *et al.* (1997) in Pakistan, Okafor *et al.* (1988) in Nigeria and Oncel (2000) in Turkey showed, 40.8%, 16.3%, 13.7%, and 4.0% prevalence of *Paramphistomum* spp., respectively. Chaudhri *et al.* (1994) in India and Islam (1969) in East Pakistan reported 33.2% and 10.0% sheep infected with *Schistosoma indicum*. The result of *Trichuris* spp. infection is similar to that of Mostafa *et al.* (1996) in Bangladesh (4.0%) and Hashem and Sayed (1997) in Egypt (4.8%) but different from Abebe and Esayas (2001) in Ethiopia (51.7%), Achi *et al.* (2003) in Cote d' Ivoire (29%) and Oncel (2000) in Turkey (28.0%). This result of *Strongyloides* spp. infection differ from Abebe and Esayas (2001) in Ethiopia (38.0%), Azad *et al.* (1997) in Pakistan (37.3%) and Achi *et al.* (2003) in Cote d' Ivoire (31.0%). The variation among the present and previous findings might be due to the differences in geographical locations, climatic conditions of the study area, rearing and management of sheep, nutritional condition of animals, sample size, and technique of sample examination.

It was revealed that age of the sheep had significant ($p < 0.01$) effect on helminth infections. Young sheep (87.0%) were the most susceptible to infection followed by adults (83.3%) and lambs (63.6%), which supported the findings of Asif *et al.* (2007) who reported the higher prevalence of helminths infection in young animals compared to adult sheep in Pakistan ($p < 0.059$). But this result differed from that of Mazid *et al.* (2006) who reported higher prevalence of helminth parasites in younger (< 1 year) and old (≥ 2 year) than in young ($\geq 1 < 2$ year) sheep in Mymensingh, Bangladesh. The higher rate of infection with *Fasciola gigantica* was in older sheep, which is similar to the earlier reports of Bhuyan (1970) in Bangladesh and Aydenizoz and Yildiz (2002) in Turkey. But the present finding is in contrast to the previous reports of Al-Bayati *et al.* (1991) who observed higher prevalence of fascioliasis in young (21.3%) than adults (12.0%) and lamb (7.3%) in Iraq. The findings of the present study is also similar to Lateef *et al.* (2005) in Pakistan with *Strongylides* infection, who observed higher prevalence in young animals but differ from Oka *et al.* (1999) in Cote d' Ivoire who stated that the highest infection was in animal below 1 year of age. This result is also dissimilar to Mazid *et al.* (2006) who reported older sheep were more susceptible to *Paramphistomum* infection than young (30.4%) in Bangladesh. The exact cause of this variation in the prevalence of helminth infection in different age groups of sheep is due to immunological status of animals. But it may be assumed that the differences in the methodology, management factors may be the reasons of this variation.

It was observed that the prevalence of helminth infections was higher in females (83.3%) than in male (79.3%) sheep. This finding is in agreement with the earlier study of Mazid *et al.* (2006) in Bangladesh who recorded higher prevalence of helminth infection in females than in male sheep (78.6%). But this report is in contrast to the previous report of Asif *et al.* (2007) who observed the sex-wise prevalence of helminths was higher in males than in females in Punjab, Pakistan. This result also differs from Okafor (1988) in Nigeria who observed that prevalence was not related to sex. The reason for higher prevalence of helminth infection in the females cannot be explained exactly but it might be assumed that the alteration in the physiological condition of the females during pregnancy, lactation and parturition as well as stresses leading to immunosuppression may be associated with this phenomenon. Higher level of prolactin and progesterone hormones makes the female individual more susceptible to any infection (Lloyd, 1983).

The present study revealed that nutritional condition of sheep had significant ($p < 0.01$) effect with helminth infection. The higher prevalence of helminth infection was recorded in poor health (75.2%) sheep than that in healthy (64.2%) sheep. The present findings are in agreement with the earlier study of Lapage (1962) who found that malnourished animals are more susceptible to any infection as they are immunocompromised. The present study also agrees with the findings of Etter *et al.* (1999) who reported that in immunocompromised animal the fecundity of parasites is usually increased.

The prevalence of helminth infection was higher in sheep reared in free ranged grazing system (82.8%) than that in sheep reared in semi-intensive system (79.1%) sheep. It was observed that fluke infection was higher in free ranged grazing sheep whereas nematode infection was higher in semi-intensively reared sheep. The present findings can not be compared due to lack of available relevant literatures. It is assumed that regular and direct contact of animals to the contaminated pastures may be associated with this phenomenon.

It was revealed that the prevalence of helminths was significantly ($p < 0.01$) higher in large flocked sheep (89.1%) than in small (81.8) and medium flocked (73.9%) sheep. The influence of flock size on the prevalence of helminths is difficult to explain exactly but it is assumed that crowding of animals, lower feed supplement and management practices may be associated with this variation.

Conclusion

The prevalence of helminth parasites of sheep at Tangail district, Bangladesh was highly susceptible to helminth infections. Age, nutritional status and flock size of sheep significantly ($p < 0.01$) influenced the prevalence of helminth infections. Identification of the helminths down to species through ova detection is very difficult, and it may be better to isolate mature or immature helminths directly from the visceral organ of the sheep. Further study should be carried out to determine the economic losses due to helminthiasis of sheep and to develop effective control measures against it.

References

- Abebe, W. and Esayas, C. 2001. Survey of ovine and caprine gastrointestinal helminthosis in eastern part of Ethiopia during the dry season of the year. *Revue de Medecine Veterinaire*, 152: 379-384.
- Achi, Y.L., Zinsstag, J., Yeo, N., Dea, V. and Dorchie, P. 2003. Epidemiology of parasites of sheep and goats from savannah area in Cote d'Ivoire. *Revue de Medecine Veterinaire*, 154(3): 179-188.
- Al-Bayati, M.M., Al-Khalidi, N.W., and Altaif, K.I. 1991. Fascioliasis in sheep in Mosul, Iraq: the infection rate and the seasonal variation. *Journal of Veterinary Parasitology*, 5(1): 45-47.
- Asif, R.M., Iqbal, Z., Jabbar, A. and Yaseen, M. 2007. Point prevalence of gastrointestinal helminthiasis in ruminants in southern Punjab, Pakistan. *Journal of Helminthology*, 81(3): 323-328.
- Aydenizoz, Z. and Yildiz, K. 2002. Prevalence of liver trematodes in sheep slaughtered in Kirikkale. *Turkiye Parazitoloji Dergisi*, 26(3): 317-319.
- Azad, E., Khan, S.A., Khan, R.M. and Khan, M.M. 1997. Prevalence of gastrointestinal helminth parasites of sheep and goats in Quetta district of the Province of Baluchistan, Pakistan. *Pakistan Journal of Zoology*, 29(3): 259-262.
- Bezubik, B.B., Uadhaya, D.S., and Juyal, P.D. 1969. The helminth parasites of Carpathian sheep in Poland. *Acta parasitology*, Poland, 17(1/19): 25-37.
- Bhatia, B.B., Uadhaya, D.S., and Juyal, P.D. 1989. Epidemiology of *Fasciola gigantica* in buffaloes, goats and sheep in Tarai region of Uttar Pradesh, India. *Journal of Veterinary Parasitology*, 3(1): 25-29.
- Bhuyan, M.A. 1970. A survey of helminths parasiting the liver of domesticated ruminants in East Pakistan. MS Thesis, Department of Parasitology, Bangladesh Agricultural University, Mymensingh.
- Celep, A., Acici, M., Cetindag, M., and Gurbuz, I. 1995. Parasito-epidemiological studies in sheep in the Samsun region, Turkey. *Turkiye Parazitoloji Dergisi*, 19(2): 290-296.
- Chaudhri, S.S., Kumar, S., and Gupta, R.P. 1994. Prevalence of *Schistosoma indicum* in Haryana and treatment trials in sheep. *Indian Veterinary Journal*, 71(3): 291-292.
- Diaz, R.P., Torres, H.G., Osorio, A. M. M., Perez, H.P., Pulido, A.A.R., Becerril, P.A.M., and Herrera, H.J.G. 2000. Resistencia genetica a parasitos gastrointestinales en ovinos Florida, Pelibuy y sus cruces en el tropico Mexicano, *Agrociencia*, 34: 13-20.
- Etter, E., Chartier, C., Hoste, H. and Borgida, L.P. 1999. The influence of nutrition on the peri parturient rise in faecal egg counts in dairy goat; Results from a two years study. *Revue de medecine Veterinaire*, 150(12): 975-980.

- FAO, 1992. Distribution and impact of helminth diseases of livestock in developing countries. *FAO Animal Production and Health Paper*, Vol. 96. FAO, Rome, p. 221.
- Gadahi, J.A., Arshed, M.J., Ali, Q., Javaid, S.B. and Shah, S.I. 2009. Prevalence of gastrointestinal parasites of sheep and goat in and around Rawalpindi and Islamabad, Pakistan. *Veterinary World*, 2(2): 51-53.
- Haq, S. and Shaikh, H. 1968. A Survey of helminth parasiting the gastrointestinal tracts of goats and sheep in East Pakistan. *East Pakistan Journal Veterinary Science*, 2 (2): 54-62.
- Hashem, M.S. and Sayed, M.H. 1997. Helminth parasites of sheep in Dakahlia province- Egypt. *Assiut Veterinary Medical Journal*, 38(75): 48-54.
- Ijaz, M., Khan, M.S., Avais, M., Ashraf, K., Ali, M.M., and Khan, M.Z.U. 2009. Infection rate and chemotherapy of various helminthes in diarrhoeic sheep in and around Lahore. *The Journal of Animal & Plant Sciences*, 19(1): 13-16.
- Islam, K.S. 1969. Incidence of schistosomiasis in domesticated ruminants in East Pakistan. M.S. Thesis, the Department of Parasitology, Bangladesh Agricultural University, Mymensingh.
- Khan, M.N., Sajid, M.S., Khan, M.K., Iqbal, Z. and Hussain, A. 2010. Gastrointestinal helminthiasis: prevalence and associated determinants in domestic ruminants of district Toba Tek Singh, Punjab, Pakistan. *Parasitology Research*, 107(4): 787-794.
- Krishnappa, K., Parimal, R., Balasubramaniam, G.A., Ramaswamy, V. and Roy, P. 1992. Studies on helminthiasis of sheep in Dharmapuri area of Tamil Nadu, India. *Indian Veterinary Medical Journal*, 16(4): 317-318.
- Lapage, G. 1962. *Monig's Veterinary Helminthology and Entomology*, 5th edition. Bailliere, Tindall and Cox Ltd. London, UK. pp: 556-723.
- Lateef, M., Iqbal, Z., Jabbar, A., Khan, M.N. and Akhter, M.S. 2005. Epidemiology of trichostrongylid nematode infections in sheep under Traditional Husbandry System in Pakistan. *International Journal of Agriculture and Biology*, 7(4): 596-600.
- Lloyd, S. 1983. Effect of pregnancy and lactation up on infection. *Veterinary immunology Immunopathology*, 4: 153-176.
- Mazid, M.A., Bhattacharjee, J., Begum, N. and Rahman, M.H. 2006. Helminth parasites of the digestive system of sheep in Mymensingh, Bangladesh. *Bangladesh Journal of Veterinary Medicine*, 4(2): 117-122.
- Mazyad, S.A., and Hl el-Nemr. 2002. The endoparasites of sheep and goats, and shepherd in North Sinai Governorate, Egypt. *Journal of the Egyptian Society of Parasitology*, 32(1): 119-126.
- Mbap, S.T., and Chiroma, B.Y. 1998. The effects of breed and environmental factors on helminth infection of sheep in bauchi, Nigeria. School of Agriculture, ATBU, PMB 0248, Bauchi, Nigeria. Proceedings of the 6th World Congress on Genetics Applied to Livestock Production, Armidale, NSW, Australia, 11-16 January, Volume 25: 261-264.
- Mir, R.A., Chishti, M.Z., Zarger, M.A., Tak, H. and Dar, F.A. 2008. Seasonal prevalence of trematode parasites of sheep (*Ovis aries* L.) in Kashmir Valley, India. *Nigerian Journal of Parasitology*, 29 (2): 80-83.
- Mostofa, M., Mckellar, Q.A. and Alam, M.N. 1996. Epidemiology of gastrointestinal helminth parasites in small ruminants in Bangladesh and their anthelmintic therapy. SPCSR, ACAIR Publication, Australia, 74: 105-108.
- Oka, C.K., Zinsstag, J., Pandey, V.S., Fofana, F. and Depo, N.A. 1999. Epidemiology of parasites of sheep in the southern forest zone of Cote d'Ivoire. LANADA, Laboratoire central de pathologie animal de Bingerville, BP 206, Bingerville, Cote d'Ivoire. *Revue d'Elevage et de Medecine Veterinaire des pays Tropicaux*, 52 (1): 39-46.
- Okafor, F.C., Mbata, G., and Anosike, J. 1988. Studies on *Paramphistomum cervi* (Schrank, 1790) infections of ruminants in Imo state, Nigeria with special reference to the role played by *Bulinius forskalii* (Ehrenberg) in there transmission. *Bulletin of Animal Health and production in Africa*, 38(2): 142-146.
- Oncel, T. 2000. The prevalence of helminth species in sheep in the southern region of Marmara, Turkey. *Turkiye Parazitoloji Dergisi*, 26(3): 317-319.
- Paz-Silva, A., Sanchez-Andrade, R., Suarez, J.L., Pedreira, J., Arias, M., Lopez, C., panadero, R., Diaz, P., Diez-Banos, P. and Morroondo, P. 2003. Prevalence of natural ovine fascioliasis shown by demonstrating the presence of serum circulating antigens. *Parasitology Research*, 91(4): 328-331.
- Rahman, M.A. and Hossain, W.I.M.A. 1997. Introductory Animal Hygiene and Management, 1st edition, Department of Microbiology and Hygiene, Bangladesh Agricultural University, Mymensingh. pp. 7-34.
- Rahman, M.H., Ahmed, S. and Mondal, M.M.H. 1996. Helminths parasites of cattle (*Bos indicus*) in Bangladesh. *Indian Journal of Parasitology*, 7(2): 173-174.
- Ratnaparkhi, M.R., Shastri, U.V., Narladkar, B.W., Degloorkar, N.M. and Digraskar, S.U. 1993. A short note on prevalence of *Fasciola* infection in domestic animals in Parbhani District, India. *Indian Journal of Animal Health*, 3(1): 83-85.
- Schlesselman, J.J. 1982. *Case Control Studies*, 1st edition, Oxford University Press, New York, pp: 174-177.
- Siddiqua and Banu 2010. Incidence of common helminth parasites in goats and sheep in Ghatampur Area of Kanpur Dehat. *Trends in Biosciences*, 3(1): 60-68.
- Soulsby, E.J.L. 1982. *Helminths, Arthropod and Protozoa of Domesticated Animals*. 7th edition, Bailliere Tindal and Cassell Ltd., London, pp. 757-759.
- Vercruysse, J. and Claerebout, E. 2001. Treatment vs. non-treatment of helminth infections in cattle: defining the thresholds. *Veterinary Parasitology*, 98: 195-214.