



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

Foreground selection through SSRs markers for the development of salt tolerant rice variety

U. Mondal*, M. S. R. Khanom, L. Hassan and S. N. Begum¹

Department of Genetics and Plant Breeding, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh and ¹Plant Breeding Division, Bangladesh Institute of Nuclear Agriculture, Mymensingh-2202, Bangladesh, *E-mail: upamamondal@gmail.com

Abstract

Salinity is a great problem for rice production worldwide incurring substantial yield loss; a great threat towards food security. Marker-assisted backcrossing is one of the feasible methods to develop a new salt tolerant rice cultivar to cope with the challenge. The study was focused on to introgress salt tolerant genes from a tolerant rice line, FL-478 to Binadhan-7, an early, agronomically suitable and susceptible variety. Backcrossing was done during boro season; where Binadhan-7 was the recurrent parent and FL-478 was the donor parent. 141 BC₁F₁ lines were developed, which were subjected to foreground selection; the first level of selection of marker assisted backcrossing program. The aim of foreground selection was to identify the introgressed lines. 141 BC₁F₁ populations were evaluated with tightly linked salt tolerant markers; AP3206f, RM3412b and RM336. A total of 47 heterozygous BC₁F₁ lines were selected finally, which have alleles of both of the parents. Those introgressed lines could be efficiently used in further development of a stable early salt tolerant rice variety.

Keywords: Foreground selection, SSRs markers, Marker-assisted backcrossing

Introduction

Climate change and food security are the two burning issues now-a-days. Agricultural production is extremely vulnerable to climate change. It is causing threatening impacts on rice production, which is the most important cereal crop for the food security worldwide. Rice is the major source of food for more than 2.7 billion people on a daily basis in South and Southeast Asia (Hossain, 2005). In Bangladesh, rice grows in all the crop-growing seasons and occupies 77% of the total cropped area of 13.9 million hectares (BBS, 2010).

Salinity is one of the major natural hazards hampering rice production. Approximately 20% of irrigated areas worldwide (about 45 million ha) are estimated to suffer from salinity problems by various degrees (FAO, 2010). 21.5 million hectares of land areas in Asia are affected by salinity and estimated to cause up to 50% loss of fertile land by the mid of 21st century (Linh *et. al.*, 2012). In Bangladesh, The coastal saline soils are distributed unevenly in 64 upazillas of 13 Districts, covering portions of 8 AEZ of the country. These areas constitute about 2.5 million hectares which amount to about 18% of the total cropland of the country (Seraj and Salam, 2000). So in the concern of food security the necessity for enhancement in salt tolerance in rice is well understood.

Molecular marker technology offers a possibility by adopting a wide range of novel approaches to improve the selection strategies in rice breeding. Molecular markers that are linked to genes controlling salt tolerance could facilitate selection and improve rice varieties with salt tolerance having high heritability and expressivity. Microsatellite markers have been used effectively to map QTLs associated with salt tolerance (Singh *et. al.* 2007). *Saltol*, A major salinity tolerance QTL on rice chromosome 1, was mapped at IRRI using a recombinant inbred line (RIL) population between tolerant Pokkali and sensitive IR29; explaining 43% of the variation for seedling shoot Na⁺ uptake (Gregorio 1997, Elahi *et. al.* 2004). This QTL confers salinity tolerance at the vegetative stage of rice.

The use of DNA markers in backcrossing greatly increases the efficiency of selection, which is known as Marker-assisted backcrossing. The basis of MABC strategy is to incorporate one or a few genes into an adapted or elite variety or to transfer a gene/QTL from a donor line to a recipient line by repeated backcrossing. This approach develops an ideal genotype within a very short time avoiding the complicated issues related to transgenic technology and conventional breeding approaches. MABC

approach is very advantageous with the following steps, (1) recombination and identification of target locus, known as 'foreground selection'; (2) minimizing linkage drag as recombinant selection; (3) harvesting maximum recurrent parent genome as background selection (Collard *et. al.* 2008). Although, the extent of effectiveness of this program is delimited by some factors, molecular breeding technologies are upgrading day by day, which has already been proven as the most effective technology for the development of salt tolerant varieties.

Hence, the attempt of this study was to introgress *Salto* QTL into a popular variety Binadhan-7 by MABC method and to identify the introgressed lines through foreground selection using SSRs markers at early generation.

Materials and Methods

Plant materials

FL-478 (IR 66946-3R-178-1-1) was used as one of the parent for transferring salt tolerant QTL as it contains *Salto* QTL on chromosome 1. It is a recombinant inbred line derived from (Pokkali X IR 29). It has a high level of seedling stage salinity tolerance in rice. Binadhan-7 is an early, high yielding transplanted aman variety was used as another parent. It is susceptible to salinity. A backcrossed program was conducted where, Binaadhan-7 was used as recurrent parent and FL-478 was the non-recurrent donor parent. From that crossing program 23 F₁ seeds were produced. The F₁ seeds were backcrossed with Binadhan-7 to make BC₁F₁ seeds.

Cultural management

For the ease of handling, the experiment was conducted on the plastic pot (bucket 10 L). Recommended cultural operations were followed as and when necessary to ensure the normal plant growth and development.

Development of backcross seeds

Crossing scheme: Synchronization of flowering time is the most important operation for MABC program. To produce BC₁F₁ seeds, F₁ seeds were seeded in two sets at 14 days interval and recurrent parents were staggered in four sets starting from 15 January, 2012. First set of recurrent parent was seeded 7 days before the seeding F₁ generation. Second and third set were seeded in same time of F₁ generation and finally fourth set was seeded in 7 days after the seeding of F₁ generation.

Raising of backcross plants: 201 mature backcrossed seeds were collected 21 days after dusting. Seeds were dried into a seed dryer at 65°C for seven days and then set for germination. Among the 201 seeds, 171 sprouted seeds were grown in the pot. The seeds were seeded in two sets in order to ease the work pressure. Among those, 141 seedlings were survived and leaves were collected for foreground selection (Fig 1).



Fig 1. BC₁F₁ generation in field

Collection of leaf sample and DNA extraction: Samples were collected from young, vigorous leaves from 25 days old seedlings (Binadhan-7, FL-478, BC₁F₁ population) to extract genomic DNA. The collected leaf samples were placed in an ice box and finally the samples were stored in (-) 80° C freezer.

Using the Cetyl Trimethyl Ammonium Bromide (CTAB) mini-prep method, DNA was extracted from the leaves collected from at least 2-3 seedling of each genotype. The simplified mini scale procedure for DNA isolation in PCR analysis developed at IRRI was followed. The quality and quantity of the isolated DNA was sufficient for PCR analysis. Quantified DNA samples from each genotype were subjected to PCR amplification with SSR primers.

Polymorphism survey: Polymorphism survey of two parents and BC₁F₁ populations were carried out using 7 foreground markers (RM585, RM336, RM3412b, RM10748, RM8094, AP3206 and AP3206f). Out of 7 markers, 3 markers (AP3206f, RM3412b and RM336) showed clear polymorphisms which were used in genotyping the foreground selection of the 141 BC₁F₁ rice lines for salt tolerant genotypes. The details of the primer are given in Table 1.

Table 1. The details of the microsatellite markers (SSRs) used for BC₁F₁ Survey

Primer Name	Chr.1 (Mb)	Nipponbare size (bp)	Notes	Sequence		Annealing temperature (°C)
				Forward	Reverse	
AP3206f	11.2	167	Foreground marker	Forward	GCAAGAATTAATCCATGTGAAAGA	55
				Reverse	ATGCTCTGGCTCCCTCAAG	
RM336	11.6	175	Foreground marker	Forward	CTTACAGAGAAACGGCATCG	55
				Reverse	GCTGGTTTGTTCAGTTTCG	
RM3412b	11.5	110	Foreground marker	Forward	TCATGATGGATCTCTGAGGTG	55
				Reverse	GGGAGGATGCACTAATCTTTC	

PCR array: The PCR cocktail had total volume of 12.75 μ L/reaction containing 1.5 μ L 10X TB buffer (containing 200mM Tris- HCl pH 8.3, 500mM KCl, 15mM MgCl₂), 0.75 μ L of 1mM dNTP, 1.0 μ L each of 5 μ M forward and reverse primers, and 0.25 μ L of *Taq* DNA polymerase with required ddH₂O. 2 μ L genomic DNA (5–25 ng of DNA Template) was added PCR cocktail. The steps of PCR reactions were: initial denaturation for 5 minutes at 94°C, then annealing at 55°C for 1 minute. Polymerization for 2 minutes at 72°C to complete a cycle and cycle was repeated for 34 times. The final extension duration was 7 minutes at 72°C. then, PCR products were mixed with 3 μ L of 2X gel loading dye. Polymorphisms in the PCR products were analyzed by electrophoresis using mini vertical polyacrylamide gels for high throughput manual genotyping. The gels were stained in ethidium bromide and documented using BIOMETRA Gel Documentation System.

Data analysis

Scoring of bands: The pattern of bands obtained after amplification with the primers was scored with reference to two parents. The band having same level of FL-478 was scored as 'T' which indicated the homozygous allele of the tolerant parent for a particular microsatellite marker. Similarly, the bands with similar level of Binadhan-7 was scored as 'S'. The heterozygous alleles having both the bands of two parents were scored as 'H'.

Allele scoring: With the help of Alpha Ease FC 5.0 software, the size (in nucleotide base pairs) of the amplified band for each SSR marker was determined based on its migration relative to 20bp DNA Ladder (a molecular weight size marker).

Analysis of SSR Data: The summary statistics including the number of alleles per locus, major allele frequency, gene diversity and Polymorphism Information Content (PIC) values were determined using POWER MARKER version 3.23 (Liu and Muse 2005), a genetic analysis software. Polymorphism Information Content (PIC) value described and modified by Anderson E. (1993) for self-pollinated species.

Estimation of gene and genotypic frequencies: The proportions of different alleles of a gene present in a Mendelian population are known as gene frequency. Gene frequencies in a population can be readily estimated by the total number of each of the alleles of the gene present in these individuals and their ratio to the total number for all the alleles of the gene is estimated. The proportions of different genotypes for a gene in a population are known as genotypic frequencies for that gene; often they are also called zygotic frequencies.

Results and Discussion

Parental polymorphism

Primer survey is very essential before starting marker-assisted backcross breeding. Monomorphic markers can not distinguish the two parental genotypes, so this type of marker bears no value in selection work. For selection of the *Saltol* locus (foreground selection), seven tightly linked SSR markers to the *Saltol* positioning at 11.2 to 11.7 Mb were used. Out of seven SSR primers, three RM336, RM3412b, and AP3206f showed polymorphisms for the parents.

Allelic information

Using the SSR markers, alleles were detected among the 141 BC₁F₁ populations along with their parents. According to Nei's (1983), the highest level of gene diversity value (0.69) was observed in loci RM3412b and the lowest level of gene diversity value (0.438) was observed in loci AP3206f with a mean diversity of 0.5260 (Table 2). The maximum number of repeats within the SSRs was also positively correlated with the genetic diversity. The PIC value was calculated to estimate the informativeness of each primer. The PIC values ranged from a low of 0.3454 (AP3206f) to a high of 0.6445 (RM3412b) with an average of 0.4460 (Table 2). So, the primer RM 3412b was found to be superior for analysis of genetic diversity among the markers in this region.

The result was consistent with the previous study of Mohammadi-nejad, *et. al.* (2010), where the PIC value varied from 0.56 to 0.88 which was also highly correlated the allele size range and the number of motif.

Table 2. Allelic informations found among 141 BC₁F₁ populations of Binadhan-7 x FL 478 along with their parents for 3 microsatellites (SSR) markers

Locus	Repeat motif	Amplicon Size ranges	Major allele frequency	PIC	Sample size	Gene Diversity
AP3206f	(GT)10	136-177	0.43	0.3454	143	0.4386
RM3412b	(TA)34	98-117	0.67	0.6445	143	0.6959
RM 336	(CTT)18	144-195	0.68	0.3482	143	0.4436
Total	-	-	-	1.3381	-	1.5755
Mean	-	-	0.59	0.4460	143	0.5260

Genotypic performance of backcross population BC₁F₁ at foreground selection

For foreground selection in the BC₁F₁ generation, individual plants that were heterozygous at the *Saltol* locus were identified reducing the population size for further screening. For each of the marker, allelic bands were scored based on the parental bands and designated as 'T' for tolerant, 'S' for susceptible and 'H' for heterozygous. 141 BC₁F₁ lines showed variation in foreground selection with 3 tightly linked salt tolerant primers (Fig 2-4). Out of 141 BC₁F₁ plants, the marker AP3206f identified 60 plants as heterozygous, RM3412b identified 59 plants and the RM336 identified 65 plants as heterozygous (Table 3).

47 BC₁F₁ plants were selected based on the heterozygous nature of all the target loci at *Saltol* region. Those lines can be used for further selection step of marker aided backcrossing. Similar result was found in the study of Rahman (2011) to introgress *Saltol* QTL into BR28 from FL-478. He found 94 heterozygous plants out of 230 BC₁F₁ rice lines using foreground markers.




Fig. 2. Banding pattern of BC₁F₁ population of Binadhan-7/FL-478 using SSR marker AP3206f




Fig. 3. Banding pattern of BC₁F₁ population of Binadhan-7/FL-478 using SSR marker RM3412b




Fig. 4. Banding pattern of BC₁F₁ population of Binadhan-7/FL-478 using SSR marker RM336

Table 3. 141 BC₁F₁ populations with respect to the alleles amplified by the microsatellite primer i.e., AP3206f, RM3412b and RM336

Name of Primers	Total number of BC ₁ F ₁ lines	Patterns of BC ₁ F ₁ population		
		Susceptible type (Binadhan-7)	Tolerant type (FL478)	Heterozygous
AP3206f	141	65	15	60
RM3412b		68	13	59
RM336		58	18	65

Gene and Genotypic Frequency

The proportions of tolerant and susceptible alleles are 0.32 and 0.67 for marker AP3206f and 0.301, 0.69 for marker RM3412b. In case of RM336 the proportions are 0.358 and 0.641. The proportions of susceptible, heterozygous and tolerant genotypes are 0.46, 0.40 and 0.1 for marker AP3206f. For RM3412b the proportions are 0.44, 0.45 and 0.09 respectively. In case of RM336 the proportion is 0.41, 0.46 and 0.127. In this study, the average proportions of susceptible and heterozygous genotypes were 0.44 and 0.43 respectively; which indicated that the results fitted to the expected 1:1 ratio of this generation (chi square value=0.48 at a significant level of 0.05).

Similar result was found in the study of Iftekharaudaula (2008), who also found 1:1 ratio at BC₁F₁ generation with a non-significant chi square value of 0.28 at a probability level of 0.05.

Conclusion

Salinity stress is the most serious factors that limit the rice (*Oryza sativa* L.) production worldwide and brought an enormous challenge for food security in developing countries. Breeding of rice varieties with in-built salt tolerance is realized as the most promising, less resource consuming, economically viable and socially acceptable approach. Marker aided backcrossing can be used as an efficient tool for this. After conducting foreground selection, 47 BC₁F₁ lines were selected for further recombinant and background selection to minimize 'linkage drag' while recovering more genetic background of recurrent parent. Three foreground markers AP3206f, RM3412b and RM336 showed polymorphism among the rice lines, could facilitate selection, mapping, cloning genes, QTL analysis and so on, which in turn increase rice production in saline environments.

Acknowledgements

The authors would like to thank the staff and lab mates at Bangladesh Institute of Nuclear Agriculture for all technical assistance. Thanks also go to all the teachers of the department of Genetics and Plant Breeding, Bangladesh Agricultural University, Mymensingh, for their valuable advice and suggestions.

References

- Anderson, E. 1993. A semi graphical method for the analysis of complex problems. Proceeding of National Academy of Science, USA, 48: 923-927.
- BBS, 2010. Agriculture Wing. Bangladesh Bureau of statistics, Ministry of planning, Government of the People's Republic of Bangladesh, Dhaka. 54.
- Collard B.C.Y and Mackill, D.J. 2008. "Marker-Assisted Selection: An Approach for Precision Plant Breeding in the 21st Century," Philosophical Transactions of the Royal Society, Vol. 363, No. 1491, pp. 557-572.
- Elahi, C.M.F., Seraj, Z.I., Rasul, N.M., Das, K.C., Biswas, K., Salam, M.A., Gomosta, A.R., Tumimbang, E., Adorada, D., Gregorio, G. and Bennett, J. 2004. Breeding rice for salinity tolerance using the Pokkali allele: finding a linked DNA marker. In: *In Vitro Culture, Transformation and Molecular markers for Crop Improvement*. Ed. Islam A.S. Science Publishers, Inc, USA. pp. 157-170.
- Food and Agriculture Organization. 2010. "Report of salt affected agriculture," <http://www.fao.org>
- Gregorio, G.B. 1997. Tagging salinity tolerance genes in rice using amplified fragment length polymorphism (AFLP). PhD. thesis, University of the Philippines, Los Baños.; 118 p.
- Hossain, M. 2005. Rice production and market: trends and outlook. Paper presented at the IFA Regional Conference for Asia and Pacific, December 6-8, Singapore.
- Iftekharaudaula, K.M. 2008. Comparison of New Selection Strategies or Marker Assisted Backcrossing for a Submergence Tolerant Gene in Rice. Ph. D. dissertation. Bangladesh agricultural University, Mymensingh, Bangladesh, pp. 1-205.
- Linh, H.L., Linh, T.H., Xuan, T.D., Ham, L.H., Ismail, M.A., Khanh, T.D. 2012. Molecular Breeding to Improve Salt Tolerance of Rice (*Oryza sativa* L.) in the Red River Delta of Vietnam. *International Journal of Plant Genomics*. 10: 1-9.
- Liu, K. and Muse, S.V. 2005. Power Marker: Integrated analysis environment for genetic marker data. *Bioinformatics*, 21: 2128-2129
- Mohammadi-Nejad, G., Singh, R.K., Arzani, A., Rezaie, A.M., Sabouri, H. and Gregorio, G.B. 2010. Evaluation of salinity tolerance in rice genotypes. *International Journal of Plant Production*, 4: 199-208.
- Nei M. 1983. Analysis of gene diversity in subdivided populations. *Proceeding of National Academy of Science, USA*, 70: 3321-3323.
- Rahman, M.A. 2011. Marker-assisted backcrossing of saltol, discovery of additional quantitative trait loci and assessment of allelic variability in *saltol* in rice germplasm. Ph. D. dissertation. Bangladesh Agricultural University, Mymensingh, Bangladesh, pp. 1-268.
- Seraj, Z.I. and Salam, M.A. 2000. Growing rice in saline soils. *The Biotechnology Directory*. Macmillan Reference Ltd., Porters South, Crinan Street, London.
- Singh, R.K., Gregorio, G.B. and Jain, R.K. 2007. QTL mapping for salinity tolerance in rice. *Physiology and Molecular Biology of Plants*, 13: 87-99.