



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

Estimation of size selectivity of fish species caught by different gears in the old Brahmaputra River

I. S. Saberin¹, M. S. Reza^{1*}, N. J. Ayon² and M. Kamal²

¹Department of Fisheries Technology, Bangladesh Agricultural University, Mymensingh 2202, Bangladesh, and

²Department of Pharmaceutical Technology, Faculty of Pharmacy, University of Dhaka, Dhaka-1000 Bangladesh,

*E-mail: rezams@gmail.com

Abstract

Over the last couple of decades, the old Brahmaputra River, one of the ancient river systems of Bangladesh, has fallen in verge of biodiversity changes affecting its range of fauna and flora. Study was, therefore, conducted to investigate the selectivity of fish species caught by different fishing gears used in the old Brahmaputra River system over a period of 1 year. $L_{50\%}$ of different gears was determined mainly for dominant species where 41 fish species were recorded. Highest $L_{50\%}$ was recorded in gill net (*fash jal*) followed by basket trap (*vair*) whereas lowest value was recorded for seine net. Seine net, cast net, *sidi bair* and hand line (*borshi*) were considered non selective gears where *fash jal* and *sidi bair* were regarded as gear specific for large fish species and *koi jal*, *ber jal*, *dharma jal*, *howra bair* and *koach* were considered specific for small species. Among all the gears, seine net, cast net, *koi jal* and *howra bair* were found to be harmful for small species of fishes. Monofilamentous gill net (*current jal*) although marked as banned gear, was found to be operated in the river and causing considerable damage to the fish stock. *Fash jal* was considered as harmful gear for catfish as it was found to be used to catch fishes below legal size. It is suggested that *fash jal* should be restricted for several months, which may allow the newly stock fingerlings to grow to market size.

Keywords: Biodiversity, Fish species, Fishing gears, The Old Brahmaputra river

Introduction

Bangladesh is topographically a deltaic plain criss-crossed by innumerable rivers and rivulets and has a great fisheries potential. Riverine fishing areas comprise nearly one fifth of the entire fishing area of 4.9 million ha of the country (DoF, 2013). Its capture fisheries in the form of common property and open access resources constitute a vital component of the agro-ecosystem of rural Bangladesh. It is also a very important source of household welfare for millions of rural poor, particularly for providing nutrition (specially the much needed protein), income and employment. Thus, the fisheries resources offer excellent opportunities for increased fish production that will effectively combat malnutrition and generate additional employment. Approximately 1.4 million people are directly engaged in fishing, 11 million in part time fishing and another 3 million in aquaculture activities (Hussain, 2010). In addition, it provides opportunities for about 70% of rural population in subsistence fishing for their own consumption. Quite a significant number of people are engaged in other related activities such as fish fry collection, fish trading, processing, net/trap and boat making, fisheries labor etc.

The old Brahmaputra River was the main flow of the Brahmaputra-Jamuna river system and rich in different fish fauna. The biodiversity of this river system has changed over the last couple of centuries, but still it is regarded as one of the most productive freshwater ecosystem of Bangladesh. The water supply of this river system is has been reported to be greatly affected due to huge withdrawal of water from the upstream parts of the river located in India (Mortuza, 2007).

Different types and form of gears have been operated in the old Brahmaputra River to exploit wild fishes. The intensity of use of any form of gear in a river is dependent on the intensity of target fish population presumed to be available in that river. Some of the gears are selective for a particular species, whereas other account for a number of species caught during operation giving multi-species nature of the fishing. Among them, many of these have been known to catch carp fingerlings before they grow to legal size and many of these responsible for sharp decline in the population of wild species from the river of the country. However, operation of all types of gear cannot be kept suspended to allow the stocked fingerlings and also wild fishes to grow. Only a few studies have been undertaken on the status of resources as well as the causes of declining resources in the old Brahmaputra River. The intensity of fishing gears, mesh sizes and catch per unit effort are good indicators of the status of exploitation level of river.

Considering the above circumstances, the present study was undertaken on the basis of the following objectives i) to identify the fishing gears operated in the old Brahmaputra River, ii) Size (length) selectivity of the species caught by different fishing gears with the potential to formulate the better management practices to conserve the resources.

Materials and Methods

Study area

The study was conducted in the area of Mymensingh sadar upzilla where the river runs from Kagdahor, Kachari, Shomvogonj, Kawatkhal Railway Bridge, BAU campus, Shesmor and Sutiakhali. The investigation have also included the Char-kalibari, the Char-gobaida, and the Char-nilakshia.

Data Collection

The catch data were collected from fish landing centers, markets and the fishing spot individually for a period starting from April, 2011 to March, 2012. Catch composition of each type of gear was recorded by physical examination of the total catch, most of the time the total catch was purchased from the fishermen on the spot. Then the samples were sorted out species wise and measurement total length of individual fish of each species. For each type of gear, the samples were collected from different spots (each collection is the catches of a fishing unit with a type of gear) to obtain the desired size of sample population (the number of fish those have been measured), i.e., at least 100 fish for a dominant species in a given gear. But less than 100 fish were considered as dominant species due to its less availability.

Data Analysis

Collected information on species composition in each type of gear was analyzed to determine the species and length selectivity. L_{50} % value and selection range ($L_{25\%}$ - $L_{75\%}$) were determined for each dominant species. To estimate size (length) selectivity of the species caught in gear, the $L_{50\%}$ value and selection range ($L_{25\%}$ - $L_{75\%}$) were calculated for each dominant species with the help of linear probit regression. The $L_{50\%}$ is the length at which 50% of the fish entering the gear are retained and 50% escape. It is also called "length at first capture". The selection range is the range of the lengths between $L_{25\%}$ (the length at which 25% of fish entering the gear are retained) and $L_{75\%}$ (the length at which 75% of fish entering the gear are retained). The 50% retention length and the selection ranges (25% and 75%) were evaluated according to Pope *et al.* (1975).

Results and Discussion

A total 19 types of fishing gears were found to be used by the fishermen for harvesting fish from old Brahmaputra River during the study period (Table 1). The types of gears used are more or less similar with the finding of BCAS (1991) in Haldi beel where the fishermen use approximately 30 different types of fishing gear. From the present study, all these types of gears recorded can be broadly classified into nets, traps and hooks and line. Besides these gears, fishermen were also found to catch fish by dewatering the water body and by hand picking. *Katha fisheries* were also popular and used extensively during dry season by the fishermen.

A total of 41 species of fishes were recorded in the catches of different gears. Data of the found gears were analyzed and results are described below:

Gill net

The $L_{50\%}$ value and selectivity range ($L_{25\%}$ - $L_{75\%}$) of *fash jal* and *koi jal* were calculated for dominant species of fish. The $L_{50\%}$ value and of selectivity range ($L_{25\%}$ - $L_{75\%}$) of *Anabas testudineus* 6.25 cm and 6.36-7.50 cm in the 2.45 cm mesh, 8.40 cm and 8.45-9.50 cm in 3.16 cm mesh, 9.24 cm and 8.50-10.22 cm in 3.6 cm mesh in June-July, respectively (Table 2). In *koi jal* $L_{50\%}$ values recorded for *Anabas testudineus* were higher than the values (8.82 cm and 8.08-9.54 cm in 2.5 cm mesh) recorded by

Rahman *et al.* (1992). These differences might be associated with the difference in the mesh sizes of the net. But in the *fash jal* the $L_{50\%}$ value and selective range ($L_{25\%}$ - $L_{75\%}$) of *Mystus aor* 21.24 cm and 21.25-22.56 cm in 6.2 cm mesh, 22.56 cm and 22.15-24.56 cm in 7.4 cm mesh, 24.25 cm and 24.50-28.26 cm in 9.3 cm mesh, respectively. The values were recorded for *Labeo rohita* 22.56 cm and 22.24-24.25 cm in 6.2 cm mesh, 24.56 cm and 24.25-26.50 cm in 7.4 cm in mesh, 27.15 cm and 27.25-30.25 cm in 9.3 cm mesh in June-July, respectively (Table 3). The $L_{50\%}$ value and selective ranges of *fash jal* (9-12 cm, mesh) recorded by Paul *et al.* (1993) in Haldi beel were 24.15 cm and 22.03-26.27 cm for catla and 23.99 cm and 21.13-26.86 cm for common carp, respectively. In the present study, the sizes of the fishes recorded in the catches of the gill net (*fash jal*) were almost similar to the sizes recorded by Paul *et al.* (1993) in Haldi beel but in Chanda beel the sizes of fish were larger due to the use of comparatively large mesh gill net. Through the sizes of carp and catfishes recorded in the present study were just within the limit of legal size, it is suggested to use relatively large mesh size as a better management strategy.

Table 1. Fishing gears used in the old Brahmaputra River

Types of gear		
Fish net	Gill net	1. <i>Koi jal</i> 2. <i>Fash jal</i> 3. <i>Current jal</i>
	Seine net	1. <i>Ber jal</i>
	Lift net	1. <i>Vassal/Khorajal</i> 2. <i>Dharma jal</i>
	Cast net	1. <i>Jhaki jal</i>
	Dragged net	1. <i>Thela jal</i> 2. <i>Moiya jal</i>
Fish Trap		1. <i>Howrabair</i> 2. <i>Sidibair</i> 3. <i>Dughair</i> 4. <i>Vair</i> 5. <i>Bitte</i>
Hook and Line		1. <i>Kobaborshi</i> 2. <i>Chip borshi</i> 3. <i>Hand borshi</i> 4. <i>Chasra</i>
Wounding Gear		1. <i>Koach</i>
Fish Aggregating Device		1. <i>Katha</i>

Table 2. The $L_{50\%}$ values and Selectivity ranges ($L_{25\%}$ - $L_{75\%}$) of gil net (*koi jal*) for major dominant species in the old Brahmaputra River

Species	Month	2.45 cm mesh		3.16 cm mesh		3.6 cm mesh	
		$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$	$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$	$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$
<i>Anabas testudineus</i> (koi)	June-July	6.25	6.36-7.50	8.40	8.45-9.50	9.24	8.50-10.22

Cast net

Length selectivity of this gear was calculated for 5 dominant species viz. *Puntius sarana*, *P. ticto*, *Chanda nama*, *Mystus tengara* and *Glossogobius giuris* caught by three different mesh size from June to October. The $L_{50\%}$ value and selective ranges of *P. sarana* were recorded 5.65 cm and 4.15-7.20 cm in 1cm mesh, 6.58 cm and 6.84-7.50 cm in 1.5 cm mesh, 8.50 cm and 7.25-10.25 cm in 2cm mesh (Table 4). The values of 10.24 cm and 9.25-12.24 cm in 1cm mesh, 12.58 cm and 6.75-18.25 cm in 2 cm mesh for *Glossogobius giuris*. The values of 6.25 cm and 4.36-6.50 cm in 1 cm mesh, 10.26 cm and 10.25-15.36 cm in 2 cm mesh for *M. tengara*. Also the values of *C. nama* were 3.36 cm and 3.75-4.50 cm, 4.85 cm and 4.50-5.26 cm, 6.24 cm and 6.15-7.25 cm in 1 cm, 1.5 cm and 2 cm mesh size, respectively. The $L_{50\%}$ value increased with the increased of mesh size in *P. sarana*, *P. ticto*, *C. nama*, *M. tengara* in July - August. For *M. tengara* value increased to change in mesh in July. But the $L_{50\%}$ value of *P. sarana* more or less similar in 1 cm and 1.5 cm mesh in July. The length selectivity value recorded for *P. sarana* and *M. tengara* in the present study were higher to the values obtained by Rahman *et al.* (1992).

Table 3. The $L_{50\%}$ values and Selectivity ranges ($L_{25\%}$ - $L_{75\%}$) of gill net (*fash ja*) for major dominant species in the old Brahmaputra River

Species	Month	6.2 cm mesh		7.4 cm mesh		9.3 cm mesh	
		$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$	$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$	$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$
<i>Mystus aor</i> (Air)	June-July	21.24	21.25-22.56	22.56	22.15-24.56	24.25	24.50-28.26
<i>Labeo rohita</i> (Rui)	June-July	22.56	22.24-24.25	24.56	24.25-26.50	27.15	27.25-30.25

Table 4. The $L_{50\%}$ values and Selectivity ranges ($L_{25\%}$ - $L_{75\%}$) of cast net for major dominant species in the old Brahmaputra River

Species	Month	1 cm mesh		1.5 cm mesh		2 cm mesh	
		$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$	$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$	$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$
<i>Glossogobius giuris</i> (Baila)	June	10.24	9.25-12.24			12.58	6.75-18.25
	July			11.25	11.36-12.50		
	August			11.50	11.45-13.25		
	September					13.25	12.36-16.50
	October					13.52	13.62-18.25
<i>Puntius sarana</i> (Sharputi)	July	5.65	4.15-7.20	6.58	6.84-7.50	8.50	7.25-10.25
	August			6.84	6.92-7.21		
	September					10.24	8.81-11.50
	October					10.32	8.75-12.36
<i>Puntius ticto</i> (Tit puti)	March	3.56	3.25-4.56	4.25	4.15-5.36		
	April					5.26	5.12-6.50
	October	2.56	2.25-3.85				
<i>Mystus tengara</i> (Tengra)	July	6.25	4.36-6.50			10.36	10.25-15.36
<i>Chanda nama</i> (Chanda)	September	3.36	3.75-4.50	4.85	4.50-5.26	6.24	6.15-7.25

Seine net

The $L_{50\%}$ value and selectivity range ($L_{25\%}$ - $L_{75\%}$) were determined for five dominant species of fish viz. baim, baila, sharputi, tengra and chanda. The $L_{50\%}$ value and selective ranges recorded were 15.42 cm and 13.65-16.36 cm for *Mastacembelus armatus*, 5.50 cm and 4.36-6.25 cm for *G. giuris*, 6.15 cm and 5.25-7.50 cm for *P. sarana*, 6.50 cm and 4.68-7.25 cm for *M. tengara*, 2.75 cm and 2.25-3.65 cm for *C. nama*, respectively (Table 5). *Colisa fasciatus* caught by seine net have less $L_{50\%}$ value then push net due to smaller mesh size of seine net. The $L_{50\%}$ value and selectivity range recorded by Rahman *et al.* (1993) in Haldi beel 3.3 cm and 2.98–3.62 cm for *C. nama* and 6.21 cm and 5.07-7.36 cm for *M. tengara*. These values almost similar with the values recorded by Rahman *et al.* (1993) but it was to much higher than the values recorded by Rahman *et al.* (2009).

Lift net

The length selectivity values of these gears were determined for 6 dominant species of fishes. The $L_{50\%}$ value and selective ranges were recorded 18.20 cm and 16.54 -21.25 cm for *Mastacembelus armatus*, 5.45 cm and 4.25-6.30 cm for *Puntius sarana* 6.17 cm and 5.43 -6.85 cm for *M. tengara*, respectively (Table 5). These values were more or less similar to the values of 5.74 cm and 4.24-7.23 cm for *P. sarana* recorded by Rahman *et al.* (2009) in Chanda beel.

Push net

The $L_{50\%}$ value and selective range were calculated for 5 dominant species. In *Colisa fasciatus* the values were 6.25 cm and 5.58-7.56 cm, in *G. giuris* 6.25 cm and 4.36-6.85 cm, in *P. sarana* 4.58 cm and 4.36-6.25 cm, in *Mystus tengara* 6.78 cm and 6.85-7.15 cm and *C. nama* the values were 3.56 cm and 3.25-4.50 cm, respectively. The $L_{50\%}$ values recorded for *G. giuris* were higher than the values recorded in the seine net due to difference of mesh size.

Table 5. The L_{50%} values and Selectivity ranges (L_{25%}-L_{75%}) of seine net, lift net and push net for major dominant species in the old Brahmaputra River

Species	Month	Seine net (1 cm mesh)		Lift net (1.2 cm mesh)		Push net(1.5 cm mesh)	
		L _{50%}	L _{25%} -L _{75%}	L _{50%}	L _{25%} -L _{75%}	L _{50%}	L _{25%} -L _{75%}
<i>Colisa fasciatus</i> (Kholisha)	June			4.25	4.36-5.84	6.25	5.58-7.56
	July	6.25	6.36-8.68			7.52	7.15-8.65
<i>Mastacembelus armatus</i> (Baim)	June	15.42	13.65-16.36	18.20	16.54-21.25		
	July			18.50	16.54-21.36		
	August	14.58	14.65-15.23				
<i>Glossogobius giuris</i> (Baila)	June	5.50	4.36-6.25			6.25	4.36-6.85
	July			5.75	5.36-6.54		
	August			5.92	5.36-6.85		
	September	5.45	5.23-5.89				
<i>Puntius sarana</i> (Sharputi)	June	6.15	5.25-7.50	5.45	4.25-6.30		
	July					4.58	4.36-6.25
	August	3.45	3.58-5.24				
<i>Mystus tengara</i> (Tengra)	June	6.50	4.68-7.25	6.17	5.43-6.85		
	July					6.78	6.85-7.15
	August	4.36	4.12-5.89				
	September			5.36	5.24-6.78		
<i>Chanda nama</i> (Chanda)	June	2.75	2.25-3.65			3.56	3.25-4.50
	July			3.89	3.56-4.87		

Traps

Most of the traps were recorded more or less selective to few species of fishes. In *howra bair* length selectivity were determined for *M. armatus*, *G. giuris* and *M. tengara*. The L_{50%} value and selective ranges recorded were 16.50 cm and 15.00-18.25 cm for *M. armatus*, 6.65 cm and 4.23-6.50 cm for *G. giuris*, 6.45 cm and 5.25-7.50 cm for *M. tengara*, respectively (Table 6).

In *sidi bair* L_{50%} value and selective ranges recorded were 4.75 cm and 4.25-6.50 cm for *P. sarana*, 15.50 cm and 14.25-18.50 cm for *Macrogathus pancalus*, 3.68 cm and 3.25-4.50 cm for *Esomus danricus*, respectively (Table 6).

Table 6. The L_{50%} values and Selectivity ranges (L_{25%}-L_{75%}) of Sidi bair, Howra bair and Vair, for major dominant species in the old Brahmaputra River

Species	Month	Fishing Trap					
		<i>Howra bair</i>		<i>Sidi bair</i>		<i>Vair</i>	
		L _{50%}	L _{25%} -L _{75%}	L _{50%}	L _{25%} -L _{75%}	L _{50%}	L _{25%} -L _{75%}
<i>Mastacembelus armatus</i> (Baim)	June	16.50	15-18.25			16.75	15.25-20.24
	July			17.25	17.12-18.52		
	August			18.12	18.05-20.25		
	September					15.36	16.25-18.65
<i>Glossogobius giuris</i> (Baila)	June	5.65	4.23-6.50			5.56	5.13-6.54
	July					6.58	6.36-7.65
	August			5.36	5.21-6.35	6.85	6.35-7.84
	September	6.45	5.65-7.85				
<i>Lepidocephalus guntea</i> (Gutum)	June-July					7.25	5.5-8.2
<i>Puntius sarana</i> (Sharputi)	June-July			4.75	4.25-6.50	4.35	4.25-6.50
<i>Mystus tengara</i> (Tengra)	June	6.45	5.25-7.50			6.25	4.20-6.75
	July			5.56	5.36-6.54		
	August	7.23	7.14-8.84				
<i>Macrogathus pancalus</i> (Guchi baim)	June			15.50	14.25-18.50		
	July					16.47	16.45-18.25
	August	15.56	15.23-16.54				
<i>Esomus danricus</i> (Darkina)	August			3.68	3.25-4.50		

Hook and line

In *nol borshi* the $L_{50\%}$ value and selective ranges recorded were for 2 species *Channa punctatus* and *C. striatus*. The values recorded were 6.72 cm and 6.57-8.64 cm for *C. punctatus*, 17.34 cm and 15.24-24.25 cm for *C. striatus* (Table 7). But the values were also determined for *L. calbasu* 16.25 cm and 16.45-20.25 cm in *koba borshi*. The values were also determined 16.45 cm and 15.65-18.25 cm for *M. armatus* in *chip borshi*. Rahman *et al.* (1999) recorded values and selective range 13.02 cm and 11.54-14.88 cm for *C. punctatus* in *nol borshi*. However, from the above finding these gears may be regarded as more or less selective and non detrimental gears in old Brahmaputra River.

Table 7. The $L_{50\%}$ values and Selectivity ranges ($L_{25\%}$ - $L_{75\%}$) of *Nol borshi*, *Koba borshi*, *Chip borshi* for major dominant species in the old Brahmaputra River

Species	Hook and Line					
	<i>Nol borshi</i>		<i>Koba borshi</i>		<i>Chip borshi</i>	
	$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$	$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$	$L_{50\%}$	$L_{25\%}$ - $L_{75\%}$
<i>Channa punctatus</i> (Taki)	6.72	6.57-8.64				
<i>Mastacembelus armatus</i> (Baim)					16.45	15.65-18.25
<i>Channa striatus</i> (Shol)	17.94	15.24-24.25				
<i>Labeo calbasu</i> (Kalibaush)			16.25	16.45-20.25		

Conclusion

Study on present status of fishing gears operated in the old Brahmaputra River revealed that some gears were selective while others were not which were used to fish in the river. *Fash jal* with a mesh size greater than 90 mm was found to be harmless to stocked carp and cat fish, therefore, nets with mesh size <90 mm should be regulated during the period June to October. Use of FADs like *katha fishery* should be prohibited and various awareness/training programs should be extended toward fishers as a rationale for the restrictions on a particular fishing gear during particular season.

References

- Bangladesh Centre for Advanced Studies (BCAS). 1991. Floodplain Production Monitoring. Initial Study Report. BCAS , Dhakha, Bangladesh. 99 pp.
- DoF. 2013. Jatiya Matshya Saptah. Department of Fisheries, Government of the Peoples Republic of Bangladesh.
- Hussain. M. G. 2010. Freshwater fishes of Bangladesh: Fisheries, Biodiversity and Habitat.
- Paul, S.K., Rahman, Razzaque, A. and Chakraborty, B.. 1993. Gear selectivity of the floodplain. Progress Report (June-September 1993). Fisheries Research Institute, TFP, Santahar, Bogra. p. 1-27.
- Pope, J., Margetts, A. and Hamely, J.M. 1975. Manual of methods for fish stock assessment. Part III. Selectivity of fishing gear. FAO Fisheries Technical Paper, 41 (Rev.1): 65.
- Rahman, S., Chakraborty, B., Razzak, A., Paul, S.K. and Chu-Fa, T.S. 1993. The fishing gear selectivity study. Preliminary Progress Report (May, 1993). Fisheries Research Institute, TFP, Santahar, Bogra. p. 1-27.
- Rahman, S., Paul, S.K., Chu-Fa, T.S. 1992. The fishing gear selectivity study. Progress Report. Fisheries Research Institute. Thrid fisheries Project, Santahar, Bogra. P.1-20.
- Rahman, S., Mazid, M.A., Kamal, M. and Hossain, M.A.1999. Studies on size selectivity caught in fishing gears BSKB beel, a floodplain in Bangladesh. Bangladesh J. Fish. 22 (2): 9-19.
- Rahman, S., Razzaque, A. and Islam, A.K.M.S. 2009. Study of Fishing Gear Selectivity of Chanda Beel, a Floodplain in Bangladesh. American-Eurasian Journal of Sustainable Agriculture, 3(4): 804-815.
- Mortuza M.R., Rashid, U.S., Rajib, M.A. and Rahman, M.M. 2007. Temporal Variation Characteristics of Flow and Water Level in the Old Brahmaputra River. In: Report on Climate Change Project No. BUET 20/11 GPN 12, p. 39.