



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

Analysis on Nutritional and Functional Components of Different *Pueraria lobata* Roots

Jinping WU¹, Fengling GUO¹, Xianqiang SHAO², Yan HU³, Jianjun ZHAO⁴, Zhengming QIU^{1*}

1. Institute Industrial Crops, Hubei Academy of Agricultural Sciences, Wuhan 430064, China; 2. Hubei Gewa Food Co. Ltd. Zhongxiang 445000, China; 3. Wuhan Gujun Plant Professional Cooperatives, Wuhan 430223, China 4. Ezhou Linjiang Township Xinchang Agricultural Products Professional Cooperatives, Ezhou 436035, China

Abstract This paper made analysis and evaluation of nutritional components and functional components of different *Pueraria lobata* roots. Nutritional components mainly include water, ash content, fat, reducing sugar, starch and cellulose; functional components mainly include flavone and polyphenol. *Pueraria lobata* root No. 1 has highest ash content, flavone, and polyphenol but lowest fat, so it is suitable for using as medical *Pueraria lobata* root resource. *Pueraria lobata* root No. 5 has starch content as high as 64.43%, and is recommended using as vegetable and processing into *Pueraria lobata* powder. *Pueraria lobata* root No. 5 has cellulose content as high as 17.79% and is recommended processing into *Pueraria lobata* tablets. Through comparison of nutritional and functional components of different *Pueraria lobata* roots, it is intended to provide reference for variety selection, breeding, production and processing of *Pueraria lobata* roots.

Key words *Pueraria lobata* roots, Nutritional components, Functional components, Analysis

1 Introduction

Pueraria lobata belongs to *Pueraria* DC. of *Leguminosae*. The plant is a member of the pea family, with reddish-purple flower clusters. It is perennial deciduous liana. Its big roots are rich in starch and essential nutritional components for human body and also active components for physiological functions. In China, there are 9 varieties and 2 variants of *Pueraria lobata*^[1]. Except Xinjiang and Tibet, most provinces and regions of China have distribution of *Pueraria lobata*. *Pueraria lobata* food functions as improving memory, suppressing postnatal diseases, and beauty care^[3]. Starch and flavone separated from *Pueraria lobata* roots can be processed into foods clearing heat, removing toxin, preventing and curing high blood pressure, high fat of blood, coronary heart disease, and diabetes, etc^[4]. In March of 1998, the Ministry of Public Health of China listed *Pueraria lobata* into Catalogue of Food and Medicine Homology. With development of China's foreign trade, especially increase in demands of international market for *Pueraria lobata* root powder and *Pueraria lobata* root flavone, the area of cultivating wild *Pueraria lobata* is gradually expanding. However, because *Pueraria lobata* roots have high ability of survival and rich wild resources, all areas input little to variety selection of *Pueraria lobata* roots, local *Pueraria lobata* roots are mainly domesticated germplasm resources, the nutritional and functional components are not well known. In order to better use and develop *Pueraria lobata* roots and provide reference for variety selection of *Pueraria lobata* roots, we made an analysis on major nutritional

and functional components of *Pueraria lobata* roots.

2 Materials and methods

2.1 Materials Samples we used are *Pueraria lobata* roots obtained from two years of growing *Pueraria lobata* roots in experimental base of Hubei Gewa Food Co. Ltd. We cleaned them with tap water, chopped and put them into 105°C baking oven for 20 minutes, dried at 80/105°C to constant weight, and ground samples to powder with pulverizer and screened with 50 mesh sieve, put into reagent bottles and affixed with labels for experiment.

2.2 Methods Water content was measured by direct drying method with reference to methods stated in GB/T5009.3–2003. Ash content was measured by methods specified in GB/T 5009.4–2003. Starch was measured by enzymic hydrolysis method with reference to methods stated in GB/T5009.9–2003. Fat was measured with acid hydrolysis method with reference to methods stated in GB/T 5009.6–2003. Total dietary fiber was measured by methods stated in GB/T 5009.10–2003. Reducing sugar was measured by direct titration with reference to methods stated in GB/T5009.7–2003. Flavone was measured by sodium nitrite - aluminum nitrate - sodium hydroxide colorimetric method. Polyphenol was measured using Folin-Ciocalteu method.

3 Results

3.1 Nutritional components of *Pueraria lobata* roots Ash content is mainly mineral salts or inorganic salts, including essential inorganic salts or minerals for human body, such as calcium (Ca), magnesium (Mg), potassium (K), sodium (Na), and sulfur (S), accounting for 80% of ash content weight, as well as trace elements such as iron (Fe), copper (Cu), zinc (Zn), manganese (Mn), selenium (Se). From Table 1, we can see that *Pueraria lobata* root No.1 has the highest ash content, next is

Received: April 10, 2015 Accepted: May 21, 2015

Supported by Special Project for Construction of Modern Agricultural Industrial Technology System of Ministry of Agriculture (nycytx-35-02-06) and Hubei Public Science and Technology Program (2014BBA183).

* Corresponding author. E-mail: 18971632353@163.com

Pueraria lobata root No. 2. With constant deepening of nutrition research and constant upgrade of diet concept, it is generally acknowledged that cardiovascular disease, high fat of blood, hypertension, obesity and cancer are connected with high intake of fat. Table 1 indicates that *Pueraria lobata* root No. 1 has the lowest fat content, next is *Pueraria lobata* root No. 2. Starch of *Pueraria lobata* roots has a series of unique features, such as transparency of starch paste, high viscosity, starch gel not easy to syneresis, and as well as unique fragrance of *Pueraria lobata* roots^[4]. Therefore, foods made of starch of *Pueraria lobata* roots are characterized by special nutrition and can be used as foods and medicine. From Table 1, we know that *Pueraria lobata* root No. 5 has starch content as high as 64.43%, followed by *Pueraria lobata* root No. 8 and

Pueraria lobata root No. 7, respectively 54.37% and 53.93%. Cellulose features high water holding capacity and chelation and can be exchanged with anion, improve bacteria microenvironment of digestive system, so it plays extremely important role in human health. Its functions include easing constipation and preventing intestinal cancer, preventing diabetes and controlling obesity, reducing blood pressure and preventing and curing coronary heart disease, preventing cancer, preventing dental caries, and protecting oral health^[5]. Besides, cellulose can be used as excellent food additives^[6-7]. From Table 1, it can be seen that *Pueraria lobata* root No. 4 has cellulose as high as 17.97%, next is *Pueraria lobata* root No. 2 and *Pueraria lobata* root No. 1.

Table 1 Nutritional components of *Pueraria lobata* roots (Unit: %)

Number of <i>Pueraria lobata</i> roots	1	2	3	4	5	6	7	8
Water content	69.06	57.95	68.12	62.24	60.56	73.79	58.37	57.85
Ash content	8.29	6.28	4.50	5.08	4.72	5.46	3.35	4.25
Fat content	2.80	3.62	6.71	5.45	5.65	4.80	5.92	4.77
Reducing sugar	16.00	9.22	12.71	7.37	19.39	14.31	17.74	17.81
Starch	43.92	14.58	27.67	9.32	64.43	35.12	53.93	54.37
Cellulose	14.03	14.16	5.30	17.97	4.44	7.15	1.37	2.74

3.2 Functional components of *Pueraria lobata* roots With constant development of modern medical and food science and technologies, use value of *Pueraria lobata* roots receives higher and higher concern^[8]. Major functional components of *Pueraria lobata* roots are flavonoids, aromatics, triterpenoids, polyphenol, etc. Flavone functions as expanding coronary artery and cerebral blood vessels, improving cardiovascular and cerebrovascular circulation, reducing myocardial consumption of oxygen, strengthening body immunity, reducing blood sugar and anti-oxidation^[9-10]. Polyphenol has high anti-oxidation and ability of removing free

radical. In addition, polyphenol has high ability of absorbing ultraviolet and far-ultraviolet, thus, it can reduce the incidence of such diseases as cancer, cardiovascular diseases, arthritis, and chronic diseases^[11]. Table 2 lists polyphenol and flavone content of 8 samples of *Pueraria lobata* roots. From Fig. 1, we can see that polyphenol content is directly proportional with flavone content. *Pueraria lobata* root No. 1 has the highest polyphenol and flavone, next is *Pueraria lobata* root No. 4, followed by *Pueraria lobata* root No. 3.

Table 2 Functional components of *Pueraria lobata* roots (Unit: %)

Number of <i>Pueraria lobata</i> roots	1	2	3	4	5	6	7	8
Polyphenol content	1.49	1.24	1.09	1.30	0.54	0.53	0.18	0.27
Flavone content	2.78	1.91	1.69	2.55	0.43	0.49	0.02	0.09





Fig. 1 Diagram for polyphenol and flavone content of *Pueraria lobata* roots

4 Conclusions

In the selective breeding of *Pueraria lobata* varieties, it is recom-

mended to select varieties with high yield, high resistance of disease and suitability according to different purposes of use. For ex-

ample, it is recommended to select varieties with high starch content to process powder of *Pueraria lobata* roots. For medical purpose, it is recommended to select varieties with high content of polyphenol and flavone. For processing into tablets of *Pueraria lobata* roots, it is recommended to select varieties with high content of cellulose. For fresh eating or making dishes, it is recommended to select varieties with low content of cellulose and few sclerotic cells. *Pueraria lobata* root is a wild natural plant resource still not developed fully. With constant deepening of extraction, separation, and detection technologies of effective components of *Pueraria lobata* roots, as well as in-depth understanding of pharmacological functions, application value of *Pueraria lobata* roots receives higher and higher concern. As a type of easily cultivated cash crop, *Pueraria lobata* has wide suitability for soil, including sandy and clay soil. In intense sunshine and high temperature, *Pueraria lobata* still can grow. Besides, resistance to aridity and barren land is also a special characteristic. Therefore, *Pueraria lobata* is suitable for planting in mountain areas, especially for economic development of poor mountain areas, it is of realistic significance for farmers shaking off poverty and setting out on the road to prosperity. For example, if planting 66700 hectare *Pueraria lobata* using standardized cultivation technologies, it is able to harvest 2 million tons of fresh *Pueraria lobata* roots, which can replace 0.652 million tons of grain. Calculating at 9000 kg/ha grain, it is equivalent to increasing 72000 hectare farmland and providing grain for 2 million people for one year^[12]. Therefore, planting and developing *Pueraria lobata* roots have prominent economic benefits and important social significance.

(From page 70)

4 Discussions and conclusions

From this experiment, it can be concluded that planting density has significant effect on the yield of *Coix Lachryma-jobi* L., while the effect of fertilizer level and number of seedlings per hole is no significant on the yield of *Coix Lachryma-jobi* L. This indicates that in the planting of *Coix Lachryma-jobi* L., planting density is a major factor influencing yield of *Coix Lachryma-jobi* L. In the actual planting, it is recommended to select appropriate planting density to ensure high yield of *Coix Lachryma-jobi* L. The plant height of *Coix Lachryma-jobi* L is as high as 1.5 – 2.0 m, mutual shading situation is serious. With increase of planting density, the tillering rate, number of grains per plant and weight of thousand grains decrease. Therefore, properly sparse planting is favorable for increasing high yield of *Coix Lachryma-jobi* L. Besides, the planting density is mutually influenced with the basic number of seedlings^[6]. When the planting density increases, although the tillering rate, number of ears per hole, and number of grains per ear decrease, since the basic number of seedlings is high, it still can realize high yield. Therefore, in the planting of

References

- [1] WU DL, CHEN ZY. Study on Chinese *Pueraria* DC. [J]. Journal of Tropical and Subtropical Botany, 1994, 2(3): 12 – 21. (in Chinese).
- [2] FENG RZ, CHEN BZ. Survey of *Pueraria lobata* resource [J]. Chinese Pharmaceutical Journal, 1993, 28(5): 273. (in Chinese).
- [3] JIANG GJ. On the development and utilization, product processing and market prospect of "wood raw *Pueraria lobata*" [J]. Bulletin of Agricultural Science and Technology, 2002(6): 38. (in Chinese).
- [4] YIN W, WANG S. The character of *pueraria lobata* starch, chemical composition and pharmacological function of *Pueraria lobata* [J]. Grain Processing, 2008, 33(1): 84 – 86. (in Chinese).
- [5] DING ZM. Discussion the effects of dietary fiber on human health [J]. Chinese Journal of Misdiagnosis, 2009, 2(5): 1070. (in Chinese).
- [6] HAN Y. Study on the degradation conditions of *Aspergillus niger* to kelp cellulose [J]. Jiangsu Agricultural Sciences, 2010(3): 358 – 360. (in Chinese).
- [7] WANG Y, ZHU DY, ZHOU JZ, *et al.* Identification and phylogenetic study of bacterial cellulose – producing strain from water kefir [J]. Jiangsu Journal of Agricultural Sciences, 2009, 25(2): 446 – 448. (in Chinese).
- [8] LIU YJ, WAN XC. On the development of radix *puerariae* germplasms and its utilization [J]. Forest By – product and Speciality in China, 1998, 45(2): 1. (in Chinese).
- [9] TANG J. The study extension of biological activities of plant polysaccharide [J]. Food Research and Development, 2006, 27(5): 130 – 132. (in Chinese).
- [10] LUYR FOOYL. Antioxidant activities of poly phenols from sage (*Salvia officinalis*) [J]. Food Chemistry, 2001, 75: 197 – 202.
- [11] JIA SP, ZENG R, DAN WD, *et al.* Study on the pharmacologic action of plant polyphenol and its application [J]. China Pharmacy, 2009, 20(12): 953 – 955. (in Chinese).
- [12] XIONG LF. Study on *Pueraria lobata* and its cultivation utilization [M]. Changsha: Hunan Science & Technology Press, 2007: 5. (in Chinese).

Coix Lachryma-jobi L., it is recommended to coordinate the relationship between planting density and basic number of seedlings and select an appropriate planting density and basic number of seedlings to ensure high yield of *Coix Lachryma-jobi* L.

References

- [1] PANG XF, ZHANG SW, QU ZC, *et al.* The growth characteristics and high – yield culture technique of *Coix* L. [J]. Shandong Agricultural Sciences, 1996, 28(3): 15 – 17. (in Chinese).
- [2] CAI ZX, LIU HJ, XIE WJ, *et al.* Research advances in *Coix* L. biology [J]. Crops, 2013, 16(2): 17 – 18. (in Chinese).
- [3] ZHAO YJ, LIU ZH. Study on the planting density of *Coix* L. [J]. Journal of Chinese Medicinal Materials, 1992, 15(12): 11 – 13. (in Chinese).
- [4] LIN YZ. Effects of different planting densities and fertilizer supplies on yield and its components in *coix* (*Coix Lachryma-jobi* L.) [J]. Chinese Agricultural Science Bulletin, 2008, 24(6): 217 – 221. (in Chinese).
- [5] TANG QY, FENG MG. Practical statistic analysis and its computer processing platform [M]. Beijing: China Agriculture Press, 1997: 77 – 90. (in Chinese).
- [6] LIN WQ, MEI XQ, SU WJ, *et al.* Studies on the cultural technique of later – ripening jobstears (*Coix lacryma-jobi*) for its high – yielding [J]. Journal of Zhejiang Agricultural Sciences, 2002(2): 57 – 59. (in Chinese).