



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

Effect of Watering Regimes and Water Quantity on the Early Seedling Growth of *Picralima nitida* (Stapf)

Gbadamosi Alaba Emmanuel¹

¹ Department of Plant Science & Biotechnology, Adekunle Ajasin University, Nigeria

Correspondence: Gbadamosi Alaba Emmanuel, Department of Plant Science & Biotechnology, Adekunle Ajasin University, Nigeria. Tel: 234-806-356-7002. E-mail: gbadamosialabae@hotmail.com

Received: July 30, 2013 Accepted: February 8, 2014 Online Published: February 24, 2014

doi:10.5539/sar.v3n2p35 URL: <http://dx.doi.org/10.5539/sar.v3n2p35>

Abstract

Competing land use pattern has necessitated planting of useful forest trees on marginal lands with varying water status. A study was conducted on the effect of watering regimes and water quantity on seedlings growth of *P. nitida*. Seedlings were raised from seeds, transplanted in polythene bags and subjected to watering at full capacity daily, once in 3 days, 5 days; 20 ml daily, 3 days, 5 days; 15 ml daily, 3 days, 5 days; 10 ml daily, 3 days, 5 days; and flooding. The growth traits assessed were plant height, collar diameter, number of leaves; shoot, root and leaf dry weights as well as leaf area. The effect of watering regime and water quantity were significant ($P \leq 0.05$) on quantitative morphological parameters of seedlings. Total dry weight and root dry weight were lower in both water-stressed and waterlogging than well-watered conditions, but the lowest values were obtained under flooding. However, specific leaf area (SLA) and leaf area ratio (LAR) were higher in water-stressed conditions than well-watered conditions. Relative growth rate of seedlings is related more to the net assimilation rate values than LAR and SLA. Also, root/shoot ratio was highest under daily full capacity watering. Thus, daily watering at full capacity with water availability or 15-20 ml of water per seedling during scarcity is recommended for raising *P. nitida* seedlings.

Keywords: watering regime, biomass, trait, *Picralima nitida*

1. Introduction

The changing weather conditions triggered by climatic change have elicited diverse reactions from plants and animals. The changing rainfall pattern with its attendant unpredictable flooding and drought conditions around the globe necessitates a new approach to research in regeneration of useful forest species.

Water is an important factor in the growth, development and productivity of plants. Hartmann et al. (2005) reported that water stress due to drought is the most significant abiotic factor limiting plant growth and development. Water stress drastically decreased fresh and dry weight, leaf number, total leaf area and stomatal conductance (Vandoorne et al., 2012). However, the present realities favoured irrigation of food crops to feed the ever increasing human population at the expense of tree species; some of which have direct impact on the food, health and general well-being of rural communities.

Plant species respond differently to water availability, also, different plant parts adapt differently to varying water stress conditions. Seeds of many crop species are sensitive to flooding stress during germination (Sesay, 2009; Wuebker et al., 2001; Sung, 1995); while prolonged flooding eliminates some species yet favour others (Van der Valk, 1994 cited in Casanova & Brock (2000)). Seabloom et al. (2001) opined that the depth of flooding can also have significant effect on species composition and biomass of establishing plants. Roots play an important role in plant survival during periods of drought (Hoogenboom et al., 1987). Hsiao and Xu (2000) reported that under water deficiency, growth is readily inhibited and growth of roots is favoured over that of leaves. Often, leaves of plants growing in water stressed environment are small both in number and size.

The study species *Picralima nitida* is a tropical tree of the family Apocynaceae. Its parts are highly medicinal, the dried seeds are used in traditional medicine throughout West Africa; the seeds are crushed or powdered and taken orally, and are mainly used for the treatment of malaria (Kapadia et al., 1993). Duwiejua et al. (2002) reported that the seed contain a mixture of alkaloids producing antipyretic and anti-inflammatory effects along with analgesia. Extracts of the seeds have been reported to have opioid activity in five alkaloids namely akuammidine,

akuammine, akuammicine, akuammigine and pseudoakuammigine (Meenzies et al., 1998).

This useful species occurs only in the wild under no form of management but with a threat of extinction, as part of the domestication strategy therefore; this study was carried out to determine the water requirements at the nursery for its subsequent inclusion in agro ecosystem.

2. Materials and Method

Mature fruits of *Picralimanitida* were collected from Benin City, Nigeria (lat. 6°23'N and long. 5° 13'E). Seeds were extracted from the fermented fruits manually and the seed coats removed to hasten germination (Gbadamosi, 2013); thereafter sown in washed river sand. The experiment was set up at the National Centre for Genetic Resources and Biotechnology, Moor plantation; Ibadan, Nigeria (lat. 7° 24'N and long. 3°49'E) in August 2011. At the two-leaf stage, young seedlings were transplanted into medium sized polythene bags measuring 12 × 14 × 16 cm and allowed to wean under shade for three weeks and watered daily.

Later, seedlings were laid out in completely randomized design in the glasshouse. The following irrigation regimes and water quantity were administered to the seedlings viz: watering at full capacity daily; watering at full capacity once every 3 days; watering at full capacity once every 5 days; 20 ml daily watering; 15 ml daily; 10 ml daily; 20 ml once every 3 days; 15 ml once every 3 days; 10 ml once every 3 days; 20 ml once every 5 days; 15 ml every 5days; 10 ml every 5days; and flooding. There were 15 single-plant replicates under each treatment. Flooding condition was accomplished by placing each seedling inside a bowl filled with water which was regularly replenished.

Growth parameters such as height, collar diameter, number of leaves and internode length were measured fortnightly for 5 months. Height of the seedlings were measured from the collar to the tip of the apical bud using a ruler calibrated in centimeter, collar diameter readings were taken using venier caliper while the number of leaves of each seedling were counted and recorded. At 3 months after transplanting, three seedlings were randomly selected every fortnight under each irrigation treatment for biomass assessment. These seedlings were separated into root and shoot components. The leaf area of each seedling was determined using the Leaf Area Meter. The root and shoot components of each seedling were put into separate envelopes for ease of identification. Each envelope was weighed using an electronic weighing balance to determine the fresh weight of the samples after which the envelopes together with their contents were oven dried for 24 h at 80 °C. The samples were reweighed to get the dry weights. The data obtained were subjected to ANOVA, using statistical analysis system (SAS 9.1) and means were separated using Duncan's Multiple Range Test (DMRT) at 5% level of significance.

The raw data were used to calculate the following growth parameters:

$$\text{NAR (Net assimilation rate)} = \frac{(\text{dry weight at end} - \text{dry weight at start})(\log_e \text{ leaf area at end} - \log_e \text{ leaf area at start})}{(\text{time at end} - \text{time at start})(\text{leaf area at end} - \text{leaf area at start})}$$

$$\text{RGR}_{\text{DW}} \text{ (Relative growth rate by dry weight)} = \frac{\ln(\text{dry weight at end}) - \ln(\text{dry weight at start})}{\text{duration of experiment}}$$

$$\text{RGR}_{\text{LA}} \text{ (Relative growth rate by leaf area)} = \frac{\ln(\text{leaf area at end}) - \ln(\text{leaf area at start})}{\text{duration of experiment}}$$

$$\text{SLA (Specific leaf area)} = \frac{\text{leaf area}}{\text{dry leaf biomass}}$$

$$\text{LAR (Leaf area ratio)} = \frac{\text{leaf area}}{\text{dry plant biomass}}$$

$$\text{RSR (Root : shoot ratio)} = \frac{\text{dry root biomass}}{\text{dry shoot biomass}}$$

3. Result

3.1 Seedlings Height

The effect of watering regime and water quantity were significant ($P \leq 0.05$) on the height of seedlings (Table 1). Overall, seedlings watered with 10 cl at 3 days interval had the highest mean value of 13.17 cm; this was

followed by seedlings which received 20 cl of water daily with a mean value of 12.85 cm while the lowest mean value of 8.5 cm was obtained from seedlings subjected to flooding. The mean height of seedlings subjected to flooding is significantly different from the mean value of seedlings under other treatments (Table 2). Height of seedlings was highly correlated with the number of leaves (Table 3).

Table 1. Summary of analysis of variance for growth variables of seedlings of *P. nitida* under different watering regimes and water quantities

SV	df	Height		Collar diameter		Number of leaves		Internode length	
		MS	F	MS	F	MS	F	MS	F
Treatment	12	207.82	7.47*	172.59	0.96 ^{ns}	195.33	13.31*	6.3	10.78*
Error	1937	27.82		179.14		14.68		0.58	
Total	1949								

*significantly different at 5% probability; ns- not significantly different at 5%.

Keys: SV- Source of variation; MS- Mean square; F- F value.

3.2 Internode Length

There were significant differences in internode length of seedlings due to watering regime and water quantity (Table 1). Seedlings which received 10 cl of water at 3 days interval had the mean value of 2.24 cm, those that were watered with 10 cl daily followed with a mean value of 2.05 cm while seedlings watered at full capacity daily had the lowest mean value of 1.60 cm. The mean internode length of seedlings that received 10 cl of water at 3 days interval was significantly different from the values of seedlings under other treatments (Table 2). There were positive correlations among growth variables assessed (Table 3).

Table 2. Means and standard errors of growth variables of seedlings of *P. nitida* under different watering regimes and watering quantities

Treatment	Height	Collar diameter	Number of leaves	Internode length
FC daily	12.16 ^a ±0.43	4.55 ^a ±0.09	9.23 ^a ±0.31	1.60 ^d ±0.05
FC 3 days	12.04 ^a ±0.47	4.39 ^a ±0.09	8.94 ^a ±0.34	1.64 ^d ±0.05
FC 5 days	12.70 ^a ±0.46	4.42 ^a ±0.1	9.38 ^a ±0.32	1.66 ^d ±0.05
20 cl daily	12.85 ^a ±0.45	4.43 ^a ±0.09	9.33 ^a ±0.32	1.65 ^d ±0.05
15 cl daily	12.72 ^a ±0.45	4.19 ^a ±0.93	8.59 ^a ±0.33	1.76 ^d ±0.08
10 cl daily	12.79 ^a ±0.42	4.38 ^a ±0.10	9.23 ^a ±0.31	2.05 ^b ±0.07
20 cl 3 days	12.31 ^a ±0.43	4.30 ^a ±0.10	9.41 ^a ±0.33	1.60 ^d ±0.06
15 cl 3 days	12.41 ^a ±0.46	4.31 ^a ±0.10	9.25 ^a ±0.32	2.01 ^b ±0.07
10 cl 3 days	13.17 ^a ±0.41	4.35 ^a ±0.09	9.39 ^a ±0.32	2.24 ^a ±0.08
20 cl 5 days	12.15 ^a ±0.47	4.14 ^a ±0.11	8.46 ^a ±0.33	1.59 ^d ±0.05
15 cl 5 days	12.84 ^a ±0.45	4.21 ^a ±0.11	9.01 ^a ±0.33	1.75 ^{cd} ±0.07
10 cl 5 days	12.73 ^a ±0.45	4.27 ^a ±0.10	9.17 ^a ±0.33	1.78 ^{cd} ±0.07
Flooding	8.5 ^b ±0.13	4.40 ^a ±0.06	5.15 ^b ±0.11	1.90 ^{bc} ±0.06

*Mean values with the same letter in a column are not significantly different at 5% probability.

Table 3. Correlation analysis of growth variables of seedlings of *P. nitida* under different watering regimes and water quantities

Parameter	Height	Collar diameter	Number of leaves	Internode length
Height	-	0.11	0.91	0.13
Collar diameter	0.11	-	0.11	0.01
Number of leaves	0.91	0.11	-	0.06
Internode length	0.13	0.01	0.06	-

3.3 Number of Leaves

Number of leaves differed significantly ($P \leq 0.05$) among seedlings subjected to different watering regimes and water quantity (Table 1). The highest number of leaves (9.41) was obtained in seedlings watered with 20 cl of water at 3 days interval; which was followed by seedlings which received 10 cl of water at 3 days interval with a mean value of 9.39 while the lowest value of 5.15 was obtained in seedlings grown under flooding conditions. The mean number of leaves of seedlings under flooding was significantly different from those of seedlings under other treatments (Table 2). Number of leaves was highly correlated with the height of seedlings (Table 3).

3.4 Collar Diameter

Collar diameter of seedlings were not significantly ($P \leq 0.05$) different under watering regime and water quantity. Seedlings watered daily at full capacity had the highest mean value of 4.55 cm; this was followed by seedlings which received 20 ml water daily (4.43 cm). The lowest mean value of 4.14 cm was obtained in seedlings which received 20 ml of water 5 days interval (4.14 cm) (Table 2). Collar diameter was positively correlated with other growth parameters assessed (Table 3).

3.5 Biomass of Seedlings

3.5.1 Shoot Dry Weight (SDW)

The effect of water quantity and watering regime were significant ($P \leq 0.05$) on SDW of *P. nitida* (Table 4). Seedlings watered daily with 20 ml water had the highest mean SDW of 2.29 g followed by seedlings under 15 ml daily and 10 ml in 3 days watering with a mean value of 2.03 g. Seedlings under flooding condition had the lowest mean value of 1.15 g. There were significant differences in mean SDW values of seedlings watered daily with 20 ml and those given 10 ml in 5 days and seedlings under flooding condition (Table 5). SDW was highly correlated with RDW, LDW and TDW (Table 6).

Table 4. Summary of analysis of variance for biomass parameters of seedlings of *P. nitida* under different watering regimes and water quantities

Parameter	SV	DF	MS	F	Pr>F
SDW	Treatment	12	1.87	2.33*	0.0073
	Error	182	1.41		
	Total	194			
RDW	Treatment	12	3.82	2.46*	0.0054
	Error	182	1.55		
	Total	194			
LDW	Treatment	12	9.64	4.58*	<0.0001
	Error	182	2.11		
	Total	194			
TDW	Treatment	12	35.32	2.65*	0.0027
	Error	182	13.32		
	Total	194			
LA	Treatment	12	121070.49	8.20*	<0.0001
	Error	182	14771.28		
	Total	194			

*Significantly different at $P \leq 0.05$

Keys: SV- Source of variation; MS- Mean square; F- F value.

3.5.2 Root Dry Weight (RDW)

There were significant ($P \leq 0.05$) differences in RDW of seedlings under varying watering regime and quantity (Table 4). The highest mean value of 2.85 g was recorded in seedlings watered daily at full capacity; seedlings which received 15 ml of water in 3 days had a mean value of 2.47 g while the lowest value of 1.25 g was obtained in seedlings under flooding. The mean RDW of seedlings watered daily at full capacity was significantly different from those under flooding, 15 ml and 10 ml in 5 days (Table 5). RDW is highly correlated with LDW and TDW of seedlings (Table 6)

3.5.3 Leaf Dry Weight (LDW)

The effect of water quantity and watering regime were highly significant ($P \leq 0.05$) on LDW of *P. nitida* seedlings (Table 4). Seedlings which received 10ml of water daily had the highest mean value of 4.09g, this was followed by seedlings watered at full capacity every 5 days with a mean value of 4.05 g. The lowest mean value of 1.58 g was obtained in seedlings under flooding condition. The mean LDW of seedlings watered at full capacity every 5 days and those watered daily with 10ml were significantly different from that watered with 10ml every 5 days and flooded seedlings (Table 5). LDW of seedlings had positive correlation with SDW, RDW and TDW.

Table 5. Means and standard errors of biomass parameters of seedlings of *P. nitida* under different watering regimes and water quantities

Treatment	SDW (g)	RDW (g)	LDW (g)	TDW (g)	LA (cm ²)
FC daily	1.86 ^{abc} ±0.23	2.85 ^a ±0.36	3.91 ^{ab} ±0.37	8.63 ^a ±0.85	363.39 ^{abc} ±33.64
FC 3 days	1.62 ^{abc} ±0.31	1.5 ^{cd} ±0.24	3.35 ^{abcd} ±0.37	6.5 ^{abcd} ±0.81	374.15 ^{ab} ±29.95
FC 5 days	1.88 ^{abc} ±0.35	2.11 ^{abcd} ±0.38	4.05 ^a ±0.54	8.05 ^{ab} ±1.24	423.3 ^a ±52.84
20 cl daily	2.29 ^a ±0.35	2.46 ^{ab} ±0.36	3.82 ^{ab} ±0.33	8.57 ^a ±1.00	410.6 ^a ±13.52
15 cl daily	2.03 ^{ab} ±0.31	1.94 ^{bcd} ±0.31	3.42 ^{abc} ±0.32	7.39 ^{abc} ±0.87	310.25 ^{bc} ±28.94
10 cl daily	1.7 ^{abc} ±0.29	1.5 ^{cd} ±0.25	4.09 ^a ±0.41	7.69 ^{abc} ±0.91	413.85 ^a ±37.30
20 cl 3 days	1.58 ^{abc} ±0.33	2.07 ^{abcd} ±0.38	2.95 ^{bcd} ±0.41	6.60 ^{abcd} ±1.06	314.43 ^{bc} ±38.89
15 cl 3 days	1.99 ^{abc} ±0.36	2.47 ^{ab} ±0.37	3.43 ^{abc} ±0.41	7.88 ^{ab} ±1.11	279.47 ^c ±21.63
10 cl 3 days	2.03 ^{ab} ±0.38	2.41 ^{abc} ±0.41	3.57 ^{ab} ±0.45	8.01 ^{ab} ±1.18	308.65 ^{bc} ±18.98
20 cl 5 days	1.81 ^{abc} ±0.38	1.63 ^{bcd} ±0.37	2.32 ^{def} ±0.38	5.75 ^{bcd} ±1.08	179.76 ^d ±25.04
15 cl 5 days	1.29 ^{bc} ±0.13	1.45 ^d ±0.18	2.48 ^{cdef} ±0.25	5.22 ^{cd} ±0.47	366.67 ^{abc} ±34.85
10 cl 5 days	1.16 ^c ±0.09	1.29 ^d ±0.16	2.17 ^{ef} ±0.20	4.61 ^d ±0.28	376.4 ^{ab} ±27.74
Flooding	1.15 ^c ±0.31	1.25 ^d ±0.30	1.58 ^f ±0.32	3.98 ^d ±0.89	125.06 ^d ±24.85

Means with the same letter in a column are not significantly different.

Table 6. Correlation analysis of biomass parameters of seedlings of *P. nitida* under different watering regimes and water quantities

Parameter	SDW	RDW	LDW	TDW	LA
SDW	-	0.81	0.79	0.92	0.24
RDW	0.81	-	0.81	0.94	0.26
LDW	0.79	0.81	-	0.94	0.5
TDW	0.92	0.94	0.94	-	0.37
LA	0.24	0.26	0.5	0.37	-

3.5.4 Total Dry Weight (TDW)

The TDW of seedlings were significantly ($P \leq 0.05$) different under varying watering regimes and water quantities (Table 4). Seedlings watered at full capacity had the highest mean value of 8.63 g followed by seedlings that received 20 ml of water daily (8.57 g). The lowest mean value of 3.98 g was obtained in seedlings under flooding conditions. The mean TDW of seedlings watered with 20 ml of water; and at full capacity daily were significantly different from seedlings given 10, 15 and 20 ml water every 5 days; and those under flooding condition (Table 5). TDW had high positive correlation with SDW, RDW and LDW (Table 6).

3.5.5 Leaf Area (LA)

The effect of water quantity and watering regime were significant ($P \leq 0.05$) on LA of seedlings (Table 4). Seedlings watered at full capacity every 5 days had the highest mean LA value of 423.3 cm² followed by seedlings which had 20 ml of water daily with a mean value of 410.6 cm². The lowest mean value of 125.06 cm² was obtained in seedlings under flooding conditions. The mean LA value of seedlings under flooding and those watered with 20 ml every 5 days were significantly different from other treatments (Table 5). Leaf area was lowly but positively correlated with SDW, RDW and TDW (Table 6).

3.5.6 Relative Growth Rate by Dry Weight (RGR_{DW}) and Leaf Area (RGR_{LA})

Relative growth rate (RGR_{DW}) varied widely among treatments. The highest value of 0.0638 g/g/wk was obtained in seedlings watered at full capacity every 5 days while the lowest value of -0.0391 g/g/wk was obtained in seedlings watered with 20 ml every 5 days. RGR_{LA} values ranges from -0.0824 g/g/wk in seedlings watered with 20 ml of water every 5 days to 0.0635 g/g/wk in seedlings which had 20 cl of water every 3 days (Table 7).

Table 7. Computed growth parameters of seedlings of *P. nitida* under different watering regimes and water quantities

Treatment	RGR _{DW}	RGR _{LA}	SLA	LAR	NAR	RSR
FC daily	0.0412	0.0466	92.94	42.11	3.139 x10 ⁻⁴	1.53
FC 3days	0.0428	0.0051	111.68	57.56	2.420 x10 ⁻⁴	0.94
FC 5days	0.0638	0.0389	104.52	52.58	3.064 x10 ⁻⁴	1.12
20 cl daily	0.0325	0.0127	107.48	47.91	1.867 x10 ⁻⁴	1.07
15 cl daily	0.0626	0.0335	90.72	41.98	0.0031	0.95
10 cl daily	0.0408	0.0571	101.19	53.82	2.862 x10 ⁻⁴	1.12
20 cl 3 days	0.0367	0.0635	106.59	47.64	2.444 x10 ⁻⁴	1.31
15 cl 3 days	0.0561	0.0026	81.48	35.47	4.049 x10 ⁻⁴	1.24
20 cl 5 days	-0.0391	-0.0824	77.48	31.26	-2.556x10 ⁻⁴	0.9
15 cl 5 days	0.0494	0.0286	147.85	70.24	2.689 x10 ⁻⁴	1.12
10 cl 5 days	0.0282	-0.0249	173.27	81.64	1.831 x10 ⁻⁴	1.1
Flooding	0.0437	-0.0661	79.14	31.42	1.737 x10 ⁻⁴	1.08

3.5.7 Root/Shoot Ratio (RSR)

The ratio of root to shoot was highest in seedlings watered at full capacity daily (1.53), while the lowest RSR value of 0.9 was obtained in seedlings which received 20ml of water every 5 days (Table 7).

3.5.8 Specific Leaf Area (SLA) and Leaf Area Ratio (LAR)

Specific leaf area (SLA) values range from 79.14 cm²/g in flooded seedlings to 173.27 cm²/g in seedlings watered with 10 ml in 5 days. LAR nearly followed the same pattern with values of 31.26 cm²/g and 81.64 cm²/g for seedlings watered with 20 and 10 ml of water every 5 days respectively (Table 7).

3.5.9 Net Assimilation Rate (NAR)

Seedlings watered daily with 15ml of water had the highest value of 0.0031 g/cm²/wk, those watered with 15 ml in 3 days had a value of 4.048 × 10⁻⁴ cm²/g while a negative value of -2.556 × 10⁻⁴ cm²/g was obtained in seedlings given 20 ml of water every 5 days (Table 7)

4. Discussion

Water is a major constituent of all living organisms which is involved in important biochemical processes including photosynthesis thus its availability in adequate quantity and at biologically tolerable interval affects productivity of plant species. In this study, total biomass and root dry weights were lower in both water stressed and water logging conditions than well-watered conditions; but the lowest values were obtained under waterlogging. Gonzales et al. (2009) made the similar observation for *Chenopodium quinoa* (Willd.). However, SLA and LAR were higher in drought conditions than well-watered conditions.

Seedlings of *P. nitida* which received the lowest water rate (10 ml) at the longest intervals of 5 days had the lowest biomass value, this agreed with the findings of several authors that drought water stress influences plant

growth and limits productions (Ky-Dembele et al., 2010; Hsiao & Xu, 2000; Busso et al., 1998; Hampson & Simpson, 1990). In this study, excess of water also had negative impact on growth and biomass production of the seedlings. This result agreed with the findings of Sakio (2005) that flooding reduces the total dry weight increment in some plant species. In flooding condition, available oxygen in the soil is usually low thus, limiting nutrient absorption for species not adapted to this condition; and as stated by Predick et al. (2009) growth of flood-intolerant species is often depressed in flooding. *P. nitida* is not a choice species for planting in flooded areas as seedlings were stunted and number of leaves and leaf expansion were very poor in line with the findings of Kozłowski (1997); this response might have reflected the species habitat and growth pattern as suggested by Sakio (2005).

It has been widely reported that extensive root growth is an adaptive feature under drought stressed conditions (Guoxiong et al., 2002); the result of the present study is contrary to this, root dry matter reduced under increased water deficit. The ratio of root to shoot was highest in seedlings under full capacity daily watering. This is in agreement with the finding of Vandoorne et al. (2012) that water stress drastically decreased fresh and dry root weight, leaf number, total leaf area and stomatal conductance in *Cichorium intybus* (var. *sativum*).

There were some wide variations in RGR of seedlings among treatments; this is often adduced to variations in SLA (Cornelissen et al., 2003; Karacic & Weih, 2006). Cornelissen et al. (1998) reported that RGR of deciduous species seems to be determined by their higher LAR, which in turn is accounted for by their higher SLA compared to evergreen. The result of the present study revealed that although *P. nitida* is a deciduous species, NAR is a better determinant of RGR as seedlings with higher SLA and LAR under water stress had very poor RGR values. Antunez et al. (2001) seems to support this by inferring that temporal variation in RGR within species was due mainly to variation in NAR.

Ordinarily, plants as part of their survival strategy to reduce water loss during periods of severe water stress may reduce leaf area or formation of new leaves; or shed old ones. Lawlor and Leach (1985) submitted that decrease in LA is a common effect of drought, in this study however, leaf area under the severest water stress tested compared reasonably with those under daily full capacity watering; this is perhaps due to the very waxy cuticle of the leaf.

5. Conclusion

The result from this study seems to suggest that *P. nitida* can be raised with some satisfactory growth under some form of water stress but cannot grow well in flooding conditions. Therefore, further investigation with longer periods of drought may help to reveal the drought resistance traits of this species. Also, these drought conditions would require field studies for verification.

Acknowledgements

This study was executed with support from International Foundation for Science (IFS) through research grant-D/4670-1 (2009).

References

- Antunez, I., Retamosa, E. C., & Villar, R. (2001). Relative growth rate in phylogenetically related deciduous and evergreen woody species. *Oecologia*, *128*, 172-180. <http://dx.doi.org/10.1007/s004420100645>
- Busso, C. A., Fernandez, O. A., & Fedorenko, D. E. F. (1998). Dry weight production and partitioning in *Medicago minima* and *Erodium cicutarium* under water stress. *Annals of Botany*, *82*, 217-227. <http://dx.doi.org/10.1006/anbo.1998.0667>
- Casanova, M. T., & Brock, M. A. (2000). How do depth, duration and frequency of flooding influence the establishment of wetland plants communities? *Plant Ecology*, *147*, 237-250. <http://dx.doi.org/10.1023/A:1009875226637>
- Cornelissen, J. H. C., Cerabolini, B., Castro-Díez, P., Villar-Salvador, P., Montserrat-Martí, G., Puyravaud, J. P., ... Aerts, R. (2003). Functional traits of woody plants: correspondence of species rankings between field adults and laboratory-grown seedlings. *Journal of Vegetation Science*, *14*(3), 311-322. <http://dx.doi.org/10.1111/j.1654-1103.2003.tb02157.x>
- Cornelissen, J. H. C., Castro-Díez, P., & Carnelli, A. C. (1998). Variation in relative growth rate among woody species: scaling up. In H. Lambers, H. Poorter, & M van. Vuuren (Eds.), *Inherent variation in plant growth: physiological mechanisms and ecological consequences* (pp. 363-392). Backhuys, Leiden.
- Duwiejua, M., Woode, E., & Obiri, D. D. (2002). Pseudo-akuammigine, an alkaloid from *Picralima nitida* seeds, has anti-inflammatory and analgesic actions in rats. *Journal of Ethnopharmacology*, *81*, 73-79.

- [http://dx.doi.org/10.1016/S0378-8741\(02\)00058-2](http://dx.doi.org/10.1016/S0378-8741(02)00058-2)
- Gbadamosi, A. E. (2013). Germination biology of *Picralimanitida* (Stapf) under pretreatments. *Greener Journal of Biological Sciences*, 3(1), 13-20.
- Gonzales, J. A., Gallardo, M., Hilal, M., Rosa, M., & Prade, F. E. (2009). Physiological responses of quinoa (*Chenopodium quinoa* Willd.) to drought and waterlogging stresses: dry matter partitioning. *Botanical Studies*, 50, 35-42.
- Guoxiong, C., Krugman, T., Fahima, T., Korol, A. B., & Nevo, E. (2002). Comparative study on morphological and physiological traits related to droughtresistance between xeric and mesic *Hordeum spontaneum* lines in Isreal. *Barley Genet. Newslett.*, 32, 22-33.
- Hampson, C. R., & Simpson, G. M. (1990). Effect of temperature, salt and osmotic potential on early growth of wheat (*Triticum aestivum* L.). I. Germination. *Can. J. Bot.*, 68, 524-528. <http://dx.doi.org/10.1139/b90-072>
- Hartmann, T., College, M., & Lumsden, P. (2005). Responses of different varieties of *Lolium perennetum* to salinity. Annual Conference of the Society for Experimental Biology, Lancashire.
- Hoogenboom, G., Huck, M. G., & Peterson, C. M. (1987). Root growth rate of soybean as affected by drought stress. *Agron J.*, 79, 607-614. <http://dx.doi.org/10.2134/agronj1987.00021962007900040003x>
- Hsiao, T. C., & Xu, L. (2000). Sensitivity of growth of roots versus leaves to water stress: biophysical analysis and relation to water transport. *Journal of Experimental Botany*, 51(350), 1595-1616. <http://dx.doi.org/10.1093/jexbot/51.350.1595>
- Kapadia, G. J., Angerhofer, C. K., & Ansa-Asamoah, R. (1993). Akuammine: an antimalarial indolemonoterpene alkaloid of *Picralimanitida* seeds. *Planta Medica*, 59(6), 565-566. <http://dx.doi.org/10.1055/s-2006-959764>
- Karacic, A., & Weih, M. (2006). Variation in growth and resource utilisation among eight poplar clones grown under different irrigation and fertilisation regimes in Sweden. *Biomass and Bioenergy*, 30(2), 115-124.
- Kozlowski, T. T. (1997). Responses of woody plants to flooding and salinity. *Tree Physiol. Monogr*, 1, 1-29.
- Ky-Dembele, C., Bayala, J., Savadogo, P., Tigabu, M., Odén, P. C., & Boussim, I. J. (2010). Comparison of growth responses of *Khayasenegalensis* seedlings and stecklings to four irrigation regimes. *Silva Fennica*, 44(5), 787-798.
- Lawlor, D. W., & Leach, J. E. (1985). Leaf growth and water deficits: biochemistry in relation to biophysics. In N. R. Baker, W. J. Davis & C. K. Ong (Eds.), *Control of Leaf Growth* (pp. 267-294). England: Cambridge University Press.
- Meenzies, J. R., Paterson, S. J., Duwiejua, M., & Corbett, A. D. (1998). Opioid activity of alkaloids extracted from *Picralimanitida* (fam. Apocynaceae). *European Journal of Pharmacology*, 350(1), 101-108. [http://dx.doi.org/10.1016/S0014-2999\(98\)00232-5](http://dx.doi.org/10.1016/S0014-2999(98)00232-5)
- Predick, K. I., Gergel, S. E., & Turner, M. G. (2009). Effect of flood regime on tree growth in the floodplain and surrounding uplands of Wisconsin River. *River Res. Applic.*, 25, 283-296. <http://dx.doi.org/10.1002/rra.1156>
- Sakio, H. (2005). Effects of flooding on growth of seedlings of woody riparian species. *J. Forest Research*, 10(4), 341-346. <http://dx.doi.org/10.1007/s10310-005-0156-9>
- Seabloom, E. W., Moloney, K. A., & van der Valk, A. G. (2001). Constraints on the establishment of plants along a fluctuating water-depth gradient. *Ecology*, 82, 2216-2232. [http://dx.doi.org/10.1890/0012-9658\(2001\)082\[2216:COTEOP\]2.0.CO;2](http://dx.doi.org/10.1890/0012-9658(2001)082[2216:COTEOP]2.0.CO;2)
- Sesay, A. (2009). Influence of flooding on Bambara groundnut (*Vigna subterranean* L.) germination: Effect of temperature, duration and timing. *African Journal of Agricultural Research*, 4(2), 100-106.
- Sung, J. M. (1995). The effect of sub-optimal O₂ on seedling emergence of soybean seeds of different size. *Seed Sci. Technol.*, 23, 807-814.
- Vandoorne, B., Mathieu, A. S., Van den Ende, W., Vergauwen, R., Perilleux, C., Javaux, M., & Lutt, S. (2012). Water stress drastically reduces root growth and inulin yield in *Cichorium intybus* (var. sativum) independently of photosynthesis. *Journal of Experimental Botany*, 63(12), 4359-4373. <http://dx.doi.org/10.1093/jxb/ers095>
- Wuebker, E. F., Mullen, R. E., & Koehler, K. (2001). Flooding and temperature effects on Soybean germination. *Crop Sci.*, 41, 1857-1861. <http://dx.doi.org/10.2135/cropsci2001.1857>

Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.

This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).