AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

Pesticides and Bees: ecological-economic modelling of bee populations on farmland.

Ciaran Ellis¹, Nick Hanley*², Adam Kleczkowski¹ and David Goulson³

¹ University of Stirling

² University of St Andrews

³ University of Sussex

Contributed Paper prepared for presentation at the 90th Annual Conference of the Agricultural Economics Society, University of Warwick, England

4 - 6 April 2016

Copyright 2016 by [author(s)]. All rights reserved. Readers may make verbatim copies of this document for non-commercial purposes by any means, provided that this copyright notice appears on all such copies.

*[Corresponding author] Email ndh3@st-andrews.ac.uk. Dept of Geography and Sustainable Development, Irvine Building, North Street, St Andrews, Fife KY16 9AL

We thank the European Investment Bank for funding part of the work on which this paper is based.

Abstract

Production of insect-pollinated crops typically relies on both pesticide use and pollination, leading to a potential conflict between these two inputs. In this paper we combine ecological modelling with economic analysis to investigate the effects of pesticide use on wild and commercial bees, whilst allowing farmers to partly offset the negative effects of pesticides on bee populations by creating more on-farm bee habitat. Farmers have incentives to invest in creating wild bee habitat to increase pollination inputs. However, the optimal allocation of on-farm habitat strongly depends on the negative effects of pesticides, with a threshold-like behaviour at a critical level of the impairment. When this threshold is crossed, the population of wild bees becomes locally extinct and their availability to pollinate breaks down. We also show that availability of commercial bees masks the decrease in pollination services which would otherwise incentivise farmers to conserve the wild pollinator population, therefore indirectly leading to local wild bee extinction. The paper demonstrates the importance of combining ecological modelling with economics to study sustainability in the provision of ecosystem services in agro-ecosystems.

Keywords: pollination, pesticides, wild bees, commercial bees, ecological-economic modelling.

1. Introduction

Globally, around three-quarters of food crops are at least partly dependent on insect pollination (Klein et al., 2007), and this share has been rising over the past 50 years (Aizen & Harder, 2009)¹. Ensuring sufficient pollination of these crops will be challenging in the future, due to adverse pressures on the supply of pollination services. Wild insect pollinator populations are threatened by both habitat loss, declines in foraging resources (Carvell et al, 2006; Winfree et al., 2009) and agricultural intensification (Biesmeijer et al., 2006, Cameron et al., 2011), leading to population declines (Cameron et al, 2011; Goulson et al. 2015). Honeybees are used to supplement or substitute wild pollinators, along with other commercial pollinators such as factory-reared bumblebees (Velthuis & Van Doorn, 2006), although the majority of crop pollination for most crops is currently delivered by wild pollinators (Breeze et al. 2011).

Commercial pollinators can be adequate substitutes for wild pollinators for many crops, (Brittain et al., 2013), but the use of commercial pollinators is not without risk. Honeybees have suffered losses in recent years due to the abandonment of hives (Colony Collapse Disorder), the impacts of the *Varroa* mite and associated diseases (Cox-Foster et al., 2007) and falling numbers of bee keepers in some countries (Potts et al, 2010). If losses of honeybees occur over a wide area, there can be an impact on the supply of these insects for pollination services, which can lead to cost increases to farmers². Given the risks associated with reliance on commercial pollination sources, maintaining viable wild pollinator populations is likely to be crucial for sustaining the production of insect-pollinated crops into the future (Winfree et al., 2007; Garibaldi et al, 2013).

¹ The data in Aizen and Harder relates to animal pollination in general, not insect pollination specifically. However, they note that the demand for pollination in agriculture has risen about 6 times more than the population of honey bees over the last 50 years.

² Prices for honeybee hire for use on almond farms doubled between 2006 and 2008 in the US (Pettis & Delaphane 2010).

One of the factors implicated in the decline of insect pollinators is the use of pesticides. There is growing evidence of negative effects of commonly used insecticides on population- determining traits such as foraging rates and navigation in bees, on the overall growth and performance of colonies, and on the pollination services that they provide (Mommaerts et al., 2010; Henry et al., 2012; Gill et al. 2012; Whitehorn et al., 2012; Goulson, 2013, 2015; Rundlöf et al. 2015; Stanley et al. 2015). Awareness of this evidence has led to the temporary banning of the use on flowering crops of a widely used group of insecticides -- neonicotinoids -- within the European Union.

Farmers of insect pollinated crops therefore face a dilemma, as one input (pesticides) is potentially dangerous to another (pollinators). One option, not investigated here, is to switch production to organic principles, and use zero pesticides. However, in the majority of global agricultural systems, abstaining from the use of all pesticides is not usually possible without sacrificing yields. Farmers must either attempt to reduce the impact of pesticides on wild pollinators, or increase the use of commercial pollinators, as these can in some cases be replenished year after year. Wild pollinators require habitat either off-farm or within the farm area. Although pollinating insects can forage over large distances, in intensive agricultural landscapes there is a decay in visitation of flowers by pollinators with increasing distance from the nearest habitat patch (Ricketts et al., 2008; Osborne et al. 2008a). To offset this, farmers can encourage wild bees to nest within foraging distance of crops by providing nesting habitat and providing alternative foraging resources on the farm for when the crop is not in flower (Carvell et al, 2007). The effect of such interventions has been found to be strongest in intensively farmed areas (Carvell et al., 2011), but depends also on the spatial location of bee-friendly habitat (Brosi et al., 2008; Keitt, 2009). Hence, local or field-scale management practices may offset the negative impacts of intensive monoculture agriculture on pollination services to some extent (Kennedy et al., 2013).

In this paper, we develop an ecological-economic model to investigate the relations between two agricultural inputs, pollination and pesticides, and two sources of pollinators with different

characteristics; commercial pollinators, which can be replaced at a cost, and wild pollinators, which rely on a population being sustained within the farm area. Dedicating some of the farm area to sustain wild pollinators (eg by cultivating wild flower strips) is assumed to be costly. The model is parameterised using farm management data for strawberries, a relatively well-studied crop on which both wild and commercial bees are used. The neonicotinoid pesticide thiacloprid is also commonly used in strawberry farming to protect the crop from destructive pests such as capsid bugs. Our modelling framework is, however, generalizable to other cropping systems where conflict occurs between pesticides, crop area and wild bee persistence, such as almonds. Our model differs from previous modelling attempts which have looked at either habitat considerations (Brosi et al., 2008; Keitt 2009) or pesticide impacts (Bryden et al., 2013) in isolation. In contrast, we combine these factors co-determining pollinator populations in a realistically-parameterised model which includes both economic and ecological behaviours.

2. Methods: the ecological economic model.

The model has three main linked components: the dynamics of the wild bee population; the production function which links bee populations and pesticide use to output, and farmers' decisions over which inputs to employ via a profit function. We assume a farm that produces a single crop; parameters are chosen to represent a typical soft-fruit production system (Nix, 2015; Ellis, 2014). The farm has an area A which is divided into a wild bee habitat conservation area, vA , and a cropping area $(1-v)A$, where v is the proportion assigned to the wild bee habitat (we vary this between 0% and 70%). Honeybees and commercially reared bumblebees are both used in fruit production. For simplicity we consider all commercial (non-wild) pollinators to have the characteristics of commercially reared bumblebees in terms of nest size and pollinating efficiency, and generate results for both a scenario where all pollinators are affected by pesticides, and a scenario where wild bees are affected but

commercial bees are not³. For simplicity we are assuming that the farm is a closed system with regard to wild or commercial bees, so that bees are not coming in from surrounding non-farmed habitat or leaving the farm. We also assume no transfer of pesticides from outside the farm.

Wild bee population

The dynamics of the wild bee population is described in terms of $N_{[t]}$ – a number of nests in a given year, t . This changes according to equation (1):

$$N_{[t]} = \min\left(R\left(N_{[t-1]} - D_{[t-1]}\right), K\right) \quad (1)$$

where $N_{[t-1]}$ is the number of nests at the beginning of year $t-1$, $D_{[t-1]}$ represents the number of nests that die during year $t-1$. $N_{[t-1]} - D_{[t-1]}$ represents the number of live nests at the end of year $t-1$ that will reproduce in the following year. R is the reproduction rate, i.e. the number of new nests that each reproducing nest produces in the following year. The carrying capacity, K , is calculated from the likely on-farm nesting densities of wild bumblebees, wN , under the assumption that wild bees nest in the conservation area only, $K = wN \vee A$.

Not all bumblebee nests will produce queens in a given year, and the likelihood of reproduction will depend in part on nest size. Pesticides can indirectly impact the likelihood of a nest reproducing by impairing the performance of foragers or increasing worker mortality and thus decreasing a nests' ability to gather and process resources. These impacts can lead to increased colony failure, either

³ The actual situation may be that commercial pollinators are affected, but to a slightly lesser extent than wild bees; efforts can be made to minimise chemical exposure to commercial nests such as shutting the bees inside the boxes before spraying, or only spraying before the placement of nest boxes. Wild nests, on the other hand, may be exposed to multiple sprays of insecticides and though both wild and commercial bumblebee nests are vulnerable to disease, wild nests are more likely to have infestations of parasites at the time spraying occurs (commercial bee boxes *should* arrive at the farms free from disease and therefore only pick up infections and parasites from that point onwards) putting wild bees at increased risk of any interactive effects between parasites and pesticides (Alaux et al., 2010).

through early colony death or by limiting the number of new queens produced (Gill et al, 2012; Whitehorn et al., 2012; Rundlöf et al. 2015). Nest success after exposure to pesticides has been found to fit a stochastic model of colony death; nests treated with imidacloprid were found to develop as predicted by a model where the probability of nest death was inversely proportional to the number of foragers adjusted for pesticide impairments (Bryden et al., 2013). In our model we use an equivalent deterministic model in which a proportion dN of nests dies in year $t-1$ so that:

$$D_{[t-1]} = dN \cdot N_{[t-1]}. \quad (2)$$

We also consider a stochastic equivalent of model (1), with nest deaths given by a random variable binomially distributed (with the maximum number of $N_{[t]}$ and probability given by dN): results are qualitatively similar to the ones presented here for the deterministic model.

Although in principle dN can depend on time, in this model we assume the constant probability of nest death following (Bryden et al., 2013),

$$dN = \frac{m}{j + wBN} \quad (3)$$

where wBN is an effective number of foraging wild bees per nest, $wBN = wF(1 - wI)$ with wF being an average number of foragers per nest and wI the impairment factor due to pesticides. If no pesticides are used, or if pesticides are used but do not affect bees, $wI = 0$; otherwise $wI > 0$, reflecting for example, the effects on the navigational ability of honeybees which reduces the number of foragers which successfully return to the nest (Henry et al., 2012).

Equation (1) can thus be rewritten

$$N_{[t]} = \begin{cases} R \times \left(1 - \frac{m}{j + wF \times (1 - wI)} \right) N_{[t-1]} & \text{if smaller than } K, \\ K & \text{otherwise.} \end{cases} \quad (4)$$

The initial condition is assumed to be $N_{[0]}=K$ for $t=0$. Under this assumption $N_{[t]}$ will stay constant for $t>0$, as long as:

$$R \times \left(1 - \frac{m}{j + wF \times (1 - wI)} \right) \geq 1 \quad (5)$$

and will decline exponentially to zero otherwise. In the following we assume such parameter values that condition (5) is always satisfied if $wI=0$, i.e. if there is no impairment due to pesticides.

Pollination and yield.

The single crop is pollinated by foragers originating from both wild and commercial nests. The total effective number of foraging wild bees is given by $wB_{[t]} = wF (1-wI) N_{[t]}$, whereas for commercial bees the number is assumed to be constant through time but proportional to the crop area, $cB=cF (1-cI) cN (1-v) A$. Here, cF is the average number of foragers per commercial nest, cI is the impairment of commercial bees due to pesticide use, cN is the number of commercial nests per ha, and $(1-v) A$ is the area under the crop (here we assume that commercial nests will only be placed where the crop is located, not in the area set aside as on-farm wild bee habitat). As for wild bees, if no pesticides are used or are used but have no effect on commercial bees, then $cI=0$.

Both wild and commercial bees are assumed to forage across the whole farm, over both crop land and the conservation area. The resulting effective density of foraging pollinators is then given by:

$$B_{[t]} = \frac{wB_{[t]} + cB}{A} = \frac{wF (1 - wI) N_{[t]} + cF (1 - cI) cN (1 - v) A}{A} . \quad (6)$$

Production.

The total farm production in year t is given by $Y_{[t]} \cdot (1 - v)A$ where $Y_{[t]}$ is the current yield (in tonnes per ha) which is assumed to be a step-wise linear function of $B_{[t]}$. We assume that without pollinators there is a set but low proportion, $aY_{\max}$, of a maximum yield ($Y_{\max}$) that can be achieved. When pollination is fully supplied, the maximum yield is given by $gY_{\max}$ with g being a maximum proportion of high quality crop (Garratt et al., 2014). For intermediate values of $B_{[t]}$ the yield per area in year t is given by:

$$Y_{[t]} = Y_{\max} \cdot \min(g, a + b B_{[t]}) \quad (7)$$

where γ is the maximum proportion of good quality, α is the proportion of good quality fruits without bees and β is the incremental effect of bee visitation. The maximum attainable yield, $Y_{\max}$, depends on pesticide use; we choose a higher value of $Y_{\max}$, $Y_{\max,p}$, if pesticides are used, and a lower value, $Y_{\max,nop}$, if they are not.

Farm economics.

There are two components to the profit function, the income from the sale of the crop and various costs. The crop is sold at price p and with commission cm so that the income is given by:

$$p \cdot (1 - cm) \cdot Y_{[t]} \cdot (1 - v)A. \quad (8)$$

Total costs for each year are the sum of variable (yield dependent) costs and other costs which include the costs of wild flower seeds, pesticides and commercial bees. Harvesting and packaging costs are assumed to be variable and calculated per tonne. We divide the costs into three components, the first one which does not directly depend on the usage of commercial bees or pesticides, given by:

$$C_{pt} \cdot Y_{[t]} \cdot (1 - v)A + C_{pa} \cdot (1 - v)A + C_{apa} \cdot A + C_{seed} \cdot vA \quad (9)$$

where C_{pt} is the cost per tonne (harvesting and packaging), C_{pa} is the cost per crop area (planting, structures, fieldwork), C_{apa} is the total cost per area regardless of whether it is cropped or not (e.g. land lease costs), and C_{seed} is the cost of maintaining the conservation area (mainly providing seed). If commercial bees are used, there is an additional cost of buying commercial nests which is proportional to the number of commercial nests per ha and the area under crop,

$$bC + cN + (1 - v)A. \quad (10)$$

If pesticides are used, there is additional cost associated with their purchase, assumed to be proportional to the area under crop,

$$pC + (1 - v)A. \quad (11)$$

We assume that the primary decision is over the proportion of on-farm wild bee habitat, v , and this is driven by profit maximisation over a decision horizon of one year. We analyse how the optimal choice of v and the resulting profit vary as pesticides are used or not, whether they affect wild or commercial bees, and whether the farmer decides to use commercial bees.

Parameters.

Although the model is generic, we calibrated it to soft fruit production in the UK (Nix, 2015; Ellis, 2014). The numerical values for parameters used are listed in Table 1. K is calculated from the likely on-farm nesting densities of wild bumblebees⁴. While actual densities will vary between locations, we assume that densities of 15 nests per ha can be found in on-farm habitat and assume that no nesting can occur within the cropped area. We follow Bryden et al. (2013) in describing the effect of pesticide impairments on the dynamics of wild nests (Table 1). Costs of seeds, pesticides and bumblebee boxes are taken from a farm survey of 25 soft-fruit farms in Scotland (Ellis, 2014). Other production costs and prices per ha are taken from farm management data from the Farm Management

⁴ Nest densities will depend on the landscape type; around 11 to 15 nests per ha were found in non-linear countryside in a large scale survey in UK habitats, with higher densities in gardens and around linear features (Osborne et al., 2008b).

Pocketbook 2016 eds. (Nix, 2015), corresponding to raised-bed June-bearing strawberries (Nix, 2015, p. 35).

3. Results

We first analyse the optimal levels of conservation area provision in the absence of pesticide use and commercial bees. The effect of pesticide on wild bees is considered next and then provision of commercial bees is considered, without and with the impact of pesticides on their ability to pollinate.

RESULT 1: When no commercial bees or pesticides are used, profits are negative without on-farm wild bee habitat, and peak at low-moderate levels of its provision. Allowing for pesticide use shifts the yield and therefore the profit upwards, but the peak remains in the same position if pesticides have no adverse impact on wild bees.

We first consider a case when pollination is provided by wild bees only. If pesticides are not used, or if they are used but do not impair pollination ability of wild bees (so that $wI=0$), the profits and the population of wild bees are stable over time (assuming that $N_{[0]} = K$). Profits peak when on-farm habitat proportion is between 10% and 20% (fig. 1a) as they depend on revenues made from the crop area balanced against the loss through providing habitat rather than growing crops on the remaining area. At low levels of on-farm habitat provision, yield is limited by pollination, fig. 1b, as

$$a + b B_{[t]} < g \supset Y_{[t]} = Y_{\max} - \left(a + b w F (1 - wI) w N v \right) \quad (12)$$

(where we used the fact that $B_{[t]} = \frac{w F (1 - wI) N_{[t]}}{A} = w F (1 - wI) w N v$ with $N_{[t]} = K = w N v A$;

see fig. 1c). Combining equations (6), (8) and (9) we see that for low values of v the leading term in the profit function is of the form $v(1-v)$, see the left hand side of fig. 1a. When v reaches the critical level

$$v = \frac{g - a}{b_w F (1 - wI) wN} \quad (13)$$

(i.e. when $a + bB_{[t]} = g$) then yield becomes independent on the wild bee population, but total production and therefore profit decreases as the area under cropping decreases with increasing v , as in figures 1a and 1b.

Profits can be negative when there is no area of the farm used for wild bee habitat and yields are low due to pesticides not being used, fig. 1a. When pesticides are used (still under assumption of no adverse effect on wild bees), the profit function is shifted upwards (thick line in fig. 1a), but this does not change the dynamics of wild bee population over time (fig. 1c) or the optimal allocation of on-farm habitat. We note that if the initial density of the wild bumblebee nests, $N_{[0]}$ is lower than K , the time projection of $N_{[t]}$ will increase towards K . Profits in this case will also increase but in the long term the behaviour is the same as that discussed above.

RESULT 2: When no commercial bees are used and wild bees are impacted by pesticides ($wI > 0$), profits are lower and peak profits occur at higher level of on-farm bee habitat.

If the pesticide-induced impairment in pollination by wild bees is relatively small (eg. $wI=0.3$ or 0.6), the wild bee population stays constant over time (assuming $N_{[0]} = K$, or increases until $N_{[t]} \simeq K$ if $N_{[0]} < K$), fig. 2a. As a result, the yield is also constant, as in figure 2c. The corresponding profits are lower and require a higher proportion of on-farm habitat to peak, see equation (13) and fig. 3a, as more nests (and therefore more habitat) are required to make up for the impairment of foragers. These results are summarised in fig. 4. Thus, with an increasing impact of pesticides on wild bees, there is a gradual increase in the optimal value of v , as shown in figure 4a (compared to figure 3a). This is associated with the gradual decrease in the maximum achievable profit, as shown in figures. 3a and 4b.

Wild bee numbers respond gradually to changes in the impairment as long as:

$$wI \leq 1 - \frac{1}{wF} \left[\frac{mR}{R-1} - j \right]; \quad (14)$$

When (14) is not satisfied, the behaviour of the population of wild bees switches from sustainability over long periods of time, $N_{[t]} = K$, to decline over time, $N_{[t]} \rightarrow 0$ with $t \rightarrow \infty$, fig. 2b. As a result, there is not enough pollination potential and production declines; in our parameterisation this occurs for $wI > 2/3 = 0.666\dots$, see figure 4. We choose $wI=0.67$ to illustrate this behaviour in fig. 2b and d. The resulting profits are significantly lower than for $wI < 2/3$ (figs. 2d and 4b). The optimal percentage of on-farm habitat changes in time and is initially ca. 50%, higher than when there is no impact of pesticides on wild bees.

The qualitative change in the long-term dynamics of wild pollinators results in a threshold-like behaviour for optimal allocation of on-farm habitat, v , fig. 4a, and the associated maximum profit, fig. 4b, both of which drop rapidly at the transition point, cf. equation (14). This points to very high sensitivity of the results to the effects of pesticides on wild bee population.

RESULT 3: The speed at which wild bumblebees decline depends on the balance of nest death relative to nest reproduction.

When wild bees are used as the sole pollination input, the likelihood of wild bee decline depends on the relationship between the impairment of foragers (and hence nest survival) and the reproductive capacity of the surviving nests each year (fig. 2b). If the impairment is high enough, the density of nests declines exponentially in time as

$$N_{[t]} = N_{[0]} \times \exp(-rt) \quad \text{with} \quad r = -\ln \left[R \times \left(1 - \frac{m}{j + wF \times (1 - wI)} \right) \right]. \quad (15)$$

Thus, the characteristic time for the decline is given by r^{-1} and sharply decreases when wI increases, fig. 5.

However, the resulting decline in the profit can initially be slow (see an example in fig. 6), effectively masking the decline in nest density (to illustrate this effect better, wN is increased by a factor of 5 so that the resulting K is higher in fig. 6 than in other figures). With higher levels of on-farm habitat, there are more wild bees per area of crop, and so there is a period where farms are over supplied with pollinators. This continues until the wild bee population drops to a level at which pollination services become limited, at which point profits begin to drop (fig. 6). Thus, the farmer might not have an incentive to change the pesticide use until it is too late.

RESULT 4: When commercial bees are used (and unaffected by pesticides), profits remain stable despite declines in wild bees, and are highest when on-farm habitat is low

When commercial bees are used at the same time as wild bees, fig. 3b and 4b, the highest profit corresponds to no on-farm habitat, i.e. $v=0$. The resulting optimal profit is higher than when pollination relies on wild bees only. The slight drop in the profit at higher values of v in fig. 3b is due to the cost of buying in commercial bees.

Profits remain stable throughout the projection period regardless of whether wild bee nests decline or not, figs. 3b, 4b and 7a, with highest yields when no farm area is set aside for habitat. Thus, when farmers can buy-in pollinators which are unaffected by pesticides, and where such commercial bees can provide a perfect substitute for wild bees in terms of their pollination delivery, this acts as a severe disincentive to conserving wild bees or to reduce pesticide use.

RESULT 5: When commercial bees are used and both these and wild bees are affected by pesticides, the optimal strategy is either to rely completely on commercial bees, or to provide a mixture of commercial bees and on-farm habitat for wild bees, depending on the level of impairment.

When both commercial and wild bees are impaired by pesticides, profits generally change little if the impairment is low and equation (14) is satisfied, as shown in figure 4. The optimal area of on-farm habitat is zero, so all pollination is provided by commercial bees. If the impairment is increased (but (14) is still satisfied) it becomes profitable to invest in a mixture of wild and commercial bees, as shown by the dash-dot line in fig. 3b and the intermediate range of wI and cI in fig. 4a (here we assume $wI=cI$). This is also associated with a drop in optimal profit as compared to the case when commercial bees are unaffected by pesticides, fig. 4b. The wild bee population remains steady for low impairment levels (if (14) is satisfied) and starts to decline when impairment becomes too high, resulting in the return to pollination based on commercial bees only, see the drop in fig. 4a. Profits continue to decline with increasing impairment, as the reduced number of commercial bee foragers cannot provide the entire pollination service, leaving crops vulnerable to pollinator decline (we assume that farmer does not change the provision of commercial bees over time: clearly, this assumption can be relaxed). However, the decline in profits at this point is smaller than if the commercial bees are not used, fig. 4b, as the commercial bees still manage to moderate the adverse impacts of pesticides.

When the impairment is high and both commercial and wild bees are affected, profit changes over time unless $v=0$, fig. 7b. Initially, when there is still sufficient number of wild bee nests, the optimal strategy is to invest in a mixture of wild and commercial bees, fig. 7b. As wild bee nests die due to pesticide impairment, the farmer starts to rely on commercial bees only, even though they are affected by pesticides.

Discussion and Conclusions.

Pollination inputs are valued by farmers as they increase the quality and quantity of a range of important crops (Klatt et al, 2014). However, commercial bee use can effectively mask declines in wild bees, reducing the private value of wild bee conservation on farms. Moreover, there may be lags

in the response of insect pollinators to pesticide use meaning that the market signal to farmers to change their management practice arrives “too late” to stop a rapid decline in pollinators. Since wild pollinators also generate ecosystem benefits for a wide range of wild plants beyond the farm from which society derives value (Hanley et al, 2015), these three factors can all drive the supply of wild bees below the social optimum.

In the modelling presented above, we consider the pollination services provided by a mix of wild and commercial bees which are inputs to a commercial crop. Farmers can “produce” more wild bees by allocating land to bee habitat, but this comes at an opportunity cost in terms of foregone profits from land allocated to cropping. Use of a third input, pesticides, contributes positively to profits through its effect on output, but negatively through any effects on bees. Farmers thus face a trade-off in the costs and benefits of pesticide use, where these costs go beyond the price paid for pesticides.

If commercial bees are unaffected by pesticides, their small cost relative to other inputs means that profits are highest when commercial bees are used and little farm area is converted to on-farm habitat for wild bees. If wild bee numbers decline under pesticide pressure, profits can remain positive, as commercial bee numbers can deliver the required pollination level for maximum yields. This is in contrast to the situation when wild bees alone are used for pollination and there is no option to use commercial bees (this is equivalent to the situation where commercial bees can substitute for wild bees). In this case there is an optimal percentage of land converted to wild bee habitat, a results which is in accordance with other studies (Brosi et al., 2008; Keitt, 2009). How big this area of land allocated to bee habitat is will depend on crop prices and the productivity of land, both for wild bees and for crops.

The outcome changes when commercial bees are impaired by pesticides along with wild bees. In this case, agricultural yields can be stable and high for a number of years and then fall suddenly, as wild pollinators decline past a particular point. High yields are maintained when there is an “over-supply”

of pollinators, but fall after wild pollinators numbers decline to a level where overall pollinator numbers limit yields.

In practice, the relative impact of pesticides on commercial and wild bees will depend on farm practices used. Farmers can reduce the impact on commercial bees by shutting the hives or boxes when spraying takes place, though systemic pesticides, by design, are likely to persist within the plant for weeks after application so bees will still be exposed through the ingestion and transport of contaminated nectar and pollen (Bonmatin et al. 2015). Wild pollinators cannot be shut inside nests while spraying takes place and so are potentially left more vulnerable, though some action can still be taken to avoid direct impact on wild pollinators such as spraying when wild bees are not active.

If declines in wild pollinators are irreversible (eg as species become extinct), and if there is uncertainty over whether wild pollinators will be more beneficial in the future, then there is an option value to maintaining this natural capital for future use (Arrow & Fisher, 1974; Kassari & Lasserre, 2004). This option value is an additional economic rationale for conserving wild pollinators, even when there are commercial pollinators present. This value, however, will depend on the time-horizon and risk-aversion of the farmer, as farm profits may be stable for years before declines are evident. If farmers are present-bias, then there may be little private benefit to conserving wild pollinators for crop production, implying that government interventions may be required given the wide range of economic and ecological benefits which wild pollinators deliver (Hanley et al, 2015; Hoehn et al, 2008).

The wild bee population modelled here will often in practice be made up of multiple populations of bee and non-bee pollinators such as hover-flies, wasps and beetles (Brittain et al, 2013b). The presence of multiple pollinator groups can buffer the system from extinction (Mommott et al., 2004; Kaiser-Bunbury et al., 2010), and we have not modelled this buffering capacity here. While different pollinators groups may respond in different ways to external pressure such as pesticide use,

the effects are likely to be negative on all groups, and may be stronger on solitary bees and non-bee pollinators as these are often smaller in size and they are not buffered by living in a social colony with numerous expendable workers (Goulson, 2013; Henry et al. 2015). There is a benefit from maintaining multiple groups of ecosystem service providers as insurance against a fluctuating environmental conditions Baumgaertner (2007), implying a role for commercial bees in providing “financial insurance” against wild bee declines.

Several simplifications made in the modelling procedure should be noted. We have assumed that all factors other than nest death (or reproductive failure) are deterministic. In reality the pollination process itself will be stochastic, and there will be an overall higher likelihood of successful pollination if more pollinators are present. We assumed that all nests which reproduce produce a set number of queens which survive until the next year, since this simplifies the actual process which will rely on perhaps a larger number of queens being produced by successful colonies, who then may or may not mate, survive until the next year and establish a nest themselves. Overall success is likely to depend on other factors such as weather conditions and the level of disturbance, so the failure rate will vary substantially between years (Bryden et al., 2013). There is evidence that pesticides can interact synergistically with diseases, poor nutrition and other chemicals, but this is not modelled either (Iwasa et al, 2004; Alaux et al, 2010; Laurino et al, 2011; reviewed in Goulson et al. 2015). Moreover, if commercial bee keepers find that their bees are being adversely affected by pesticides, then supply may decline, leading to a future rise in the prices charged for commercial pollinator services.

Finally, note that we have based model parameters on a specific crop, strawberries. As Keitt (2009) concluded, the actual form of the production relationship between pollinators and profits is likely to vary across and within crops, depending on the yield response to both pesticides and bees, and the landscape in which the farmers are working. However our model is applicable for a range of crops with similar or higher dependency on bees which also benefit from applications of pesticides, and which are grown within intensive agricultural environments, including other soft-fruits and almonds.

We show that pesticide use is not only an externality, affecting wild bees in the vicinity of the farm, but part of an internal trade-off decision for farmers of insect pollination-dependent crops. In the presence of commercial bees, farmers have little incentive to support wild bees around their farms; while bees might be important to crop yields, the availability of cheap substitutes means that high profits can be maintained in the short-term. This is despite a longer term risk of declining profits which can threaten the ability of farmers to maintain production. Safeguarding farmland pollinators may therefore require monetary incentives to encourage the creation of on-farm habitat so that future pollination options are not reduced.

References

- AIZEN, M.A. and HARDER, L.D., 2009. The global stock of domesticated honey bees is growing slower than agricultural demand for pollination. *Current Biology*, **19**(11), pp. 915-918.
- ALAUX, C., BRUNET, J., DUSSAUBAT, C., MONDET, F., TCHAMITCHAN, S., COUSIN, M., BRILLARD, J., BALDY, A., BELZUNCES, L.P. and LE CONTE, Y., 2010. Interactions between *Nosema* microspores and a neonicotinoid weaken honeybees (*Apis mellifera*). *Environmental microbiology*, **12**(3), pp. 774-782.
- ARROW, K.J. and FISHER, A.C., 1974. Environmental preservation, uncertainty, and irreversibility. *Quarterly Journal of Economics*, **88**(2), pp. 312-319.
- BAUMGAERTNER, S., 2007. The insurance value of biodiversity in the provision of ecosystem services. *Natural Resource Modeling*, **20**(1), pp. 87-127.
- BIESMEIJER, J.C., ROBERTS, S.P.M., REEMER, M., OHLEMUELLER, R., EDWARDS, M., PEETERS, T., SCHAFFERS, A.P., POTTS, S.G., KLEUKERS, R., THOMAS, C.D., SETTELE, J. and KUNIN, W.E., 2006. Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science*, **313**(5785), pp. 351-354.
- BONMATIN J-M., GIORIO C., GIROLAMI V., GOULSON D., KREUTZWEISER D., KRUPKE C., LIESS M., LONG E., MARZARO M., MITCHELL E., NOOME D., SIMON-DELSON N., TAPPARO A. 2015. Environmental fate and exposure; neonicotinoids and fipronil. *Environmental Science & Pollution Research* **22**, 35-67.
- BREEZE, T.D., BAILEY, A.P., BALCOMBE, K.G. and POTTS, S.G., 2011. Pollination services in the UK: How important are honeybees? *Agriculture Ecosystems & Environment*, **142**(3-4), 137-143.
- BRITAIN, C., KREMEN, C. and KLEIN, A., 2013a. Biodiversity buffers pollination from changes in environmental conditions. *Global Change Biology*, **19**(2), pp. 540-547.

BRITTAİN, C., WILLIAMS, N., KREMEN, C. and KLEIN, A., 2013b. Synergistic effects of non-*Apis* bees and honey bees for pollination services. *Proceedings of the Royal Society B-Biological Sciences*, **280**(1754), pp. 20122767.

BROSI, B.J., DAILY, G.C., SHIH, T.M., OVIEDO, F. and DURAN, G., 2008. The effects of forest fragmentation on bee communities in tropical countryside. *Journal of Applied Ecology*, **45**(3), 773-783.

BRYDEN, J., GILL, R.J., MITTON, R.A.A., RAINE, N.E. and JANSEN, V.A.A., 2013. Chronic sublethal stress causes bee colony failure. *Ecology Letters*, **16**(12), pp. 1463-1469.

CAMERON, S.A., LOZIER, J.D., STRANGE, J.P., KOCH, J.B., CORDES, N., SOLTER, L.F. and GRISWOLD, T.L., 2011. Patterns of widespread decline in North American bumble bees. *Proceedings of the National Academy of Sciences of the United States of America*, **108**(2), pp. 662-667.

CARVALHEIRO, L.G., KUNIN, W.E., KEIL, P., AGUIRRE-GUTIERREZ, J., ELLIS, W.N., FOX, R., GROOM, Q., HENNEKENS, S., VAN LANDUYT, W., MAES, D., VAN DE MEUTTER, F., MICHEZ, D., RASMONT, P., ODE, B., POTTS, S.G., REEMER, M.,

CARVELL, C., MEEK, W.R., PYWELL, R.F., GOULSON, D. and NOWAKOWSKI, M., 2007. Comparing the efficacy of agri-environment schemes to enhance bumble bee abundance and diversity on arable field margins. *Journal of Applied Ecology*, **44**(1), pp. 29-40.

CARVELL, C., OSBORNE, J.L., BOURKE, A.F.G., FREEMAN, S.N., PYWELL, R.F. and HEARD, M.S., 2011. Bumble bee species' responses to a targeted conservation measure depend on landscape context and habitat quality. *Ecological Applications*, **21**(5), pp. 1760-1771.

CARVELL, C., ROY, D.B., SMART, S.M., PYWELL, R.F., PRESTON, C.D. and GOULSON, D., 2006. Declines in forage availability for bumblebees at a national scale. *Biological Conservation*, **132**(4), pp. 481-489.

COX-FOSTER, D.L., CONLAN, S., HOLMES, E.C., PALACIOS, G., EVANS, J.D., MORAN, N.A., QUAN, P., BRIESE, T., HORNIG, M., GEISER, D.M., MARTINSON, V., VANENGELSDORP, D., KALKSTEIN, A.L., DRYSDALE, A., HUI, J., ZHAI, J., CUI, L., HUTCHISON, S.K., SIMONS, J.F., EGHOLM, M., PETTIS, J.S. and LIPKIN, W.I., 2007. A metagenomic survey of microbes in honey bee colony collapse disorder. *Science*, **318**(5848), pp. 283-287.

ELLIS, C., 2014. Valuing wild pollinators for sustainable crop production, PhD thesis, University of Stirling.

FAOSTAT, 2013. *Honey producer prices 2000 to 2010, United Kingdom*. [Online]. Available from: <http://faostat3.fao.org/home/E>

FELTHAM, H., PARK, K. and GOULSON, D., 2014. Field realistic doses of pesticide imidacloprid reduce bumblebee pollen foraging efficiency. *Ecotoxicology*, **23**(3), pp. 317-323.

GARIBALDI, L.A., STEFFAN-DEWENTER, I., WINFREE, R., AIZEN, M.A., BOMMARCO, R., CUNNINGHAM, S.A., KREMEN, C., CARVALHEIRO, L.G., HARDER, L.D., AFIK, O., BARTOMEUS, I., BENJAMIN, F., BOREUX, V., CARIVEAU, D., CHACOFF, N.P., DUDENHOFFER, J.H., FREITAS, B.M., GHAZOUL, J., GREENLEAF, S., HIPOLITO, J., HOLZSCHUH, A., HOWLETT, B., ISAACS, R., JAVOREK, S.K., KENNEDY, C.M., KREWENKA, K.M., KRISHNAN, S., MANDELIK, Y., MAYFIELD, M.M., MOTZKE, I., MUNYULI, T., NAULT, B.A., OTIENO, M., PETERSEN, J., PISANTY, G., POTTS, S.G., RADER, R., RICKETTS, T.H., RUNDLOF, M., SEYMOUR, C.L., SCHUEPP, C., SZENTGYORGYI, H., TAKI, H., TSCHARNTKE, T., VERGARA, C.H., VIANA, B.F., WANGER, T.C., WESTPHAL, C.,

WILLIAMS, N. and KLEIN, A.M., 2013. Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science (New York, N.Y.)*, **339**(6127), pp. 1608-11.

GARRATT, M.P.D., BREEZE, T.D., JENNER, N., POLCE, C., BIESMEIJER, J.C. and POTTS, S.G., 2014. Avoiding a bad apple: Insect pollination enhances fruit quality and economic value. *Agriculture Ecosystems & Environment*, **184**, pp. 34-40.

GILL, R.J., RAMOS-RODRIGUEZ, O. and RAINE, N.E., 2012. Combined pesticide exposure severely affects individual- and colony-level traits in bees. *Nature*, **491**(7422), pp. 105-119.

GOULSON, D., 2013. REVIEW: An overview of the environmental risks posed by neonicotinoid insecticides. *Journal of Applied Ecology*, **50**(4), pp. 977-987.

GOULSON D. 2015. Neonicotinoids impact bumblebee colony fitness in the field; a reanalysis of the UK's Food & Environment Research Agency 2012 experiment. *PeerJ* **3**:e854.

GOULSON, D., NICHOLLS E., BOTÍAS C., & ROTHERAY, E.L. 2015. Combined stress from parasites, pesticides and lack of flowers drives bee declines. *Science*, **347**, 1435-+.

HANLEY N., BREEZE T., ELLIS C. and GOULSON D. (2015) Measuring the economic value of pollination services: principles, evidence and knowledge gaps. *Ecosystem Services. Volume 14*, Pages 124-132.

HENRY, M., BEGUIN, M., REQUIER, F., ROLLIN, O., ODOUX, J., AUPINEL, P., APTEL, J., TCHAMITCHIAN, S. and DECOURTYE, A., 2012. A common pesticide decreases foraging success and survival in honey bees. *Science*, **336**(6079), pp. 348-350.

HENRY, M., CERRUTTI, N., AUPINEL, P., DECOURTYE, A., GAYRARD, M., ODOUX, J., PISSARD, A., RUGER, C., BRETAGNOLLE, V. 2015. Reconciling laboratory and field assessments of neonicotinoid toxicity to honeybees. *Proceedings of the Royal Society B*, 282: 20152110.

HOEHN, P., TSCHARNTKE, T., TYLIANAKIS, J.M. and STEFFAN-DEWENTER, I., 2008.

Functional group diversity of bee pollinators increases crop yield. *Proceedings of the Royal Society B-Biological Sciences*, **275**(1648), pp. 2283-2291.

KAISER-BUNBURY, C.N., MUFF, S., MEMMOTT, J., MUELLER, C.B. and CAFLISCH, A., 2010. The robustness of pollination networks to the loss of species and interactions: a quantitative approach incorporating pollinator behaviour. *Ecology Letters*, **13**(4), pp. 442-452.

KASSAR, I. and LASSERRE, P., 2004. Species preservation and biodiversity value: a real options approach. *Journal of Environmental Economics and Management*, **48**(2), pp. 857-879.

KEITT, T.H., 2009. Habitat conversion, extinction thresholds, and pollination services in agroecosystems. *Ecological Applications*, **19**(6), pp. 1561-1573.

KENNEDY, C.M., LONSDORF, E., NEEL, M.C., WILLIAMS, N.M., RICKETTS, T.H., WINFREE, R., BOMMARCO, R., BRITTAIN, C., BURLEY, A.L., CARIVEAU, D., CARVALHEIRO, L.G., CHACOFF, N.P., CUNNINGHAM, S.A., DANFORTH, B.N., DUDENHOEFFER, J., ELLE, E., GAINES, H.R., GARIBALDI, L.A., GRATTON, C., HOLZSCHUH, A., ISAACS, R., JAVOREK, S.K., JHA, S., KLEIN, A.M., KREWENKA, K., MANDELIK, Y., MAYFIELD, M.M., MORANDIN, L., NEAME, L.A., OTIENO, M., PARK, M., POTTS, S.G., RUNDLOF, M., SAEZ, A., STEFFAN-DEWENTER, I., TAKI, H., VIANA, B.F., WESTPHAL, C., WILSON, J.K., GREENLEAF, S.S. and KREMEN, C., 2013. A global quantitative synthesis of local and landscape effects on wild bee pollinators in agroecosystems. *Ecology Letters*, **16**(5), pp. 584-599.

KLATT, B.K., HOLZSCHUH, A., WESTPHAL, C., CLOUGH, Y., SMIT, I., PAWELZIK, E. and TSCHARNTKE, T., 2014. Bee pollination improves crop quality, shelf life and commercial value. *Proceedings of the Royal Society B-Biological Sciences*, **281**(1775), pp. 20132440.

- KLEIN, A., BRITTAIN, C., HENDRIX, S.D., THORP, R., WILLIAMS, N. and KREMEN, C., 2012. Wild pollination services to California almond rely on semi-natural habitat. *Journal of Applied Ecology*, **49**(3), pp. 723-732.
- LAURINO, D., PORPORATO, M., PATETTA, A. and MANINO, A., 2011. Toxicity of neonicotinoid insecticides to honey bees: laboratory tests. *Bulletin of Insectology*, **64**(1), pp. 107-113.
- MEMMOTT, J., WASER, N.M. and PRICE, M.V., 2004. Tolerance of pollination networks to species extinctions. *Proceedings of the Royal Society B-Biological Sciences*, **271**(1557), pp. 2605-2611.
- MOMMAERTS, V., REYNDERS, S., BOULET, J., BESARD, L., STERK, G. and SMAGGHE, G., 2010. Risk assessment for side-effects of neonicotinoids against bumblebees with and without impairing foraging behavior. *Ecotoxicology*, **19**(1), pp. 207-215.
- NIX, J., ed., 2015. John Nix Farm Management Pocketbook. 46th ed. Melton Mowbray: Agro Business Consultants Ltd.
- OSBORNE, J.L., MARTIN, A.P., CARRECK, N.L., SWAIN, J.L., KNIGHT, M.E.; GOULSON, D., HALE, R.J. & SANDERSON, R.A. 2008a. Bumblebee flight distances in relation to the forage landscape. *Journal of Animal Ecology* **77**, 401-415.
- OSBORNE, J.L., MARTIN, A.P., SHORTALL, C.R., TODD, A.D., GOULSON, D., KNIGHT, M.E., HALE, R.J. and SANDERSON, R.A., 2008b. Quantifying and comparing bumblebee nest densities in gardens and countryside habitats. *Journal of Applied Ecology*, **45**(3), pp. 784-792.
- PETTIS, J.S. and DELAPLANE, K.S., 2010. Coordinated responses to honey bee decline in the USA. *Apidologie*, **41**(3), pp. 256-263.

- POTTS, S.G., ROBERTS, S.P.M., DEAN, R., MARRIS, G., BROWN, M.A., JONES, R., NEUMANN, P. and SETTELE, J., 2010. Declines of managed honey bees and beekeepers in Europe. *Journal of Apicultural Research*, **49**(1), 15-22.
- RICKETTS, T.H., REGETZ, J., STEFFAN-DEWENTER, I., CUNNINGHAM, S.A., KREMEN, C., BOGDANSKI, A., GEMMILL-HERREN, B., GREENLEAF, S.S., KLEIN, A.M., MAYFIELD, M.M., MORANDIN, L.A., OCHIENG, A. and VIANA, B.F., 2008. Landscape effects on crop pollination services: are there general patterns? *Ecology Letters*, **11**(5), 499-515.
- ROBERTS, S.P.M., SCHAMINEE, J., WALLISDEVRIES, M.F. and BIESMEIJER, J.C., 2013. Species richness declines and biotic homogenisation have slowed down for NW-European pollinators and plants. *Ecology Letters*, **16**(7), pp. 870-878.
- RUNDLÖF, M., ANDERSSON, G.K.S., BOMMARCO, R., et al (2015) Seed coating with a neonicotinoid insecticide negatively affects wild bees. *Nature* 521, 77–80.
- STANLEY, d.a., Garratt, M.P.D., Wickens, J.B., Wickens, V.J., Potts, S.G. & Raine, N.E. 2015. Neonicotinoid pesticide exposure impairs crop pollination services provided by bumblebees. *Nature* doi:10.1038/nature16167
- VELTHUIS, H.H.W. and VAN DOORN, A., 2006. A century of advances in bumblebee domestication and the economic and environmental aspects of its commercialization for pollination. *Apidologie*, **37**(4), pp. 421-451.
- WHITEHORN, P.R., O'CONNOR, S., WACKERS, F.L. and GOULSON, D., 2012. Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science*, **336**(6079), pp. 351-352.
- WINFREE, R., AGUILAR, R., VAZQUEZ, D.P., LEBUHN, G. and AIZEN, M.A., 2009. A meta-analysis of bees' responses to anthropogenic disturbance. *Ecology*, **90**(8), pp. 2068-2076.

WINFREE, R. and KREMEN, C., 2009. Are ecosystem services stabilized by differences among species? A test using crop pollination. *Proceedings of the Royal Society B-Biological Sciences*, **276**(1655), pp. 229-237.

WINFREE, R., WILLIAMS, N.M., DUSHOFF, J. and KREMEN, C., 2007. Native bees provide insurance against ongoing honey bee losses. *Ecology Letters*, **10**(11), pp. 1105-1113.

Table 1: Key parameters in the model (modelled after soft fruit production). Source is (Ellis, 2014) unless otherwise specified.

Parameter	Interpretation	Value	Source/comments
v	Proportion in conservation area	0-0.7	Key variable
A	Farm area	100ha	
R	Nest reproduction ratio	4	Incorporates the relatively small chance of queens mating and overwintering
wN	Wild bees nesting density	15	Osborne et al. (2008b)
cN	Commercial bees nesting density	4	
μ	Nest death parameter	55	Bryden et al., (2013)
ϕ	Nest death parameter	40	Bryden et al., (2013)
wF	Avg. number of wild foragers per nest	100	
cF	Avg. number of commercial foragers per nest	100	Same as wF
wI	Impairment due to pesticides, wild bees	0 if no impairment; variable	Key variable
cI	Impairment due to pesticides, commercial bees	0 if no impairment; variable	Key variable
$Y_{\max, \text{nop}}$	Maximum attainable yield when pesticides are not used	11.5 tonne per ha	Estimated from Nix (2015) as 50% of max yield
$Y_{\max, \text{p}}$	Maximum attainable yield when pesticides are used	23 tonne per ha	Max yield in Nix (2015)

γ	maximum proportion of good quality fruits	0.9	
α	proportion of good quality fruits without bees	0.35	
β	incremental effect of bee visitation	0.0024	Combined visitation and efficiency in (Ellis, 2014)
p	Price per tonne	3445	Nix (2015)
cm	Commission	0.09	Nix (2015)
Cpt	Cost per tonne (harvesting and packaging)	£1650 per tonne	Nix (2015)
Cpa	Cost per crop area (planting structures, fieldwork)	£18700 per ha	Nix (2015)
Capa	Total cost per area (land lease)	£150 per ha	
Cseed	Cost of maintaining the conservation area (mainly seed)	£100 per ha	
bC	Cost of commercial nests, per nest	£60 per nest	
pC	Cost of pesticide use, per ha of crop area	£10 per ha

Figure 1: Total profit (a), yield (b), and the number of wild bee nests, $N_{[t]}$ as functions of the proportion of on-farm habitat proportion, v . Thin line: no pesticides; thick line: with pesticides. No commercial bees are used and when pesticides are used, they do not affect wild bees. Parameters as in Table 1.

Fig. 2: Total profit as a function of the on-farm habitat proportion, v , for (a) no commercial bees, (b) with commercial bees but with small impact of pesticides, and (c) with commercial bees but with large impact of pesticides. Horizontal line represents zero profit. In (a), solid line corresponds to $wI=1$, dashed line to $wI=0.3$ and dotted line to $wI=0.6$. In (b) dotted line corresponds to no impact of pesticides on wild or commercial bees ($wI=cI=0$), and dash-dot line corresponds to $wI=cI=0.6$ (solid line from (a) is redrawn for comparison). All other parameters as in Table 1.

Fig. 3: Dependence of (a) and (b): the number of wild bee nests $N_{[t]}$, and (c) and (d): total profit, on the on-farm habitat proportion, v and time (between 0 and 200 years), when pesticides are used but commercial bees are not. In (a) and (c), there is no effect of pesticides on wild bees, $wI=0$, and in (b) and (d), $wI=0.67$. Other parameters as in Table 1.

Fig. 4: Dependence of the optimal on-farm habitat proportion (a) and the corresponding total profit (b) on the wild and commercial bee impairment due to pesticides. Thin solid line corresponds to the case without commercial bees; dashed line corresponds to the case with commercial bees, but with no impairment of their performance, $cI=0$. For the thick solid line, commercial bees are used and affected by pesticides in the same way as wild bees, $cI=wI$. Other parameters as in Table 1.

Fig. 5: Dependence of the characteristic time of decay for the wild bee nests, τ^{-1} , in response to the impairment, wI .

Fig. 6: Examples of time projections for profit over 200 years. Pesticides are used, but no commercial bees; high impact of pesticides on wild bees ($wI=0.67$). For illustration, the carrying capacity for wild bees is doubled so that the effect of overpollination is more pronounced. Solid line: $\nu=0.22$ (optimal), thick line: $\nu=0.52$, dashed line: $\nu=0.7$. Other parameters as in Table 1.

Fig. 7: Comparison of dependence of the profit on time and on-farm habitat proportion for the case when pesticides and commercial bees are used and pesticides strongly affect (a) wild bees only ($wI=0.67$, $cI=0$) and (b) both wild and commercial bees ($wI=cI=0.67$). Other parameters as in Table 1.