



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

Papers downloaded from AgEcon Search may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

Evaluation on Potential Ecological Risk of Heavy Metals in Soil from Penglai Fairyland Park

Zhen YANG¹, Di WU¹, Zhun XIANG², Chao KANG², Fang WANG², Zhu HUANG², Lin ZHANG^{2*}

1. Key Laboratory of Mountain Area and Protection of Ecological Environment of Guizhou Province/Guizhou Normal University, Guiyang 550001, China; 2. Guizhou Institute of Biology, Guiyang 550009, China

Abstract This study aims at evaluating the heavy metals pollution in soil collected from Penglai Fairyland Park in Guiyang, and predicting the harm of heavy metals to human health. Based on the background value of soil in Guizhou and *Soil Environment Quality Standard* (GB15618 – 1995), by using the geoaccumulation index method and Hakanson potential ecological risk assessment method, we research pollution characteristics and ecological risk of heavy metals in soil from the study area. The results show that from the index of geoaccumulation, the pollution level of heavy metals in agricultural soil from Penglai Fairyland Park is in the order of Ni > Cu > Hg > Cd > As > Zn > Cr > Pb. From the potential ecological harm index evaluation, soil is at slight ecological risk level in the park and the potential ecological hazard degree is in the order of Hg > Cd > Ni > As > Cu > Cr > Pb > Zn. The soil ecological environment in research area is only mildly damaged. Thus, the soil in research area is at the level of security, and it is suitable for cultivation of crops and less harmful to human health.

Key words Soil, Heavy metals, Ecological damage, Evaluation

1 Introduction

Soil is the means of production for human survival and development, and is also one of the most important natural resources in human society^[1]. Among all the pollutants in the soil, the heavy metal brings the greatest harm to human body, and heavy metal pollution of the soil has captured more and more attention of scholars at home and abroad^[2–6]. Heavy metal is a kind of important pollutant with potential hazards, and it is hidden and irreversible^[7]. It can lead to human cancer and chronic toxicity, mutagenicity and teratogenicity. Heavy metals can enter the food chain^[8–9] through the water-soil-crop ecosystem. The heavy metals in the soil can be absorbed by plants and can not be degraded by biological degradation. When the heavy metal content is too high, it will affect the normal growth, development and reproduction and other physiological activities of the plant^[10–11]. In order to understand the pollution situation of heavy metals in soil, it is necessary to predict the harm of heavy metals to human health and the impact on the ecosystem. This paper takes the soil from Penglai Fairyland Park in Baiyun District of Guiyang City as the research object, and based on the soil background value of Guizhou Province and *Soil Environment Quality Standard* (GB15618 – 1995), this paper analyzes the content of 8 kinds of heavy metals in the soil from Penglai Fairyland Park, and uses the method of

geoaccumulation index and potential ecological risk index to evaluate and analyze. It aims to provide scientific basis for the safe use of soil and the safety of agricultural products in the park.

2 Materials and methods

2.1 Soil sampling Global positioning system (GPS) is used for the positioning of soil sampling points in Penglai Fairyland Park of Baiyun District, and the quincunx sampling method is used for collecting 1 kg of soil samples (0 – 20cm) in sampling bag. After the soil samples are brought back to the lab, soil samples are cooled and put on white paper for indoor natural air drying. Then some stones and residual plant roots are removed to retain about 200 g. The air dried soil sample is ground and sifted with a mortar and 100 nylon mesh, then it is packaged for later use.

2.2 Sample handling and monitoring The total amount of Pb, Cr, Cd, Cu, Zn, Ni elements is digested by $\text{HNO}_3 - \text{HClO}_4 - \text{HF}$ wet method, Hg, As as elements with 1:1 aqua regia digestion. Heavy metal Cd is determined by atomic absorption spectrometry and graphite furnace method. Zn, Cu and Ni are determined by atomic absorption spectrometry. Hg is determined by Atomic Fluorescence Spectrometry, and Pb, Cr and As are determined by inductively coupled plasma atomic emission spectrometry. All reagents are GR, and ultra pure water is used for water analysis in determination process. The samples include blank sample, and the second parallel samples, coupled with national soil standard reference material in accordance with GBW – 070010 and GBW – 07430 quality control. The relative deviation of second parallel kind is less than 5%, the samples with the recovery are in 96.4% – 115.1% and standard sample determination results are within the allowable error range. EXCEL software and SPSS18 software are used for analysis and processing.

Received: March 14, 2016 Accepted: May 25, 2016

Supported by Agricultural Science and Technology Achievement Transformation Fund Project of the Ministry of Science and Technology (2013F-2001007); Project of Edible Fungus Engineering Center of Guizhou Province (G20144002); Guiyang Science and Technology Project (201410014); Project of Baiyun District Science and Technology Bureau (201419).

* Corresponding author. E-mail: 953827773@qq.com

2.3 Evaluation methods

2.3.1 Cumulative pollution index method. Land accumulation index method is an environmental assessment method widely used in the assessment of heavy metal pollution in soil or sediment^[12-15], proposed by scientist Muller from Sediment Research Institute of German University of Heidelberg in 1969. The method not only reflects the characteristics of natural changes in the distribution of heavy metals, but also intuitively reflects the exogenous heavy metals in the sediment concentration. The formula is as follows:

$$I_{geo} = \log_2 \left[\frac{C_j}{1.5 \cdot BE_j} \right]$$

where I_{geo} represents the value of element j in the product; BE_j represents the background concentration of element j ; C_j represents the measured value of element j in the sample; 1.5 is modified exponential, which takes into account element background values change caused by diagenesis and human activities^[16].

Table 1 The evaluation criterion classification of geoaccumulation index

Grade	Geoaccumulation index	Class of pollution
0	$I_{geo} < 0$	Non-pollution
1	$0 \leq I_{geo} < 1$	Slight pollution
2	$1 \leq I_{geo} < 2$	Moderate pollution
3	$2 \leq I_{geo} < 3$	Moderate to heavy pollution
4	$3 \leq I_{geo} < 4$	Heavy pollution
5	$4 \leq I_{geo} < 5$	Heavy to extreme pollution
6	$I_{geo} \geq 5$	Extreme pollution

Geoaccumulation index has the advantages of comprehensive consideration of the sedimentary rock and other natural geological processes and human activities, and geochemical background values will have a greater impact on the geological accumulation index^[17]. Integrated pollution index method is used for embedding I_{geo} . Based on the index of Geoaccumulation advantages, combined with the geological structure features of Penglai Fairyland Park, Nemerow comprehensive pollution index formula and cumulative index are used to replace the single factor index. The embedded formula of the comprehensive pollution index of Geoaccumulation (I_{geo}) is as follows:

Table 3 Classification standard of potential ecological risk of heavy metals in soils

Potential ecological risk	Light	Moderate	A bit strong	Very strong	Extremely strong
Er^i	< 40	40-80	80-160	160-320	> 320
R_j	< 150	150-300	300-600	600-1200	> 1200

Table 4 Coefficient of heavy metal toxicity in soil

	Pb	Cd	Cr	Cu	Zn	Hg	As	Ni
Toxicity Coefficient Tr^i ^[18]	5	30	2	5	1	40	10	5

3 Results and discussions

3.1 Soil heavy metal content

The content of heavy metals in

$$P_j = \sqrt{\frac{(I_{max})^2 + (I_{ave})^2}{2}}$$

where P_j represents the comprehensive pollution index of element j ; I_{max} indicates the maximum value of the ground accumulation index; I_{ave} indicates the average value of the cumulative index. The integrated pollution index of the embedded I_{geo} is shown in Table 2.

Table 2 The evaluation criterion classification of embedded I_{geo} composite index

Grade	Embedded I_{geo} comprehensive pollution index	Class of pollution
0	$P_j < 0$	Non-pollution
1	$0 \leq P_j < 0.5$	Slight pollution
2	$0.5 \leq P_j < 1$	Mild to moderate pollution
3	$1 \leq P_j < 2$	Moderate pollution
4	$2 \leq P_j < 3$	Moderate to heavy pollution
5	$3 \leq P_j < 4$	Heavy pollution
6	$4 \leq P_j < 5$	Heavy to extreme pollution
7	$P_j \geq 5$	Extreme pollution

2.3.2 Hakanson potential ecological risk assessment method. Swedish scholar Lars Hakanson's potential ecological risk index is used to evaluate heavy metal potential ecological hazard in soil from Penglai Fairyland Park. The potential ecological risk index reflects the heavy metal content, species, toxic level and the sensitivity to heavy metal pollution in the soil of the cultivated layer.

$$Er^i = T_f^i \times C_f^i \quad (1)$$

where Er^i is the potential ecological risk coefficient in the soil of a certain region; T_f^i is the toxic coefficient of heavy metal i , which reflects the level of toxicity and the sensitivity of biological to the pollution.

$$R_j = \sum_{i=1}^n Er^i \quad (2)$$

where R_j is a potential ecological risk factor for a variety of heavy metals in sediment.

The potential ecological hazard classification standards of Er^i and R_j are shown in Table 3, and the toxicity coefficient of heavy metal in soil is shown in Table 4.

the soil from Penglai Fairyland Park is shown in Table 5. Table 5 shows that the average content of Cu, Hg, Ni is slightly higher

than that of Guizhou's soil background value, while the average content of the remaining heavy metals is lower than that of the background value. The average values of Cu, Hg and Ni are 2.82, 3.82 and 2.05, respectively, and the average content is in

the order of Hg > Cd > Cu > Cr > Ni > Zn > Pb > As > Hg > Cd. It shows that the content of Cu, Hg and Ni in the soil of this area has been affected by human activities.

Table 5 The content of heavy metals in soil from Penglai Fairyland Park (mg/kg)

Heavy Metals	Range	Average value	Standard deviation	Coefficient of variation	Soil background values in Guizhou Province ^[19]	Multiple of mean value	Exceeding standard rate // %
Pb	11.05 – 35.09	30.66	6.94	22.64	35.20	0	0
Cd	0.12 – 0.63	0.37	0.17	45.95	0.66	0	0
Cr	48.25 – 96.90	84.42	12.21	14.46	95.90	0	0
Hg	0.26 – 0.49	0.42	0.07	16.67	0.11	3.82	100
As	12.63 – 19.59	16.30	2.20	13.50	20.00	0	0
Cu	75.80 – 99.70	90.08	6.80	7.55	32.00	2.82	100
Zn	34.47 – 98.09	73.78	18.40	24.94	99.50	0	0
Ni	54.48 – 96.96	80.04	12.94	16.17	39.10	2.05	100

3.2 Correlation of heavy metal content in soil From Table 6, we can see that the correlation coefficient of Zn and Cd is 0.433, there is a very significant positive correlation; the correlation coefficient of Zn and Cr, Cu is 0.443, and there is a significant positive correlation at 5% level; the correlation coefficients of

Ni and Cr and Zn are 0.424 and 0.428, respectively, and there is a significant positive correlation at the 5% level. The results show that Zn, Cr, Cd, Cu, Ni in the park soil may have the same source of pollution, and have similar geochemical behavior.

Table 6 The correlation matrix of heavy metal content in soil from Penglai Fairyland Park

	pH	Pb	Cd	Cr	Hg	As	Cu	Zn	Ni	mg/kg
pH	1									
Pb	-0.109	1								
Cd	.620 **	-0.371	1							
Cr	-0.199	-0.231	0.095	1						
Hg	-0.307	-0.157	0.007	-0.182	1					
As	-0.028	0.330	-0.188	-0.395	0.305	1				
Cu	-0.076	-0.090	0.345	0.214	-0.035	-0.248	1			
Zn	0.119	-0.415	0.433 *	0.443 *	0.164	-0.033	0.443 *	1		
Ni	0.311	0.018	-0.031	0.424 *	-0.297	-0.221	0.072	0.428 *	1	

Note: * * p < 0.01, * p < 0.05.

3.3 Principal component analysis of heavy metal elements in soil

Principal component analysis (PCA) can be used to determine the source of trace elements in soil^[20-21]. Principal component analysis is performed on heavy metal element content in soil, and multiple factors can be selected from correlation factors, so as to provide a basis^[22].

Table 7 The decomposition of total variance for heavy metals in soil

Principal component factor	Initial eigenvalue and contribution rate		
	Total characteristic value	Contribution rate // %	Cumulative contribution rate // %
1	2.600	32.501	32.501
2	1.589	19.865	52.366
3	1.113	13.914	66.281
4	0.955	11.939	78.219
5	0.697	8.709	86.928
6	0.479	5.984	92.912
7	0.414	5.178	98.090
8	0.153	1.910	100.000

Table 7 shows that the total variance value of first three principal components accounts for 66.281% of cumulative contribution rate, and the first three principal components have the feature information of 8 kinds of heavy metals contained in the soil. The first principal component includes 32.501% of general information; the second principal component contains 19.865% of general information; the third principal component contains 13.914% of general information. So, we choose the first three factors as the main components, to represent the 8 kinds of heavy metals in the soil characteristic index. Based on the principal component analysis principle, the load value is a reflection of the principal component and variable correlation coefficient of heavy metal elements in the soil. The greater the load value, the stronger the correlation between the variable and the principal component. The element is the main component of the important characteristic index^[23]. The principal component matrix of soil heavy metal content is shown in Table 8.

Table 8 The matrix of principal component loading for heavy metals in soil

Principal component factors	Pb	Cd	Cr	Hg	As	Cu	Zn	Ni
First principal component	-0.555	0.556	0.691	-0.154	-0.553	0.569	0.774	0.503
Second principal component	-0.414	0.456	-0.334	0.747	0.349	0.175	0.299	-0.546
Third principal component	0.294	-0.283	0.166	0.316	0.597	-0.045	0.467	0.494

Table 8 shows Cd, Cr, Cu load, and Zn, Ni value in 8 kinds of heavy metal elements, and this indicates that the sources of 5 kinds of heavy metals are similar. By the variation coefficient of Cd, Zn in Table 5, we can see that the source is not stable, and may be influenced by people. Zn also can be used as identification element of traffic pollution, and the first principal component is mainly human activity and traffic pollution. Cd and Hg load is highest in 8 kinds of heavy metal elements of the second principal components, and it indicates that the sources of 2 kinds of heavy metals are similar, the variation coefficient of Hg is higher, and

Guizhou is a typical karst province, affected by the topography and soil factors^[24]. The background values of heavy metals in soil are generally high. The second principal components may be subject to a combined effect of soil parent material and human activities. As, Zn, Ni load value is largest in 8 principal components of the third heavy metal elements, and it indicates that the sources of 3 kinds of heavy metals are similar, probably from soil parent material, less affected by human activities. It can be seen that the result of principal component analysis and correlation analysis are consistent.

Table 9 The geoaccumulation index value of heavy metals in soil

Detection item order	Evaluation index range	Number of test points	Geo-accumulation index I_{geo}									
			$I_{geo} < 0$		$0 \leq I_{geo} < 1$		$1 \leq I_{geo} < 2$		$2 \leq I_{geo} < 3$	Moderate	$3 \leq I_{geo} < 4$	
			Non-pollution	Number	Slight pollution	Number	Moderate pollution	Number	to heavy pollution	Number	Heavy pollution	
Pb	-5.08 -- 3.42	22	22	100	0	0	0	0	0	0	0	
Cd	-1.91 - 0.49	15	68.18	7	31.82	0	0	0	0	0	0	
Cr	-2.96 - -1.95	22	100	0	0	0	0	0	0	0	0	
Hg	-0.79 - 0.12	12	54.55	10	45.45	0	0	0	0	0	0	
As	-1.83 - -1.20	22	100	0	0	0	0	0	0	0	0	
Cu	0.01 - 0.41	0	0	22	100	0	0	0	0	0	0	
Zn	-3.12 - -1.61	22	100	0	0	0	0	0	0	0	0	
Ni	-0.14 - 0.69	1	4.55	21	95.45	0	0	0	0	0	0	
P_j Integrated pollution index of embedded I_{geo}	Number of test points		$P_j < 0$ (non-pollution)		$0 \leq P_j < 0.5$ (Slight pollution)		$0.5 \leq P_j < 1$ (Mild to moderate pollution)		$1 \leq P_j < 2$ (Moderate pollution)		$2 \leq P_j < 3$ (Moderate to heavy pollution)	
			Number	Percentage %	Number	Percentage %	Number	Percentage %	Number	Percentage %	Number	Percentage %
		22	0	0	0	0	19	86.36	3	13.64	0	0

The pollution status of heavy metals in the park is evaluated by using Geoaccumulation index method. We consider the sedimentary rock natural geological process and human activities, such as the influence of various factors on the soil environment. As an important parameter to measure the degree of the impact of human activities, the cumulative index can comprehensively reflect the natural characteristics of the soil heavy metals and the impact of human activities on the soil environment^[25-26]. The calculation results are shown in Table 9. Geoaccumulation index calculation results show that soil Cu accumulation index is 0.01 - 0.41, which belongs to the light pollution; soil Cd accumulation index is -1.91 - 0.49, which belongs to non-pollution to light pollution; soil Hg accumulation index is -0.79 - 0.12, which belongs to non-pollution to light pollution; soil Ni accumulation index is -0.14 - 0.69, which belongs to non-pollution to the light pollution; soil Pb, Cr, as, Hg, Zn accumulation index is less than 0, belonging to non-pollution. Thus, there is light pollution or no pollu-

tion in the soil of Penglai Fairyland Park, and the pollution index of eight kinds of soil heavy metal is in the order of Ni > Cu > Cd > Hg > As > Zn > Cr > Pb.

3.4 Potential ecological risk assessment of heavy metals in soils

One of the commonly used methods in the study of soil heavy metals is the quantitative classification of the possible ecological damage degree of heavy metal elements^[27]. Using potential ecological risk index method, we use the toxicity analysis of heavy metal elements in soil and sediment migration transformation rule analysis in order to make the evaluation, excluding the effects of regional differences, which can indicate that the heavy metals have potential harm to ecological environment^[28]. It can be seen from Table 10 that the value of Pb, Cd, Cr, Cu, Zn, Hg, As, Ni and Er^i is less than 40, and the ecological risk is slight. From the composite ecological risk hazard index evaluation, the hazard index is less than 150, and the degree of ecological harm is relatively slight. The potential ecological harm degree is in the order of

$Hg > Cd > Ni > As > Cu > Cr > Pb > Zn$. Based on potential ecological harm evaluation results, using cumulative index method, it is found that the heavy metals in agricultural soil from Penglai Fairyland Park are in the state of non-pollution to light pollution, and the accumulation pollution index size is in the order of $Ni > Cu > Hg > Cd > As > Zn > Cr > Pb$. Potential ecological hazard index evaluation shows that the park soil is at the level of slight ecological damage, and the potential ecological harm degree is in the order of $Hg > Cd > Ni > As > Cu > Cr > Pb > Zn$. The cumulative index method and the potential ecological risk index method show

that the soil is mainly affected by Cd , Cu , Ni , Hg , and the factor analysis method is also used in the 22 sampling points. Potential ecological risk index method not only considers the types of heavy metal elements in terms of the synergistic effect, heavy metal toxicity, pollution concentration, but also considers the ecological environment's sensitivity to heavy metals, to make up for the defects of accumulation index method. Therefore, the concentration of heavy metals in the soil should be combined with its potential ecological risk index, in order to better evaluate the extent of heavy metal pollution in the soil^[29].

Table 10 Indexes of single factor potential ecological risk (E^i_r) and compound ecological risk of heavy metals (R_j) in soil

Detection item order	Evaluation index range	Number of test points	Single element potential ecological risk factor E^i_r									
			$E^i_r < 40$ (Slight ecological harm)		$40 \leq E^i_r < 80$ (Moderate ecological risk)		$80 \leq E^i_r < 160$ (Strong ecological hazard)		$160 \leq E^i_r < 320$ (Stronger ecological hazard)		$E^i_r > 320$ (Extremely strong ecological hazard)	
			Number	Percentage %	Number	Percentage %	Number	Percentage %	Number	Percentage %	Number	Percentage %
Pb	0.20–0.60	22	22	100	0	0	0	0	0	0	0	0
Cd	6.00–31.50	22	100	0	0	0	0	0	0	0	0	0
Cr	0.48–0.97	22	100	0	0	0	0	0	0	0	0	0
Hg	20.80–39.20	22	100	0	0	0	0	0	0	0	0	0
As	4.21–6.53	22	100	0	0	0	0	0	0	0	0	0
Cu	3.80–5.00	22	100	0	0	0	0	0	0	0	0	0
Zn	0.14–0.39	22	100	0	0	0	0	0	0	0	0	0
Ni	5.45–9.70	22	100	0	0	0	0	0	0	0	0	0
RI (Potential ecological risk index)		Number of test points	$RI < 150$ (Slight ecological harm)		$150 \leq RI < 300$ (Moderate ecological risk)		$300 \leq RI < 600$ (Strong ecological hazard)		$RI \geq 600$ (Strong ecological hazard)			
			Number	Percentage %	Number	Percentage %	Number	Percentage %	Number	Percentage %	Number	Percentage %
			22	22	100	0	0	0	0	0	0	0

4 Conclusions

(i) The heavy metals, Cu , Hg , Ni in soil from Penglai Fairyland Park have greater value than Guizhou's soil background value in varying degrees, and the pollution level is in the order of $Cu > Cr > Ni > Zn > Pb > As > Hg > Cd$. (ii) Factor analysis results show that Cd , Cu , Ni , Zn , Cr pollution mainly comes from human activities and traffic pollution, Hg pollution comes mainly from soil parent material and human activities, and As , Zn , Ni are from a natural source, less affected by human. (iii) The cumulative pollution index shows that the heavy metals in agricultural soil from Penglai Fairyland Park are in the state of non-pollution to light pollution, and the accumulation pollution index size is in the order of $Ni > Cu > Hg > Cd > As > Zn > Cr > Pb$. (iv) The assessment of potential ecological risk index shows that the soil in the park is at the level of slight ecological harm, and the potential ecological harm degree is in the order of $Hg > Cd > Ni > As > Cu > Cr > Pb > Zn$.

References

- LI GF, WANG JY. Analysis and evaluation of heavy metal pollution in soil around a gold mine drainage ditch [J]. Gold, 2012(8):50–52.
- MACHENDER G, DHAKATE R, PRASANNA L, et al. Assessment of heavy metal contamination in soils around Bala Nagar industrial area, Hyderabad, India [J]. Environment Earth Sciences, 2010(10): 325–334.
- ZHUANG P, ZOU B, LI NY, et al. Heavy metal contamination in soils and food crops around Dabaoshan mine in Guangdong, China: implication for human health [J]. Environmental Geochemistry and Health, 2009 (31): 707–715.
- BHATTACHARYYA P, TRIPATHY S, CHAKRABARTI K, et al. Fractionation and bioavailability of metals and their impacts on microbial properties in sewage irrigated soil [J]. Chemosphere, 2008(72): 543–550.
- SORIANO – DISLA JM, SPEIR TW, GOMEZ I, et al. Evaluation of different extraction methods for the assessment of heavy metal bioavailability in various soils [J]. Water, Air and soil Pollution, 2010, 213: 471–483.
- LI J, YU TM, ZHOU J, et al. Health risk assessment of lead, zinc, cadmium and copper in lead zinc mining area and surrounding soil [J]. Environmental Science, 2008, 29(8):2327–2330.
- LI GF. Contamination analysis of Mn, Cu, Zn and Pb four kinds of heavy metals in the soil around gold mine drainage ditch [J]. Journal of Shaanxi University of Technology (Natural Science Edition), 2012(4): 60–63.
- GAO YX, FENG JG, TANG L, et al. Distribution and risk assessment of heavy metals in the soil of the metal mining area in the upper reaches of Miyun Reservoir [J]. Environmental Science, 2012(5): 1707–1717.
- ZHANG G, WANG N, WANG Y, et al. Environmental mercury pollution in the gold mining area of China [J]. Environmental Science and Management, 2012(11):54–60.
- ZHONG LY, GUO LZ. Soil heavy metal pollution status and dynamic change characteristics of agricultural land in different ways: a case study of Xuwen County, Guangdong Province [J]. Journal of Ecological Environment, 2011, 20(12):1934—1939.
- ZHANG GY, WANG LY, SUN SY, et al. Heavy metal pollution of soil and its prevention and cure [J]. Hebei Agricultural Sciences, 2003, 7 (5):59–62.
- WAN JB, WANG JY, WU D. Le'an river sediment heavy metal pollution evaluation [J]. Environmental Science and Technology, 2008, 31 (11): 130 – 133.

respectively, that is, the water consumption due to soil evaporation and crop transpiration accounted for 46.4% and 24.1% of total irrigation water and precipitation, respectively, and leakage accounted for 30.3% and 60.6%, respectively. The annual irrigation water loss in GT-TR1 and GT-TR2 of Guanting irrigation area in 2013 was also simulated. Results show that the annual actual evapotranspiration in GT-TR1 and GT-TR2 of Guanting irrigation area was 632.6 mm and 646.9 mm, respectively, and leakage was only 14.9 mm and 6.7 mm, respectively, that is, leakage accounted for 2.6% and 1.2% of total irrigation water and precipitation, respectively, indicating that the leakage was very weak. RMSE of the simulation results of the groundwater depth in Daxia irrigation area during the two periods was 92.3 mm and 27.7 mm, respectively. And RMSE of the simulation results of the moisture content of soil profile in the two monitoring sites of Guanting irrigation area was 2.04% and 5.81%, respectively, indicating that the simulation results were reliable.

References

[1] CAO JD. Taking the water as the source, promoting ecological protection and benefiting people[J]. *Qinghai Science and Technology*, 2013 (2): 54–56. (in Chinese).

(From page 79)

[13] ZHANG YJ, SUN B, ZHAO XL. Heavy metal pollution assessment and resource utilization in rural inland river sediments [J]. *Journal of Agro-environment Science*, 2008, 27 (4): 1398–1402.

[14] GU XJ, HOU YQ. Shaanxi Province Jing Hui canal irrigation area soil heavy metals geological accumulation index evaluation [J]. *Earth and Environmental Science Journal*, 2010 (3): 288–291.

[15] LIU Y, YE LL, LI JC. Taiyuan city soil heavy metal pollution and the potential ecological risk assessment [J]. *Journal of Environmental Science*, 2011, 31 (6): 1285–1293.

[16] LI ZK, WANG LD, LI Y, *et al.* Research progress of method for evaluation of heavy metal pollution of soil mineral resources and geology [J]. *Journal of Environmental Science*, 2011, 25 (2): 172–176.

[17] DONG DL, LI HJ, ZHANG J, *et al.* Removal of heavy metals from mine water by cyanobacterial calcification [J]. *Mining Science and Technology*, 2010, 20(4): 566–570.

[18] ZHUANG SY. Heavy metal pollution and potential ecological risk of heavy metals in vegetable soils [J]. *China*, 2008, 24 (7): 284–287.

[19] National Environmental Protection Bureau. Soil element background value of China [S]. Beijing: China Environmental Science Press, 1990.

[20] MARTIN JAR, ARIAS ML, CORBI JMG. Heavy metals contents in agricultural top soils in the Ebro basin(Spain). Application of the multivariate geo-statistical methods to study spatial variations [J]. *Environmental Pollution*, 2006, 144(3) : 1001–1021.

[21] FACCINELLI A, SACCHI E, MALLEN L. Multivariate statistical and GIS-based approach to identify heavy metal sources in soils [J]. *Environmental Pollution*, 2001, 114: 313–324.

[2] BAIER W, ROBERTSON GW. A new versatile soil moisture budget[J]. *Canadian Journal of Plant Science*, 1996, 46: 299–315.

[3] DYER JA, BAIER W. Weather-based estimation of field workdays in fall [J]. *Canadian Agricultural English*, 1979, 21: 119–222.

[4] DYER JA. A new infiltration sub-model for soil moisture budgeting [R]. In: *Proceedings of Manitoba Society of Soil Science 27th Annual Meeting*. University of Manitoba. Winnipeg, Manitoba, Canada. January 1984: 79–93.

[5] DYER JA, WILKIE KI. Estimation of vertical transport of inert water soluble compounds in soil[J]. *Canadian Water Resources Journal*, 1988, 13: 63–73.

[6] BROUGHTON, RS, FOROUD N. A model to predict water table depths for flat lands[J]. *Canadian Agricultural English*, 1978, 20: 81–86.

[7] BELOMANS CJ, FEYEN D. An attempt at experimental validation of macroscales models of soil moisture extraction by roots[J]. *Soil Science*, 1979, 127: 174–186.

[8] ROBERTSON GW. Calculation of soil water in an oil palm plantation in a humid tropical area[Z]. UNDP/FAO Technical Assistance to the Federal Land Development Authority, Tun Razak Agriculture Research Centre, Sungai Tekam, Jerantut, Pahang, Malaysia. Project Field Report No. A-5, 1976: 35.

[9] DYER JA, KELBE BE, DE JAGER JM. Lysimetric calibration of a Canadian soil moisture budget model under bare soil in southern Africa[J]. *Climatological Bulletin*, 1988, 22: 33–47.

[10] BOISVERT JB, DYER JA, LAGACE R. Estimating watertable fluctuations with a daily weather – based budget approach[J]. *Canadian Agricultural English*, 1992, 34: 115–124.

[22] GUO Q. Principal component analysis method of software – land ecosystem health evaluation of Yuci district based on [J]. *Journal of Shandong Agricultural University (Natural Science Edition)*, 2012, 32 (1): 58–62.

[23] YOSHIHOGI M, FANG L, ZHANG J, *et al.* Analysis of operation method in SPSS software and the application in the evaluation of river water quality [J]. *Environmental Research and Monitoring of Principal Components*, 2012, 25 (4): 68–73.

[24] RUAN YL, LI XD, LI TY, *et al.* Karst area of farmland soil heavy metal pollution and human health hazards [J]. *Earth and Environment*, 2015, 43 (1): 92–97.

[25] LI SS. Application of geo – accumulation index method in the assessment of heavy metal and fluorine pollution in soil of Yima mining area [J]. *Henan Science*, 2011, 29 (5): 614–618.

[26] ZHENG GZ. Heavy metal pollution and potential ecological risk assessment of heavy metals in farmland soils in the Loess Hilly and gully region of Northern Shaanxi Province [J]. *Soil Bulletin*, 2013, 49 (6): 1491–1494.

[27] RKLAL R. Changes in physical chemical and Shrestha properties of soil after surface mining and reclamation [J]. *Geoderma*, 2011, 161 (3–4): 168–176.

[28] HE YF, ZHU GW, CHEN YX, *et al.* The Grand Canal (Hangzhou section) the potential ecological risk of heavy metals in sediments [J]. *Journal of Zhejiang University: agricultural and Life Science Edition*, 2002, 28 (6): 669–674.

[29] DOU L, ZHOU YZ, WANG XR, *et al.* According to the mathematical model of fuzzy evaluation of heavy metal pollution in soil improvement and application of [J]. *Journal of Soil Science*, 2007, 38 (1): 101–105.