AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

Economic growth and global particulate pollution concentrations

CCEP Working Paper 1604
Feb 2016

David I. Stern

Crawford School of Public Policy, The Australian National University

Jeremy van Dijk

Australian Bureau of Agricultural and Resource Economics and Sciences, Australia

Abstract

Though the environmental Kuznets curve (EKC) was originally developed to model the ambient concentrations of pollutants, most subsequent applications focused on pollution emissions. Yet, previous research suggests that it is more likely that economic growth could eventually reduce the concentrations of local pollutants than emissions. We examine the role of income, convergence, and time related factors in explaining changes in PM_{2.5} pollution in a global panel of 158 countries between 1990 and 2010. We find that economic growth has positive but relatively small effects, time effects are also small but larger in wealthier and formerly centrally planned economies, and, for our main dataset, convergence effects are small and not statistically significant. There is no in-sample income turning point for regressions that include both the convergence variables and a set of control variables.

Keywords:

Air pollution; economic growth; environmental Kuznets curve

JEL Classification:

O44, Q53, Q56

Suggested Citation:

Stern, D.I. and van Dijk, J. (2016), Economic growth and global particulate pollution concentrations, CCEP Working Paper 1604, Feb 2016. Crawford School of Public Policy, The Australian National University.

Address for Correspondence:

David I. Stern
Professor
Crawford School of Public Policy
The Australian National University
132 Lennox Crossing
Acton
ACT 2601
Australia
Tel: +61 2 61250176
Email: david.stern@anu.edu.au

The Crawford School of Public Policy is the Australian National University's public policy school, serving and influencing Australia, Asia and the Pacific through advanced policy research, graduate and executive education, and policy impact.

[The Centre for Climate Economics & Policy](#) is an organized research unit at the Crawford School of Public Policy, The Australian National University. The working paper series is intended to facilitate academic and policy discussion, and the views expressed in working papers are those of the authors. Contact for the Centre: Dr Frank Jotzo, frank.jotzo@anu.edu.au

Economic Growth and Global Particulate Pollution Concentrations

David I. Stern

Crawford School of Public Policy, The Australian National University, 132 Lennox Crossing, Acton, ACT 2601, Australia. david.stern@anu.edu.au

Jeremy van Dijk

Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra, ACT 2601, Australia.

29 February 2016

Abstract

Though the environmental Kuznets curve (EKC) was originally developed to model the ambient concentrations of pollutants, most subsequent applications focused on pollution emissions. Yet, previous research suggests that it is more likely that economic growth could eventually reduce the concentrations of local pollutants than emissions. We examine the role of income, convergence, and time related factors in explaining changes in PM_{2.5} pollution in a global panel of 158 countries between 1990 and 2010. We find that economic growth has positive but relatively small effects, time effects are also small but larger in wealthier and formerly centrally planned economies, and, for our main dataset, convergence effects are small and not statistically significant. There is no in-sample income turning point for regressions that include both the convergence variables and a set of control variables.

Keywords: Air pollution, economic growth, environmental Kuznets curve

JEL Codes: O44, Q53, Q56

Acknowledgements: We thank Rohan Best, Zsuzsanna Csereklyei, Paul Burke, Astrid Kander, and Luis Sanchez for helpful comments.

1. Introduction

Particulate pollution, especially PM_{2.5}, is thought to be the form of pollution with the most serious human health impacts (WHO, 2013). It is estimated that PM_{2.5} exposure causes 3.1 million deaths a year, globally, and any level above zero is deemed unsafe, i.e. there is no threshold above zero below which negative health effects do not occur (WHO 2013). Black carbon is an important fraction of PM_{2.5} pollution (Vidanoja *et al.*, 2002) that may contribute significantly to anthropogenic radiative forcing (Bond *et al.*, 2013) and, therefore, there may be significant co-benefits to reducing its concentration (Victor *et al.*, 2015). Though the environmental Kuznets curve (EKC) was originally developed to model the ambient concentrations of pollutants, most subsequent applications focused on pollution emissions. Yet, it would seem more likely that economic growth could eventually reduce the concentrations of local pollutants than emissions (Selden and Song, 1994; Stern *et al.*, 1996). Here, we examine the role of income, convergence, and time related factors in explaining changes in PM_{2.5} particulate pollution in a global panel of countries between 1990 and 2010. We use a recently developed model that integrates the EKC and convergence approaches. We find that economic growth has positive but relatively small effects, time effects are also small but larger in wealthier and formerly centrally planned economies, and, for our main dataset, convergence effects are small and not statistically significant. The surprising finding is that there isn't an EKC even for local pollution concentrations, though the effects of economic growth are much smaller than they are for emissions of carbon and sulfur dioxide.

The environmental Kuznets curve (EKC) has been the dominant approach among economists to modeling ambient pollution concentrations and aggregate emissions since Grossman and Krueger (1991) introduced it a quarter of a century ago. The EKC is characterized by an income turning point – the level of GDP per capita after which economic growth reduces rather than increases environmental impacts. Though the EKC was originally developed to model the ambient concentrations of pollutants, most subsequent applications have focused on pollution emissions and in particular carbon dioxide and sulfur dioxide (Carson, 2010). Recent studies using global data sets find that, in fact, income has a monotonic positive effect on the emissions of both these pollutants (Wagner, 2008; Vollebergh *et al.*, 2009; Stern, 2010; Anjum *et al.*, 2014).

Both Selden and Song (1994) and Stern *et al.* (1996) noted that ambient concentrations of pollutants were likely to fall before emissions did. Stern (2004) suggests that this may be due to both the decline in urban population densities and the decentralization of industry that tend

to accompany economic growth. Furthermore, actions through which governments can try to reduce local air pollution include moving industry outside of populated areas and building taller smokestacks. The latter reduced urban air pollution in developed countries in the 20th Century at the expense of increasing acid rain in neighboring countries and the formation of sulfate aerosols (Wigley and Raper, 1992). Additionally, pollutants with severe and obvious human health impacts such as particulates are more likely to be controlled earlier than pollutants with less obvious impacts such as carbon dioxide (Shafik, 1994). Despite this, relatively little recent research has attempted to apply the EKC to pollution concentrations rather than emissions.

More recently, it has become popular to model the evolution of emissions using convergence approaches. Pettersson *et al.* (2013) provide a review of the literature on convergence of carbon emissions. There are three main approaches to testing for convergence: sigma convergence, which tests whether the dispersion of the variable in question declines over time using either just its variance or its full distribution (e.g. Ezcurra, 2007); stochastic convergence, which tests whether the time series for different countries cointegrate; and beta convergence, which tests whether the growth rate of a variable is negatively correlated to the initial level. We are not aware of attempts to test for convergence in pollution concentrations rather than emissions. Yet, it seems reasonable that high concentrations of pollution would encourage defensive action to reduce that pollution (Ordás Criado *et al.*, 2011).

Sanchez and Stern (2016) propose a regression model that nests both the EKC and beta convergence models, which can be seen as an extension of Ordás Criado *et al.*'s (2011) model to also include the EKC effect. The model allows us to test the contributions of economic growth, convergence, and time effects to the evolution of pollution.

Our main results use population-weighted estimates of national average concentrations of PM_{2.5} pollution from the *World Bank Development Indicators*. These data are based on Brauer *et al.* (2016), who used satellite observations of aerosol optical depth, pollution emissions data to obtain estimates, which were then regressed on the available ground-based observations. The resulting coefficients were used to project calibrated PM_{2.5} for all parts of the world. To check robustness we also use the *Environmental Performance Index* dataset. These data are based on Boys *et al.* (2014) and van Donkelaar *et al.* (2015). Neither of these latter studies uses ground-based ambient observations in deriving their estimates. More details are provided in the Appendix. Because both these datasets are weighted by population exposure, they most reflect the concentrations of these pollutants in densely populated areas such as cities. Thus,

though obviously particulates travel between cities and countries in the wind, we are capturing local pollution to a large extent with this data set.

The next section of the paper reviews previous research on modeling particulate pollution concentrations. The third section presents our modeling approach, the fourth our data, and the fifth our results. The sixth section presents our conclusions.

2. Previous Research

Grossman and Krueger (1991) estimated the first EKC models as part of a study of the potential environmental impacts of NAFTA. They estimated EKCs for SO₂, dark matter (fine smoke), and suspended particles (SPM) using the GEMS dataset. This dataset is a panel of ambient measurements from a number of locations in cities around the world. Each regression involved a cubic function in levels (not logarithms) of PPP (Purchasing Power Parity adjusted) per capita GDP, various site-related variables, a time trend, and a trade intensity variable. The turning points for SO₂ and dark matter were at around \$4,000-5,000 while the concentration of suspended particles appeared to decline even at low income levels. However, Harbaugh *et al.* (2002) re-examined an updated version of Grossman and Krueger's data. They found that the locations of the turning points for the various pollutants, as well as even their existence, were sensitive both to variations in the data sampled and to reasonable changes in the econometric specification.

Shafik's (1994) study was particularly influential, as its results were used in the 1992 *World Development Report*. Shafik estimated EKCs for ten different indicators using three different functional forms. She found that local air pollutant concentrations conformed to the EKC hypothesis with turning points between \$3,000 and \$4,000. Selden and Song (1994) estimated EKCs for four emissions series: SO₂, NO_x, SPM, and CO. The estimated turning points were all very high compared to the two earlier studies. For the fixed effects version of their model they are (in 1990 US dollars): SO₂, \$10,391; NO_x, \$13,383; SPM, \$12,275; and CO, \$7,114. This showed that the turning points for emissions were likely to be higher than for ambient concentrations.

There has been little recent EKC research on particulate pollution. Keene and Deller (2015) recently published an EKC analysis of PM_{2.5} concentrations for a cross-section of U.S. counties. The model includes state dummies and various control variables and they use OLS and spatial econometric estimators. They find that the peak of the EKC occurs at between

US\$24,000 and US\$25,500, depending on the estimator used.

Some recent research focuses on Chinese cities. Brajer *et al.* (2011) investigate ambient concentrations of SO₂, NO₂, and total suspended particulates and also construct indices of total air pollution using the Nemerow approach and an alternative proposed by Khanna (2000). Their data cover the period 1990-2006 for 139 Chinese cities. They use a logarithmic EKC model with city random effects and a linear time trend with the addition of population density variable. Using the quadratic EKC model, they estimate the turning point for TSP at RMB 3,794, not controlling for population density, and at RMB 6,253, controlling for population density. However, the regression coefficient of the cube of log income in a cubic EKC model is statistically significantly greater than zero. This second turning point occurs around RMB 125k. Hao and Liu (2016) estimate EKC models for PM_{2.5} concentrations and the official Air Quality Index in a cross-section of 73 Chinese cities in 2013. They find an inverted U shape curve with highly significant parameter estimates for OLS and SEM estimates, with turning points of RMB 9k to 40k and PM_{2.5}, respectively.

3. Models

Our model combines the three main approaches in the literature and includes other possible drivers of change in concentrations by nesting these existing specifications in a single regression equation:

$$\hat{C}_i = \alpha + \beta_1 \hat{G}_i + \beta_2 G_{i0} \hat{G}_i + \beta_3 G_{i0} + \beta_4 C_{i0} + \sum_j \psi_j X_{ji} + \varepsilon_i$$

where i indexes countries, 0 indicates the initial year of the sample, and ε_i is a random error term. $\hat{C}_i$ and $\hat{G}_i$ are the long-run growth rates of concentrations and income, respectively. G_{i0} is the log of income per capita in the first year in the sample in each country and C_{i0} is the log of concentrations in the initial year. X is a vector of additional explanatory or “control” variables. We also estimate models that exclude the control variables, and which exclude the control and levels variables, G_{i0} and C_{i0} . The latter model is analogous to the traditional EKC model, but estimated using differences rather than levels of the variables.

We compute long-run growth rates using: $\hat{Y}_i = (Y_{iT} - Y_{i0})/T$, where Y is the logarithm of concentrations or per capita income and $T+1$ is the number of years in the sample. By formulating our model in long-run growth rates we avoid most of the econometric problems

troubling the existing literature on the environmental Kuznets curve, which are discussed in several recent contributions (Wagner, 2008, 2015; Vollebergh *et al.*, 2009; Stern, 2010; Anjum *et al.*, 2014).

We subtract the means of all the continuous levels variables (as opposed to growth rates or dummy variables) prior to estimation. Therefore, the first term on the RHS of the equation, α , is the growth rate of concentrations when there is no economic growth and all the other continuous levels variables are at their sample means. This can, therefore, be interpreted as the average time effect. β_1 is an estimate of the income-concentrations elasticity at the sample mean. The third term tests for the EKC effect. If β_2 is statistically significantly negative and β_1 is positive, then there is a level of income after which concentrations reduce with growth. We can find the EKC turning point, μ , by estimating the regression without demeaning G_{i0} prior to computing $G_{i0}\hat{G}_i$ and then computing $\mu = \exp(-\beta_1/\beta_2)$ using the estimated coefficients. If this turning point is within the sample range of income then there is an environmental Kuznets curve. If β_2 is negative, but the turning point is out of sample, we can still say that there is an EKC effect so that growth has a reduced effect on concentrations at higher income levels.

The fourth term tests whether concentrations change at a different rate in richer countries in the absence of growth and the fifth term is intended to model convergence by allowing us to test for convergence in concentrations using the beta convergence approach. If $\beta_4 < 0$, $t_{\beta_4} > 0$ concentrations converge across countries so that concentrations growth is slower in countries that commence the period with higher pollution concentrations and *vice versa*.

A wide variety of “control variables” have been considered in the EKC literature. Some of these are genuinely exogenous or predetermined, whereas others are variables that typically change in the course of economic development and might be seen as factors through which the development process drives concentrations changes. Examples of the latter are democracy, free press, good governance, lack of corruption, or industrial structure. We are interested in testing the overall effect of income and economic growth on pollution growth and so we only include variables that are pre-determined or exogenous to the development process and found in previous research to be potentially relevant.

Stern (2005) first noted that English speaking OECD countries seemed to abate sulfur emissions less and Germanic and Scandinavian countries more. Stern (2012) related this to

differences in legal origins (La Porta *et al.*, 2008) and found that energy intensity was lower in non-English legal origin countries, *ceteris paribus*. Fredriksson and Wollscheid (2015) present evidence that legal origin has a significant effect on environmental policy. Here, we include dummies for French and German legal origin. We also control for whether a country was a formerly centrally planned country using a dummy variable. We expect that market reform would reduce the level of pollution, *ceteris paribus*.

We also control for the effect of climate, by using historical country averages of temperatures over the three summer months and the three winter months, annual precipitation, and average elevation above sea level. The latter two variables are converted to logarithms. We also control for landlocked status, as landlocked countries are less likely to have the higher wind speeds seen in coastal regions. Indonesia, Singapore and Malaysia experienced very high levels of pollution in 1990 associated with the periodic haze episodes due to forest fires in the region (Osterman and Brauer, 2001). We add a dummy for these three countries. Finally, we include the average of the log of population density, which might be expected to increase the concentration of pollution, *ceteris paribus*. Also, the higher population is, the more people will be exposed to pollution and the more likely that action might be taken (Ordás Criado *et al.*, 2011).

When observations on variables are aggregated into regions – here countries - of different sizes, it is likely that much of the local variation across individual locations is cancelled out in the larger regions while more idiosyncratic variation remains in smaller regions. This means that the error terms in a regression using such aggregated data are likely to be heteroskedastic with the error variance proportional to the district size (Maddala, 1977; Stern, 1994). As our data consists of population-weighted concentrations by country, the appropriate measure of region size is population. To address this grouping heteroskedasticity we estimate the models using weighted least squares, where the weights are the square root of population, and heteroskedasticity-robust standard errors. Using weighted least squares (WLS) can result in large efficiency gains over using ordinary least squares (OLS) even when the model for reweighting the data is misspecified. But in case there is misspecification, heteroskedasticity robust standard errors should be used to ensure correct inference (Romano and Wolf, 2014). We measure goodness of fit using Buse's (1973) R-squared, weighting the squared deviations by population.

We assume that the explanatory variables in our regressions are exogenous. Clearly, there can be no reverse causality from growth rates to initial values. There may potentially be apparent feedback from the growth rate of concentrations to the growth rate of income. This is because pollution growth may be correlated with the growth of energy use and energy use contributes to economic growth. Csereklyei and Stern (2015) argue that this bias will be fairly small even when the dependent variable is energy use and so the estimated energy-income elasticity will not be far from the causal effect size of an exogenous change in income. Here, the potential bias should be smaller still. Omitted variables bias is an important issue as there are many variables that may be correlated with the level of GDP or GDP growth, and which may help explain concentrations growth. Our differenced approach should help reduce this bias (Angrist and Pischke, 2010). Finally, measurement error is a significant issue in the estimation of GDP and pollution concentrations. Obviously there are significant uncertainties in the concentrations data, which are modeled based on satellite and ground-based measurements. Measurement error is likely greater for some of the smaller economies. Weighted least squares can, therefore, help reduce the effects of this measurement error. A common approach to dealing with reverse causality, omitted variables bias, and measurement error is to use instrumental variables. However, it is hard to find plausible instrumental variables in the macro-economic context (Bazzi and Clemens, 2013).

4. Data

Details of the data sources are presented in the Appendix.

Table 1 presents some descriptive statistics for our variables. These are the variables sourced from the *World Bank Development Indicators* used in the main analysis. The PM_{2.5} concentration exposure level in 1990 is mostly above the WHO guideline of 10 $\mu\text{g}/\text{m}^3$ with a mean and median of 18-19 $\mu\text{g}/\text{m}^3$. The range of observations is quite large, from below 1 $\mu\text{g}/\text{m}^3$ to over 76 $\mu\text{g}/\text{m}^3$ in Mauritania. In 1990, Singapore had the second highest level of PM_{2.5} at 49.8 $\mu\text{g}/\text{m}^3$. But this was anomalously high as discussed above. As of 2005, 89% of the world's population was exposed to annual mean PM_{2.5} concentration levels higher than the WHO concentration guideline of 10 $\mu\text{g}/\text{m}^3$, while approximately two thirds of countries were in this category (Brauer *et al.*, 2012). This difference is due to the large populations in East and South Asia, which have high PM_{2.5} concentration levels. In the base year of the study, 1990, 67% of countries in the sample had exposure levels higher than the WHO recommendation.

The level of initial per capita GDP has a wide range from \$365 to \$114,519 in constant 2011 PPP Dollars. While mean income per capita is \$11,895, the median value is only half the mean, at \$6,440. The descriptive statistics for the continuous control variables exhibit the wide range that would be expected in a globally representative sample.

Table 1 also presents the annual growth rates of income per capita and pollution. GDP per capita grows at an average rate of 1.76% p.a. over the period 1990 to 2010. The median is only 0.13 percentage points lower. The income growth rates are mostly positive, however 24 countries had negative growth over the period. There are two outliers with GDP per capita growth rates larger than 9% p.a. – China and Equatorial Guinea. Compared to GDP growth rates, the growth rates of pollution exposure are mostly modest. The mean rate of decline was -0.35% p.a., and the median 0.17% p.a. 78 countries had positive growth in PM_{2.5} exposure over the period. The highest growth was in Micronesia, averaging 6.5% p.a., while the most rapid decrease was observed in Singapore averaging -6.9% p.a. But in both these countries there was a large change in one decade but not the other. In fact, while the mean annual rate of decline of PM_{2.5} in the 1990s was 0.88%, PM_{2.5} concentrations on average increased in the following decade with an average annual growth rate of 0.18%.

Figure 1 presents the data in growth rates form. There would not be much point in presenting the actual concentrations of pollution as the mean levels are swept out when growth rates are computed and much of the variation in levels reflects idiosyncrasies of geography. The size of the bubbles is proportional to population in 1990, which is used to weight the observations in the regression analysis. The large circle to the right is, of course, China, with India to its left. The USA is the largest circle among the countries with negative pollution growth rates. Indonesia is to its lower right. As we can see, both pollution and GDP per capita rose quite strongly in the World's two most populous countries. This and the negative pollution growth rates in many of the countries with moderate growth suggest that economic growth should have significant effects on pollution growth. OLS estimates are likely to be influenced by some of the small outlier countries such as Equatorial Guinea on the far right of the Figure, which is mitigated by using WLS to estimate our main models.

5. Results

Table 2 presents the main results, which use World Bank pollution and GDP data. The simple EKC model has a turning point at \$3,336, which is statistically significantly different from zero.

The concentrations-GDP elasticity at the sample mean is -0.18, though not very precisely estimated. It is negative because the income turning point is below the mean income in the sample. The time effect is small and not statistically significant. These results would seem to strongly support the environmental Kuznets curve story and the hypothesis that the income turning point is lower for concentrations of local pollutants than it is for emissions of pollutants such as sulfur dioxide. The second column adds the two initial levels terms. The EKC turning point is now much higher, but very imprecisely estimated. In addition, the concentrations-GDP elasticity at the sample mean is now positive. As expected, the coefficient of initial pollution is negative indicating beta convergence. The coefficient of initial income is also negative and very significant. This implies that concentrations fall faster in richer countries, *ceteris paribus*.

When we add the control variables, the concentrations-income elasticity rises to 0.21 at the sample mean and is highly significant and the interaction term, which tests the EKC hypothesis, is significant at the 10% level. However the EKC turning point rises further to \$66,728 so that the EKC is effectively monotonic. The effects of the initial levels are reduced in strength and statistical significance. Of the control variables, concentrations rise faster (or fall slower) in countries with higher summer, lower winter temperatures, and higher precipitation and rise slower (or fall faster) in formerly centrally planned countries as we would expect. Of these, the effect of precipitation is unexpected, as higher precipitation would be expected to clear the air. Many of the countries where concentrations fell strongly are in Europe and have moderate levels of rainfall around 500-1000mm, while many of the countries where concentrations rose most strongly happen to be in areas of heavy rainfall in the tropics. This effect might be related, therefore, to deforestation. The Malaysia, Singapore, and Indonesia dummy has a highly significant and negative effect on concentrations growth.

Results are, therefore, similar to those found by Anjum *et al.* (2014) for sulfur and carbon emissions, but the effect of economic growth is far smaller and even smaller than that for non-industrial greenhouse gas emissions (Sanchez and Stern, 2016). The convergence effect is also weaker than for industry related emissions. When we control for other relevant variables there is not even an environmental Kuznets curve for particulate concentrations.

We also present results for the following variations, to test robustness to different data sources, time periods, and estimation methods:

1. Use OLS instead of WLS.
2. Split the data used in Table 2 into two time periods – 1990-2000 and 2000-2010.

3. Use income from the Penn World Table instead of the World Bank.
4. Use pollution data from EPI instead of the World Bank.

We report these results for the full model in Table 3. Looking first at the OLS results, the main differences are that both income terms are much smaller and not significant, the convergence effect is highly significant, the effects of elevation and legal origin are larger and much more significant, and the effects of centrally planned status are smaller. On the other hand, these results are not dramatically different from the WLS results. One of the advantages of the latter are that they are much more robust to changes in the sample of countries, as we go to the remaining analyses.

The results from splitting the sample into 1990-2000 and 2000-2010 periods in the Columns 2 and 3 differ in somewhat expected ways from the 1990-2010 estimates in Table 2. Again the EKC is effectively monotonic but in one case there is an out of sample turning point and in the other a minimum near zero. The income elasticity at the sample mean is higher in the second period. One reason for this is that income increases from the first to second period. The effects of central planning and the Malaysia, Singapore, and Indonesia dummy decrease in the second period, as we would expect. Unexpected results are that elevation has a positive effect in the first period and precipitation only in the second period. Though these results show a stronger effect of growth in the second period, the effect of growth on concentrations is still relatively small compared to estimates for emissions of other pollutants related to industrial activity but about the same as for non-industrial greenhouse gas emissions, which are primarily from land-use change (Sanchez and Stern, 2016).

Results using income data from the Penn World Table in Column 4 are very similar to those for the World Bank income data in Table 2 but there are more statistically significant coefficients including for elevation, landlockedness, and French legal origin. However, central planning is not statistically significant here.

The results in the final column using EPI data for 2000-2010 and World Bank income data are similar in some respects to the 2000-2010 World Bank pollution data estimates in Column 3. The concentrations-income elasticity is 0.38 and very statistically significant. In contrast to the World Bank pollution data, the convergence effect is quite large and statistically significant. Also, landlockedness and French legal origin now have significant negative effects and population density a significant positive effect.

6. Conclusions

The evidence presented in this article shows that economic growth has positive though relatively small effects on the growth in PM_{2.5} concentrations. For our models that include convergence terms and control variables there is no sign of in-sample income turning point. However, when we estimate a model analogous to the classic EKC model we find a turning point of around \$3,000 per capita. Our results suggest that prior studies that find a relatively low income turning point for the environmental Kuznets curve for particulate concentrations (e.g. Grossman and Krueger, 1991; Shafik, 1994; Brajer *et al.*, 2011; Hao and Liu, 2016) suffer from omitted variables bias. Our results are more similar to Keene and Deller (2015) who found a much higher, but still in-sample, turning point for U.S. counties.

On the other hand, the negative time effect is stronger in richer countries, but this is unrelated to increases in income. What is clear is that this behavior is very different from emissions of sulfur or industrial greenhouse gases where typically a strong positive effect of economic growth is found at the sample mean income (Anjum *et al.*, 2014; Sanchez and Stern, 2016). That there is not even an EKC for particulate pollution, which is a classic example of a mostly local pollutant that impacts human health, casts further doubt on the general usefulness of the EKC model.

References

- Amante, C. and Eakins, B. W. 2009. *Etopo1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis*, NOAA Technical Memorandum NESDIS NGDC-24, National Geophysical Data Center, NOAA.
- Angrist, J. D. and J.-S. Pischke. 2010. The credibility revolution in empirical economics: how better research design is taking the con out of econometrics. *Journal of Economic Perspectives*, 24(2): 3-30.
- Anjum Z., P. J. Burke, R. Gerlagh, and D. I. Stern. 2014. Modeling the emissions-income relationship using long-run growth rates, *CCEP Working Papers* 1403.
- Bazzi, S. and M. A. Clemens. 2013. Blunt instruments: Avoiding common pitfalls in identifying the causes of economic growth. *American Economic Journal: Macroeconomics* 5(2): 152–186.
- Bond, T. C., S. J. Doherty, D. W. Fahey, P. M. Forster, T. Berntsen, B. J. DeAngelo, M. G. Flanner, S. Ghan, B. Kärcher, D. Koch, S. Kinne, Y. Kondo, P. K. Quinn, M. C. Sarofim, M. G. Schultz, M. Schulz, C. Venkataraman, H. Zhang, S. Zhang, N. Bellouin, S. K. Guttikunda, P. K. Hopke, M. Z. Jacobson, J. W. Kaiser, Z. Klimont, U. Lohmann, J. P. Schwarz, D. Shindell, T. Storelvmo, S. G. Warren, C. S. Zender. 2013. Bounding the role of black carbon in the climate system: a scientific assessment, *Journal of Geophysical Research-Atmospheres*

118(11): 5380–5552.

Boys, B., Martin, R. V., van Donkelaar, A., MacDonald, R., Hsu, N. C., Cooper M. J., *et al.* 2014. Fifteen year global time series of satellite-derived fine particulate matter. *Environmental Science and Technology* 48: 11109–11118.

Brajer, V., Mead, R. W. Xiao, F. 2011. Searching for an Environmental Kuznets Curve in China's air pollution, *China Economic Review* 22, 383–397.

Brauer, M., Amann, M., Burnett, R. T., Cohen, A., Dentener, F., Ezzati, M., Henderson, S. B., Krzyzanowski, M., Martin, R. V., Van Dingenen, R., Van Donkelaar, A., and Thurston, G. D. 2012. Exposure assessment for estimation of the global burden of disease attributable to outdoor air pollution, *Environmental Science and Technology* 46(2): 652-660.

Brauer, M., G. Freedman, J. Frostad, A. van Donkelaar, *et al.* 2016. Ambient Air Pollution Exposure Estimation for the Global Burden of Disease 2013, *Environmental Science and Technology* 50(1), 79-88.

Buse, A. 1973. Goodness of fit in generalized least squares estimation. *American Statistician* 27(3): 106-108.

Carson, R. T. 2010. The environmental Kuznets curve: Seeking empirical regularity and theoretical structure. *Review of Environmental Economics and Policy* 4(1): 3-23.

Csereklyei, Z. and D. I. Stern. 2015. Global energy use: Decoupling or convergence? *Energy Economics* 51: 633-641.

Ezcurra, R. 2007. Is there cross-country convergence in carbon dioxide emissions? *Energy Policy*, 35, 1363-1372.

Feenstra, R. C., R. Inklaar, and M. P. Timmer. 2015. The next generation of the Penn World Table. *American Economic Review* 105(10): 3150-82.

Fredriksson, P. G. and Wollscheid, J. R. 2015. Legal origins and climate change policies in former colonies, *Environmental and Resource Economics* 62(2), 309-327.

Grossman, G. M. & Krueger, A. B. 1991. Environmental impacts of a North American Free Trade Agreement. *NBER Working Papers*, 3914.

Hao, Y. & Liu, Y.-M. (2016) The influential factors of urban PM 2.5 concentrations in China: a spatial econometric analysis, *Journal of Cleaner Production* 112, 1443-1453.

Harbaugh, W., Levinson, A., & Wilson, D. 2002. Re-examining the empirical evidence for an environmental Kuznets curve. *Review of Economics and Statistics*, 84(3), 541–551.

Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H. 2014. Updated high-resolution grids of monthly climatic observations - the CRU Ts3.10 Dataset, *International Journal of Climatology* 34(3): 623-642.

Keene, A. & Deller, S. C. 2015. Evidence of the environmental Kuznets' curve among US counties and the impact of social capital, *International Regional Science Review* 38(4), 358-387.

Khanna, N. 2000. Measuring environmental quality: An index of pollution, *Ecological Economics* 35, 191–202.

La Porta, R., F. Lopez-de-Silanes, and A. Shleifer. 2008. The economic consequences of legal origins. *Journal of Economic Literature* 46(2): 285-332.

Maddala, G. S. 1977. *Econometrics*. McGraw-Hill, Singapore.

Ordás Criado, C., Valente, S., & Stengos, T., 2011. Growth and pollution convergence: Theory and evidence. *Journal of Environmental Economics and Management*, 62, 199-214.

Osterman, K. and M. Brauer. 2001. Air quality during haze episodes and its impact on health, in: P. Eaton and M. Radojevic (eds.) *Forest Fires and Regional Haze in Southeast Asia*, Nova Science Publishers, New York. 195-226.

Pettersson, F., Maddison, D., Acar, S., & Söderholm, P. 2013. Convergence of carbon dioxide emissions: A review of the literature. *International Review of Environmental and Resource Economics*, 7, 141–178.

Romano, J. P. and M. Wolf. 2014. Resurrecting weighted least squares. *University of Zurich, Department of Economics, Working Paper Series* 172.

Sanchez, L. F. & Stern, D. I. 2016. Drivers of industrial and non-industrial greenhouse gas emissions. *Ecological Economics* 124, 17-24.

Selden, T. M. & Song, D. 1994. Environmental quality and development: Is there a Kuznets curve for air pollution? *Journal of Environmental Economics and Environmental Management*, 27, 147–162.

Shafik, N. 1994. Economic development and environmental quality: an econometric analysis. *Oxford Economic Papers*, 46, 757–773.

Stern, D. I. 1994. Historical path-dependence of the urban population density gradient. *Annals of Regional Science* 28: 197-223.

Stern, D. I. 2004. The rise and fall of the environmental Kuznets curve. *World Development* 32(8): 1419-39.

Stern, D. I. 2005. Beyond the environmental Kuznets curve: Diffusion of sulfur-emissions-abating technology. *Journal of Environment and Development* 14(1): 101-24.

Stern, D. I. 2010. Between estimates of the emissions-income elasticity. *Ecological Economics* 69: 2173-2182.

Stern, D. I. 2012. Modeling international trends in energy efficiency. *Energy Economics* 34: 2200–2208.

Stern, D. I., Common, M. S., and Barbier, E. B. 1996. Economic growth and environmental degradation: the environmental Kuznets curve and sustainable development. *World Development*, 24, 1151–1160.

van Donkelaar, A., R. V. Martin, M. Brauer and B. L. Boys. 2015. Use of Satellite Observations for Long-Term Exposure Assessment of Global Concentrations of Fine Particulate Matter, *Environmental Health Perspectives* 123(2): 135-143.

Victor, D. G., Zaelke, D., & Ramanathan, V. (2015) Soot and short-lived pollutants provide political opportunity, *Nature Climate Change* 5, 796–798.

Viidanoja, J., Sillanpää, M., Laakia, J., Kerminen, V.-M., Hillamo, R., Aarnio, P., Koskentalo, T. 2002. Organic and black carbon in PM_{2.5} and PM₁₀: 1 year of data from an urban site in Helsinki, Finland, *Atmospheric Environment* 36(19): 3183–3193.

Vollebergh, H. R. J., B. Melenberg, and E. Dijkgraaf. 2009. Identifying reduced-form relations with panel data: The case of pollution and income. *Journal of Environmental Economics and Management* 58(1): 27-42.

Wagner, M. 2008. The carbon Kuznets curve: A cloudy picture emitted by bad econometrics. *Resource and Energy Economics* 30: 388-408.

Wagner, M. 2015. The environmental Kuznets curve, cointegration and nonlinearity. *Journal of Applied Econometrics* 30(6): 948-967.

WHO. 2013. *Health Effects of Particulate Matter*, World Health Organisation, Copenhagen, Denmark.

Wigley, T. M. L. & Raper, S. C. B. 1992. Implications for climate and sea level of revised IPCC emissions scenarios, *Nature* 357, 293-300.

Appendix

The pollution datasets used in this paper have slightly different methodologies and data sources. Both datasets used provide population-weighted mean annual exposure to PM_{2.5} in micrograms per m³ for all countries across the globe.

The *Environmental Performance Index* (EPI) dataset derives estimates from the studies of van Donkelaar *et al.* (2015) and Boys *et al.* (2014). Both these studies used the GEOS-Chem chemical transport model (CTM) to relate satellite observations of Aerosol Optical Depth (AOD) to ground-level PM_{2.5} concentration levels. The two papers used the satellite instruments named MISR and SeaWiFS, while van Donkelaar *et al.* additionally utilized MODIS. The spatial resolution of the concentration data differed from grids of 10x10km in van Donkelaar *et al.* and 1x1 degree in Boys *et al.* While the latter reported concentration values for each grid, the former additionally calculated national population-weighted exposure levels, as the EPI reported, using population data from the Global Rural Urban Mapping Project database. van Donkelaar *et al.* additionally compared the estimates with ground-based observations from trusted established networks in North America and Europe and 210 other global sites from other publications. The satellite observations of North America closely matched the ground-based findings, with a regression slope of 0.96 where the ground-based data are the dependent variable. Globally, however, their estimates had a poorer fit, with a regression slope of 0.68.

The *World Bank Development Indicators* dataset is based on the study of Brauer *et al.* (2016). As with van Donkelaar *et al.* and Boys *et al.*, this study used AOD data obtained from the MODIS, MISR and SeaWiFS satellite instruments, with the additional use of the CALIOP instrument. As above, Brauer *et al.* utilized the GEOS-Chem CTM to relate the satellite AOD observations to ground-level PM_{2.5} concentrations. This study, however, additionally used the TM5-FASST model to provide estimates of PM_{2.5} concentrations from pollutant emissions and meteorological data. The mean of the satellite and TM5 values for each grid were then regressed on the available ground-based observations, and the resulting coefficients used to produce ‘calibrated’ PM_{2.5} estimates across the globe based on the means of the satellite and TM5 data. The spatial resolution used by Brauer *et al.* is 0.1x0.1 degree. To calculate national population-weighted exposure levels, population data was used from the Gridded Population of the World (GPW) v3.

These data were downloaded as follows:

EPI (Environmental Performance Index) (2015), 2014 EPI Downloads, Yale University, viewed 11/2015, at <<http://www.epi.yale.edu/downloads>>.

World Bank (2015), *PM2.5 air pollution, mean annual exposure (micrograms per cubic meter)*, viewed 12/2015, at <<http://data.worldbank.org/indicator/EN.ATM.PM25.MC.M3>>.

GDP, population, and area data are from the *World Bank Development Indicators*. We also used income and population data from the *Penn World Table* (Feenstra *et al.*, 2015) as a robustness check. The dummy variables for legal origin came from LaPorta *et al.* (2008). The dummy variables for whether or not a country is landlocked and for centrally planned economies came from the NYU Development Research Institute (2009):

NYU Development Research Institute 2009, *Global Development Network Growth Database*, viewed 07/2015 <<http://www.nyudri.org/resources/global-development-network-growth-database/%3E>

The climate variables used in this study come from the Climate Research Unit of the University of East Anglia (Harris *et al.* 2014). The temperatures are provided as monthly means for each country, which is then averaged for each season used (winter and summer) and over the time period of the study. The temperatures are given in degrees Celsius. Precipitation is given as total annual precipitation level in millimeters for each country and is then averaged over 1990-2010. Amante and Eakins (2009) provided the observations of each nation's mean elevation above sea level in meters.

Table 1: Descriptive Statistics

Variable	Mean	Median	Min	Max
PM _{2.5} exposure 1990 ($\mu\text{g}/\text{m}^3$)	19.35	18.06	1.16	76.51
GDP per capita 1990 (2011 \$PPP)	11,895	6,440	375	114,519
Growth rate of PM _{2.5} concentrations 1990- 2010 (% p.a.)	-0.35%	-0.17%	-6.92%	6.50%
Growth rate of GDP per capita 1990-2010 (% p.a.)	1.76%	1.63%	-3.77%	17.80%
Mean Summer Temperature ($^{\circ}\text{C}$)	24.0	25.3	8.5	36.9
Mean Winter Temperature ($^{\circ}\text{C}$)	14.1	18.6	-22.6	28.6
Annual Precipitation (mm)	1,221	1,054	41	3,653
Mean national elevation (masl)	625	442	9	3,186
Population Density (people/ km^2)	357	62	1	19,890

Table 2. Main Results: World Bank Pollution and GDP 1990-2010

Variable	EKC	EKC & Convergence	Full Model
Constant	-0.0009 (0.0040)	-0.0021 (0.0028)	-0.0030 (0.0022)
$\hat{G}_t$	-0.1760 (0.1479)	0.0637 (0.1487)	0.2089*** (0.0694)
$\hat{G}_t \times G_{0i}$	-0.2880*** (0.0861)	-0.0892 (0.1102)	-0.0876* (0.0482)
G_{0i}		-0.0069*** (0.0024)	-0.0043** (0.0018)
C_{0i}		-0.0078* (0.0044)	-0.0048 (0.0032)
Mean Summer Temperature			0.0020*** (0.0004)
Mean Winter Temperature			-0.0005** (0.0002)
Log Annual Precipitation			0.0063*** (0.0024)
Log Mean Elevation			0.0018 (0.0016)
Landlocked			-0.0003 (0.0027)
French Legal Origin			-0.0004 (0.0028)
German Legal Origin			-0.0033 (0.0033)
Centrally planned			-0.0085** (0.0036)
Malaysia, Singapore, and Indonesia			-0.0346*** (0.0046)
Log Population Density			-0.0022 (0.0015)
EKC income per capita turning point (\$PPP)	3,336*** (1,172)	12,557 (31,442)	66,728 (125,076)
n	158	158	132
Adjusted Buse R ²	0.5296	0.6318	0.8575

Notes: All variables demeaned (except LR growth rates and dummies). Heteroskedasticity-robust standard error in parentheses. Significance Levels: * 10%, ** 5%, *** 1%

Table 3. Robustness Checks

Variable	OLS WB Data 1990-2010	WLS WB Data 1990-2000	WLS WB Data 2000-2010	WLS PWT Income 1990-2010	WLS EPI Pollution 2000-2010
Constant	0.0004 (0.0021)	-0.0054*** (0.0018)	-0.0033 (0.0037)	-0.0025 (0.0026)	0.0073 (0.0047)
$\widehat{G}_t$	0.0307 (0.0540)	0.1838*** (0.0510)	0.4309*** (0.1398)	0.2343*** (0.0702)	0.3729*** (0.1101)
$\widehat{G}_t \times G_{0i}$	-0.0247 (0.0412)	-0.0292 (0.0509)	0.1006 (0.0740)	0.1453** (0.0651)	0.0265 (0.0971)
G_{0i}	-0.0021* (0.0011)	-0.0056*** (0.0019)	-0.0052* (0.0028)	-0.0106*** (0.0021)	-0.0118*** (0.0030)
C_{0i}	-0.0056*** (0.0018)	-0.0039 (0.0031)	-0.0028 (0.0050)	-0.0051 (0.0031)	-0.0140*** (0.0029)
Mean Summer Temperature	0.0013*** (0.0004)	0.0021*** (0.0005)	0.0019*** (0.0007)	0.0023*** (0.0004)	0.0030*** (0.0007)
Mean Winter Temperature	-0.0002 (0.0002)	-0.0004* (0.0003)	-0.0003 (0.0004)	-0.0008*** (0.0003)	-0.0013*** (0.0003)
Log Annual Precipitation	0.0016 (0.0018)	0.0048 (0.0031)	0.0083*** (0.0032)	0.0096*** (0.0031)	0.0034 (0.0030)
Log Mean Elevation	0.0040*** (0.0008)	0.0069*** (0.0021)	0.0006 (0.0022)	0.0049*** (0.0016)	0.0028 (0.0020)
Landlocked	0.0011 (0.0023)	-0.0037 (0.0028)	-0.0030 (0.0037)	-0.0081*** (0.0024)	-0.0117** (0.0052)
French Legal Origin	-0.0034* (0.0019)	-0.0012 (0.0021)	-0.0020 (0.0043)	-0.0046* (0.0026)	-0.0194*** (0.0046)
German Legal Origin	-0.0097*** (0.0034)	-0.0054 (0.0053)	0.0013 (0.0055)	-0.0042 (0.0040)	-0.0049 (0.0053)
Centrally planned	-0.0047 (0.0032)	-0.0087* (0.0053)	-0.0072 (0.0057)	0.0028 (0.0040)	-0.0013 (0.0060)
Malaysia, Singapore, and Indonesia	-0.0237*** (0.0029)	-0.0556*** (0.0030)	-0.0193** (0.0095)	-0.0331*** (0.0043)	0.0044 (0.0053)
Log Population Density	0.0004 (0.0007)	0.0021 (0.0025)	-0.0019 (0.0029)	0.0016 (0.0016)	0.0068*** (0.0017)
EKC income per capita turning point (\$PPP)	21,386 (74,564)	3,352,581 (40,156,293)	97 (249)	963 (549)	0.01 (0.26)
n	132	132	149	142	150
Adjusted Buse R ²	0.5071	0.8777	0.8730	0.8258	0.8630

Notes: All variables demeaned (except LR growth rates and dummies). Heteroskedasticity-robust standard error in parentheses. Significance Levels: * 10%, ** 5%, *** 1%

Figure 1. Growth Rates of PM 2.5 Pollution and GDP per Capita 1990-2010

Notes: Size of bubbles is proportional to population in 1990. Data source is *World Bank Development Indicators*.