



*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

*No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.*

# Effect of Vector Density and Competence on Macromolecular Vector Transformation Efficiency

Hongni Qin<sup>1\*</sup>, Lanlan ZHANG<sup>2</sup>

1. Suzhou Industrial Park Institute of Services Outsourcing, Suzhou 215123, China; 2. Genewiz (Suzhou) Biotechnology Co., Ltd., Suzhou 215123, China

**Abstract** To research the effects of vector quantity and competence on the positive cloning rate, with a known gene sequence but in the absence of DNA template, we artificially designed 26 primers to synthesize a target gene of 835 bp in vitro using overlapping PCR technique. The whole experiment design with two factors and six levels (36 combinations) was applied to study the effects of the vector density and competent cells on the macromolecular vector transformation efficiency. Based on the 1 500 ng target gene, the vector density grades were designed (50, 100, 150, 200, 250, 300 ng), and then the recombinant plasmids were transformed into Top10F<sup>+</sup>, DH5, Stbl3, Epi400, JM108, SCS1. Results showed that the positive cloning rates of different vector amount from big to small were in the order of 200,250,300,150,100 and 50 ng. The maximum positive cloning rate of 200 ng reached 75%; and the average value was 28. 5%. The positive cloning rates of different competent cells from big to small were in the order of stbl3, Top10F<sup>+</sup>, DH5, JM108, Epi400 and SCS1. Stbl3 was higher than other competent cells under any vector density, and its average positive cloning rate was 42.4%. Both the vector density and competent cells had significant effects on the macromolecular vector transformation efficiency. The optimal combination was C4 with 200 ng vector density and Stbl3, the positive cloning rate of which could reach 75%.

**Key words** Macromolecular vector, Vector density, Competence, Transformation efficiency

## 1 Introduction

The plasmid vector is the small DNA molecule having autonomous replication ability in cells, and the most common recombinant DNA carrier in genetic engineering. It can use the recombinant DNA technology to transport the purposefully strong DNA fragments into recipient cells for propagation and expression<sup>[1-7]</sup>. The DNA length of plasmid vector varies from a few thousand base pairs to hundreds of thousands of base pairs. Compared with the small molecular weight vector, the macromolecular vector has low extracellular recombination and connection efficiency, low transformation efficiency in competence, and low replication rate in the host cells, so it is easy to produce random mutation and deletion of target DNA and unstable expression in the passage<sup>[8-10]</sup>. Therefore, it needs an optimized recombination and connection system and an optimized competence transformation system, so as to improve the recombination and transformation efficiency of macromolecular vector<sup>[11-13]</sup>. In this study, the PCR (Polymerase Chain Reaction) technology was used to design a 835bp target gene synthesized from 26 primers, and this gene was connected with about 20 kb macromolecular vector pCMV5-ClaI-RyR1\_WT to form the recombinant plasmid. On the basis of 1500 ng target gene substrate, the recombinant reaction of 6-gradient vector amount (50 ng, 100 ng, 150 ng, 200 ng, 250 ng and 300 ng) was carried out, and the recombinant reaction products obtained were transformed into 6 different kinds of competence (Top10F<sup>+</sup>, DH5 $\alpha$ , St-


bl3, Epi400, JM108, SCS1) to form 36 different combinations. Then the rate of positive clones from the 36 different combinations was analyzed, respectively, in order to get the optimal vector amount and best competence in the recombinant cloning of macromolecular vector.

## 2 Materials and methods

### 2.1 Materials

**2.1.1 Target gene.** In this study, we selected one 835 bp target gene (Fig. 1). The overall GC content was 56.17%, it was not the gene with high GC content, and there was slight fluctuation in the GC content. Overall, it was easy to be amplified (Fig. 2). The target gene had no significant direct, inverted repeat, or auto-reduplication, without difficulty in primer design.

GGGGCTGCTGACCTGGCTCATGTCCATCGATGTCAAGTACCAAGTCTGGAAGTTCGGGGTTCATCTTCACGGACAACCTCGTTCTGTAACCTGGGCTGGTATGATGTCCCTCTGGGCACTACAACTCTCTTTTGGCCGCCACTGCTGGACATGGCATGGGGTCAAGACGCTGCGTACCATCTCTCTCTGTCAACCCACAATGGGAAACAGCTGGTGATGACTGTGGGCTCTCTGGCCGTGCTGGTCTACTGTACATCTGGTGGCTTCAACTTCTCCGCAAGTCTACAACAAGAGCGAGGACGAGGACGAGCGGGACATGAAGTGGATGACATGATGACGTGCTACTGTTCACATGTACGTGGCGCTCCGGCTGGCGGAGGATCGGGGACGAGATCGAGGACCCAGCGGGCGATGAATACGAGCTCTACCGGGTGGTCTTCGACATCACTTCTTCTCTGTCATTGTGATCTCTGCTGGCCATCATCCAGGGTCTGATTATCGCCGCTTCGCGGAGCTCCGAGACGAGGACGAGTGAAGGAAGATATGGAGACCAATGCTTCATCTGCGGAGTGGCAGTGACTACTTCGATACCAACGCGGACGGCTTCGAGACCCACAGCTAGAGGAGGACCAATCTGGCCATTATGTTCTTCTTGATGATCTGATAAAACAGGACGAGACGAGACACGCGGACAGGAGTCTCTACGTCTGGAAGATGATACGAGAGAGGTGCTGGGACTTCTTCCCGCCGCGACTGCTTCCGACAGCATACGAGGACAGCTGAGCTGAGAAGCTTGTCATGCTGAGGTGACATAGAGGATCCGGGTGGCATCCCTGTGAC



Received: September 8, 2016 Accepted: November 18, 2016  
Supported by Teaching and Research Reform Project of Suzhou Industrial Park Institute of Services Outsourcing (JG-201601).  
\* Corresponding author. E-mail: qhn224@126.com



tection product were judged as follows: electrophoretic bands were matched with ladder; 1149 bp meant that the positive clones were denoted as 1; 412 bp meant that the empty vector was denoted as

0; the primer dimer meant that the variegated bacteria sample was denoted as 0.

Table 3 Transformation combination

Top10F <sup>+</sup>	DH5α	Stb13	Epi400	JM108	SCSI
A1	B1	C1	D1	E1	F1
A2	B2	C2	D2	E2	F2
A3	B3	C3	D3	E3	F3
A4	B4	C4	D4	E4	F4
A5	B5	C5	D5	E5	F5
A6	B6	C6	D6	E6	F6

3 Results

3.1 Synthesis of target gene As shown in Fig. 4, PCR products of the target gene were 750 – 1000 bp, consistent with the theoretical value of 855 bp.





Fig. 4 Electrophoretogram of PCR products

3.2 Linearization of vector As shown in Fig. 5, the theoretical size after enzyme digestion was 19043 bp and 98 bp, and the minimum value of 5000 Ladder was 100 bp, so 98 bp could not be displayed in the figure. The enzyme digestion result was consistent with the theoretical value.



Note: Band 1, 2, 3 represented 5000 bp Ladder, enzyme digestion sample and 10 kb Ladder, respectively.

Fig. 5 Electrophoretogram of vector enzyme digestion

3.3 Growth of transformed colonies As shown in Fig. 6, in 6 kinds of competence, there were a total of 36 plates (A<sub>1</sub>-F<sub>6</sub>), and the colonies with vector amount of 50, 100, 150, 200, 250 and 300 ng were added, respectively. It was clearly found that the colonies on the 36 plates grew well.





Fig. 6 Colony growth in 6 kinds of competence

**3.4 Rate of positive clones** From Table 4 and Fig. 7, it was found that different vector density had a significant impact on the rate of positive clones of macromolecular vectors, and in terms of the average rate of positive clones, 200 ng > 250 ng > 300 ng > 150 ng > 100 ng > 50 ng. In all kinds of competence, the rate of positive clones was highest at 200 ng, the highest rate was up to 75%, and the average rate reached 28.5%. If the vector density was less than 100 ng, the transformation efficiency was less than 10%. Different

competence had a significant effect on the rate of positive clones of macromolecular vector, and the average rate of positive clones was in the order of stb3 > Top10F<sup>+</sup> > DH5 $\alpha$  > Jm108 > Epi400 > SCS1. The best competence was stb3, and its rate of positive clones was higher than that of other kinds of competence under any vector density, and the average rate of positive clones was 42.4%. The average rate of positive clones was below 10% for Jm108, Epi400 and SCS1.

**Table 4** The rate of positive clones

Concentration/ng	A (Top10F <sup>+</sup> )	B (DH5 $\alpha$ )	C (Stb3)	D (Epi400)	E (JM108)	F (SCS1)	Average
50	0.0	0.0	4.2	0.0	0.0	0.0	0.7
100	4.2	4.2	12.5	0.0	4.2	0.0	4.2
150	8.3	12.5	41.7	4.2	8.3	4.2	13.2
200	33.3	29.2	75.0	8.3	16.7	8.3	28.5
250	29.2	20.8	66.7	4.2	8.3	4.2	22.2
300	20.8	12.5	58.3	0.0	4.2	4.2	16.7
Average	16.0	13.2	42.4	2.8	6.9	3.5	—



**Fig. 7** The rate of positive clones

## 4 Conclusions and discussions

In this study, for the 20 kb macromolecular vector, on the basis of 1500 ng target gene substrate, we set the vector amount with six gradients (50 ng, 100 ng, 150 ng, 200 ng, 250 ng, 300 ng), and the recombinant reaction products obtained were transformed into six different kinds of competence (Top10F<sup>+</sup>, DH5 $\alpha$ , Stb3, Epi400, JM108, SCS1) to form 36 test combinations. The results showed that both the vector amount and competence significantly affected the rate of positive clones. When the vector amount in the reaction system was too low, the recombinant reaction could not get enough positive clones, and the screening efficiency was very low. When the vector amount in the reaction system was too high, many empty vectors might be selected, affecting the rate of positive clones. For the 20 kb vector, when the vector molecular weight was about 200 ng, the rate of positive clones was highest. Compared with other kinds of competence, Stb3 had a greater ability to copy large vectors. The optimal combination was 200 ng vector density and Stb3, and the rate of positive clones could reach 75%. For the common recombinant test, the vector sample amount was far less than 200 ng, and taking the common puc57 series vector for example, the vector sample amount for recombination reaction was only about 30 ng<sup>[1, 3, 4]</sup>. When there was a large vector in the laboratory, the rate of positive clones might be low and even 0. Due to different genotypes for different competence, there were large differences in copying rate of different sizes and different

types of plasmid<sup>[14-18]</sup>. In this study, the vector used was close to 20 kb, and it was difficult to clone compared with the commonly used 3–4 kb vector. The macromolecular vector had features of low recombination efficiency, low transformation efficiency, low copying rate, unstable expression, difficult extraction, separation and purification<sup>[8-10]</sup>, so many domestic DNA companies were not willing to accept cloning of macromolecular vector, and even if there are corresponding services, the service price was high due to high failure rate and experimentation cost. In this study, with 20kb vector as example, we selected the vector amount with six gradients and six kinds of competence, indicating that the vector amount and competence indeed had a significant impact on the rate of positive clones. We obtained the optimal combination which can be used as reference for improving the cloning efficiency of macromolecular vector.

## References

- [1] ZHONG X, ZHAI C, CHEN L, *et al.* Construction of directional T vector for gene cloning and expression[J]. Chinese Journal of Biotechnology, 2013, 29(4): 510–519. (in Chinese).
- [2] YU Y, JIANG SC, WANG KY, *et al.* Research progress of large fragment DNA cloning vector[J]. Heilongjiang Agricultural Science, 2015(2): 147–151. (in Chinese).
- [3] WU YF, LIANG DC, GUO G, *et al.* Construction of pUC-T vector[J]. Tianjin Medical Journal, 2005, 33(3): 159–160. (in Chinese).
- [4] LI J, ZHANG JF, ZHEN YL, *et al.* Construction of recombinant plasmid pCMV-Myc-PIASx and its protein expression[J]. Journal of Jilin University: Med Ed, 2009, 35(3): 415–418. (in Chinese).
- [5] YAN F, ZHAO XY, DENG HX, *et al.* Construction and expression of a novel bisbicistronic expression vector: pCMV-Myc-IRES-EGFP[J]. Chinese Journal of Biotechnology, 2007, 23(3): 423–428. (in Chinese).
- [6] WANG HZ, ZHOU XY, SONG ZX, *et al.* Degenerate PCR and its application in gene cloning[J]. Hereditas, 2003, 25(2): 201–204. (in Chinese).
- [7] ZHENG BQ, WANG Y, PENG ZH, *et al.* Cloning of ACC oxidase gene from Cattleya flower and construction of its plant antisense expression vector[J]. Acta Agriculturae Nucleatae Sinica, 2009, 23(3): 442–446. (in Chinese).

cause behind the big data, there is the privacy of the group information. The cloud computing concerns the network security, and the standard for implementation of the platform economy is the high degree of customer information security. Mobile Internet also imposes higher requirements on the network carrier, of which some parts need establishing support system of intellectual property right, to guarantee the security of operating environment. On the other hand, as maker groups, their innovative and creative inspiration has characteristic of originality. At present, there is serious problem of plagiarism at current domestic market, which poses a great challenge for protection of intellectual property rights. Therefore, at the time of building "Maker" urban culture in Shanghai, it is required to firstly make innovation on the original systems, in other words, the institution should go first.

## References

[illegible]

( From page 84 )

[illegible]

## About KIT

The Royal Tropical Institute (KIT) in Amsterdam is an independent centre of knowledge and expertise in the areas of international and intercultural cooperation, operating at the interface between theory and practice and between policy and implementation. The Institute contributes to sustainable development, poverty alleviation and cultural preservation and exchange.