



**AgEcon** SEARCH  
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search  
<http://ageconsearch.umn.edu>  
[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

# THE STATA JOURNAL

## Editors

H. JOSEPH NEWTON  
Department of Statistics  
Texas A&M University  
College Station, Texas  
editors@stata-journal.com

NICHOLAS J. COX  
Department of Geography  
Durham University  
Durham, UK  
editors@stata-journal.com

## Associate Editors

CHRISTOPHER F. BAUM, Boston College  
NATHANIEL BECK, New York University  
RINO BELLOCCO, Karolinska Institutet, Sweden, and  
University of Milano-Bicocca, Italy  
MAARTEN L. BUIS, WZB, Germany  
A. COLIN CAMERON, University of California–Davis  
MARIO A. CLEVES, University of Arkansas for  
Medical Sciences  
WILLIAM D. DUPONT, Vanderbilt University  
PHILIP ENDER, University of California–Los Angeles  
DAVID EPSTEIN, Columbia University  
ALLAN GREGORY, Queen’s University  
JAMES HARDIN, University of South Carolina  
BEN JANN, University of Bern, Switzerland  
STEPHEN JENKINS, London School of Economics and  
Political Science  
ULRICH KOHLER, University of Potsdam, Germany

FRAUKE KREUTER, Univ. of Maryland–College Park  
PETER A. LACHENBRUCH, Oregon State University  
JENS LAURITSEN, Odense University Hospital  
STANLEY LEMESHOW, Ohio State University  
J. SCOTT LONG, Indiana University  
ROGER NEWSON, Imperial College, London  
AUSTIN NICHOLS, Urban Institute, Washington DC  
MARCELLO PAGANO, Harvard School of Public Health  
SOPHIA RABE-HESKETH, Univ. of California–Berkeley  
J. PATRICK ROYSTON, MRC Clinical Trials Unit,  
London  
PHILIP RYAN, University of Adelaide  
MARK E. SCHAFFER, Heriot-Watt Univ., Edinburgh  
JEROEN WEESIE, Utrecht University  
IAN WHITE, MRC Biostatistics Unit, Cambridge  
NICHOLAS J. G. WINTER, University of Virginia  
JEFFREY WOOLDRIDGE, Michigan State University

## Stata Press Editorial Manager

LISA GILMORE

## Stata Press Copy Editors

DAVID CULWELL and DEIRDRE SKAGGS

The *Stata Journal* publishes reviewed papers together with shorter notes or comments, regular columns, book reviews, and other material of interest to Stata users. Examples of the types of papers include 1) expository papers that link the use of Stata commands or programs to associated principles, such as those that will serve as tutorials for users first encountering a new field of statistics or a major new technique; 2) papers that go “beyond the Stata manual” in explaining key features or uses of Stata that are of interest to intermediate or advanced users of Stata; 3) papers that discuss new commands or Stata programs of interest either to a wide spectrum of users (e.g., in data management or graphics) or to some large segment of Stata users (e.g., in survey statistics, survival analysis, panel analysis, or limited dependent variable modeling); 4) papers analyzing the statistical properties of new or existing estimators and tests in Stata; 5) papers that could be of interest or usefulness to researchers, especially in fields that are of practical importance but are not often included in texts or other journals, such as the use of Stata in managing datasets, especially large datasets, with advice from hard-won experience; and 6) papers of interest to those who teach, including Stata with topics such as extended examples of techniques and interpretation of results, simulations of statistical concepts, and overviews of subject areas.

The *Stata Journal* is indexed and abstracted by *CompuMath Citation Index*, *Current Contents/Social and Behavioral Sciences*, *RePEc: Research Papers in Economics*, *Science Citation Index Expanded* (also known as *SciSearch*), *Scopus*, and *Social Sciences Citation Index*.

For more information on the *Stata Journal*, including information for authors, see the webpage

<http://www.stata-journal.com>

Subscriptions are available from StataCorp, 4905 Lakeway Drive, College Station, Texas 77845, telephone 979-696-4600 or 800-STATA-PC, fax 979-696-4601, or online at

<http://www.stata.com/bookstore/sj.html>

Subscription rates listed below include both a printed and an electronic copy unless otherwise mentioned.

| U.S. and Canada                   |       | Elsewhere                         |       |
|-----------------------------------|-------|-----------------------------------|-------|
| <b>Printed &amp; electronic</b>   |       | <b>Printed &amp; electronic</b>   |       |
| 1-year subscription               | \$ 98 | 1-year subscription               | \$138 |
| 2-year subscription               | \$165 | 2-year subscription               | \$245 |
| 3-year subscription               | \$225 | 3-year subscription               | \$345 |
| 1-year student subscription       | \$ 75 | 1-year student subscription       | \$ 99 |
| 1-year institutional subscription | \$245 | 1-year institutional subscription | \$285 |
| 2-year institutional subscription | \$445 | 2-year institutional subscription | \$525 |
| 3-year institutional subscription | \$645 | 3-year institutional subscription | \$765 |
| <b>Electronic only</b>            |       | <b>Electronic only</b>            |       |
| 1-year subscription               | \$ 75 | 1-year subscription               | \$ 75 |
| 2-year subscription               | \$125 | 2-year subscription               | \$125 |
| 3-year subscription               | \$165 | 3-year subscription               | \$165 |
| 1-year student subscription       | \$ 45 | 1-year student subscription       | \$ 45 |

Back issues of the *Stata Journal* may be ordered online at

<http://www.stata.com/bookstore/sjj.html>

Individual articles three or more years old may be accessed online without charge. More recent articles may be ordered online.

<http://www.stata-journal.com/archives.html>

The *Stata Journal* is published quarterly by the Stata Press, College Station, Texas, USA.

Address changes should be sent to the *Stata Journal*, StataCorp, 4905 Lakeway Drive, College Station, TX 77845, USA, or emailed to [sj@stata.com](mailto:sj@stata.com).



Copyright © 2014 by StataCorp LP

**Copyright Statement:** The *Stata Journal* and the contents of the supporting files (programs, datasets, and help files) are copyright © by StataCorp LP. The contents of the supporting files (programs, datasets, and help files) may be copied or reproduced by any means whatsoever, in whole or in part, as long as any copy or reproduction includes attribution to both (1) the author and (2) the *Stata Journal*.

The articles appearing in the *Stata Journal* may be copied or reproduced as printed copies, in whole or in part, as long as any copy or reproduction includes attribution to both (1) the author and (2) the *Stata Journal*.

Written permission must be obtained from StataCorp if you wish to make electronic copies of the insertions. This precludes placing electronic copies of the *Stata Journal*, in whole or in part, on publicly accessible websites, fileservers, or other locations where the copy may be accessed by anyone other than the subscriber.

Users of any of the software, ideas, data, or other materials published in the *Stata Journal* or the supporting files understand that such use is made without warranty of any kind, by either the *Stata Journal*, the author, or StataCorp. In particular, there is no warranty of fitness of purpose or merchantability, nor for special, incidental, or consequential damages such as loss of profits. The purpose of the *Stata Journal* is to promote free communication among Stata users.

The *Stata Journal* (ISSN 1536-867X) is a publication of Stata Press. Stata, **STATA**, Stata Press, Mata, **MATA**, and NetCourse are registered trademarks of StataCorp LP.

## Stata tip 116: Where did my $p$ -values go? (Part 3)

Maarten L. Buis  
Wissenschaftszentrum Berlin für Sozialforschung (WZB)  
Berlin, Germany  
maarten.buis@wzb.eu

In a previous Stata tip (Buis 2007), I discussed how to recover  $t$  statistics,  $p$ -values, and confidence intervals for regression parameters by using the results that are returned by an estimation command. In a subsequent Stata tip (Buis 2011), I discussed how to recover parameter estimates for parameters that were estimated on a transformed scale. For example, if a likelihood function contains a standard deviation or a correlation, then many Stata commands will maximize the likelihood with respect to  $\ln(\text{standard deviation})$  and the Fisher's  $z$  transformation of the correlation. In this tip, I will discuss how to recover the standard errors for the back-transformed parameters, that is, the standard errors of the standard deviation and the correlation.

Stata often displays the back-transformed parameters and their standard errors, but it leaves behind only the estimates of the transformed parameters and their standard errors. In those cases, the delta method (for example, Feiveson 2005) was used to compute the standard errors of the back-transformed parameters. In its simplest form, the delta method means that if we apply a transformation  $G(\cdot)$  to a parameter  $b$ , then we can approximate the standard error of the transformed parameter as

$$\text{se}\{G(b)\} \approx \text{se}(b) \times G'(\hat{b})$$

where  $G'(\hat{b})$  is the first derivative of  $G(b)$  with respect to  $b$  evaluated at  $\hat{b}$ . If Stata returned  $\ln(\text{standard deviation})$  and we wanted the standard deviation and its standard error, then  $G(b) = \exp(b)$  and  $G'(\hat{b}) = \exp(\hat{b})$ . If Stata returned Fisher's  $z$  transformation of a correlation and we wanted the correlation and its standard error, then  $G(b) = \tanh(b)$  and  $G'(\hat{b}) = \cosh(\hat{b})^{-2}$ . This is illustrated below using a model estimated with `heckman` (see [R] `heckman`). This model was chosen because it returns transformed parameters of both types.

```
. use http://www.stata-press.com/data/r13/womenwk
. heckman wage educ, select(married children educ) nolog
Heckman selection model                Number of obs    =    2000
(regression model with sample selection) Censored obs      =     657
                                          Uncensored obs    =    1343
                                          Wald chi2(1)      =    403.39
Log likelihood = -5250.348              Prob > chi2       =     0.0000
```

| wage          | Coef.     | Std. Err. | z      | P> z  | [95% Conf. Interval] |          |
|---------------|-----------|-----------|--------|-------|----------------------|----------|
| <b>wage</b>   |           |           |        |       |                      |          |
| education     | 1.099506  | .0547435  | 20.08  | 0.000 | .9922102             | 1.206801 |
| _cons         | 7.042147  | .8423253  | 8.36   | 0.000 | 5.39122              | 8.693074 |
| <b>select</b> |           |           |        |       |                      |          |
| married       | .5420304  | .0657798  | 8.24   | 0.000 | .4131044             | .6709564 |
| children      | .4409418  | .0276093  | 15.97  | 0.000 | .3868286             | .495055  |
| education     | .0722993  | .0105096  | 6.88   | 0.000 | .0517007             | .0928978 |
| _cons         | -1.473038 | .1465476  | -10.05 | 0.000 | -1.760266            | -1.18581 |
| /athrho       | .8081049  | .1108545  | 7.29   | 0.000 | .5908341             | 1.025376 |
| /lnsigma      | 1.807547  | .0291035  | 62.11  | 0.000 | 1.750506             | 1.864589 |
| rho           | .6685435  | .061308   |        |       | .5304953             | .772047  |
| sigma         | 6.095479  | .1773995  |        |       | 5.757513             | 6.453283 |
| lambda        | 4.075093  | .4690025  |        |       | 3.155865             | 4.994321 |

```
LR test of indep. eqns. (rho = 0):  chi2(1) = 47.02  Prob > chi2 = 0.0000
. tempname gprime
. scalar `gprime' = exp(_b[lnsigma:_cons])
. display "se of sigma = " _se[lnsigma:_cons]*`gprime'
se of sigma = .17739954
. scalar `gprime' = cosh(_b[athrho:_cons])^-2
. display "se of rho = " _se[athrho:_cons]*`gprime'
se of rho = .06130802
```

Alternatively, one can use `nlcom` (see [R] `nlcom`) to compute these standard errors, as follows:

```
. nlcom (rho: tanh(_b[athrho:_cons])) (sigma: exp(_b[lnsigma:_cons]))
      rho:  tanh(  _b[athrho:_cons]  )
      sigma: exp(  _b[lnsigma:_cons]  )
```

| wage  | Coef.    | Std. Err. | z     | P> z  | [95% Conf. Interval] |          |
|-------|----------|-----------|-------|-------|----------------------|----------|
| rho   | .6685435 | .061308   | 10.90 | 0.000 | .548382              | .788705  |
| sigma | 6.095479 | .1773995  | 34.36 | 0.000 | 5.747782             | 6.443175 |

Notice that the confidence intervals do not correspond with those in the output of `heckman`. This is because `heckman` first computes the bounds of the confidence intervals for the transformed parameters and then back-transforms those bounds to the original metric, while `nlcom` uses the standard errors for the back-transformed parameters for computing these bounds. In most cases, computing the bounds on the transformed scale

and then back-transforming those bounds to the original scale results in somewhat better bounds. This is because the sampling distribution of the transformed parameters is likely to be better approximated by a normal distribution than is the sampling distribution of the back-transformed parameters. For more discussion, see Sribney and Wiggins (2009). You can use the techniques discussed in Buis (2007) to recover the confidence intervals reported by `heckman`.

```
. display "confidence interval for rho: ["
> tanh( _b[athrho:_cons] - invnormal(.975)*_se[athrho:_cons] ) ", "
> tanh( _b[athrho:_cons] + invnormal(.975)*_se[athrho:_cons] ) "]"
confidence interval for rho: [.53049526, .77204703]

. display "confidence interval for sigma: ["
> exp( _b[lnsigma:_cons] - invnormal(.975)*_se[lnsigma:_cons] ) ", "
> exp( _b[lnsigma:_cons] + invnormal(.975)*_se[lnsigma:_cons] ) "]"
confidence interval for sigma: [5.7575126, 6.4532831]
```

Also note that `nlcom` returns the  $z$  statistic and  $p$ -value for the test of the null hypothesis that the standard deviation and the correlation are 0, which were not reported by `heckman`. This test is problematic in the case of the standard deviation, because this is a test “on the boundary of the parameter space”. A standard deviation can only take values larger than or equal to 0. Thus the hypothesis that the standard deviation is equal to 0 is on the boundary of the possible values for the standard deviation, and standard tests do not tend to behave well in this extreme area (for example, Gutierrez, Carter, and Drukker [2001]).

## References

- Buis, M. L. 2007. Stata tip 53: Where did my  $p$ -values go? *Stata Journal* 7: 584–586.
- . 2011. Stata tip 97: Getting at  $\rho$ 's and  $\sigma$ 's. *Stata Journal* 11: 315–317.
- Feiveson, A. H. 2005. FAQ: What is the delta method and how is it used to estimate the standard error of a transformed parameter?  
<http://www.stata.com/support/faqs/statistics/delta-method/>.
- Gutierrez, R. G., S. Carter, and D. M. Drukker. 2001. sg160: On boundary-value likelihood-ratio tests. *Stata Technical Bulletin* 60: 15–18. Reprinted in *Stata Technical Bulletin Reprints*, vol. 10, pp. 269–273. College Station, TX: Stata Press.
- Sribney, W., and V. Wiggins. 2009. FAQ: Standard errors, confidence intervals, and significance tests for ORs, HRs, IRRs, and RRRs.  
<http://www.stata.com/support/faqs/statistics/delta-rule/>.