



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*



**CARIBBEAN
FOOD
CROPS SOCIETY**

22

**Twenty Second
Annual Meeting 1986**

St. Lucia

Vol. XXII

STUDIES ON THE PRODUCTION AND UTILIZATION OF OILSEEDS IN BELIZE

B.K. Rai and A.K. Sinha

CARDI
Belmopan, Belize

ABSTRACT

To conserve foreign exchange and increase income of farmers, studies for oilseed production and utilization were started in 1981. The limitations to annual crop production were: high rainfall during crop growth especially at harvesttime for the June planted crop; difficulty of land preparation and less and erratic rainfall for the November to January planted crop. These difficulties have been overcome to a great extent by forming land into raised beds, minimum tillage technology and selection of crops and their varieties. Soybean varieties yielding up to 3.5 tons per ha have been identified. High yielding sesame varieties have been selected. Low erucic varieties of mustard have been tested.

Yield of sunflower was considered low for commercialization. To keep the capital and recurring expenses of oil mill low, and to get high protein and high energy feed, oilseeds should be mechanically pressed rather than solvent-extracted. Results of a trial indicated that broilers could be raised on whole fat soybean. Whole fat soybean is being mixed with corn by a tortilla factory, for improved human nutrition.

INTRODUCTION

Diversification of monoculture economies is of importance not only because of fluctuations in the price of the commodities exported, but also to provide increased income to farmers and to conserve foreign exchange by import substitution. Belize imports food worth \$45 million dollars per annum. Simultaneously, conventional crops like corn and bean tended to be over produced and could not be sold. Cultivation of oilseeds and extraction of oil would provide oils and fats for human consumption and feed for livestock. Oils and fats and feed were imported at a cost of about \$9 million annually (Table 1). Further, additional amounts of proteinaceous feed would be required for shrimp culture for export and to produce milk and milk products and meat, imported to the tune of \$14 million annually. Thus, cultivation of oilseeds is pivotal for improvement of an agrarian economy. With the increased use of broilers, globally, the vegetable oils have become a by-product of the feed industry. Thus quality of feed produced must be considered while selecting oilseed crops.

Existing vegetable oil processing facilities in the CARICOM region were importing more and more soybean, to keep the plants running and to be competitive with the extra-regional oil as the regional price of copra was 2-3 times the world price. CARICOM countries imported soybean, and soybean oil and cake worth \$44 million annually (Table 2). Further, other vegetable oils and ready made feed were imported into the CARICOM countries.

Table 1. Quantities and value of edible oils and fats and feed imported into Belize

| Item | 1982 | | 1983 | | 1984 | | 1985 | |
|--------------------------------------|-----------------|-------|-------|-------|-------|--------|-------|-------|
| | a ^{1/} | b | a | b | a | b | a | b |
| A. Oils and Fats | | | | | | | | |
| Lard | 1,384 | 3,182 | 1,801 | 3,412 | 1,570 | 3,680 | 1,291 | 2,774 |
| Margarine | 487 | 1,371 | 376 | 1,127 | 379 | 1,236 | 467 | 1,400 |
| Oils | 248 | 310 | 486 | 417 | 89 | 343 | 93 | 341 |
| Sub total | 2,119 | 4,863 | 2,663 | 4,956 | 2,038 | 5,529 | 1,851 | 4,515 |
| B. Feed (Protein Concentrate) | | | | | | | | |
| For poultry | 3,580 | 2,249 | 4,401 | 3,977 | 4,226 | 4,450 | 4,552 | 3,186 |
| For cattle | 193 | 107 | 204 | 176 | 371 | 273 | 791 | 284 |
| For pigs | 16 | 11 | 58 | 44 | 38 | 41 | 96 | 35 |
| Sub total | 3,789 | 3,367 | 4,663 | 4,197 | 4,635 | 4,764 | 5,439 | 3,505 |
| Grand total | | 8,230 | | 9,153 | | 10,023 | | 8,020 |

^{1/} a = Quantity in tons; b = Belize dollars in thousands.

Table 2. Import of soybean and soybean products into CARICOM countries^{1/}

| Country | Quantity (MT) | | | Value (US \$'000) | | | Value (US\$/MT) |
|---------------------|---------------|---------|---------|-------------------|--------|--------|-----------------|
| | 1982 | 1983 | 1984 | 1982 | 1983 | 1984 | 1984 |
| Soybean | | | | | | | |
| Jamaica | 54,859 | 94,194 | 50,813 | 14,455 | 19,577 | 15,312 | 301 |
| Soybean cake | | | | | | | |
| Barbados | 10,276 | 10,000 | 8,500 | 3,052 | 3,100 | 2,550 | 300 |
| Trinidad | 42,058 | 30,402 | 33,901 | 16,466 | 8,903 | 8,228 | 243 |
| Guyana | 5,303 | 3,000 | 4,100 | 2,968 | 1,800 | 2,400 | 585 |
| Soybean oil | | | | | | | |
| Barbados | 3,681 | 2,600 | 1,500 | 1,568 | 1,800 | 1,050 | 700 |
| Jamaica | 2,778 | 3,549 | 4,962 | 2,091 | 2,853 | 4,393 | 885 |
| St. Lucia | 519 | 8 | | 350 | 7 | | |
| Trinidad, etc. | 7,603 | 7,811 | 8,196 | 5,169 | 5,678 | 7,549 | 921 |
| Total | 127,077 | 151,609 | 111,972 | 46,119 | 43,718 | 41,482 | |

^{1/} Source: FAO Trade Year Book, 1984.

Table 3. Rainfall data for various districts

| Month | Corozal | | | | o/Walk | | | | Belize | | | | Cayo | | | | Stann Creek | | | | Toledo | | | |
|-----------|------------|----|-----------|----|--------|----|---------|-----|---------------|-----|---------|-----|--------|-----|------------|---|-------------|---|---|---|--------|---|---|---|
| | San Andres | | San Pablo | | San | | Lemonal | | Roaring Creek | | Ignacio | | Ponona | | Middle Sex | | P | | G | | | | | |
| | a | b | a | b | a | b | a | b | a | b | a | b | a | b | a | b | a | b | a | b | a | b | a | b |
| January | 64 | 7 | 43 | 7 | 78 | 6 | 101 | 9 | 107 | 10 | 150 | 13 | 167 | 15 | 227 | | | | | | | | | |
| February | 33 | 5 | 64 | 3 | 41 | 3 | 78 | 5 | 50 | 6 | 86 | 9 | 89 | 9 | 129 | | | | | | | | | |
| March | 18 | 2 | 20 | 2 | 39 | 2 | 56 | 3 | 40 | 5 | 62 | 6 | 58 | 6 | 99 | | | | | | | | | |
| April | 41 | 3 | 25 | 2 | 51 | 2 | 51 | 3 | 39 | 4 | 86 | 6 | 75 | 6 | 99 | | | | | | | | | |
| May | 122 | 5 | 80 | 4 | 104 | 4 | 79 | 4 | 104 | 6 | 152 | 10 | 176 | 10 | 249 | | | | | | | | | |
| June | 232 | 10 | 242 | 9 | 264 | 9 | 221 | 12 | 199 | 12 | 320 | 18 | 425 | 18 | 619 | | | | | | | | | |
| July | 174 | 11 | 63 | 4 | 193 | 9 | 256 | 14 | 186 | 15 | 332 | 20 | 435 | 24 | 710 | | | | | | | | | |
| August | 150 | 10 | 135 | 6 | 210 | 11 | 201 | 12 | 136 | 11 | 257 | 19 | 316 | 20 | 633 | | | | | | | | | |
| September | 237 | 12 | 181 | 10 | 275 | 12 | 220 | 12 | 175 | 14 | 339 | 19 | 369 | 18 | 567 | | | | | | | | | |
| October | 164 | 10 | 179 | 9 | 189 | 9 | 228 | 13 | 182 | 13 | 314 | 17 | 322 | 17 | 363 | | | | | | | | | |
| November | 100 | 7 | 46 | 5 | 125 | 8 | 162 | 10 | 148 | 11 | 216 | 14 | 200 | 15 | 256 | | | | | | | | | |
| December | 71 | 6 | 71 | 6 | 151 | 8 | 169 | 9 | 139 | 11 | 177 | 13 | 190 | 15 | 229 | | | | | | | | | |
| Total | 1,406 | 88 | 1,149 | 67 | 1,720 | 83 | 1,822 | 106 | 1,505 | 118 | 2,491 | 164 | 2,822 | 173 | 4,140 | | | | | | | | | |

1/ a = Rainfall in mm. b = Number of days with more than 1 mm rainfall.

CLIMATIC CONDITIONS IN BELIZE FOR OILSEED PRODUCTION

Because of the rainfall pattern, about 11,500 and 5,400 ha are planted to annual crops in June and in November to December, respectively. Thus 53 per cent of the area planted in June remains fallow during the second planting season. The weed load on fallow land increases and cultivation of even one crop per year becomes difficult and expensive.

June Planting Season

At the beginning of the rainy season it is possible to prepare land and plant crops in the first half of June, in the Corozal, Orange Walk, Belize and Cayo Districts. In the Stann Creek District, June has more rainfall (Table 3), and land preparation and planting have to be done earlier. Similarly in the Toledo District, though in some years dry land preparation is not possible because of no dry period. June planted crops are not expected to experience shortage of water. On the other hand, the success of a crop depends on its ability to withstand high rainfall especially at harvest. A 120-day duration crop, planted in mid-June, should be ready for harvest in mid-October. Because of the amount of rainfall and number of rainy days in October, the best chances of successful harvest of a 120 days crop are in the Corozal and Orange Walk districts, followed by Belize and Cayo, and the least in the Stann Creek and Toledo districts.

November to January Planting Season

Rainfall data for San Ignacio, Cayo District, for 1966 to 1979 were subjected to pentagonal probabilities to determine the chances of land preparation. When any five days received less than 2.5 cm rainfall, that period was considered dry. A period of five dry days preceded by another five dry days was considered suitable for land preparation. The probabilities thus worked out are presented in Table 4.

Table 4. Probability of land preparation in Cayo

| Dates | November | December | January |
|---------|----------|----------|---------|
| 1 - 5 | 0.2449 | 0.4082 | xxx |
| 5 - 10 | 0.2143 | 0.1224 | 0.1837 |
| 11 - 15 | 0.3367 | 0.1633 | 0.2755 |
| 16 - 20 | 0.1224 | 0.3061 | 0.4490 |
| 21 - 25 | 0.2006 | 0.4490 | 0.5510 |
| 26 - 30 | 0.3214 | 0.7857 | 0.6122 |
| Mean | 0.2401 | 0.3727 | 0.4143 |

The probability of land preparation was low in November and early December, but was good from December 16 to December 30, and again from January 16. Corozal, Orange Walk and Belize Districts received less while Stann Creek and Toledo Districts received more rainfall than Cayo and thus chances of land preparation would either be more or less, respectively, over the same period.

The data on the amount of rainfall received by various growth stages of a 95 day soybean crop, presented in Table 5, indicated that, as planting is delayed from November to January, the crop receives less rainfall. Based on the water requirement of various crops, the time of planting can be determined. It should be mentioned that rainfall from November onwards is erratic. For assured success of crops irrigation is necessary.

Average monthly maximum temperatures from Cayo District from January to December were 28, 29, 30, 31, 32, 31, 30, 30, 30, 29, 28, and 30°C, and average monthly minimum were 17, 18, 18, 20, 20, 22, 21, 21, 21, 18, 18, and 20°C. The "absolute" minimum temperatures recorded in January, February and March were 4, 4, and 8°C. Similar temperature conditions prevail in all the other districts. These data suggested that some semi-tropical crops could be planted in November and December.

Thus, major limitations to the production of annual crops are high rainfall during crop growth and at harvesttime for the June planted crop and difficulties of land preparation, and less and erratic rainfall for November to January planted crops. A system of farming on raised beds developed at ICRISAT, India, using animal power, has been adapted for tractor use at CARDI's Field Station, Belmopan, Belize. For the last few years, crops of corn, soybean, peanut and some vegetables have been successfully raised on the beds by planting in June.

Manual tools and specialized harvesting equipment are needed to harvest crops in October so that the beds are not damaged. Crops are planted from November to January, on the same beds without land preparation, using minimum tillage. Crops that have been successfully raised on beds from November to January are: corn, soybean, peanut, sesame, chickpea, dry pea, pigeon pea, sunflower, mustard, and some vegetables.

OILSEED CROPS AND THEIR VARIETIES

Since 1981, work has been in progress on oilseed crops such as soybean (*Glycine max*), sesame (*Sesamum indicum*), rape and mustard (*Brassica* spp.), sunflower (*Helianthus annuus*), peanut (*Arachis hypogaea*), safflower (*Carthamus tinctorius*), and castor (*Ricinus communis*). Five varieties of safflower failed to grow when planted in November and January. Safflower requires drier conditions than those prevailing in Belize, and work on safflower was suspended in 1982. Peanut production systems have been developed, and varieties with various characteristics have been identified, but the cost of production is high for use as oil and feed. Work on castor was initiated in 1986. Work on soybean, sesame, mustard and sunflower is discussed in this paper.

Soybean

Variety selection: From 1982 to the present, 188 varieties from INTSOY (USA), AVRDC (Taiwan), IITA (Nigeria) and from Colombia were test-planted in June and in November to December. Selection criteria included yield, plant and lowest pod height, and grain quality.

Based on previous tests, four varieties were tested in a thrice replicated trial planted in June, 1985. Establishment was uneven because of low

rainfall immediately after planting. IPB 213-81 yielded the most (3,586 kg ha⁻¹) but had 27 per cent discoloured grains (Table 5). PR 140 (11) yielded 2,384 kg per ha, with 12 per cent discoloured grains, but apparently healthy and discoloured grains germinated 97 and 65 per cent, respectively. Varieties AGS-29 and TAC 23-1385 yielded less than PR 140 (11), had higher percentages of discoloured grains and that germinated poorly. In 1984, apparently healthy and discoloured grains of AGS-29 were analyzed for aflatoxin but none was found. The effect of discolouration on quality of oil needs to be determined. As the quality of seed obtained from the June planted crop was poor, except for PR 140 (11), the effect of planting in November was tested. IPB 213-81, PR 140 (11) and IAC 73-1385, along with UFV-1, another variety selected for planting in June 1985, were planted in November, 1985. The data presented in Table 6 indicate that the plants were very short in height, and that per cent discoloured grains and seed germination ranged from 5 to 18 and 92 to 95, respectively. Yield per unit area could not be assessed as the plant stand was not uniform due to poor quality seed from the June planted crop. Four of these five varieties had been planted in November 1984 when yield varied from 1.6 to 2.1 ton per ha.

The seed harvested in March could be used for planting a commercial crop in June and a seed crop in November. Such a system would reduce expenses on cold storage of seed. Seeds of varieties IPB 213-81, PR 140 (11), AGS-29, and IAC 73-1185 were analyzed and the results were 19, 23, 18, and 21 per cent, respectively, of oil; and 37, 34, 40, and 37, respectively, of protein.

Twenty varieties were tested in a thrice replicated trial planted in November 1985. Ten varieties were rejected because of susceptibility to diseases, grain quality, lodging, shattering, green leaves at harvest stage or 10 percentile plant height less than 45 cm. There were no significant differences in yield of the ten varieties not rejected (Table 7). Longer duration and greater plant height have been considered important characters (Spehar et al., 1983). Greater height of lowest pod was an advantage for a crop to be combined.

Seed longevity: Seed longevity during storage, especially under high ambient temperatures and relative humidity (RH), is important for good establishment. The International Institute of Tropical Agriculture (IITA), Nigeria, has developed techniques to test varieties for this character and has developed varieties with good seed longevity. Dr. E. A. Keuneman, during a visit to CARDI-Belize, suggested that while testing varieties for adaptability, seed longevity should also be considered. Dried seeds of 15 varieties harvested in February/March were exposed to 40°C and 60 per cent RH in a cabinet. At various time intervals, 100 seeds of each variety were planted to determine seed viability. Corrected per cent viability was calculated using the modified Abbot's formula. Variety TGX 814-230 developed at IITA was the most viable (Table 8).

Time of Planting: Data presented in Table 6 indicate that the same varieties planted in June yield more than when planted in November. Further, yields decreased on planting delayed from November to January (Table 9).

Table 5. Rainfall received by various growth stages of a 95-day duration crop when planted at various times

| District | Date of Planting | Amount of rainfall (mm) | | | |
|----------------------------------|------------------|---------------------------|-------------------------------------|---------------------------------|-----------------------------------|
| | | Seedling (0-10 dys) | Vegetative growth (11-50 dys) | Pod formation (51-80 dys) | Grain hardening (81-95 dys) |
| Corozal | Nov. 1 | 68 | 132 | 78 | 33 |
| | Dec. 1 | 48 | 107 | 51 | 17 |
| | Jan. 1 | 39 | 74 | 31 | 15 |
| Orange Walk | Nov. 1 | 62 | 120 | 85 | 38 |
| | Dec. 1 | 45 | 114 | 63 | 22 |
| | Jan. 1 | 42 | 90 | 36 | 16 |
| Big Falls, Belize District | Nov. 1 | 79 | 149 | 91 | 45 |
| | Dec. 1 | 61 | 119 | 79 | 38 |
| | Jan. 1 | 45 | 109 | 85 | 37 |
| San Ignacio, Cayo District | Nov. 1 | 81 | 185 | 115 | 47 |
| | Dec. 1 | 68 | 158 | 74 | 24 |
| | Jan. 1 | 67 | 107 | 44 | 20 |
| Stan Creek | Nov. 1 | 134 | 298 | 168 | 68 |
| | Dec. 1 | 108 | 235 | 109 | 37 |
| | Jan. 1 | 83 | 157 | 62 | 27 |

Table 6. Performance of varieties selected for planting in June

| Variety | No. plants ha ⁻¹ ('000) | No. days to harvest | Plant height (cm) | Yield (kg ha ⁻¹) | Percent dis- coloured grains | % germination | |
|-------------------------------------|------------------------------------|---------------------|-------------------|------------------------------|------------------------------|---------------|---------------|
| | | | | | | Heal- thy | Dis- coloured |
| <u>Planted on June 11, 1985</u> | | | | | | | |
| IPB 213-81 | 217 | 125 | 79 | 3,586 | 27 | 38 | 35 |
| PR 140(11) | 200 | 130 | 73 | 2,384 | 12 | 97 | 65 |
| AGS-29 | 242 | 98 | 69 | 2,115 | 20 | 97 | 30 |
| IAC 73-1385 | 115 | 135 | 75 | 1,892 | 50 | 48 | 18 |
| <u>Planted on November 20, 1985</u> | | | | | | | |
| IPB 213-81 | | | 26 | 2,169 ^{1/} | 18 | | 92 |
| PR 140(11) | | | 54 | 1,737 | 11 | | 93 |
| AGS-29 | | | 60 | 1,974 | 15 | | 92 |
| IAC 73-1385 | | | 34 | 1,643 | 13 | | 93 |
| UFV-1 | | | 28 | | 5 | | 95 |

^{1/} Yield data in this case were obtained from a November 1985 planting.

Table 7. Data on ten soybean varieties planted in November

| Days to harvest | Plant | Lowest pod | Lodging score | Percent grain discoloured | Variety | Yield (kg ha ⁻¹) |
|-----------------|--------|------------|---------------|---------------------------|---------------|------------------------------|
| 97 | 49-77 | 10-19 | 1 | 0 | TGX 814-230 | 2,159 |
| | 52-75 | 10-19 | 1 | 9 | Jupiter-R | 1,812 |
| 104 | 55-85 | 10-20 | 2 | 14 | M-98 | 2,045 |
| 110- | 65-75 | 10-20 | 3 | 11 | IPB 189-81 | 2,056 |
| 111 | 69-80 | 10-20 | 3 | 12 | IAC 73-5115 | 1,899 |
| | 65-90 | 10-19 | 2 | 17 | PR 21-35-2x4 | 2,078 |
| | 72-90 | 13-20 | 2 | 15 | ICAL-139 | 2,259 |
| | 66-100 | 13-25 | 2 | 13 | PR 30-38-3x8 | 1,918 |
| | 76-97 | 10-25 | 2 | 9 | PR 15-100-4B8 | 2,157 |
| | 70-105 | 15-24 | 1 | 9 | PR 30-38-3x3 | 1,926 |

Table 8. Comparative seed viability of soybean varieties exposed to high temperatures and relative humidity

| Variety | Viability of seed at 0 day exposure | Corrected per cent viability after | | | |
|-----------------------------|-------------------------------------|------------------------------------|----|----|----|
| | | No. of days of exposure | | | |
| | | 23 | 28 | 33 | 36 |
| Varieties November planting | | | | | |
| TGX 814-230 | 92 | 95 | 80 | 39 | 36 |
| M-98 | 92 | 93 | 69 | 15 | 18 |
| ICAL-139 | 96 | 64 | 72 | 12 | 10 |
| IAC 73-5115 | 88 | 59 | 45 | 4 | 12 |
| AGS-29 | 96 | 62 | 54 | 2 | 0 |
| PR 21-35-2x4 | 96 | 52 | 52 | 16 | 0 |
| Tulumayo-2 | 92 | 58 | 34 | 0 | 0 |
| PR 15-100-4B8 | 94 | 46 | 42 | 0 | 0 |
| TGX 239-520 | 100 | 32 | 24 | 0 | 0 |
| PR 30-38-3x3 | 92 | 28 | 15 | 0 | 0 |
| IPB 189-81 | 94 | 23 | 19 | 0 | 0 |
| PR 30-38-3x8 | 92 | 15 | 13 | 0 | 0 |
| Varieties for June planting | | | | | |
| IPB 213-81 | 88 | 68 | 59 | 9 | 8 |
| UFV-1 | 96 | 70 | 56 | 0 | 0 |
| PR-140(11) | 92 | 39 | 28 | 4 | 0 |
| IAC 73-1385 | 98 | 20 | 16 | 0 | 0 |

Plant spacing: A spacing trial indicated that at closer spacing there was a greater reduction in plant number from germination to harvest, the lowest pod height tended to increase, but there was no significant difference in yield (Table 10).

Fertilizer trials: Two rates of basal fertilizer applied at planting and three rates of fertilizer application at R₅ stage of growth were tested on three varieties. One variety (Baru) lodged, and data on the other two are presented in Table 11.

There was no significant difference between basal fertilizer application rates, nor did side dressing or foliar application of fertilizer increase yields over the control.

Sesame

Cultivation of sesame was the least capital intensive but was labour intensive, and is thus suited to the 5,000 or so slash and burn farmers in Belize. Some farmers already grow a few plants for use in confectionery. In 1983, a large number of varieties were imported from FAO, Mexico, Brazil, Panama, Nicaragua, Guatemala, Colombia and Venezuela, and some were collected locally. In all, 80 varieties were screened for suitability. The varieties found suitable, up to 1985, are Yuamar, Acarigua, Aceitera, Local 1, Chino Rojo, Phutil No. 1 and FAO 59.833 (Table 12). However, during multiplication of these varieties in 1986, Aceitera was observed to suffer from a physiological condition and was rejected. In a variety trial harvested in March 1986, the other varieties which have been observed to be promising are Glauca, Inamar and Venezuela 52.

Young sesame plants grow slowly and compete poorly with weeds. Successful production will require development of an effective herbicide treatment. Schroder and Newson (1984) tested pre- and post-emergence herbicides in a series of experiments in Queensland. Alachlor at 2.25 kg a.i. ha⁻¹ was the most effective treatment. Alachlor at three different rates was tested pre-emergence and its effectiveness observed up to 30 days after application. Control plots were hand weeded 16 days after planting. Three fixed 1 m² quadrats per plot, were counted for numbers of monocot and dicot weeds. Up to 30 days, weed size in treated plots was small.

Alachlor at 3.33 a.i. ha⁻¹ resulted in the least numbers of weeds (Table 13). Fatty acid composition of the Yuamar variety of sesame grown in Belize was determined using a New Port Analyzer MK 111A at the Oilseeds Laboratory, Agriculture Canada, Saskatoon, with the assistance of Dr. D. Woods. On the basis of poly-unsaturated fatty acid content, sesame oil was next to safflower, sunflower, and corn oil and better than soybean, cotton, and peanut oil.

Mustard

Rape and mustard seeds were obtained from Canada, England, Sweden, France, Federal Republic of Germany, and India, between 1981 and 1985. The species of *Brassica* tested were *B. napus* (15 varieties), *B. campestris* (4 varieties), *B. juncea* (15 varieties), *B. carinata* (2 varieties) and

Table 9. Data on varieties of soybeans planted at different times

| Variety | Plant height (cm) | | | Yield (kg/ha ⁻¹) | | |
|----------------|-------------------|----|----|------------------------------|-------|-------|
| | Date of planting | | | Date of planting | | |
| | 1 | 2 | 3 | 1 | 2 | 3 |
| <u>1983-84</u> | | | | | | |
| AGS-29 | 66 | 60 | 48 | 2,708 | 2,182 | 1,284 |
| AGS-129 | 67 | 53 | 40 | 2,735 | 2,161 | 932 |
| AGS-17 | 65 | 58 | 50 | 2,409 | 2,280 | 687 |
| <u>1984-85</u> | | | | | | |
| AGS-29 | 68 | 50 | 64 | 1,972 | 1,665 | 1,555 |
| AGS-129 | 62 | 43 | 50 | 1,958 | 1,504 | 1,380 |
| AGS-17 | 64 | 30 | 56 | 1,742 | 1,211 | 1,286 |

1/ 1, 2, and 3 are November, December and January, respectively.

Table 10. Soybean spacing trial planted in November 1983

| Variety | No. plants/ha at harvest ('000) | Per cent reduction in no. plants up to harvest | Plant height (cm) | Height of lowest pod (cm) | Yield kg/ha |
|---------|---------------------------------|--|-------------------|---------------------------|-------------|
| AGS-29 | 386 | 18 | 61 | 15 | 2,647 |
| | 368 | 7 | 58 | 14 | 2,630 |
| | 306 | 2 | 53 | 13 | 2,644 |
| AGS-2 | 392 | 19 | 110 | 15 | 2,418 |
| | 380 | 5 | 119 | 15 | 2,168 |
| | 311 | 0 | 109 | 14 | 2,372 |

Table 11. Data on fertilizer trial (soybeans)

| Variety | Fertilizer rate at $R_{5/5}$ growth stage | Yield (kg ha ⁻¹) | | |
|-------------|--|--------------------------------------|-------|-------|
| | | Basal fertilizer rates ^{2/} | | |
| | | F1 | F2 | Mean |
| IAC 73-5115 | A | 2,499 | 2,436 | 2,468 |
| | B | 2,295 | 2,380 | 2,338 |
| | C | 2,400 | 2,334 | 2,367 |
| | D | 2,239 | 2,013 | 2,126 |
| AGS-29 | A | 2,103 | 2,120 | 2,111 |
| | B | 1,902 | 2,000 | 1,951 |
| | C | 2,065 | 2,015 | 2,040 |
| | D | 2,036 | 2,207 | 2,122 |
| Mean | | 2,192 | 2,188 | |

^{1/} A = nil; B = ammonium sulphate sidedressed (150 gk ha⁻¹); C = urea (70 kg ha⁻¹) and Emgeo (80 kg ha⁻¹) sidedressed; D = Nitrophoska (4.5 l ha⁻¹) as foliar spray.

^{2/} F₁ = 18.46.0 + K-Mag + micronutrient mixture at, respectively, 100, 100 and 10 kg ha⁻¹. F₂ = as F₁ plus triple superphosphate and potassium sulphate both at 50 kg ha⁻¹. Applied at planting.

Table 12. Data on selected varieties of sesame

| Variety | Yield (kg/ha) | | | | | Percent | |
|---------------|---------------|-------|------|-------|-------|---------------|----------------|
| | 1983 | | 1984 | | 1985 | | |
| | Nov. | Dec. | Jan. | Dec. | Jan. | Mean yield | oil content |
| Yuamar | - | - | - | 1,261 | - | 1,261 | 51 |
| Aceitera | - | - | 719 | 1,169 | 1,520 | 1,136 | 52 |
| Acarigua | - | - | - | 1,055 | - | 1,055 | 50 |
| Local I | 1,238 | 1,346 | 569 | 655 | - | 955 | 51 |
| Chino Rojo | 813 | - | - | 828 | 870 | 837 | 53 |
| Phultil No. 1 | - | - | 664 | 709 | 898 | 757 | 52 |
| FAO 59.833 | 290 | 1,219 | 263 | 622 | 949 | 599 | 55 |

Table 13. Effectiveness of alachlor applied post-plant to sesame

| Rate of alachlor (kg a.i. ha ⁻¹) | No. days after sowing | | | | | |
|---|-----------------------|----|--------------------|---|----|----|
| | 16 | | 23 | | 30 | |
| | a ^{1/} | b | a | b | a | b |
| 3.33 | 22 | 1 | 21 | 1 | 24 | 4 |
| 2.66 | 52 | 3 | 50 | 3 | 38 | 22 |
| 2.00 | 45 | 5 | 34 | 6 | 33 | 15 |
| 0.00 | 104 | 17 | Hand weeded weekly | | | |

^{1/} a and b are number of dicot and monocot weeds respesping m⁻².

B. hirta (1 variety). All except **B. juncea** performed poorly. Some varieties of **B. juncea** yielded up to 1.5 ton/ha⁻¹, but all were high in erucic acid and glucosinolates. The FAO/WHO Joint Expert Consultation on Dietary Fats and Oils in Human Nutrition, in 1980, suggested: "In view of the present knowledge from animal studies, it seems prudent to recommend for populations in which fat constitutes a high proportion of dietary energy (a) the reduction of erucic acid in **Brassica** oils and/or (b) the blending or use of **Brassica** oils with other fats and oils". However, high erucic acid **Brassica** oils have been consumed for centuries on the Indian sub-continent, China and Japan, with no apparent adverse effects.

During a visit to Agriculture Canada, Saskatoon, Canada, Dr. R. K. Sowney and Dr. Don L. Woods indicated that they were breeding varieties of **B. juncea** with low erucic acid. In November 1985, two such varieties were received and planted (Table 14).

Table 14. Data on two mustard varieties

| Variety | Days to | | Plant height (cm) | Seed colour | Yield |
|------------|------------------|---------|-------------------------|-------------------|-------|
| | 50% flowering | harvest | | | |
| T 931 R70 | 43 | 105 | 180 | yellow + brown | Well |
| T 931 R139 | 89 | 145 | 210 | yellow | Well |

Plant breeding has not yet met with success in breeding varieties low in glucosinolates. Some varieties predominantly contain allyl-, others butenyl-glucosinolates. McGregor et al. (1963) have demonstrated that allyl-glucosinolates could be detoxified by ammoniation, but not butenyl-glucosinolates. Ammoniated mustard meal contains 21 per cent less available

lysine, about 3 per cent more digestible energy and 8 per cent more digestible crude protein than the low glucosinolate canola meal (Bell et al., 1984). In view of this, emphasis is now being placed on selecting varieties containing low amounts of erucic acid and predominantly allyl glucosinolates.

Sunflower

Thirty nine hybrid varieties of sunflower from commercial firms in the USA, from FAO, and from the Research Institute for Cereals and Industrial Crops in Romania, were tested between 1982 and 1985. Varieties were planted in June/July 1982, in Cayo and Corozal districts. The plants grew well, flowered, set seed, but flower heads rotted at both locations due to high rainfall at maturity. Trials planted in November and January indicated that the higher yielding varieties were RO-29, RO-130 and Ag-S49, but the maximum yield obtained was only 2,048 kg per ha. The shelling percentage of high yield varieties was 70. The yield was considered too low to warrant commercialization.

UTILIZATION OF OILSEEDS

Trial to Raise Broilers on Whole Fat Soybean

Whole fat soybean was roasted to destroy urease and trypsin inhibitors. Composition of various rations used is given in Table 15. Starter ration was given for the first 42 days. The trial was started on March 19, 1986 with one day old chicks and the birds were processed on May 18, 1986.

Table 15. Composition of broiler rations

| Ingredient | Starter Ration (g) | | | Finisher (g) | | |
|----------------------------------|--------------------|--------|--------|--------------|--------|--------|
| | Ration No. | | | Ration No. | | |
| | 1 | 2 | 3 | 1 | 2 | 3 |
| Dicalcium phosphate | 750 | - | - | - | - | - |
| Limestone | 750 | - | - | - | - | - |
| d L methionene | 100 | 100 | - | 100 | 100 | - |
| Mineral and vit. mix | 120 | 120 | - | 120 | 120 | - |
| Common salt | 200 | 200 | - | 200 | 200 | - |
| Meat and bone meal ^{1/} | - | 5,000 | - | - | 5,000 | - |
| Roasted soybean | 23,000 | 15,580 | - | 18,480 | 14,480 | - |
| 48% protein commercial feed | - | - | 16,666 | - | - | 14,280 |
| Corn | 25,000 | 26,000 | 33,334 | 30,200 | 30,500 | 35,720 |
| Total | 50,000 | 50,000 | 50,000 | 50,000 | 50,000 | 50,000 |
| Per cent protein in feed | 22 | 22 | 22 | 20 | 20 | 20 |
| Per cent fat in feed | 10 | 10 | 3 | 8 | 9 | 3 |

^{1/} Per cent content in meat and bone meal, roasted soybean, corn and 48% protein feed were: 35, 37.6, 9, and 48, respectively, crude protein, and 17, 17, 3.5, and 2.5, respectively.

Water and feed were provided *ad libitum*. Data on live and dressed birds and feed consumed, are presented in Table 16, indicated that 2.72 kg and 2.34 to 2.36 kg feed were consumed per kg dressed bird weight when feed was prepared from 48 per cent crude protein (CP) content imported feed and 41 per cent CP content whole fat soybean, respectively. The birds raised on whole fat soybean had slightly more body fat. Thus ration prepared from whole fat soybean should be mixed with ration prepared from seed containing 48 per cent CP, in a proportion, so that the mixture contained about 5 per cent fat. It has been calculated that by following this system, about 700 ha of soybean could be produced and utilized, without having an oil mill in the country.

Table 16. Data on broiler production

| Ration No. | No. birds | | Live weight (kg) | | Dressing per cent | | Kg feed consumed | kg feed consumed per kg of wt. | |
|------------|-----------|----|------------------|------|-------------------|----|------------------|--------------------------------|---------|
| | ♂ | ♀ | ♂ | ♀ | ♂ | ♀ | | Live | dressed |
| 1 | 16 | 24 | 41.4 | 54.5 | 84 | 86 | 190 | 1.99 | 2.34 |
| 2 | 17 | 18 | 47.3 | 42.3 | 83 | 85 | 177 | 1.98 | 2.36 |
| 3 | 11 | 27 | 30.6 | 62.6 | 82 | 81 | 207 | 2.22 | 2.72 |

Cheaper mechanical press oil mill was preferable to capital and high technology intensive solvent extraction mill, as the meal would contain about 7 per cent fat so very necessary for better growth of livestock. The meal obtained from solvent extraction method contained almost no fat.

Improved Human Nutrition

Soybean is being mixed with corn, 12 per cent by weight, by one tortilla making factory in Belize for improved corn protein efficiency ratio, increased protein content, and better texture. Another entrepreneur is considering production of soy milk.

REFERENCES

- Boll, J.M., Keith, M.D., Blake, J.A., and McGregor, D.I. 1984. Nutritional evaluation of ammoniated mustard meal for use in swine feed. *Can. J. Anim. Sci.* 64: 1023-1033.
- McGregor, D.I., Blake, J.A., and Pickard, M.D. 1983. Detoxification of *Brassica juncea* with ammonia. *Intl. Rapeseed Conf.*, Paris, 17-19 May, 1983; 1425-1431.
- Paredes-Lopes and Saharopoulos-Paredes, M.E. 1983. A review of tortilla production technology. *Bakers Digest*, Sept. 1983: 16-25.

Schroder, G.N. and Rawson, J.E. 1984. Herbicide evaluation studies in sesame. *Australian Weeds*, 3(2): 47-49.

Spehar, C.R., Souza, P.I.M., and Urban Filho, S. 1983. Soybean in the Brazilian tropics. In: "Soybean in the Tropical and Subtropical Cropping Systems", *Proc. Symp. Taikuba, Japan*. Sept. 25 to Oct. 1, 1983. Eds. S. Shanmugasundaram and E.W. Sulzberger. AVRDC, Taiwan, China.