



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.


Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*



JOINT PROCEEDINGS



TROPICAL REGION

**21st Annual Meeting
of the Caribbean Food Crops Society
and
32nd Annual Meeting of the American Society for
Horticultural Science — Tropical Region**

technology for agricultural development

**Hilton Hotel, Port of Spain, Trinidad
8 - 13 September 1985**

Host Institutions

- Caribbean Agricultural
Research and Development
Institute
- Ministry of Agriculture, Lands
and Food Production, Trinidad
& Tobago
- Faculty of Agriculture,
University of the West Indies

Published by the Caribbean Food Crops Society, Box 506, Isabela, Puerto Rico 00662

CARBON DIOXIDE ENRICHMENT ON GROWTH AND YIELD OF SWEET POTATOES (*IPOMOEA BATATAS*, cv GEORGIA JET)

P.R. Biswas, D.R. Hileman, J.N. Mc Crimmon, P.P. Ghosh, N.C. Bhattacharya and M.E.M. Tolbert

Carbon Dioxide Research Laboratory, Tuskegee Institute, Alabama, 36088

ABSTRACT

This study was conducted to obtain field data on the growth and yield of sweet potatoes in elevated levels of carbon dioxide (CO₂). Sweet potato plants were planted in open-top chambers as well as in open field plots. The plants were grown at ambient CO₂ with and without chambers, ambient +75, +150, and +300 ppm (90 days). Enriched CO₂ concentration increased the number of tubers and the percentage of dry matter of the tubers. The density, length and diameter of the tubers did not vary significantly in enriched CO₂ levels.

RESUMEN

Este estudio se llevó a cabo con el objeto de obtener información sobre los ensayos relacionados con el desarrollo y la producción del camote con niveles altos de dióxido de carbono (CO₂). Los camotes fueron plantados en cámaras sin techos, como también, se los cultivó en parcelas de campo. Las plantas fueron cultivadas al ambiente CO₂, en cámaras y fuera de ellas, ambiente +75, +150 y +300 ppm (noventa días). Una concentración más fuerte de CO₂ incrementó el número de tubérculos. La densidad, el largo y el diámetro de los tubérculos no varió significativamente en niveles altos de CO₂.

The global carbon dioxide (CO₂) concentration is gradually increasing due to fossil fuel consumption, rapid advancement in industrialization and deforestation. It has been estimated that atmospheric CO₂ concentration will be doubled by 2025 (Clark *et al.*, 1982). Extensive literature is now available to demonstrate that elevated CO₂ results in the increase of dry matter accumulation in both vegetation and reproductive components of plants (Wittwer, 1980, 1983; Kimball, 1983).

Although there are a number of reports on the influence of atmospheric CO₂ enrichment on photosynthesis and short-term growth, very few reports are available on crops which have been subjected to CO₂ enrichment during their entire growing season. Furthermore, little information is available on plant responses to CO₂ enrichment under field conditions (Kramer, 1981; Strain *et al.*, 1984). It is therefore, imperative to have a better understanding of the effects of CO₂ enrichment on crop plants.

Reports on the influence of elevated CO₂ levels on tuber crops like radish (*Raphanus sativus* L. 'Whitetip'; Knecht, 1975) and potato (*Solanum tuberosum*, 'Kennebec'; Collins, 1976; Goudriaan and deRuiter, 1983) did not consider the effects of long-term exposure to elevated CO₂. Therefore, more information is needed to understand the overall effects of CO₂ enrichment on growth, development, and yield potential of root crops.

Sweet potatoes (*Ipomoea batatas* L.) were selected as an experimental crop in the southern United States. In addition, sweet potatoes are one of the world's major food crops. Currently, little is known about the responses of sweet potatoes to elevated CO₂ (Strain *et al.*, 1984). Sweet potatoes have widespread growth ranges and, consequently, can be easily grown for experimental purposes. In the present study, CO₂ enrichment has been applied to sweet potatoes in order to investigate its effect on their growth, physiology, and yield under field conditions.

In the summer of 1984, under the sponsorship of DOE and Tuskegee University, experiments were conducted to study the physiological and biochemical effects of enriched CO₂ on sweet potatoes in open top chambers at the George Washington Carver Agricultural Experiment Station, Tuskegee, Alabama.

The techniques for the generation of large scale test atmospheres in the field for the purpose of obtaining dose response relationships of crop plants were developed at the USDA Air Quality Laboratory at Raleigh. The main task of the 1984 study at Tuskegee University was to assemble equipment to generate test atmospheres of CO₂ in the field, using open top chambers as the basic exposure unit for studying the responses of sweet potatoes and cowpeas to enriched CO₂.

Plant responses to CO₂ in open top chambers have been demonstrated by several investigators in a variety of crops. This study focused on growth and development of sweet potatoes at levels of CO₂ ranging from the ambient level of 354 ppm to 659 ppm. The effects of CO₂ on leaf and stem weights, stem length, leaf area, and stomatal number and conductance were studied. Additional studies on sweet potatoes included the effects of CO₂ on the weight, chemical content and quality of tubers.

In addition to the field studies, a series of experiments by Tuskegee University scientists conducted at the Duke University Phytotron were completed in 1984. These experiments focused on the biochemical and physiological effects of elevated CO₂ on sweet potatoes grown in controlled environments.

Sweet potatoes grown in open top chambers, at ambient CO₂ concentrations had fewer leaves, less total runner length, and lower fresh and dry weights of shoots, leaves and tubers as compared to sweet potatoes grown in the open field without chambers. These results emphasize the need to quantify more carefully the environmental differences between open top chambers and the open field. These chamber effects need to be considered when drawing conclusions about the effects of elevated CO₂ on the growth and yield of plants grown in open top chambers.

While shoot growth in sweet potatoes increased with increasing CO₂, few of the effects were large enough to be significant. However, the percentages of nitrogen and protein nitrogen in sweet potato leaves decreased significantly at higher CO₂ concentrations. The total fresh weight of tubers increased significantly at the higher levels of CO₂, due primarily to the increase in the number of tubers. This suggests a shift in the partitioning of photoassimilates toward

tubers with increasing CO₂. There were no differences in the density of stomates or in stomatal conductances in sweet potato leaves. Analyses of tubers indicated that protein, total carotenoids and insoluble dietary fiber all decreased with increasing CO₂, while dry matter content increased with increasing CO₂. Taste panel tests indicated small but significant preferences in some test categories for potatoes grown at the highest CO₂ level.

In phytotron studies with pot-grown sweet potatoes, plants grown at 675 or 1000 ppm CO₂ showed increases in the length of the main stem, total branch length, the number of branches and leaf area as compared to those grown in 350 ppm CO₂. At each harvest interval the production of total dry matter increased in response to increases in the level of CO₂. Specific leaf weight also increased with increased CO₂ concentrations. At the final harvest, the dry weights of roots and tubers increased 1.8 and 2.6 times in plants grown at 675 and 1000 ppm CO₂, respectively, compared to those grown at 350 ppm. Carbon dioxide enrichment resulted in early tuber maturation in sweet potatoes.

References

- Bhattacharya, N.C., Biswas, P.K., Bhattacharay, S., Sionit, N. and Strain, B.R. (1985) Growth and yield response of sweet potato (*Ipomoea batatas*) to atmospheric CO₂ enrichment. *Crop Sci.* (in press).
- Clark, W.C., Cook, K.H., Moreland, G., Weinberg, A.M., Totty, R.M., Bell, P.R., Cooper, L.J., and Cooper, C.L. (1982) The carbon dioxide question: A perspective for 1982. In: W.C. Clark (ed.) *Carbon Dioxide Review*, Clarendon Press, Oxford. pp. 3 - 43.
- Collins, W.B. (1976) Effect of carbon dioxide enrichment on growth of potato plant, *Hortscience*. 11 467 - 469.
- Kimball, B.A. (1983) Carbon dioxide and agricultural yield: an assemblage and analysis of 430 prior observations. *Agron. J.* 75 779 - 788.
- Knecht, G.N. (1975) Response of radish to high carbon dioxide, *Hortscience* 10 271 - 275.
- Kramer, P.J. (1981) Carbon dioxide concentration, photosynthesis and dry matter production. *Bioscience* 31 29 - 33.
- Strain, B.R., Sionit, N., and Cure, J.D. (1984) Direct effects of carbon dioxide on plants and communities: a bibliography. Dept. of Botany, Duke University, Durham, North Carolina, 70 pp.
- Wittwer, S.H. (1980a) Carbon dioxide and climatic change: an agricultural perspective. *J. Soil Water Conserv.* 35 116 - 120.
- Wittwer, S.H. (1980b) Environmental and society consequence of a possible CO₂ induced climate change on agriculture. Paper presented at annual meeting of Am. Assoc. Adv. Sci., San Francisco, Jan. 5.
- Wittwer, S.H. (1983) Rising atmospheric CO₂ and crop productivity. *Hort. Sci.* 18 667 - 673.