



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.


Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*



JOINT PROCEEDINGS



TROPICAL REGION

**21st Annual Meeting
of the Caribbean Food Crops Society
and
32nd Annual Meeting of the American Society for
Horticultural Science — Tropical Region**

technology for agricultural development

**Hilton Hotel, Port of Spain, Trinidad
8 - 13 September 1985**

Host Institutions

- Caribbean Agricultural
Research and Development
Institute
- Ministry of Agriculture, Lands
and Food Production, Trinidad
& Tobago
- Faculty of Agriculture,
University of the West Indies

Published by the Caribbean Food Crops Society, Box 506, Isabela, Puerto Rico 00662

SOLUBLE SALTS INTERPRETATION FOR ORNAMENTAL FOLIAGE CROP PRODUCTION

R. T. Poole, C. A. Conover and A. R. Chase

University of Florida, IFAS
Agricultural Research and Education Centre,
2807 Binion Road, Apopka, FL 32703

ABSTRACT

During the last few years, experiments have been conducted with potting media and fertilizer as variables. Soluble salts of the media and plant ratings were sometimes determined. Comparisons of the levels of soluble salts and plant growth indicated soluble salts readings are a poor indicator of the potting media ability to produce satisfactory foliage plants.

RESUMEN

Durante los últimos años, se ha conducido investigaciones con diferentes mezclas de tierra y abonos. De vez en cuando se determinó los niveles de sales solubles en las mezclas de tierra y el comportamiento de las plantas. La comparación de los niveles de sales solubles y el crecimiento de las plantas indicaron que el nivel de sales solubles no es un indicador seguro de la capacidad de una mezcla de tierra para producir plantas ornamentales satisfactorias.

Satisfactory plant growth requires macro and micro-elements, normally supplied by specific fertilizers. Addition of fertilizer increases soluble salts levels, actually soluble ions, in the medium. To determine soluble salts levels, a conductivity meter is used since dissolved salts increase electrical conductivity. Horticulturists use soluble salts levels to indicate fertility level desired for plant production. If a plant is chlorotic or necrotic, roots are usually examined and sometimes soluble salts of the soil determined. If levels are too low, additional fertilizer is suggested and if levels are too high removal of fertilizer (soluble salts) by heavy applications of water to the soil is suggested.

Four methods to determine soluble salts levels are: (1) 1:2 — soil:water, by dry weight; (2) 1:2 — soil: water, by volume; (3) saturated paste and (4) leachate. The dry weight method should not be used for potting media because of the variability of bulk density of the different mixtures used for growing plants. The volume procedure is commonly used because of its simplicity. One volume of soil is added to 2 volumes of water, mixed, and usually allowed to stand for about 15 minutes with occasional stirring. The electrode of the conductivity meter can be placed directly in the solution, or the solution can be filtered prior to measuring conductivity. The saturated paste procedure had been recommended because readings obtained by this method are said to

approximate those in the root environment, but variations in water holding capacity of potting mixtures suggests this may not be much of an advantage. The procedure for saturated paste is the most complex since the soil sample is moistened until a glistening paste is obtained. Obtaining the same degree of glistening each time can lead to errors in reproduction. After glistening is obtained, vacuum is applied to extract the solution and conductivity determined. Although the leachate method is not commonly used, it is the easiest to conduct. Water is poured through the medium in the container which is in a beaker. Fifty to 100 ml of leachate is collected and conductivity determined. This method allows frequent sampling without removal of soil or disturbing the root system.

Unfortunately, many types of electrical conductivity meters are used and units of measurement differ. The usual unit is mhos, but the reading could be micromhos (mhos $\times 10^{-6}$) millimhos (mhos $\times 10^{-3}$) or mhos $\times 10^{-5}$. Numerous tables have been printed with ranges for the interpretation of these readings, (Table 1).

During the last few years, experiments have been conducted at AREC-Apopka, Florida, with potting media and fertilizer as variables. Soluble salts of the media and plant ratings were sometimes determined. A summary of soluble salts of the media and con-

Table 1. Interpretation^a of soluble salts readings (mhos $\times 10^{-5}$) for 1:2 air dry soil (peat or light weight mix) to water by volume (A) and saturated paste (B) methods (5).

Mohs $\times 10^{-5}$	Rating	Remarks
A 0-50 B 0-100	Low	Need fertilizer
A 51-100 B 101-132	Medium	Satisfactory
A 101-175 B 133-164	High	Desirable
A 176 + B 165 +	Very High	Leach

^a There is no guide for leachate soluble salts.

dition of the plants is contained in Table 2.

An examination of Table 2 clearly reveals the difficulty of determining the adequacy of a potting medium or fertilizer program by soluble salts readings. Media containing plants with the worst indicators (grade or height) had 11 ratings in the low range, 3 in the medium range, 4 in high and 4 in the

very high range. Media containing plants with the best indicators had an almost identical rating: 12 in the low range, 3 medium, 6 high and 1 in the very high range. When comparing the worst and best ratings of the same plant, ratings were the same 11 times, went from a high to a low rating 6 times and from a low to a high rating 5 times. Plants with a

Table 2. Comparisons of growth indicators with soluble salts of potting media.

Species	Reference	Mohs \times 10		Indicator		Method	Time of reading	Rating Table 1	
		worst	best	worst	best			worst	best
<i>Aglaonema</i>	4	6	5	3.5 ^p	3.9	2:1, vol	Initial	Low	Low
<i>commutatum</i>	9	149	218	2.2 ^p	4.6	Sat. paste	Initial	Med	High
<i>Alphelandra</i>	1	12	27	1.9 ^p	3.1	2:1, vol	Final	Low	Low
<i>squarrosa</i>	2	27	45	3.2 ^p	3.9	2:1, vol	Final	Low	Low
<i>Brassaia</i>	3	242	161	1.8 ^p	3.9	2:1 vol	Final	V.high	High
<i>actinophylla</i>	6	205	51	14 ^c	23	2:1, vol	Initial	V.high	Med
	9	149	156	3.4 ^p	4.6	Sat. paste	Initial	High	High
	11	515	185	15 ^w	24	Leachate	Final		
	11	690	300	122 ^w	182	Leachate	Final		
	12	310	155	3.8 ^p	4.1	Leachate	Initial		
	12	85	70	3.8 ^p	4.1	Leachate	Final		
<i>Calathea</i>									
<i>makoyana</i>	9	158	155	2.0 ^p	4.4	Sat. paste	Initial	High	High
<i>Chamaedorea</i>									
<i>elegans</i>	11	730	225	6	8	Leachate	Final		
<i>Chrysalidocarpus</i>									
<i>lutescens</i>	7	0	19	1.2 ^p	4.1	2:1, vol	Final	Low	Low
<i>Dieffenbachia</i>	1	24	45	26.3 ^c	28	2:1, vol	Final	Low	Low
<i>maculata</i>	2	47	80	26 ^c	28	2:1, vol	Final	Low	Med
	3	164	142	2.2 ^p	4.4	2:1, vol	Final	High	High
	10	34	208	2.1 ^p	3.9	2:1, vol	Final	Low	High
	12	110	50	3.7 ^p	3.9	Leachate	Initial		
	12	40	40	3.7 ^p	3.9	Leachate	Final		
	12	150	165	3.6 ^p	4.8	Leachate	Initial		
	12	40	60	3.6 ^p	.8	Leachate	Final		
<i>Ficus</i>									
<i>benjamins</i>	13	171	262	3.8 ^p	4.5	2:1, vol	Initial	High	V.High
	13	51	53	3.8 ^p	4.5	2:1, vol	Final	Med	Med
<i>Maranta</i>									
<i>leuconeura</i>									
<i>erythroneura</i>	9	705	200	13 ^w	15 ^w	Leachate	Final		
<i>Monstera</i>									
<i>deliciosa</i>	4	207	14	33 ^c	38	2:1, vol	Initial	V. high	Low
<i>Nephrolepis</i>	1	5	29	26 ^c	31	2:1, vol	Final	Low	Low
<i>exalata</i>	2	66	21	30 ^c	33	2:1, vol	Final	Med	Low
	4	3	3	3.7 ^p	3.9	2:1, vol	Final	Low	Low
<i>Pellionia</i>									
<i>pulchra</i>	8	50	59	14 ^c	17	2:1, vol	Final	Low	Med
<i>Peperomia</i>									
<i>obtusifolia</i>	8	50	59	14 ^c	59	2:1, vol	Final	Low	Med
	10	234	22	2.4 ^p	22	2:1, vol	Final	V. high	Low
	11	680	158	9 ^w	158	Leachate	Final		
	11	621	172	116 ^w	172	Leachate			
<i>Philodendron</i>	1	20	18	24 ^c	26	2:1, vol	Final	Low	Low
<i>scandens</i>	2	64	49	21 ^c	23	2:1, vol	Final	Med	Low
<i>oxycardium</i>	11	225	690	8 ^w	10	Leachate	Final		
<i>Pilea</i>									
<i>involucrata</i>	8	48	59	12 ^c	13	2:1, vol	Final	Low	Med

^c Centimeters

^p Plant grade, 1= poor, 5=excellent

^w Weight in g

grade of 3.0 or above (considered salable) were growing in 8 soil mixes with low soluble salts, 2 with medium levels, 8 with high salts and 1 with very high soluble salts.

Summary

Results of this compilation show that soluble salts readings are a poor indicator of the potting media ability to produce satisfactory, salable foliage plants.

Literature Cited

1. Conover, C.A. and R.T. Poole. 1977. Influence of irrigation method and fertilizer source and level on growth of four foliage plants. *Proc. Fla. State Hort. Soc.* 90: 312-313.
2. Conover, C.A. and R.T. Poole. 1977. Influence of potting media and fertilizer source and level on growth of four foliage plants on capillary mats. *Proc. Fla. State Hort. Soc.* 90: 316-318.
3. Conover, C.A. and R.T. Poole. 1981. Influence of light and fertilizer levels and fertilizer sources on foliage plants maintained under interior environments for one year. *J. Amer. Soc. Hort. Sci.* 106 (5): 517-574.
4. Conover, C.A. and R.T. Poole. 1983. Utilization of *Melaleuca quinquenervia* as a potting medium component for greenhouse production of foliage plants. *HortScience* 18 (6): 886-888.
5. Joiner, J.N. ed. 1981. *Foliage Plant Production*. Prentice-Hall, Inc. Englewood Cliffs, N.J. 07632. 613 pp.
6. Poole, R.T. and W.E. Waters. 1972. Evaluation of various potting media for growth of foliage plants. *Proc. Fla. State Hort. Soc.* 85: 395-398.
7. Poole, R.T. and C.A. Conover. 1975. Media, shade and fertilizer influence production of the Areca palm, *Chrysalidocarpus lutescens* Wendl. *Proc. Fla. State Hort. Soc.* 88: 603-605.
8. Poole, R.T. and C.A. Conover. 1977. Influence of medium, container size and water regime on growth of *Pellionia pulchra* N.E. Br. and *Pilea involucreata* (Sims) Urb. *Proc. Fla. State Hort. Soc.* 90: 319-320.
9. Poole, R.T. and C.A. Conover. 1979. Melaleuca bark and solite as potential potting ingredients for foliage plants. *Proc. Fla. State Hort. Soc.* 92: 327-329.
10. Poole, R.T. and R.W. Henley. 1980. Fertilization and water use of *Dieffenbachia maculata* and *Peperomia obtusifolia*. *Proc. Fla. State Hort. Soc.* 93: 162-164.
11. Poole, R.T. and R.W. Henley. 1981. Constant fertilization of foliage plants. *J. Amer. Soc. Hort. Sci.* 106 (1): 61-63.
12. Poole, R.T. and C.A. Conover. 1982. Influence of leaching, fertilizer source and rate, and potting media on foliage plant growth, quality, and water utilization. *J. Amer. Soc. Hort. Sci.* 107 (5): 793-797.
13. Poole, R.T. and C.A. Conover. 1985. Growth of *Ficus benjamina* in combinations of peat, sand and *Melaleuca*. *HortScience* 20 (3): 383-385.
14. Waters, W.E. and W. Llewellyn. 1968. Effects of coated-slow-release fertilizer on growth responses, chemical composition and soil salinity levels for foliage plants. *Proc. Fla. State Hort. Soc.* 81: 380-388.