



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

CANTER

8804 ✓

Department of Economics
UNIVERSITY OF CANTERBURY

CHRISTCHURCH, NEW ZEALAND



GIANNINI FOUNDATION OF
AGRICULTURAL ECONOMICS
LIBRARY

FEB 13 1989

**PRE-TESTING FOR LINEAR RESTRICTIONS
IN A REGRESSION MODEL WITH
STUDENT- t ERRORS**

By Judith A. Clarke

Discussion Paper

No. 8804

Department of Economics, University of Canterbury
Christchurch, New Zealand

Discussion Paper No. 8804

September 1988

**PRE-TESTING FOR LINEAR RESTRICTIONS
IN A REGRESSION MODEL WITH
STUDENT- t ERRORS**

By Judith A. Clarke

*This paper is circulated for discussion and comments. It should not be quoted without the prior approval of the author.

The Author's Electronic Mail Address is:

D.Giles%Canterbury.AC.NZ@Relay.CS.NET

ABSTRACT

In this paper, we derive the exact risk (under quadratic loss) of pre-test estimators of the prediction vector and error variance of a linear regression model whose errors are assumed to be normally distributed but in fact follow a multivariate Student-t distribution. The pre-test in question is one of the validity of a set of exact linear restrictions on the model's coefficient vector. We demonstrate how the known results for the model with normal disturbances can be extended to this broader case. Numerical evaluations of the risk expressions suggest that misspecifying the error distribution in this way does not, qualitatively, affect the risk properties of the estimators.

*This work forms part of the author's Ph.D. research. My thanks to David Giles, Tony Rayner, Robin Carter, George Judge, John Knight, Peter Morgan, Kazuhiro Ohtani, Mike Veall and Victoria Zinde-Walsh for their constructive comments and suggestions. I am also extremely grateful to Robert Davies for suggesting the use of his algorithm AS155 and for providing his software.

1. INTRODUCTION

The sampling properties of the estimators of the parameters of the linear regression model, after a pre-test for linear restrictions on the coefficient vector, have widely been examined (see, for example, Clarke *et al.* [5,6], Giles [12], Giles and Clarke [13], Judge and Bock [17], Mittelhammer [22], Ohtani [24,25,26], Wallace [37]). All have assumed that the regression errors are normally distributed. There is a large body of literature, however, which suggests that some economic data series may be generated by processes whose error distributions have thick tails, or even infinite variances. Examples include price-change analysis in the stock, financial and commodity markets (Fama [9,10,11], Sharpe [29]), cash flow analysis (Granger and Orr [14]), and demand analysis (Coursey and Nyquist [7]). Such possibilities have led to studies of the robustness of the traditional estimators and searches for 'robust' estimators. Huber [15], Koenker [21], and Judge *et al.* [18] provide surveys of this literature.

One distribution receiving considerable attention is the multivariate Student-t (Mt) with zero mean vector. For this error distribution, as shown by Singh [30,31], Thomas [33], and Zellner [38], for example, the marginal distributions of the errors are univariate Student-t and the errors are uncorrelated but are not independent. For small values of γ , the degrees of freedom of the distribution, the marginal distributions have thick tails; as $\gamma \rightarrow \infty$, the pdf approaches a normal form; and when $\gamma = 1$, the pdf is Cauchy.

Linear regression models with M_t errors have been considered by some authors, including Judge *et al.* [19], King [20], Nimmo-Smith [23], Singh [30,31], Ullah and Zinde-Walsh [36], and Zellner [38]. Judge *et al.* [19] establish sampling properties of the James-Stein [16] estimator of the location parameter vector (and its positive part counterpart) under a squared error loss measure and a M_t error density. They compare, via a Monte Carlo experiment, the finite sample behaviour (empirical risks) for their Stein-like, and some conventional robust, estimators. In general, the risk characteristics for their Stein-like estimators when the errors are nonnormal are found to be the same as for the normal errors case.

In this paper we consider the question of the robustness, in terms of risk under squared error loss, of the usual pre-test estimators of the prediction vector¹ and the error variance of a linear regression model, after a pre-test for exact linear restrictions on the coefficient vector using the traditional F-ratio, when the errors follow a M_t distribution with γ degrees of freedom. In the next section we detail the model and give the unrestricted and restricted estimators of the model's parameters. Sections 3 and 4 present, discuss and evaluate the finite sample risk functions of the various estimators of the prediction vector and the error variance respectively. Some concluding remarks are given in the final section, and the proofs of the theorems appear in an appendix.

2. THE MODEL AND THE PARAMETER ESTIMATORS

Consider the linear regression model

$$y = X\beta + \epsilon, \quad \epsilon \sim Mt(0, \sigma_\epsilon^2 I_T), \quad (1)$$

where y is a $(T \times 1)$ vector of observations on the dependent variable, X is a $(T \times k)$ non-stochastic design matrix of rank $k (< T)$, β is a $(k \times 1)$ vector of unknown parameters. The $(T \times 1)$ vector of disturbances, ϵ , is assumed to follow the multivariate Student-t (Mt) distribution

$$f(\epsilon) = \frac{C^*}{\sigma^T} \left[1 + \frac{\gamma-1}{\sigma^2} \epsilon' \epsilon \right]^{-(T+\gamma)/2}, \quad \begin{array}{l} \sigma > 0, \gamma > 0 \\ -\infty < \epsilon < \infty \end{array} \quad (2)$$

where $C^* = \Gamma\left(\frac{T+\gamma}{2}\right) \left[(\pi\gamma)^{T/2} \Gamma(\gamma/2) \right]^{-1}$

is a normalising constant, γ and σ^2 are the degrees of freedom and scale parameters of the distribution, and $\sigma_\epsilon^2 = \gamma\sigma^2/(\gamma-2)$ is the common variance of the ϵ_i 's, $i = 1, \dots, T$; $\gamma > 2$.

Consider m independent linear restrictions on β , summarized by the hypothesis

$$H_0: R\beta = r \text{ vs. } H_1: R\beta \neq r \quad (3)$$

where R and r are $(m \times k)$ and $(m \times 1)$ matrices of known constants, and rank $R = m (< k)$.

The unrestricted least squares estimator of β , $\bar{\beta} = (X'X)^{-1}X'y$, is also the unrestricted maximum likelihood estimator (see, Singh [30,31], Thomas [33], Zellner [38]). Furthermore, it is easy to show that the restricted least squares estimator of β , $\beta^* = \bar{\beta} - (X'X)^{-1}R'[R(X'X)^{-1}R']^{-1}(R\bar{\beta}-r)$, is also the restricted maximum likelihood estimator of β .

The usual F-ratio for testing the linear restrictions (3) is

$$F = \frac{v(R\bar{\beta}-r)' [R(X'X)^{-1}R']^{-1} (R\bar{\beta}-r)}{m(y-X\bar{\beta})'(y-X\bar{\beta})}, \quad (4)$$

where $v = (T-k)$. Thomas [33], and Ullah and Phillips [35] show that under Mt errors, F has a density function given by

$$f(F) = \left[\Gamma\left(\frac{v}{2}\right) \Gamma\left(\frac{\gamma}{2}\right) \right]^{-1} \sum_{r=0}^{\infty} \frac{\Gamma\left(\frac{m+v}{2} + r\right) \Gamma\left(\frac{\gamma}{2} + r\right)}{\Gamma\left(\frac{m}{2} + r\right) r!} \cdot \left(\frac{m}{v}\right)^{m/2+r} \cdot \frac{F^{m/2+r-1}}{\left(1 + \frac{m}{v}F\right)^{(m+v)/2+r}} \cdot \frac{(2\theta/\gamma)^r}{\left(1 + \frac{2\theta}{\gamma}\right)^{\gamma/2+r}} \quad (5)$$

where $\theta = (R\beta - r)' [R(X'X)^{-1}R']^{-1} (R\beta - r) / 2\sigma^2$, and is a measure of the hypothesis error. Note, first, that when H_0 is true $F \sim F_{m,v}$ (see, Zellner [38, p.401]). Secondly, when $\gamma \rightarrow \infty$, $\epsilon \sim MN(0, \sigma^2 I_T)$, and (5) reduces to the non-central F density, $F_{m,v,\theta}$.

In deriving their result, Ullah and Phillips [35] use the fact that if $\epsilon \sim Mt(0, \frac{\gamma\sigma^2}{\gamma-2} I_T)$, then we can write

$$\epsilon = \sqrt{\gamma}a/q \quad (6)$$

where a and q^2 are independently distributed as $MN(0, \sigma^2 I_T)$ and χ_{γ}^2 respectively. This decomposition enables one to condition on q^2 and work with normal random variates to obtain the conditional density $f(F|q^2)$. This is found to be non-central $F_{(m,v,\lambda)}$ with non-centrality parameter

$$\lambda = \frac{q^2\theta}{\gamma} = \frac{q^2(R\beta - r)' [R(X'X)^{-1}R']^{-1} (R\beta - r)}{2\sigma^2\gamma} \quad (7)$$

The unconditional density of F, given by equation (5), is then obtained by noting that

$$f(F) = \int_0^{\infty} f(F|q^2) f(q^2) dq^2, \quad (8)$$

$$\text{where } f(q^2) = 2^{-\gamma/2} \left[\Gamma(\gamma/2) \right]^{-1} e^{-q^2/2} (q^2)^{\gamma/2-1}.$$

This approach contrasts with that employed by some other authors,

including Zellner [38] and Singh [30,31]. These authors proceed by regarding the error vector ϵ as being randomly drawn from a multivariate normal distribution with a random standard deviation generated from the inverted gamma distribution. It does not matter which approach is used, given the relationship between the χ^2 , inverted gamma and gamma pdf's (see, for example, Zellner [39, pp.369-373]). In the following sections, we employ the Ullah and Phillips decomposition (6) to derive the risk expressions, under an invariant quadratic loss function, of the component and the pre-test estimators of $X\beta$ and σ_ϵ^2 .

3. THE RISKS OF ALTERNATIVE ESTIMATORS OF $E(y)$

We define the pre-test estimator for $E(y) = X\beta$ as

$$\hat{X\beta} = \begin{cases} X\bar{\beta} & ; \text{ if } F > c \\ X\beta^* & ; \text{ if } F \leq c \end{cases}$$

$$= X\bar{\beta}I_{(c, \infty)}(F) + X\beta^*I_{[0, c]}(F) \quad (9)$$

where $c = c(\alpha)$ satisfies $\int_0^c dF_{(m, \nu)} = (1-\alpha)$ and $I_{[a, b]}(F)$ is an indicator function with value unity if $F \in [a, b]$, zero otherwise. Further, if Xb is any estimator of $E(y)$, then its risk function, under quadratic loss, is

$$\rho[Xb, E(y)] = E[Xb - E(y)]' [Xb - E(y)] / \sigma^2, \quad (10)$$

and is the relative mean squared error of Xb . We now derive the risk expressions for the various estimators of $E(y)$. Notice that we are considering a situation involving model misspecification: the F-ratio is being used to test H_0 because the errors are wrongly being assumed to be normally distributed.

3.1 Finite Sample Risk Functions

THEOREM 1.

Under the stated assumptions,

$$\rho \left[X\bar{\beta}, E(y) \right] = k\gamma/(\gamma-2) \quad (11)$$

$$\rho \left[X\beta^*, E(y) \right] = [\gamma(k-m) + 2\theta(\gamma-2)]/(\gamma-2) \quad (12)$$

$$\rho \left[X\hat{\beta}, E(y) \right] = \left[2\theta(\gamma-2)(2P_{202} - P_{402}) + \gamma(k-mP_{201}) \right]/(\gamma-2) \quad (13)$$

where

$$P_{a_1 a_2 a_3} = \sum_{r=0}^{\infty} \frac{(2\theta/\gamma)^r \Gamma(\frac{\gamma}{2} + r + a_3 - 2)}{r! (1+2\theta/\gamma)^{\gamma/2 + r + a_3 - 2} \Gamma(\frac{\gamma}{2} + a_3 - 2)} \cdot I_u \left[\frac{1}{2}(m+a_1) + r; \frac{1}{2}(v+a_2) \right], \quad (14)$$

$I_u(\cdot; \cdot)$ is Pearson's incomplete beta function with $u = cm/(v+cm)$.

PROOF. $\bar{\beta}$ is unbiased, so,

$$\begin{aligned} \rho \left[X\bar{\beta}, E(y) \right] &= \text{tr.} [V(X\bar{\beta})]/\sigma^2 \\ &= \text{tr.} [\gamma\sigma^2 X(X'X)^{-1}X'] / [\sigma^2(\gamma-2)] \end{aligned}$$

where $V(X\bar{\beta})$ is the variance-covariance matrix of $X\bar{\beta}$. (11) then follows immediately.

Turning to the restricted estimator, we have

$$\text{Bias } (X\beta^*) = -X(X'X)^{-1}R' [R(X'X)^{-1}R']^{-1}(R\beta-r)$$

$$\text{and } V(X\beta^*) = \gamma\sigma^2 XG(X'X)^{-1}X' / (\gamma-2), \quad G = I - (X'X)^{-1}R' [R(X'X)^{-1}R']^{-1}R.$$

So,

$$\rho \left[X\beta^*, E(y) \right] = (R\beta-r)' [R(X'X)^{-1}R']^{-1}(R\beta-r)/\sigma^2 + \gamma(k-m)/(\gamma-2)$$

and (12) follows directly, as $(R\beta-r)' [R(X'X)^{-1}R']^{-1}(R\beta-r)/\sigma^2 = 2\theta$.

The proof of (13) is given in the Appendix. ■

Comparing (11), (12) and (13) we note directly:

(i) When $\gamma \rightarrow \infty$, $\epsilon \sim MN(0, \sigma^2 I_T)$, $P_{a_1 a_2 a_3} \rightarrow P_{a_1 a_2}$, where

$$P_{a_1 a_2} = \Pr. \left[F_{(m+a_1, v+a_2; \theta)} \leq \left[\frac{cm(v+a_2)}{v(m+a_1)} \right] \right] \quad (15)$$

(11), (12) and (13) reduce to the well known expressions (see, for example, Judge and Bock [17], Wallace [37]).

(ii) When $\alpha \rightarrow 1[0]$, $c \rightarrow 0[\infty]$, $P_{a_1 a_2 a_3} \rightarrow 0[1]$ for all

$$a_1, a_2, a_3 \text{ and } \rho \left[X\hat{\beta}, E(y) \right] \rightarrow \rho \left[X\bar{\beta}, E(y) \right] \left[\rho \left[X\beta^*, E(y) \right] \right].$$

(iii) When the null hypothesis is true ($\theta = 0$),

$$\rho \left[X\beta^*, E(y) \right] - \frac{\gamma(k-m)}{\gamma-2} < \rho \left[X\hat{\beta}, E(y) \right] - \frac{\gamma(k-mF_{20}^*)}{\gamma-2} < \rho \left[X\bar{\beta}, E(y) \right] - \frac{\gamma k}{\gamma-2}$$

$$\text{where } F_{a_1 a_2}^* = \Pr. \left[F_{(m+a_1, v+a_2)} \leq \left[\frac{cm(v+a_2)}{v(m+a_1)} \right] \right].$$

(iv) As $\theta \rightarrow \infty$, the risk of $X\beta^*$ is unbounded, while

$$\rho \left[X\hat{\beta}, E(y) \right] \rightarrow \rho \left[X\bar{\beta}, E(y) \right] \text{ as } P_{a_1 a_2 a_3} \rightarrow 0 \text{ for all } a_1, a_2, a_3.$$

(v) When $\theta = m\gamma/(\gamma-2)$, $\rho \left[X\bar{\beta}, E(y) \right] - \rho \left[X\beta^*, E(y) \right] \leq \rho \left[X\hat{\beta}, E(y) \right]$.

Equality of the three risk functions results when $2P_{202} -$

$$P_{402} - P_{201} = 0.$$

3.2 Numerical Evaluations of the Risk Functions

The risk function of the pre-test estimator depends on the data and the unknown parameters only through θ and so we have evaluated the risk expressions for various choices of γ , α , m , k and T .² Some representative results, for various γ values, appear in Figures 1 to 4.

Comparing the figures, we see that a decrease in the value of γ from the normal errors case ($\gamma = \infty$) causes the estimator risk

functions to shift upwards, a decrease in the rate at which the risk of the pre-test estimator approaches that of the unrestricted estimator, and an increase in the risk gain of the restricted estimator over the unrestricted estimator for all θ such that $\rho[X\beta^*, E(y)] < \rho[X\bar{\beta}, E(y)]$. For the unrestricted and the restricted estimators these changes occur because of the increase in the estimators' variances as γ decreases (the marginal distribution has fatter tails). For the pre-test estimator, the increase in its variance and its absolute bias (for relatively large θ) both contribute to the observed differences. Our numerical evaluations suggest that, in general, the difference between an estimator's risk under normality and Mt errors is relatively insignificant for a γ value of at least 100.

Comparing the risk functions of the pre-test estimator and its component estimators for a given γ , we find that the conclusions observed when the errors are normally distributed hold for all values of γ . When the null hypothesis is true the pre-test estimator is risk inferior to the restricted estimator but superior to the unrestricted estimator. However, the pre-test estimator is dominated by the unrestricted estimator over a wide range of θ , and by both of its component estimators over part of the parameter space.³ Hence, no one of the estimators strictly dominates the other two. This latter feature suggests choosing an estimator according to some optimality criterion. Such a study is beyond the scope of this paper (see, for example, Brook [3], Ohtani and Toyoda [27], Ohtani [24], Toyoda and Wallace [34]). So, aside from appropriate scaling, we find that the risk properties of the estimators are robust to the choice of γ , and

hence to the misspecification of the error term's distribution being considered here.

4. THE RISKS OF ALTERNATIVE ESTIMATORS OF σ_ϵ^2 .

Let the pre-test estimator of σ_ϵ^2 be given as

$$\hat{\sigma}_\epsilon^2 = \begin{cases} \bar{\sigma}_\epsilon^2 & ; \text{ if } F > c \\ \sigma_\epsilon^{*2} & ; \text{ if } F \leq c \end{cases}$$

$$= \bar{\sigma}_\epsilon^2 I_{(c, \infty)}(F) + \sigma_\epsilon^{*2} I_{[0, c]}(F)$$

where c , F and $I_{[a, b]}(F)$ are as previously defined. The risk function, under quadratic loss, of any estimator $\bar{\sigma}_\epsilon^2$ of σ_ϵ^2 is

$$\rho(\bar{\sigma}_\epsilon^2, \sigma_\epsilon^2) = E(\bar{\sigma}_\epsilon^2 - \sigma_\epsilon^2)^2 / \sigma_\epsilon^4,$$

and is the relative mean squared error of $\bar{\sigma}_\epsilon^2$. We now consider the risks of the various estimators of σ_ϵ^2 .

4.1 Finite Sample Risk Functions

THEOREM 2. *Under the stated assumptions,*

$$\rho(\bar{\sigma}_\epsilon^2, \sigma_\epsilon^2) = 2(v+\gamma-2)/[v(\gamma-4)] \quad (15)$$

$$\rho(\sigma_\epsilon^{*2}, \sigma_\epsilon^2) = 2[\gamma^2(v+m)(v+m+\gamma-2) + 4\theta\gamma(\gamma-2)(\gamma-4) + 2\theta^2(\gamma-2)^2(\gamma-4)] / [(v+m)^2\gamma^2(\gamma-4)] \quad (16)$$

$$\rho(\hat{\sigma}_\epsilon^2, \sigma_\epsilon^2) = \left\{ 2\gamma^2(v+m)^2(v+\gamma-2) + m\gamma^2 \left[- (v+2)(m+2v)(\gamma-2)P_{040} + v(\gamma-2)(m+2)P_{400} + 2v^2(\gamma-2)P_{220} + 2(\gamma-4)(v+m)v(P_{021} - P_{201}) \right] + 4\theta\gamma v(\gamma-2)(\gamma-4) \left[(m+2)P_{601} + vP_{421} - (v+m)P_{402} \right] + 4\theta^2 v(\gamma-2)^2(\gamma-4)P_{802} \right\} / \left[\gamma^2(\gamma-4)v(v+m)^2 \right]. \quad (17)$$

PROOF. See the Appendix. ■

Remarks:

- (i) When $\gamma \rightarrow \infty$, $\epsilon \sim \text{MN}(0, \sigma^2 I_T)$, $P_{a_1 a_2 a_3} \rightarrow P_{a_1 a_2}$, $\sigma_\epsilon^2 = \sigma^2$ and so, (15), (16) and (17) collapse to the risk functions derived by Clarke *et al.* [6] when the errors are normally distributed.
- (ii) $\rho(\hat{\sigma}_\epsilon^2, \sigma_\epsilon^2)$ converges to $\rho(\tilde{\sigma}_\epsilon^2, \sigma_\epsilon^2)$ when $\alpha \rightarrow 1$, and to $\rho(\sigma_\epsilon^{*2}, \sigma_\epsilon^2)$ when $\alpha \rightarrow 0$.
- (iii) $\rho(\hat{\sigma}_\epsilon^2, \sigma_\epsilon^2) \rightarrow \rho(\tilde{\sigma}_\epsilon^2, \sigma_\epsilon^2)$ as $\theta \rightarrow \infty$, while $\rho(\sigma_\epsilon^{*2}, \sigma_\epsilon^2)$ is unbounded.
- (iv) $\rho(\sigma_\epsilon^{*2}, \sigma_\epsilon^2) < \rho(\tilde{\sigma}_\epsilon^2, \sigma_\epsilon^2)$ when the restrictions are valid.

4.2 Numerical Evaluations of the Risk Functions

As in Section 3.2 we have evaluated the risk expressions, as functions of θ , for the same values of γ , α , m , and v as before. Figures 5 to 8 illustrate typical cases. Note that Figure 6, which considers the risk functions when $\gamma = 5$, is drawn on a different scale from that of Figures 5, 7 and 8, so as to enable the features for all four cases to be distinguishable. We have included the pre-test estimator with a critical value of 1 for two reasons. First, under the null, the risk of the pre-test estimator attains a local minimum when $c = 1$ (for all γ), and, in some situations is also the global minimum. We discuss this further below. Secondly, if one undertook the Stein-like strategy of using an estimator which always selects the minimum of $\tilde{\sigma}_\epsilon^2$ or σ_ϵ^{*2} then this corresponds to the pre-test estimator with $c = 1$.

Consider first Figure 5, which illustrates the risk functions of the estimators when the errors are normally distributed. We see that there exists a family of pre-test estimators, with $c \in (0, 1]$ which strictly dominate the unrestricted estimator for all

θ , and the restricted estimator over part of this parameter space.⁴ This feature is not observed in the evaluations undertaken by Clarke *et al.* [6] but is noted in subsequent work by Ohtani [24]. Ohtani [24] considers the question of the optimal significance level for the pre-test problem examined by Clarke *et al.* [6] when the component estimators are based on the minimum mean squared error principle. He compares the sampling properties of the pre-test estimator with those of the Stein [32] estimator (extended to the linear regression case) and shows first, that the Stein estimator can be written as a pre-test estimator with critical value $v/(v+2)$, and secondly, that the numerical evaluations suggest that the "Stein pre-test" estimator is optimal in some sense. Further, Clarke [4] shows that in certain situations the pre-test estimator, appropriately chosen, can strictly dominate both of its component estimators for all θ . She finds this typically occurs for small m , say 1 or 2, but does not appear to depend on the value of v . In these cases, the risk of the pre-test estimator with critical value unity corresponds to the global minimum risk at the origin and her numerical evaluations suggest this feature also holds for $\theta \neq 0$.⁵

Turning to the consequences of decreasing the value of γ from infinity, we find that the risk functions change in a similar way to that observed when estimating the prediction vector. That is, the estimator risk functions shift upwards, there is a decrease in the rate at which the risk of the pre-test estimator approaches that of the unrestricted estimator, and there is an increase in the risk gain of the restricted estimator over the unrestricted estimator for all θ such that $\rho(\sigma_{\epsilon}^{*2}, \sigma_{\epsilon}^2) < \rho(\bar{\sigma}_{\epsilon}^2, \sigma_{\epsilon}^2)$. These effects

occur because of the increase in the variances of all the estimators when γ decreases and changes in the bias functions of the restricted and pre-test estimators; the bias of the restricted estimator decreases for all θ while the bias function of the pre-test estimator shifts down for relatively small θ (which may increase or decrease absolute bias) but becomes unbiased at a slower rate.

In Section 3.2, when comparing the risk functions of an estimator of the prediction vector for different values of γ , we found that, in general, there was little difference between the normal risk functions and those when $\gamma = 100$. However, when estimating the error variance we find that relatively large differences may still be evident for this gamma value, and even for much larger values of γ , say 5,000 or 10,000.

Nevertheless, when comparing the risk functions of the pre-test estimator and its component estimators for a given γ , the conclusions for when the errors are normally distributed continue to hold for all values of γ . Namely, there exists a family of pre-test estimators, with $c \in (0,1]$ which strictly dominate the unrestricted estimator for all θ . Further, some members of this family,⁶ for some γ , also strictly dominate the restricted estimator. The numerical evaluations suggest that the pre-test estimator with critical value 1 strictly dominates all other members of this family.⁷

Moreover, for some γ , the restricted estimator is also strictly dominated by pre-test estimators with $1 < c < \infty$. Comparing equations (16) and (17) this will depend on m and v as well as γ . For the cases analyzed we found, in general, that the

restricted estimator is strictly dominated by all pre-test estimators, except for those with c around 0 and $c = \infty$, if γ is at most 15.

So, we have shown that, apart from appropriate scaling, the risk properties of the estimators are robust to the choice of γ . In particular, regardless of the value of γ , our recommendation is to pre-test rather than to impose the restrictions without testing their validity. Further, when using the least squares component estimators, a critical value of 1 seems to be the appropriate choice for the pre-test.

5. CONCLUSIONS

In this paper we have considered the sampling properties of various estimators of the parameters of the linear regression model, after a preliminary test of restrictions on the coefficients, when the error vector follows a multivariate Student-t distribution, but normality is wrongly assumed. The results suggest that, qualitatively, the estimator properties found for the model with normal errors carry over to this wider case. So, these properties are robust to such possible misspecification of the error distribution. These results are of interest to applied researchers working with data likely to follow fat-tailed empirical distributions.

FOOTNOTES

1. We consider estimation of the prediction vector rather than the location vector so that our results are independent of the design matrix.
2. We numerically evaluated the risk functions for $v = 10, 16, 20, 30$, $m = 1, 2, 3, 4, 5$, $\alpha = 0.01, 0.05, 0.25, 0.30, 0.50, 0.75, 0.90$ and that value associated with a critical value of unity, $\gamma = 5, 10, 50, 100, 500, 1000, 5000, 10,000, 100,000, \infty$, and $\theta \in [0,3(0.1);3,20(0.5)]$. Full results are available on request. All evaluations were carried out using double-precision FORTRAN on an AT computer. Davies' [8] algorithm was used to evaluate the $P_{a_1 a_2}$'s and the subroutines GAMMLN and BETAI from Press *et al.* [28] were utilized to obtain the $P_{a_1 a_2 a_3}$'s. Using these programs we found that the risk expressions were efficiently evaluated with no observed convergence problems.
3. Except when $2P_{202} - P_{402} - P_{201} = 0$. See Figure 2 for example.
4. That the risk of the pre-test estimator can dominate both of its components over any or all of the parameter space may seem counter intuitive. We may believe that as the pre-test estimator is a weighted sum of its component estimators then its risk function should be enveloped by those of its components. This, however, confuses the distinction between a weighted sum of the moments of the component estimators and the moments of their weighted sum.
5. The dominance of the pre-test estimator, for suitably chosen c , over the unrestricted estimator for all θ , and over the

restricted estimator for some or all θ , also occurs when estimating the error variance after a pre-test for homogeneity in the two-sample linear regression model. See, for example, Bancroft [1], Bancroft and Han [2], Clarke [4], Ohtani and Toyoda [27], and Toyoda and Wallace [34].

6. The exceptions are those pre-test estimators with critical value around the neighbourhood of $c = 0$.
7. It is straightforward to show that this feature holds under the null for any γ , but the proof for $\theta \neq 0$ is not obvious.

APPENDIX

The Proofs of THEOREMS 1. and 2. require the following lemma.

LEMMA A.1. Let $q^2 = \chi_\gamma^2$, $\lambda = q^2\theta/\gamma$, $\theta = (R\beta-r)' [R(X'X)^{-1}R']^{-1}(R\beta-r)/2\sigma^2$, and let $P_{a_1 a_2}$, given q^2 , be defined by

$$\begin{aligned} P_{a_1 a_2} &= \text{Pr.} \left[F' (m+a_1, v+a_2; \lambda) \leq \left[cm(v+a_2) \right] / \left[v(m+a_1) \right] \right] \\ &= \sum_{r=0}^{\infty} \lambda^r e^{-\lambda} (r!)^{-1} I_u \left[\frac{1}{2}(m+a_1)+r; \frac{1}{2}(v+a_2) \right], \end{aligned}$$

$I_u(\cdot; \cdot)$ is Pearson's incomplete beta function with $u = cm/(v+cm)$, and $a_1, a_2 = 0, 1, 2, \dots$. Then, for any real a_4, a_5 :

$$\begin{aligned} E \left[(q^2)^{a_4} \lambda^{a_5} P_{a_1 a_2} \right] &= \sum_{r=0}^{\infty} \frac{(2\theta/\gamma)^{r+a_5} \cdot 2^{a_4} \cdot \Gamma(\gamma/3+t+a_4+a_5)}{r! (1+2\theta/\gamma)^{\gamma/2+r+a_4+a_5} \Gamma(\gamma/2)} \\ &I_u \left[\frac{1}{2}(m+a_1)+r; \frac{1}{2}(v+a_2) \right] \end{aligned} \quad (\text{A.1})$$

PROOF.

$$\begin{aligned} E \left[(q^2)^{a_4} \lambda^{a_5} P_{a_1 a_2} \right] &= \int_0^{\infty} (q^2)^{a_4} \lambda^{a_5} \sum_{r=0}^{\infty} \frac{\lambda^r e^{-\lambda}}{r!} I_u \left[\frac{1}{2}(m+a_1)+r; \frac{1}{2}(v+a_2) \right] \\ &f(q^2) dq^2 \\ &= \sum_{r=0}^{\infty} (\theta/\gamma)^{r+a_5} \left[r! 2^{\gamma/2} \Gamma(\gamma/2) \right]^{-1} I_u \left[\frac{1}{2}(m+a_1)+r; \frac{1}{2}(v+a_2) \right] \\ &\int_0^{\infty} (q^2)^{\gamma/2+r+a_4+a_5-1} e^{-q^2(\gamma+2\theta)/2\gamma} dq^2, \end{aligned}$$

because $\lambda = q^2\theta/\gamma$, $I_u(\cdot; \cdot)$ does not depend on q^2 , and

$$f(q^2) = \left[2^{\gamma/2} \Gamma(\gamma/2) \right]^{-1} e^{-q^2/2} (q^2)^{\gamma/2-1}.$$

Applying the transformation $t = q^2(\gamma+2\theta)/(2\gamma)$, collecting terms

and noting that $\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt$, the result follows directly.

Now, to establish (13) of THEOREM 1, we have:

PROOF. $I_{(c,\infty)}(F) = 1 - I_{[0,c]}(F)$,

$$(X\hat{\beta}^* - X\bar{\beta}) = -X(X'X)^{-1}R' [R(X'X)^{-1}R']^{-1}(R\bar{\beta}-r),$$

$$\text{so, } X\hat{\beta} - X\beta = X\bar{\beta} - X\beta - I_{[0,c]}(F) \left[X(X'X)^{-1}R' [R(X'X)^{-1}R']^{-1}(R\bar{\beta}-r) \right]$$

$$\text{and, } \rho \left[X\hat{\beta}, E(y) \right] = \rho \left[X\bar{\beta}, E(y) \right] + E(I^*), \quad (\text{A.2})$$

$$\text{where } I^* = I_{[0,c]}(F) \left\{ 2(R\bar{\beta}-r)' [R(X'X)^{-1}R']^{-1}(R\beta-r) \right.$$

$$\left. - (R\bar{\beta}-r)' [R(X'X)^{-1}R']^{-1}(R\bar{\beta}-r) \right\} / \sigma^2.$$

Following Ullah and Phillips [35], we note that $R\bar{\beta}-r = R\beta-r + R(X'X)^{-1}X'\epsilon = R(\beta-\beta_0) + R(X'X)^{-1}X'\epsilon$ where $\beta_0 = R\bar{r}$ is any solution of $R\beta_0 = r$. Further, let

$$B = [R(X'X)^{-1}R']^{-1}, \quad A = R(X'X)^{-1}X', \quad C = A'BA - C^2,$$

$$M = I_T - X(X'X)^{-1}X' = M^2, \quad \text{and } \delta = X(\beta-\beta_0). \quad (\text{A.3})$$

So, noting that $AX = R$ and $MC = 0$, we have $F = v(\epsilon+\delta)'C(\epsilon+\delta)/(m\epsilon'M\epsilon)$ and

$$I^* = I_{[0,c]} \left[v(\epsilon+\delta)'C(\epsilon+\delta)/(m\epsilon'M\epsilon) \right] \left\{ 2(\epsilon+\delta)'C\delta - (\epsilon+\delta)'C(\epsilon+\delta) \right\} / \sigma^2 \\ - I_{[0,c]} \left[(vb'Cb)/(mb'Mb) \right] \left\{ \gamma(2b'Cb - b'Cb)/q^2 \right\} / \sigma^2, \quad (\text{A.4})$$

where $b = a + \delta_1$, $\delta_1 = q\delta/\sqrt{\gamma}$ and use is made of decomposition (6), $\epsilon = \sqrt{\gamma}a/q$. Now, observe that, given q^2 ,

$$b = MN(\delta_1, \sigma^2 I_T), \quad \frac{b'Mb}{\sigma^2} = \frac{a'Ma}{\sigma^2} = \chi_v^2, \quad \text{and } \frac{b'Cb}{\sigma^2} = \chi_m^2; \lambda \quad (\text{A.5})$$

where $\lambda = \delta_1^* C \delta_1 / \sigma^2 = q^2 / \gamma$. Further, given q^2 , the quadratic forms $b' M b / \sigma^2$ and $b' C b / \sigma^2$ are independent and so, taking expectations of (A.4), conditional on q^2 , we have

$$\begin{aligned} E(I^* | q^2) &= \gamma \left\{ 2E \left[I_{[0,c]} \left[(v b' C b) / (m b' M b) \right] (b' C \delta_1) | q^2 \right] \right. \\ &\quad \left. - E \left[I_{[0,c]} \left[(v b' C b) / (m b' M b) \right] (b' C b) | q^2 \right] \right\} / q^2. \end{aligned} \quad (A.6)$$

First, using Lemma 1. of Clarke *et al.* [5] gives,

$$E \left[I_{[0,c]} \left[(v b' C b) / (m b' M b) \right] (b' C b) | q^2 \right] = \sigma^2 (m P_{20} + 2 \lambda P_{40}). \quad (A.7)$$

Secondly, we can write

$$\begin{aligned} E \left[I_{[0,c]} \left[(v b' C b) / (m b' M b) \right] (b' C \delta_1) | q^2 \right] &= \\ E \left[I_{[0,c]} \left[(v b^* b^*) / (m b' M b) \right] (b^* \delta_1^*) | q^2 \right] & \end{aligned}$$

where $b^* = L' b$, $\delta_1^* = L' \delta_1$ and $LL' = C$. Note that as C is symmetric and idempotent, $L'L = I_m$ and so, given q^2 , $b^* = MN(\delta_1^*, \sigma^2 I_m)$. Then, using Theorem 1 of Judge and Bock [17, p.321], noting that $\lambda = \delta_1^* \delta_1^* / 2\sigma^2$, we have

$$E \left[I_{[0,c]} \left[(v b' C b) / (m b' M b) \right] (b' C \delta_1) | q^2 \right] = 2\sigma^2 \lambda P_{20}. \quad (A.8)$$

Substituting (A.7) and (A.8) appropriately in (A.6) yields

$$E(I^* | q^2) = \gamma [(4\lambda - m) P_{20} - 2\lambda P_{40}] / q^2.$$

The unconditional expectation then follows as

$$\begin{aligned} E(I^*) &= \int_0^\infty E(I^* | q^2) f(q^2) dq^2 \\ &= \gamma \left\{ 4E \left[(q^2)^{-1} \lambda P_{20} \right] - mE \left[(q^2)^{-1} P_{20} \right] - 2E \left[(q^2)^{-1} \lambda P_{40} \right] \right\} \\ &= \left[2\theta(\gamma - 2)(2P_{202} - P_{402}) - m\gamma P_{201} \right] / (\gamma - 2) \end{aligned} \quad (A.9)$$

using LEMMA 1 repeatedly and defining

$$P_{a_1 a_2 a_3} = \sum_{r=0}^{\infty} \frac{(2\theta/\gamma)^r \Gamma(\gamma/2+r+a_3-2)}{r!(1+2\theta/\gamma) \gamma^{2+r+a_3-2} \Gamma(\gamma/2+a_3-2)} \cdot I_u \left[\frac{1}{2}(m+a_1)+r; \frac{1}{2}(v+a_2) \right]. \quad (A.10)$$

Substituting (A.8) in (A.2) yields $\rho \left[\hat{X}\beta, E(y) \right]$ as stated. \blacksquare

Turning now to THEOREM 2 and considering first $\rho(\tilde{\sigma}_\epsilon^2, \sigma_\epsilon^2)$ we have

$$\tilde{\sigma}_\epsilon^2 = \epsilon' M \epsilon / v - \gamma a' M a / (v q^2)$$

So,

$$\rho(\tilde{\sigma}_\epsilon^2, \sigma_\epsilon^2) = E \left[(\gamma-2)^2 (a' M a / \sigma^2)^2 + v^2 (q^2)^2 - 2(\gamma-2) v q^2 (a' M a / \sigma^2) \right] / [v^2 (q^2)^2].$$

Now, given q^2 , $a' M a / \sigma^2 = \chi_v^2$ and, therefore, the risk of $\tilde{\sigma}_\epsilon^2$ conditional on q^2 is

$$\rho(\tilde{\sigma}_\epsilon^2, \sigma_\epsilon^2 | q^2) = \left[(\gamma-2)^2 (v+2) + v (q^2)^2 - 2(\gamma-2) v q^2 \right] / [v (q^2)^2].$$

The unconditional risk is then obtained by noting that

$$\rho(\tilde{\sigma}_\epsilon^2, \sigma_\epsilon^2) = \int_0^\infty \rho(\tilde{\sigma}_\epsilon^2, \sigma_\epsilon^2 | q^2) f(q^2) dq^2,$$

$$E(1/q^2) = (\gamma-2)^{-1}, \text{ and } E(1/q^2)^2 = [(\gamma-2)(\gamma-4)]^{-1}. \quad (A.11)$$

Secondly, by a similar argument,

$$\begin{aligned} \sigma_\epsilon^{*2} &= [\epsilon' M \epsilon + (R\bar{\beta} - r)' [R(X'X)^{-1}R']^{-1} (R\bar{\beta} - r)] / (v+m) \\ &\quad - \gamma [b' M b + b' C b] / [(v+m)q^2] \end{aligned}$$

using previous notation. So,

$$\begin{aligned} \rho(\sigma_\epsilon^{*2}, \sigma_\epsilon^2) &= E \left\{ (\gamma-2)^2 [(b' M b + b' C b) / \sigma^2]^2 + (q^2)^2 (v+m)^2 - \right. \\ &\quad \left. 2(v+m)q^2(\gamma-2) [(b' M b + b' C b) / \sigma^2] \right\} / [(q^2)^2 (v+m)^2] \end{aligned}$$

and, as $(b' M b + b' C b) / \sigma^2 = \chi_{m+v; \lambda}^2$, given q^2 , we have

$$\begin{aligned} \rho(\sigma_\epsilon^{*2}, \sigma_\epsilon^2 | q^2) &= \left\{ (\gamma-2)^2 [(v+m)(v+m+2) + 4\lambda(v+m+2) + 4\lambda^2] \right. \\ &\quad \left. + (q^2)^2 (v+m)^2 - 2(v+m)q^2(\gamma-2)(v+m+2\lambda) \right\} / [(q^2)^2 (v+m)^2]. \end{aligned}$$

The unconditional risk is then obtained by integrating over q^2 , noting that $\lambda = q^2\theta/\gamma$.

Thirdly, to derive $\rho(\hat{\sigma}_\epsilon^2, \sigma_\epsilon^2)$ we write,

$$\begin{aligned} \hat{\sigma}_\epsilon^2 - \bar{\sigma}_\epsilon^2 + (\sigma_\epsilon^{*2} - \bar{\sigma}_\epsilon^2) I_{[0, c]}(F) \\ - \gamma \left\{ (v+m)(b'Mb) + [vb'Cb - mb'Mb] I_{[0, c]} \left[(vb'Cb)/(mb'Mb) \right] \right\} / \\ [q^2 v(v+m)] \end{aligned}$$

and so,

$$\begin{aligned} \rho(\hat{\sigma}_\epsilon^2, \sigma_\epsilon^2) - E \left\{ (\gamma-2)^2 (v+m)^2 (b'Mb/\sigma^2)^2 + (q^2)^2 v^2 (v+m)^2 \right. \\ - 2q^2 v(v+m)^2 (\gamma-2) (b'Mb/\sigma^2) + [(\gamma-2)^2 [v^2 (b'Cb/\sigma^2)^2 \\ + 2v^2 (b'Mb/\sigma^2)(b'Cb/\sigma^2) - m(m+2v)(b'Mb/\sigma^2)^2] \\ \left. - 2q^2 v(v+m)(\gamma-2)(vb'Cb/\sigma^2 - mb'Mb/\sigma^2) \right\} I_{[0, c]} \left[(vb'Cb)/(mb'Mb) \right] \\ / [q^2 v(v+m)]^2. \end{aligned}$$

Then, using (A.5) and the results from the Appendix of Clarke *et al.* [5], we have

$$\begin{aligned} \rho(\hat{\sigma}_\epsilon^2, \sigma_\epsilon^2 | q^2) - \left\{ (v+m)^2 [(v+2)(\gamma-2)^2 + (q^2)^2 v - 2q^2 v(\gamma-2)] \right. \\ + (\gamma-2)^2 [mv(m+2)P_{40} - m(v+2)(m+2v)P_{04} + 2mv^2 P_{22}] \\ - 2mvq^2 (v+m)(\gamma-2)(P_{20} - P_{02}) + 4v\lambda(\gamma-2)[(m+2)(\gamma-2)P_{60} \\ \left. + v(\gamma-2)P_{42} - (v+m)q^2 P_{40}] + 4v\lambda^2 (\gamma-2)^2 P_{80} \right\} / \\ [(q^2)^2 v(v+m)^2] \end{aligned}$$

and $\rho(\hat{\sigma}_\epsilon^2, \sigma_\epsilon^2)$ follows by integrating over q^2 using LEMMA 1, (A.10)

and (A.11) appropriately. ■

REFERENCES

1. Bancroft, T.A. On biases in estimation due to the use of preliminary tests of significance. *Annals of Mathematical Statistics* 15 (1944): 190-204.
2. Bancroft, T.A. & C. Han. A note on pooling variances. *Journal of the American Statistical Association* 78 (1983): 981-983.
3. Brook, R.J. On the use of a regret function to set significance points in prior tests of estimation. *Journal of the American Statistical Association* 71 (1976): 126-131.
4. Clarke, J.A. The optimality of pre-testing. Department of Economics: University of Canterbury, Christchurch, 1988.
5. Clarke, J.A., D.E.A. Giles & T.D. Wallace. Estimating the error variance in regression after a preliminary test of restrictions on the coefficients. *Journal of Econometrics* 34 (1987): 293-304.
6. Clarke, J.A., D.E.A. Giles, and T.D. Wallace. Preliminary test estimation of the error variance in linear regression. *Econometric Theory* 3 (1987): 299-304.
7. Coursey, D. & H. Nyquist. Applications of robust estimation techniques in demand analysis. *Applied Economics* 20 (1988): 595-610.
8. Davies, R.B. The distribution of a linear combination of χ^2 random variables (Algorithm AS 155). *Applied Statistics* 29 (1980): 323-333.
9. Fama, E.F. Mandelbrot and the stable Paretian hypothesis. *Journal of Business* 36 (1963): 420-449.

10. Fama, E.F. The behaviour of stock market prices. *Journal of Business* 38 (1965): 34-105.
11. Fama, E.F. Efficient capital markets: a review of theory and empirical work. *Journal of Finance* 25 (1970): 383-417.
12. Giles, D.E.A. Preliminary-test estimation in mis-specified regressions. *Economics Letters* 21 (1986): 325-328.
13. Giles, D.E.A. and J.A. Clarke. Preliminary-test estimation of the scale parameter in a mis-specified regression model. Department of Economics: University of Canterbury, Christchurch, 1987.
14. Granger, C.W.J. & D. Orr. Infinite variance and research strategy in time series analysis. *Journal of the American Statistical Association* 67 (1972): 275-285.
15. Huber, P.J. *Robust Statistical Procedures*. Philadelphia: SIAM, 1977.
16. James, W. & C. Stein. Estimation with quadratic loss. In *Proceedings of the Fourth Berkeley Symposium*, Vol.1, pp.361-379. Berkeley: University of California, 1961.
17. Judge, G.G. & M.E. Bock. *The Statistical Implications of Pre-Test and Stein-Rule Estimators in Econometrics*. Amsterdam: North-Holland, 1978.
18. Judge, G.G., W. Griffiths, C. Hill, H. Lutkepohl, & T.C. Lee. *The Theory and Practice of Econometrics*, 2nd ed. New York: Wiley, 1985.
19. Judge, G.G., S. Miyazaki, & T. Yancey. Minimax estimators for the location vectors of spherically symmetric densities. *Econometric Theory* 1 (1985): 409-417.

20. King, M.L. Robust tests for spherical symmetry and their application to least squares regression. *Annals of Statistics* 8 (1980): 1265-1271.
21. Koenker, R.W. Robust methods in Econometrics. *Econometric Reviews* 1 (1982): 213-255.
22. Mittelhammer, R.C. Restricted least squares, pre-test, OLS and Stein rule estimators: risk comparisons under model mis-specification. *Journal of Econometrics* 15 (1984): 151-164.
23. Nimmo-Smith, I. Linear regressions and sphericity. *Biometrika* 66 (1979): 390-392.
24. Ohtani, K. Optimal levels of significance of a pre-test in estimating the disturbance variance after the pre-test for a linear hypothesis on coefficients in a linear regression. Faculty of Economics: Kobe University, Kobe, 1988.
25. Ohtani, K. Some small sample properties of a pre-test estimator of the disturbance variance in a mis-specified linear regression. *Journal of the Japan Statistical Society* 17 (1987): 81-89.
26. Ohtani, K. Preliminary test predictor in the linear regression model including a proxy variable. *Journal of the Japan Statistical Society* 13 (1983): 11-19.
27. Ohtani, K. & T. Toyoda. Minimax regret critical values for a preliminary test in pooling variances. *Journal of the Japan Statistical Society* 8 (1978): 15-20.
28. Press, W.H., B.P. Flannery, S.A. Teukolsky, & W.T. Vetterling. *Numerical Recipes: The Art of Scientific Computing*. New York: Cambridge University Press, 1986.

29. Sharpe, W.F. Mean-absolute-deviation characteristic lines for securities and portfolios. *Management Science* 18 (1971): B1-B13.
30. Singh, R.S. A family of improved estimators in linear regression models with errors having multivariate Student-t distribution. Department of Econometrics: Monash University, Melbourne, 1987.
31. Singh, R.S. Estimation of error variance in linear regression models with errors having multivariate Student-t distribution with unknown degrees of freedom. Department of Econometrics: Monash University, Melbourne, 1987.
32. Stein, C. Inadmissibility of the usual estimator for the variance of a normal distribution with unknown mean. *Annals of the Institute of Statistical Mathematics* 16 (1964): 155-160.
33. Thomas, D.H. Some contributions to radial probability distributions, statistics, and the operational calculi. Unpublished Ph.D. thesis. Wayne State University, Michigan.
34. Toyoda, T. & T.D. Wallace. Estimation after a preliminary test of homogeneity and optimal levels of significance for the pre-test. *Journal of Econometrics* 3 (1975): 395-404.
35. Ullah, A. & P.C.B. Phillips. Distribution of the F-ratio. *Econometric Theory* 2 (1986): 449-452.
36. Ullah, A. & V. Zinde-Walsh. On the robustness of LM, LR and W tests in regression models. *Econometrica* 52 (1984): 1055-1066.

37. Wallace, T.D. Pre-test estimation in regression: a survey.
American Journal of Agricultural Economics 59 (1977):
431-444.
38. Zellner, A. Bayesian and non-Bayesian analysis of the
regression model with multivariate Student-t error terms.
Journal of the American Statistical Association 71 (1976):
400-405.
39. Zellner, A. *An Introduction to Bayesian Inference in
Econometrics*. New York: Wiley, 1971.




FIGURE 1. Risk functions for the unrestricted, restricted and pre-test estimators of $E(y)$ when $T = 30$, $k = 5$, $m = 3$, and $\gamma = \omega$.




FIGURE 2. Risk functions for the unrestricted, restricted and pre-test estimators of $E(y)$ when $T = 30$, $k = 5$, $m = 3$ and $\gamma = 5$.




FIGURE 3. Risk functions for the unrestricted, restricted and pre-test estimators of $E(y)$ when $T = 30$, $k = 5$, $m = 3$ and $\gamma = 10$.




FIGURE 4. Risk functions for the unrestricted, restricted and pre-test estimators of $E(y)$ when $T = 30$, $k = 5$, $m = 3$, and $\gamma = 100$.




FIGURE 5. Risk functions for the unrestricted, restricted and pre-test estimators of σ_ϵ^2 when $T = 20, k = 4, m = 2,$ and $\gamma = \infty$.




FIGURE 6. Risk functions for the unrestricted, restricted and pre-test estimators of σ_ϵ^2 when $T = 20, k = 4, m = 2,$ and $\gamma = 5$.




FIGURE 7. Risk function for the unrestricted, restricted and pre-test estimators of σ_ϵ^2 when $T = 20, k = 4, m = 2, \text{ and } \gamma = 10$.




FIGURE 8. Risk functions for the unrestricted, restricted, and pre-test estimators of σ_ϵ^2 when $T = 20, k = 4, m = 2, \text{ and } \gamma = 100$.

LIST OF DISCUSSION PAPERS*

- No. 8401 Optimal Search, by Peter B. Morgan and Richard Manning.
- No. 8402 Regional Production Relationships During the Industrialization of New Zealand, 1935-1948, by David E. A. Giles and Peter Hampton.
- No. 8403 Pricing Strategies for a Non-Replenishable Item Under Variable Demand and Inflation, by John A. George.
- No. 8404 Alienation Rights in Traditional Maori Society, by Brent Layton.
- No. 8405 An Engel Curve Analysis of Household Expenditure in New Zealand, by David E. A. Giles and Peter Hampton.
- No. 8406 Paying for Public Inputs, by Richard Manning, James R. Markusen, and John McMillan.
- No. 8501 Perfectly Discriminatory Policies in International Trade, by Richard Manning and Koon-Lam Shea.
- No. 8502 Perfectly Discriminatory Policy Towards International Capital Movements in a Dynamic World, by Richard Manning and Koon-Lam Shea.
- No. 8503 A Regional Consumer Demand Model for New Zealand, by David E. A. Giles and Peter Hampton.
- No. 8504 Optimal Human and Physical Capital Accumulation in a Fixed-Coefficients Economy, by R. Manning.
- No. 8601 Estimating the Error Variance in Regression After a Preliminary Test of Restrictions on the Coefficients, by David E. A. Giles, Judith A. Mikolajczyk and T. Dudley Wallace.
- No. 8602 Search While Consuming, by Richard Manning.
- No. 8603 Implementing Computable General Equilibrium Models: Data Preparation, Calibration, and Replication, by K. R. Henry, R. Manning, E. McCann and A. E. Woodfield.
- No. 8604 Credit Rationing: A Further Remark, by John G. Riley.
- No. 8605 Preliminary-Test Estimation in Mis-Specified Regressions, by David E. A. Giles.
- No. 8606 The Positive-Part Stein-Rule Estimator and Tests of Linear Hypotheses, by Aman Ullah and David E. A. Giles.
- No. 8607 Production Functions that are Consistent with an Arbitrary Production-Possibility Frontier, by Richard Manning.
- No. 8608 Preliminary-Test Estimation of the Error Variance in Linear Regression, by Judith A. Clarke, David E. A. Giles and T. Dudley Wallace.
- No. 8609 Dual Dynamic Programming for Linear Production/Inventory Systems, by E. Grant Read and John A. George.
- No. 8610 Ownership Concentration and the Efficiency of Monopoly, by R. Manning.
- No. 8701 Stochastic Simulation of the Reserve Bank's Model of the New Zealand Economy, by J. N. Lye.
- No. 8702 Urban Expenditure Patterns in New Zealand, by Peter Hampton and David E. A. Giles.
- No. 8703 Preliminary-Test Estimation of Mis-Specified Regression Models, by David E. A. Giles.
- No. 8704 Instrumental Variables Regression Without an Intercept, by David E. A. Giles and Robin W. Harrison.

(continued on back cover)

- No. 8705 Household Expenditure in Sri Lanka: An Engel Curve Analysis, by Mallika Dissanayake and David E. A. Giles.
- No. 8706 Preliminary-Test Estimation of the Standard Error of Estimate in Linear Regression, by Judith A. Clarke.
- No. 8707 Invariance Results for FIML Estimation of an Integrated Model of Expenditure and Portfolio Behaviour, by P. Dorian Owen.
- No. 8708 Social Cost and Benefit as a Basis for Industry Regulation with Special Reference to the Tobacco Industry, by Alan E. Woodfield.
- No. 8709 The Estimation of Allocation Models With Autocorrelated Disturbances, by David E. A. Giles.
- No. 8710 Aggregate Demand Curves in General-Equilibrium Macroeconomic Models: Comparisons with Partial-Equilibrium Microeconomic Demand Curves, by P. Dorian Owen.
- No. 8711 Alternative Aggregate Demand Functions in Macro-economics: A Comment, by P. Dorian Owen.
- No. 8712 Evaluation of the Two-Stage Least Squares Distribution Function by Imhof's Procedure by P. Cribbett, J. N. Lye and A. Ullah.
- No. 8713 The Size of the Underground Economy: Problems and Evidence, by Michael Carter.
- No. 8714 A Computable General Equilibrium Model of a Fisherine Method to Close the Foreign Sector, by Ewen McCann and Keith McLaren.
- No. 8715 Preliminary-Test Estimation of the Scale Parameter in a Mis-Specified Regression Model, by David E. A. Giles and Judith A. Clarke.
- No. 8716 A Simple Graphical Proof of Arrow's Impossibility Theorem, by John Fountain.
- No. 8717 Rational Choice and Implementation of Social Decision Functions, by Manimay Sen.
- No. 8718 Divisia Monetary Aggregates for New Zealand, by Ewen McCann and David E. A. Giles.
- No. 8719 Telecommunications in New Zealand: The Case for Reform, by John Fountain.
- No. 8801 Workers' Compensation Rates and the Demand for Apprentices and Non-Apprentices in Victoria, by Pasquale M. Sgro and David E. A. Giles.
- No. 8802 The Adventures of Sherlock Holmes, the 48% Solution, by Michael Carter.
- No. 8803 The Exact Distribution of a Simple Pre-Test Estimator, by David E. A. Giles.
- No. 8804 Pre-testing for Linear Restrictions in a Regression Model With Student-t Errors, by Judith A. Clarke.

* Copies of these Discussion Papers may be obtained for \$4 (including postage, price changes occasionally) each by writing to the Secretary, Department of Economics, University of Canterbury, Christchurch, New Zealand.