AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

CANTER

8907✓

Department of Economics
UNIVERSITY OF CANTERBURY

CHRISTCHURCH, NEW ZEALAND

GRANTING FOUNDATION OF
AGRICULTURAL ECONOMICS
LIBRARY

1 1990

**UNBIASED ESTIMATION OF THE
MEAN SQUARED ERROR
FEASIBLE GENERALISED RIDGE REGRESSION ESTIMATOR**

V. K. SRIVASTAVA AND D. E. A. GILES

Discussion Paper

No.8907

Department of Economics University of Canterbury
Christchurch, New Zealand

Discussion Paper No. 8907

October 1989

**UNBIASED ESTIMATION OF THE
MEAN SQUARED ERROR
FEASIBLE GENERALISED RIDGE REGRESSION ESTIMATOR**

**V. K. SRIVASTAVA
University of Lucknow
and
D. E. A. GILES
University of Canterbury**

*This paper is circulated for discussion and comments. It should not be quoted without the prior approval of the author.

Address for Correspondence:

Professor D. E. A. Giles, Department of Economics, University of Canterbury, Christchurch 1, New Zealand.

UNBIASED ESTIMATION OF THE
MEAN SQUARED ERROR OF THE
FEASIBLE GENERALISED RIDGE REGRESSION

ESTIMATOR

V.K. Srivastava
Department of Statistics
Lucknow University
Lucknow 226007
INDIA

AND

D.E.A. Giles
Department of Economics
University of Canterbury
Christchurch 1
NEW ZEALAND

Abstract

The paper derives an exact unbiased estimator of the mean squared error of the feasible generalised ridge regression estimator for a linear regression coefficient. This provides the basis for calculating the standard error of such an estimator in a meaningful way.

Keywords:

Ridge regression; estimated efficiency; multicollinearity

1. INTRODUCTION

Hoerl and Kennard (1970) suggest the method of ridge regression to deal with the problem of multicollinear data. Subsequently, a variety of ridge and ridge-type estimators have been proposed and their properties derived and compared. However, the practical application of these estimators has been limited by the fact that they produce only point estimates - in general, appropriate formulae for computing the associated "standard errors" are unavailable.

The reason for this is that the expressions for the exact first and second moments of these estimators are extremely complex, and obtaining unbiased estimators of these moments (as are needed to derive standard errors which are meaningful in finite samples) seems to be a non-trivial task.

Although asymptotic standard errors for the feasible ridge regression estimator are easily obtained, they are of limited use. In particular, as they coincide with the least squares standard errors, no distinction can be drawn between the relative estimated efficiencies of the two estimators if this approach is pursued.

In the spirit of results obtained for the Stein-rule estimator by Carter et al. (19**), this paper derives an unbiased estimator of the exact mean squared error of the feasible ridge estimator. The result obtained is of a simple form and is readily computed. This facilitates the computation of meaningful standard errors which have a proper finite-sample justification. We deal only with the generalised feasible ridge estimator, but in principle the same analytical approach could be applied to other related estimators.

2. GENERALISED RIDGE REGRESSION

Consider the model

$$(2.1) \quad y = X\beta + u ; \quad u \sim N(0, \sigma^2 I_n)$$

where X is $(n \times p)$, and for convenience (and without loss of generality) we take the model to be in canonical form, so that $X'X = \Lambda = \text{diag.}(\lambda_i)$. The Ordinary Least Squares (OLS) estimator of β is

$$(2.2) \quad b = (X'X)^{-1}X'y = \Lambda^{-1}X'y$$

and this unbiased estimator has covariance matrix

$$(2.3) \quad V(b) = \sigma^2 \Lambda^{-1},$$

for which an unbiased estimator is

$$(2.4) \quad \hat{V}(b) = s\Lambda^{-1}/\nu,$$

where $\nu = (n-p)$ denotes degrees of freedom, and $s = (y - Xb)'(y - Xb)$ is the residual sum of squares.

The generalised ridge regression estimator of β is

$$(2.5) \quad \hat{\beta} = (\Lambda + K)^{-1}X'y,$$

where $K = \text{diag.}(k_i)$, and the k_i 's are biasing parameters. The i th element of $\hat{\beta}$ is

$$(2.6) \quad \hat{\beta}_i = \lambda_i b_i / (\lambda_i + k_i),$$

where b_i is the i th element of b .

If k_i is fixed then $\hat{\beta}_i$ has bias given by

$$(2.7) \quad B(\hat{\beta}_i) = - \left(\frac{k_i \beta_i}{\lambda_i + k_i} \right)$$

and Mean Squared Error (MSE):

$$(2.8) \quad M(\hat{\beta}_i) = \frac{\lambda_i \sigma^2 + k_i^2 \beta_i^2}{(\lambda_i + k_i)^2}.$$

In this case, the choice of k_i which minimizes the MSE of $\hat{\beta}_i$ is $k_i = (\sigma^2/\beta_i^2)$. Hoerl and Kennard (1970) suggest a feasible version of this optimal ridge estimator:

$$(2.9) \quad \hat{\beta}_i^* = \left[\lambda_i b_i^2 / (\lambda_i b_i^2 + s/\nu) \right] b_i .$$

Clearly, $\hat{\beta}_i^*$ no longer minimises MSE - some of its exact finite sample properties are derived by Dwivedi et al. (1980).

For applied work, a major limitation of $\hat{\beta}_i^*$ is that it provides only a point estimate of the regression coefficient. This problem is resolved in various ad hoc ways in practice (e.g. Judge et al. (1988, pp.878-882)). One possibility is to use the OLS standard error $(s/\nu\lambda_i)^{\frac{1}{2}}$ as a standard error for $\hat{\beta}_i^*$. This is legitimate asymptotically, but has no finite-sample justification and provides no information about the gains or losses in measured efficiency when ridge regression is used in preference to OLS. Another approach that is sometimes adopted involves using estimators of β_i and σ^2 to estimate k_i (as in (2.9)) and hence (2.7) and (2.8). Although "standard errors" for the $\hat{\beta}_i^*$ estimators can be generated in this way, their obvious limitation is that (2.7) and (2.8) are no longer the correct expressions for the estimator's bias and mean squared error once k_i is estimated.

Complete inference based on the feasible ridge estimator requires knowledge of its sampling distribution. Dwivedi et al. (1980) derive the first two moments of $\hat{\beta}_i^*$, but the expressions concerned are complex and unobservable. However, if unbiased estimators of these two moments can be derived, then we have a legitimate basis for constructing ridge regression standard

errors. This task is considered in the next section. Our objective is to provide exact finite sample measures, and hence resolve the difficulties currently facing an applied researcher using feasible generalised ridge regression.

3. UNBIASED ESTIMATION OF THE BIAS AND MEAN SQUARED ERROR

From (2.9), we observe that the bias of $\hat{\beta}_i^*$ is

$$\begin{aligned} (3.1) \quad B(\hat{\beta}_i^*) &= E \left[(b_i - \beta_i) - \frac{sb_i}{\nu\lambda_i b_i^2 + s} \right] \\ &= -E \left[\frac{sb_i}{\nu\lambda_i b_i^2 + s} \right], \end{aligned}$$

from which it follows that an unbiased estimator of the bias is

$$(3.2) \quad \hat{B}(\hat{\beta}_i^*) = - \left(\frac{b_i}{\nu f_i + 1} \right)$$

where $f_i = \left(\frac{\lambda_i b_i^2}{s} \right)$. It may be noticed that unbiased estimation of the bias does not require normality of the disturbances.

Similarly, the mean squared error of $\hat{\beta}_i^*$ is

$$\begin{aligned} (3.3) \quad M(\hat{\beta}_i^*) &= E(b_i - \beta_i)^2 + E \left[\frac{sb_i}{\nu\lambda_i b_i^2 + s} \right]^2 - 2E \left[\frac{sb_i(b_i - \beta_i)}{\nu\lambda_i b_i^2 + s} \right] \\ &= \frac{\sigma^2}{\lambda_i} + E \left[\frac{sb_i}{\nu\lambda_i b_i^2 + s} \right]^2 - 2E \left[\frac{sb_i(b_i - \beta_i)}{\nu\lambda_i b_i^2 + s} \right]. \end{aligned}$$

An unbiased estimator of the first term on the right hand side of (3.3) is

$$(3.4) \quad \left(\frac{s}{\nu \lambda_i} \right) ,$$

while an unbiased estimator for the second term is

$$(3.5) \quad \left(\frac{s}{\nu \lambda_i} \right) \left[\frac{\nu f_i}{(\nu f_i + 1)^2} \right].$$

For the third term, we observe that b_i has a normal distribution with mean β_i and variance (σ^2/λ_i) while (s/σ^2) has a χ^2 -distribution with ν degrees of freedom, independent of b_i . Using this result and integrating by parts:

$$(3.6) \quad E \left[\frac{s b_i (b_i - \beta_i)}{\nu \lambda_i b_i^2 + s} \right] = \frac{\sigma^2}{\lambda_i} E \left[s \frac{\partial}{\partial b_i} \left(\frac{b_i}{\nu \lambda_i b_i^2 + s} \right) \right]$$

$$= - \frac{\sigma^2}{\lambda_i} E \left[\frac{s (\nu \lambda_i b_i^2 - s)}{(\nu \lambda_i b_i^2 + s)^2} \right].$$

Now suppose that $g(s, b_i)$ is absolutely continuous in s and is such that

$$(3.7) \quad E[s g(s, b_i)] = E \left[\frac{s b_i (b_i - \beta_i)}{\nu \lambda_i b_i^2 + s} \right].$$

Integrating the left side of (3.7) by parts and using (3.6), we get

$$\sigma^2 E \left[\nu g(s, b_i) + 2s \frac{\partial g(s, b_i)}{\partial s} \right] = E \left[\frac{s b_i (b_i - \beta_i)}{\nu \lambda_i b_i^2 + s} \right]$$

$$= - \frac{\sigma^2}{\lambda_i} E \left[\frac{s (\nu \lambda_i b_i^2 - s)}{(\nu \lambda_i b_i^2 + s)^2} \right]$$

or

$$E \left[\nu g(s, b_i) + 2s \frac{\partial g(s, b_i)}{\partial s} + \frac{s (\nu \lambda_i b_i^2 - s)}{\lambda_i (\nu \lambda_i b_i^2 + s)^2} \right] = 0.$$

As (s, b_i) are jointly complete sufficient statistics for (σ^2, β_i) , this last equation suggests

$$(3.9) \quad \nu g(s, b_i) + 2s \frac{\partial g(s, b_i)}{\partial s} + \frac{s (\nu \lambda_i b_i^2 - s)}{\lambda_i (\nu \lambda_i b_i^2 + s)^2} = 0.$$

This is a linear differential equation of first order and first degree, with solution

$$(3.10) \quad g(s, b_i) = - \left(\frac{1}{2} \right) s^{-\frac{\nu}{2}} \int_0^s \frac{(\nu \lambda_i b_i^2 - x) x^{\frac{\nu}{2}}}{\lambda_i (\nu \lambda_i b_i^2 + x)^2} dx$$

$$= - \left(\frac{1}{2 \lambda_i} \right) \int_0^1 \frac{(\nu f_i - x) x^{\frac{\nu}{2}}}{(\nu f_i + x)^2} dx,$$

which provides the functional form of $g(s, b_i)$.

Substituting (3.10) in (3.7) we observe that an unbiased estimator of the third term on the right hand side of (3.3) is

$$(3.11) \quad - \left(\frac{s}{\lambda_i} \right) \int_0^1 \left[\frac{(\nu f_i - x) x^{\frac{\nu}{2}}}{(\nu f_i + x)^2} \right] dx.$$

Using (3.4), (3.5) and (3.11), an unbiased estimator of the MSE of $\hat{\beta}_i^*$ is

$$(3.12) \quad \hat{M}(\hat{\beta}_i^*) = \left(\frac{s}{\nu \lambda_i} \right) \left[1 + \frac{\nu f_i}{(\nu f_i + 1)^2} + \nu J(\nu, f_i) \right]$$

$$\text{where } J(\nu, f_i) = \int_0^1 \left[\frac{(\nu f_i - x) x^{\frac{\nu}{2}}}{(\nu f_i + x)^2} \right] dx.$$

The square root of $\hat{M}(\hat{\beta}_i^*)$ gives a meaningful standard error for $\hat{\beta}_i^*$. This expression is simple, especially when compared with its population counterpart (see Dwivedi et al. (1980, p.206)). The value of $J(\nu, f_i)$ is easily determined and $\hat{M}(\hat{\beta}_i^*)$ is expressed in terms of quantities readily obtained from conventional regression output.

Finally, (3.12) depends on the regressors through λ_i and is derived on the assumption that $X'X$ has been diagonalised. However, considering the orthogonal transformations that would be used to achieve such a diagonalisation it is easily seen that $\hat{V}(b)$ in (2.4) depends on the data in a corresponding way; that "t-ratios" based on (2.9) and (3.12) are independent of this assumption; and that the comparative evaluations considered in the next section are perfectly general in the sense that they are not in fact limited to the case of a diagonalised design matrix.

4. NUMERICAL EVALUATION

The numerical evaluation of (3.12) for any choice of ν and f_i is straightforward - we have used the FORTRAN routines QROMB, TRAPZD and POLINT given by Press et al. (1986) to implement Romberg's method to determine $J(\nu, f_i)$. The relative estimated efficiency of b_i to $\hat{\beta}_i^*$ is $e = \hat{M}(\hat{\beta}_i^*)/\hat{V}(b_i)$, where $\hat{V}(b_i) = (s/\nu\lambda_i)$, and values of e are given in Table 1 for various choices of ν and $t_i = (\nu f_i)^{\frac{1}{2}}$. The latter quantity is the t-ratio for testing if $\beta_i = 0$, so this parameterisation facilitates practical prescriptions.

We see that $e < 1$ for combinations of small values of ν and t_i . Small values of t_i are characteristic of the application of OLS in the context of collinear data, and our results indicate that in such cases the OLS standard errors understate the precision of the estimates. If $t_i \leq 0.5$ then $\hat{\beta}_i^*$ is estimated to be more efficient than b_i for any ν ; and conversely if $t_i \geq 0.8$ then b_i is estimated to be more efficient than $\hat{\beta}_i^*$ for any ν .

From (2.9), $\hat{\beta}_i^* = b_i t_i^2 / (1 + t_i^2)$, so as $\nu \rightarrow \infty$ or $f_i \rightarrow \infty$, $\hat{\beta}_i^* \rightarrow b_i$ and $e \rightarrow 1$. This is supported by the results in Table 1 and also when e is tabulated in terms of ν and f_i values. The results in Table 1 also indicate that for the majority of combinations of ν and t_i , b_i is estimated to be efficient relative to $\hat{\beta}_i^*$, and the maximum such efficiency gain is 46%. Regardless of ν , this maximum gain arises for t_i between 1.2 and 1.6, and generally for $t_i = 1.4$ ($t_i^2 \approx 2$).

As noted already, in practice $\hat{V}(b_i)$ is often used to produce standard errors to use in association with $\hat{\beta}_i^*$. This can be justified asymptotically, but in finite samples this can now be improved upon through the use of $\hat{M}(\hat{\beta}_i^*)$. Comparing these two

approaches to measuring the precision of $\hat{\beta}_i^*$, our values for e show that generally the use of OLS standard errors in conjunction with ridge regression point estimates involves an overstatement of the precision of these estimates, relative to that indicated by the more appropriate measure derived in this paper. The figures in Table 1 may be viewed as correction factors to compensate for this overstatement - multiplying an OLS standard error by $e^{\frac{1}{2}}$ yields an appropriate standard error to use with the ridge estimator.

It is interesting to note that, in terms of actual MSE, $\hat{\beta}_i^*$ is efficient relative to b_i if $(\lambda_i \beta_i^2 / 2\sigma^2) < 1$. While it may be tempting to estimate the left hand side of this expression to see if the inequality holds empirically, it is clear that this strategy can be very misleading. For example, substituting the least squares estimators of β_i and σ^2 gives the condition $t_i^2 < 2$. Adjusting for the bias introduced by estimating in this way yields the condition $(\nu-2)t_i^2/\nu < 3$. That this is an unsatisfactory procedure can be seen by noting that these last two inequalities involve the estimation of a condition relating to the true relative efficiencies of the two estimators. In contrast, the information in Table 1 relates to the relative estimated efficiencies of $\hat{\beta}_i^*$ and b_i . Dealing with the inequality relating to the actual MSE's of $\hat{\beta}_i^*$ and b_i properly involves testing if $(\lambda_i \beta_i^2 / 2\sigma^2) < 1$ and choosing the estimator accordingly. This "preliminary-test" estimation strategy is discussed by Srivastava and Giles (1984).

Another way of considering the practical implications of our results is to compare t_i with $t_i^* = \hat{\beta}_i^* / (M(\hat{\beta}_i^*))^{\frac{1}{2}}$, the ridge-regression analogue of the usual t-ratio. The quantity $\gamma_i^* =$

(t_i^*/t_i) is the factor by which an OLS t -ratio should be scaled to convert it to its ridge regression counterpart. Of course, t_i^* is not t -distributed, but it is still a measure of some practical interest. Values of γ_i^* appear in Table 2. We see that, especially for small values of ν and/or t_i , a t -ratio based on OLS is misleading in the sense that it suggests substantially greater estimation precision than would be inferred from the value of t_i^* .

5. CONCLUDING REMARKS

This paper solves the problem of presenting a meaningful measure of the precision associated with a point estimate obtained by feasible generalised ridge regression. The practices generally adopted by applied workers are unsatisfactory because either they involve the estimation of a quantity which does not actually measure the population precision; or they are based on expressions which are uninformative because they are valid only asymptotically. In contrast, we present a simple expression for an unbiased estimator of the ridge estimator's mean squared error which is exact in finite samples and whose square root provides a meaningful standard error to be associated with a ridge regression coefficient. The tabulated evaluations indicate the extent to which standard errors and "t-ratios" based on least squares results may differ from those based on the procedure proposed here. The calculations needed to compute the proposed standard error are trivial, and can easily be incorporated into ridge regression packages.

October 1989

Table 1: Values of $e = \hat{M}(\hat{\theta}_i^2)/\hat{V}(b_i)$

v	t _i	0.50	0.60	0.70	0.80	0.90	1.00	1.20	1.40	1.60	1.80	2.00	2.50	3.00	4.00	5.00
1	0.97	1.14	1.25	1.32	1.37	1.39	1.40	1.37	1.34	1.30	1.27	1.20	1.15	1.09	1.06	
2	0.77	1.01	1.17	1.29	1.36	1.41	1.44	1.42	1.39	1.35	1.32	1.24	1.18	1.11	1.07	
3	0.64	0.90	1.10	1.25	1.34	1.40	1.45	1.44	1.41	1.38	1.34	1.26	1.20	1.12	1.08	
4	0.55	0.83	1.05	1.21	1.32	1.39	1.45	1.45	1.43	1.39	1.35	1.27	1.21	1.13	1.09	
5	0.49	0.78	1.01	1.18	1.30	1.37	1.45	1.46	1.44	1.40	1.36	1.28	1.21	1.13	1.09	
6	0.44	0.74	0.98	1.15	1.28	1.36	1.44	1.46	1.44	1.41	1.37	1.28	1.22	1.14	1.09	
7	0.41	0.71	0.95	1.13	1.26	1.35	1.44	1.46	1.44	1.41	1.37	1.29	1.22	1.14	1.09	
8	0.38	0.69	0.93	1.12	1.25	1.34	1.44	1.46	1.45	1.41	1.38	1.29	1.23	1.14	1.09	
9	0.36	0.67	0.91	1.10	1.24	1.34	1.44	1.46	1.45	1.42	1.38	1.29	1.23	1.14	1.10	
10	0.35	0.65	0.90	1.09	1.23	1.33	1.43	1.46	1.45	1.42	1.38	1.30	1.23	1.14	1.10	
12	0.32	0.63	0.88	1.07	1.22	1.32	1.43	1.46	1.45	1.42	1.39	1.30	1.23	1.15	1.10	
14	0.30	0.61	0.86	1.06	1.21	1.31	1.42	1.46	1.45	1.42	1.39	1.30	1.24	1.15	1.10	
16	0.29	0.60	0.85	1.05	1.20	1.30	1.42	1.46	1.45	1.42	1.39	1.31	1.24	1.15	1.10	
18	0.28	0.59	0.84	1.04	1.19	1.30	1.42	1.46	1.45	1.42	1.39	1.31	1.24	1.15	1.10	
20	0.27	0.58	0.83	1.04	1.19	1.29	1.42	1.45	1.45	1.43	1.39	1.31	1.24	1.15	1.10	
25	0.26	0.56	0.82	1.02	1.18	1.29	1.41	1.45	1.45	1.43	1.39	1.31	1.24	1.15	1.10	
30	0.25	0.55	0.81	1.01	1.17	1.28	1.41	1.45	1.45	1.43	1.40	1.31	1.24	1.15	1.10	
35	0.24	0.55	0.80	1.01	1.16	1.28	1.41	1.45	1.45	1.43	1.40	1.31	1.24	1.15	1.10	

Table 2: Values of $\gamma_i^* = (t_i^*/t_i)$

v	t_i	0.50	0.60	0.70	0.80	0.90	1.00	1.20	1.40	1.60	1.80	2.00	2.50	3.00	4.00	5.00
1		0.20	0.28	0.37	0.45	0.52	0.59	0.70	0.78	0.83	0.87	0.90	0.94	0.97	0.98	0.99
2		0.18	0.27	0.36	0.44	0.52	0.59	0.71	0.79	0.85	0.89	0.92	0.96	0.98	0.99	1.00
3		0.16	0.25	0.35	0.44	0.52	0.59	0.71	0.80	0.86	0.90	0.93	0.97	0.98	0.99	1.00
4		0.15	0.24	0.34	0.43	0.51	0.59	0.71	0.80	0.86	0.90	0.93	0.97	0.99	1.00	1.00
5		0.14	0.23	0.33	0.42	0.51	0.59	0.71	0.80	0.86	0.90	0.93	0.97	0.99	1.00	1.00
6		0.13	0.23	0.32	0.42	0.51	0.58	0.71	0.80	0.86	0.91	0.94	0.98	0.99	1.00	1.00
7		0.13	0.22	0.32	0.42	0.50	0.58	0.71	0.80	0.86	0.91	0.94	0.98	0.99	1.00	1.00
8		0.12	0.22	0.32	0.41	0.50	0.58	0.71	0.80	0.86	0.91	0.94	0.98	1.00	1.00	1.00
9		0.12	0.22	0.31	0.41	0.50	0.58	0.71	0.80	0.86	0.91	0.94	0.98	1.00	1.00	1.00
10		0.12	0.21	0.31	0.41	0.50	0.58	0.71	0.80	0.87	0.91	0.94	0.98	1.00	1.00	1.00
12		0.11	0.21	0.31	0.40	0.49	0.57	0.71	0.80	0.87	0.91	0.94	0.98	1.00	1.00	1.00
14		0.11	0.21	0.31	0.40	0.49	0.57	0.70	0.80	0.87	0.91	0.94	0.98	1.00	1.00	1.00
16		0.11	0.20	0.30	0.40	0.49	0.57	0.70	0.80	0.87	0.91	0.94	0.98	1.00	1.00	1.00
18		0.11	0.20	0.30	0.40	0.49	0.57	0.70	0.80	0.87	0.91	0.94	0.99	1.00	1.00	1.00
20		0.10	0.20	0.30	0.40	0.49	0.57	0.70	0.80	0.87	0.91	0.94	0.99	1.00	1.00	1.00
25		0.10	0.20	0.30	0.39	0.49	0.57	0.70	0.80	0.87	0.91	0.94	0.99	1.00	1.00	1.00
30		0.10	0.20	0.30	0.39	0.48	0.57	0.70	0.80	0.87	0.91	0.94	0.99	1.00	1.00	1.00
35		0.10	0.20	0.29	0.39	0.48	0.57	0.70	0.80	0.87	0.91	0.95	0.99	1.00	1.00	1.00

REFERENCES

- Carter, R.A.L., Srivastava, M.S., Srivastava, V.K. and Ullah, A. (19**). Unbiased Estimation of the MSE Matrix of Stein-Rule Estimators, Confidence Ellipsoids and Hypothesis Testing. Econometric Theory (forthcoming).
- Dwivedi, T.D., Srivastava, V.K. and Hall, R.L. (1980). Finite Sample Properties of Ridge Estimators. Technometrics, 22, 205-121.
- Hoerl, A.E. and Kennard, R.W. (1970). Ridge Regression: Biased Estimation for Non-Orthogonal Problems. Techometrics, 12, 55-67.
- Judge, G.G., Hill R.C., Griffiths, W.E. and Lee T-C. (1988). Introduction to the Theory and Practice of Econometrics (Wiley, New York).
- Press, W.H., Flannery, B.P., Teukolsky, S.A. and Vetterling, W.T. (1986). Numerical Recipies: The Art of Scientific Computing (Cambridge University Press, Cambridge).
- Srivastava, V.K. and Giles, D.E.A. (1984). A Pre-Test General Ridge Regression Estimator: Exact Finite-Sample Properties. Australian Journal of Statistics, 26, 323-336.

LIST OF DISCUSSION PAPERS*

- No. 8401 Optimal Search, by Peter B. Morgan and Richard Manning.
- No. 8402 Regional Production Relationships During the Industrialization of New Zealand, 1935-1948, by David E. A. Giles and Peter Hampton.
- No. 8403 Pricing Strategies for a Non-Replenishable Item Under Variable Demand and Inflation, by John A. George.
- No. 8404 Alienation Rights in Traditional Maori Society, by Brent Layton.
- No. 8405 An Engel Curve Analysis of Household Expenditure in New Zealand, by David E. A. Giles and Peter Hampton.
- No. 8406 Paying for Public Inputs, by Richard Manning, James R. Markusen, and John McMillan.
- No. 8501 Perfectly Discriminatory Policies in International Trade, by Richard Manning and Koon-Lam Shea.
- No. 8502 Perfectly Discriminatory Policy Towards International Capital Movements in a Dynamic World, by Richard Manning and Koon-Lam Shea.
- No. 8503 A Regional Consumer Demand Model for New Zealand, by David E. A. Giles and Peter Hampton.
- No. 8504 Optimal Human and Physical Capital Accumulation in a Fixed-Coefficients Economy, by R. Manning.
- No. 8601 Estimating the Error Variance in Regression After a Preliminary Test of Restrictions on the Coefficients, by David E. A. Giles, Judith A. Mikolajczyk and T. Dudley Wallace.
- No. 8602 Search While Consuming, by Richard Manning.
- No. 8603 Implementing Computable General Equilibrium Models: Data Preparation, Calibration, and Replication, by K. R. Henry, R. Manning, E. McCann and A. E. Woodfield.
- No. 8604 Credit Rationing: A Further Remark, by John G. Riley.
- No. 8605 Preliminary-Test Estimation in Mis-Specified Regressions, by David E. A. Giles.
- No. 8606 The Positive-Part Stein-Rule Estimator and Tests of Linear Hypotheses, by Aman Ullah and David E. A. Giles.
- No. 8607 Production Functions that are Consistent with an Arbitrary Production-Possibility Frontier, by Richard Manning.
- No. 8608 Preliminary-Test Estimation of the Error Variance in Linear Regression, by Judith A. Clarke, David E. A. Giles and T. Dudley Wallace.
- No. 8609 Dual Dynamic Programming for Linear Production/Inventory Systems, by E. Grant Read and John A. George.
- No. 8610 Ownership Concentration and the Efficiency of Monopoly, by R. Manning.
- No. 8701 Stochastic Simulation of the Reserve Bank's Model of the New Zealand Economy, by J. N. Lye.
- No. 8702 Urban Expenditure Patterns in New Zealand, by Peter Hampton and David E. A. Giles.
- No. 8703 Preliminary-Test Estimation of Mis-Specified Regression Models, by David E. A. Giles.
- No. 8704 Instrumental Variables Regression Without an Intercept, by David E. A. Giles and Robin W. Harrison.
- No. 8705 Household Expenditure in Sri Lanka: An Engel Curve Analysis, by Mallika Dissanayake and David E. A. Giles.
- No. 8706 Preliminary-Test Estimation of the Standard Error of Estimate in Linear Regression, by Judith A. Clarke.
- No. 8707 Invariance Results for FIML Estimation of an Integrated Model of Expenditure and Portfolio Behaviour, by P. Dorian Owen.
- No. 8708 Social Cost and Benefit as a Basis for Industry Regulation with Special Reference to the Tobacco Industry, by Alan E. Woodfield.
- No. 8709 The Estimation of Allocation Models With Autocorrelated Disturbances, by David E. A. Giles.
- No. 8710 Aggregate Demand Curves in General-Equilibrium Macroeconomic Models: Comparisons with Partial-Equilibrium Microeconomic Demand Curves, by P. Dorian Owen.
- No. 8711 Alternative Aggregate Demand Functions in Macro-economics: A Comment, by P. Dorian Owen.
- No. 8712 Evaluation of the Two-Stage Least Squares Distribution Function by Imhof's Procedure by P. Cribbitt, J. N. Lye and A. Ullah.
- No. 8713 The Size of the Underground Economy: Problems and Evidence, by Michael Carter.

(Continued on back cover)

- No. 8714 A Computable General Equilibrium Model of a Fisherine Method to Close the Foreign Sector, by Ewen McCann and Keith McLaren.
- No. 8715 Preliminary-Test Estimation of the Scale Parameter in a Mis-Specified Regression Model, by David E. A. Giles and Judith A. Clarke.
- No. 8716 A Simple Graphical Proof of Arrow's Impossibility Theorem, by John Fountain.
- No. 8717 Rational Choice and Implementation of Social Decision Functions, by Manimay Sen.
- No. 8718 Divisia Monetary Aggregates for New Zealand, by Ewen McCann and David E. A. Giles.
- No. 8719 Telecommunications in New Zealand: The Case for Reform, by John Fountain.
- No. 8801 Workers' Compensation Rates and the Demand for Apprentices and Non-Apprentices in Victoria, by Pasquale M. Sgro and David E. A. Giles.
- No. 8802 The Adventures of Sherlock Holmes, the 48% Solution, by Michael Carter.
- No. 8803 The Exact Distribution of a Simple Pre-Test Estimator, by David E. A. Giles.
- No. 8804 Pre-testing for Linear Restrictions in a Regression Model With Student-t Errors, by Judith A. Clarke.
- No. 8805 Divisia Monetary Aggregates and the Real User Cost of Money, by Ewen McCann and David Giles.
- No. 8806 The Management of New Zealand's Lobster Fishery, by Alan Woodfield and Pim Borren.
- No. 8807 Poverty Measurement: A Generalization of Sen's Result, by Prasanta K. Pattanaik and Manimay Sen.
- No. 8808 A Note on Sen's Normalization Axiom for a Poverty Measure, by Prasanta K. Pattanaik and Manimay Sen.
- No. 8809 Budget Deficits and Asset Sales, by Ewen McCann.
- No. 8810 Unorganized Money Markets and 'Unproductive' Assets in the New Structuralist Critique of Financial Liberalization, by P. Dorian Owen and Otton Solis-Fallas.
- No. 8901 Testing for Financial Buffer Stocks in Sectoral Portfolio Models, by P. Dorian Owen.
- No. 8902 Provisional Data and Unbiased Prediction of Economic Time Series by Karen Browning and David Giles.
- No. 8903 Coefficient Sign Changes When Restricting Regression Models Under Instrumental Variables Estimation, by David E. A. Giles.
- No. 8904 Economies of Scale in the New Zealand Electricity Distribution Industry, by David E. A. Giles and Nicolas S. Wyatt.
- No. 8905 Some Recent Developments in Econometrics: Lessons for Applied Economists, by David E. A. Giles.
- No. 8906 Asymptotic Properties of the Ordinary Least Squares Estimator in Simultaneous Equations Models, by V. K. Srivastava and D. E. A. Giles.
- No. 8907 Unbiased Estimation of the Mean Squared Error of the Feasible Generalised Ridge Regression, by V. K. Srivasatva and D. E. A. Giles.

* Copies of these Discussion Papers may be obtained for \$4 (including postage, price changes occasionally) each by writing to the Secretary, Department of Economics, University of Canterbury, Christchurch, New Zealand.