



**AgEcon** SEARCH  
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search  
<http://ageconsearch.umn.edu>  
[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

CANTER

DISCUSSION PAPER 91/07

Department of Economics  
UNIVERSITY OF CANTERBURY  
CHRISTCHURCH, NEW ZEALAND



GIANNINI FOUNDATION OF  
AGRICULTURAL ECONOMICS  
LIBRARY

WITHDRAWN

AUG 26 1991

**$L_p$ -NORM CONSISTENCIES OF NONPARAMETRIC  
ESTIMATES OF REGRESSION, HETEROSKEDASTICITY  
AND VARIANCE OF REGRESSION ESTIMATE  
WHEN DISTRIBUTION OF REGRESSOR IS KNOWN**

Radhey S. Singh  
(University of Guelph)

***Discussion Paper***

No. 9107

This paper is circulated for discussion and comments. It should not be quoted without the prior approval of the author. It reflects the views of the author who is responsible for the facts and accuracy of the data presented. Responsibility for the application of material to specific cases, however, lies with any user of the paper and no responsibility in such cases will be attributed to the author or to the University of Canterbury.

Department of Economics, University of Canterbury  
Christchurch, New Zealand

***Discussion Paper No. 9107***

**April 1991**

**$L_p$ -NORM CONSISTENCIES OF NONPARAMETRIC  
ESTIMATES OF REGRESSION, HETEROSKEDASTICITY  
AND VARIANCE OF REGRESSION ESTIMATE  
WHEN DISTRIBUTION OF REGRESSOR IS KNOWN**

**Radhey S. Singh**

$L_p$ -NORM CONSISTENCIES OF NONPARAMETRIC ESTIMATES

OF REGRESSION, HETEROSKEDASTICITY

AND VARIANCE OF REGRESSION ESTIMATE

WHEN DISTRIBUTION OF REGRESSOR IS KNOWN<sup>1,2</sup>

by

RADHEY S. SINGH

UNIVERSITY OF GUELPH

1. Part of the research was carried out during the author's visit to the Department of Economics, University of Canterbury, Christchurch, New Zealand. The author thanks David Giles, Alan Woodfield, Judith Giles and their Department for the invitation to visit and providing the research support.
2. The paper is to be presented at the "Symposium on Nonparametric Statistical Inference and Related Topics" to be held during May 5 - 8, 1991 at Carleton University, Ottawa, Canada.

### ABSTRACT

When dealing with heteroskedastic models  $Y = \mu(X) + \epsilon$  in econometrics and other disciplines, situations often arise (especially with structural models) where the probability distribution of the ( $\mathbb{R}^d$ -valued) regressor vector  $X$  is known, but postulations about the functional form of the regression  $\mu(x)$ , the heteroskedasticity  $\sigma^2(x) = \text{var}(\epsilon|X=x)$  and the distribution of the disturbance term  $\epsilon$  are made. These three postulations generally lead to misspecification of the models. This paper, based on a data set on  $(Y, X)$ , considers nonparametric kernel estimators  $\hat{\mu}(x)$ ,  $\hat{\sigma}^2(x)$  and  $\hat{V}(\hat{\mu}(x))$ , respectively, of the regression  $\mu(x)$ , the heteroskedasticity  $\sigma^2(x)$  and the asymptotic variances  $V(\hat{\mu}(x))$  of the regression estimate  $\hat{\mu}(x)$  for situations where only the probability distribution of  $X$ , say  $\lambda$  is known. For an arbitrary subset  $A$  in the interior of the support of  $\lambda$  and for  $1 \leq p < \infty$ , we establish convergences to zero, as the data set gets large, of the  $L_p$ -norms  $\|\hat{\mu} - \mu\|^p = \int_A E|\hat{\mu}(x) - \mu(x)|^p d\lambda(x)$  (with  $A \equiv \mathbb{R}^d$  for  $p = 1$ ),  $\|\hat{\sigma}^2 - \sigma^2\|^p$  (with  $A \equiv \mathbb{R}^d$  for  $p = 1$ ) and  $\|\hat{V}(\hat{\mu}) - V(\hat{\mu})\|^p$  under certain moment conditions on  $Y$  but with no assumptions on the joint distribution of  $(Y, X)$  or the continuity of  $\mu(x)$ ,  $\sigma^2(x)$  or the density of  $X$ .

## 1. INTRODUCTION

Let  $(Y, X)$  be a  $\mathbb{R}^1 \times \mathbb{R}^d$  - valued random vector defined on a probability space  $(\Omega, \mathcal{B}, P)$ . Then the general heteroskedastic regression model is given by  $Y = \mu(X) + \epsilon$ . A measurement on response  $Y$  at a value  $x$  of the regressor  $X$  is the value of the unknown regression  $\mu(x)$  at  $X = x$  contaminated with an unobservable disturbance term  $\epsilon$ , which is random on  $(\Omega, \mathcal{B}, P)$  with conditional mean  $E(\epsilon|X) = 0$  and unknown conditional variance  $\text{Var}(\epsilon|X) = \sigma^2(x)$ , called the heteroskedasticity of the regression model. Estimations of the regression function  $\mu(x) = E(Y|X=x)$  and the heteroskedasticity function  $\sigma^2(x) = \text{var}(Y|X=x)$  are invariably handled in various sciences by postulating a certain fixed model (functional form) for the regression, by assuming a constant conditional variance,  $\sigma^2(x) \equiv \sigma^2$  (the homoskedasticity), for the response variable  $Y$  at any value  $x$  of the regressor  $X$  and by assuming a knowledge of the functional form of the distribution of the disturbance term  $\epsilon$ . However any postulations regarding the functional form of the regression  $\mu(x)$ , the heteroskedasticity  $\sigma^2(x)$  or the distributions of disturbances  $\epsilon$  are all questionable, and often lead to misspecifications of the models, which thereby lead to serious impact on decisions and plannings.

This problem of misspecification of the models, however, can be avoided by assuming no specific parametric form for the regression or heteroskedasticity function; and by estimating them completely nonparametrically.

Whereas a vast literature on nonparametric estimation of regression  $\mu$  is at hand, (for example, Watson (1964), Rosenblatt (1969), Schuster (1972), Schuster and Yakowitz (1979), Noda (1976), Greblicki and Krzyzak (1980), Härdle (1984), among others, discussed pointwise consistency; and

Nadarya (1964, 1970), Deheuvels (1974), Hall (1981), Mack and Silverman (1987), Härdle and Luckhans (1984) Singh and Ahmad (1987), among others discussed uniform consistency of nonparametric kernel estimators of regression), nonparametric estimation of heteroskedasticity function  $\sigma^2(x)$  has drawn little attention. Hildreth and Houck (1968), Fraehlich (1973), Box and Hill (1974), Jobson and Fuller (1980) and Judge et al (1988) adopted a sort of parametric approach in which they assumed that  $\sigma^2(x)$  has a known functional form involving a finite number of unknown parameters. Fuller and Rao (1978), White (1980), Carroll (1982) and MacKinnon and White (1985) took a somewhat semi-non-parametric approach in which they assumed that the regression function  $\mu(x)$  has a known functional form involving a finite number of unknown parameters, estimated these parameters and then used the residuals and the estimated variance-covariance of the parameter estimates to estimate  $\sigma^2(x)$  nonparametrically. In the latter works, the assumed functional form of  $\mu(x)$  is a linear regression. Müller and Stadtmüller (1987) considered estimation of  $\sigma^2(x)$  with nonstochastic ordered regressors assuming, among others, that the density of Y and  $\sigma^2(x)$  satisfy certain order of Lipschitz condition. Singh and Ullah (1985) and Singh et al. (1987) discussed estimation of  $\sigma^2(x)$  under the continuity of  $\sigma^2(x)$ , the density of X and the joint density of (Y,X) at x.

In most of the literature on the nonparametric estimation of regression or heteroscedasticity (e.g. the references cited in the preceding paragraph), it is invariably assumed, particularly when discussing consistencies like weak, strong, mean square or asymptotic normality that the regression  $\mu(x)$  and heteroskedasticity  $\sigma^2(x)$  (in case of its estimation) are continuous, and the joint density of (Y,X), the density of Y and the density of X (and hence the conditional density of Y given X = x) not only exist but are also continuous. Recently Singh (1989)



considered kernel estimations of  $\mu$  and  $\sigma^2$  under no condition on the distribution of  $(Y,X)$  other than that the distribution function of  $X$  is absolutely continuous w.r.t. the Lebesgue measure on  $\mathbb{R}^d$  so that the probability density function (pdf) of  $X$  exists. He established the weak and strong consistencies as well as the asymptotic normality of the estimators  $\mu(x)$  and  $\sigma^2(x)$  only under some appropriate moment conditions on  $Y$ . Thus for his results the joint density of  $(Y,X)$ , the density of  $Y$  or the conditional density of  $Y$  given  $X$  need not exist and  $\mu(x)$ ,  $\sigma^2(x)$  or the density of  $X$  need not be continuous at  $x$ . Under similar weaker conditions he has also established weak and strong consistencies of the estimates of the variances of his regression and heteroskedasticity estimators by establishing similar properties for the estimators of a general function, namely  $\mu_r(x) = E(Y^r | X=x)$  for  $r \geq 0$ .

Greblicki and Krzyzak (1980), Johnston (1982) and Härdle (1986), among others considered Watson-Nadaraya type kernel estimators of regression when the pdf of the regressor  $X$ , say  $\phi$ , is known. Under the assumption that  $\phi(x)$  and  $\mu(x)$  are continuous and  $EY^2 < \infty$ , pointwise weak consistency of such estimators  $\hat{\mu}(x)$  is established by Greblicki and Krzyzak, while Johnston and Härdle obtained results, respectively, on the asymptotic distribution of the maximal deviation of  $\hat{\mu}(x)$  from  $\mu(x)$  and an  $L_2(0,1)$ -norm  $\int_0^1 E|\hat{\mu}(x) - \mu(x)| d\lambda(x)$ , where  $\lambda$  is the probability measure of  $X$ . Considering situations where the distributions of the regressor vectors are known is not quite unrealistic. For example, in structural regression models in econometrics quite often the regression vector is consisted of endogenous variables generated from some known functions of exogenous variables plus independent multivariate normal errors with mean zero and known variances.

In this note we further look at the situation considered in the preceding paragraph, i.e. when the pdf of  $X$  is known. We estimate the regression as well as the heteroskedasticity and the variance of the regression estimator, and examine the  $L_p$ -norm ( $p \geq 1$ ) consistencies of these estimators. Weak and strong pointwise consistencies of these estimators follow from those established in Singh (1989) under weaker conditions stated in preceding paragraph. However, note that weak or strong consistency of an estimator does not imply  $L_p$ -norm consistency. Without any continuity assumption on  $\mu(\cdot)$ ,  $\sigma^2(\cdot)$  or the p.d.f.  $\phi(\cdot)$  of  $X$ , and without making any assumption on the distribution of  $(Y, X)$ , the distribution of  $Y$  or the conditional distribution of  $Y$  given  $X$ , we establish the convergence to zero, as the sample size gets large, of the  $L_p$ -norm distances,  $1 \leq p < \infty$ , of the estimators from their respective true values.

In Section 2, we introduce some notations, give a proper definition of " $L_p$ -norm convergence" considered here and define estimators  $\hat{\mu}_r(x)$  of the regression function  $\mu_r(x) = (EY^r | X=x)$  of  $Y^r$  on  $X$  for  $r \geq 0$ , estimators  $\hat{\sigma}^2(x)$  of the heteroskedasticity  $\sigma^2(x)$  and estimators  $\hat{V}(\hat{\mu}(x))$  of the asymptotic variance of the regression estimate  $\hat{\mu}(x) = \hat{\mu}_1(x)$ . In Section 3 we establish  $L_p$ -norm consistencies of the estimator  $\hat{\mu}_r$  (Theorem 3.1),  $\hat{\sigma}^2$  (Theorem 3.2) and  $\hat{V}(\hat{\mu})$  (Theorem 3.3). The paper is concluded with a few remarks in Section 4.

## 2. NOTATIONS, DEFINITIONS AND ESTIMATORS

Let  $\mathbb{R}^d$  denote the  $d$ -dimensional Euclidean space with the usual norm  $\|z\| = (z'z)^{1/2} = (\sum_1^d z_i^2)^{1/2}$  for  $z = (z_1, \dots, z_d)' \in \mathbb{R}^d$ . Let  $(Y, X)$ ,  $(Y_1, X_1)$ , ...,  $(Y_n, X_n)$  be independent identically distributed  $\mathbb{R}^1 \times \mathbb{R}^d$ -valued random vectors defined on a common probability space  $(\Omega, \mathcal{B}, P)$ . We assume that the regression  $\mu(x) = E(Y|X)$  and the heteroskedasticity  $\sigma^2(x) = \text{var}(Y|X)$  are properly defined and exist a.e.  $(\lambda)$ , where  $\lambda$  is the probability measure of  $X$ . We assume nothing about the joint distribution of  $(Y, X)$ , the distribution of  $Y$  or the conditional distribution of  $Y$  given  $X$ . We do however assume that the probability measure  $\lambda$  of  $X$  is known and is dominated by the Lebesgue measure on  $\mathbb{R}^d$  so that the probability density, say  $\phi$ , of  $X$  exists and is known. Thus the joint density of  $(Y, X)$ , the density of  $Y$  or the conditional density of  $Y$  given  $X$  need not exist and the functions  $\mu(x)$ ,  $\sigma^2(x)$  or  $\phi(x)$  need not be continuous. When  $\phi$  is unknown but the response variable  $Y$  is bounded w.p.1, estimators of the regression and heteroskedasticity functions, which are some retracted versions of those considered in Singh (1989), have been recently studied by Singh and Giles (1991) and  $L_1$ -norm consistencies (which also imply  $L_p$ -norm consistencies for  $p \geq 1$  in their case) have been established.

All real-valued functions on  $\mathbb{R}^d$  in this paper carrying an argument  $x$  (explicitly or implicitly) are only defined a.e.  $\lambda$ , i.e. on the support of the pdf  $\phi$ . Also all convergences of sequences (in  $n$ ) of functions carrying an argument  $x$  (explicitly or implicitly) are only a.e.  $\lambda$ .

For  $r \geq 0$ , define

$$\mu_r(x) = E(Y^r | X=x)$$

(so that  $\mu_1(x) = \mu(x)$ ). In the heteroscedastic model  $Y = \mu(X) + \varepsilon$ , with

$E(\varepsilon|X) = 0$ , the regression function is  $\mu(x) = E(Y|X=x) = (\mu_1(x)$ , according to our notation) and the heteroskedasticity function is

$$\sigma^2(x) = \text{var}(\varepsilon|X=x) = E(\varepsilon^2|X=x),$$

which can be expressed as  $\sigma^2(x) = \mu_2(x) - \mu_1^2(x)$ .

Define  $\psi_r = \mu_r \phi$  so that

$$(2.1) \quad \mu_r(x) = \frac{\psi_r(x)}{\phi(x)}.$$

Notice that  $\psi_0 = \phi$ .

Let  $K$  be an arbitrary Borel-measurable function on  $\mathbb{R}^d$  with

$\int_{\mathbb{R}^d} K(u)du = 1$ . Let  $\{h_n\}$  be a sequence of positive real numbers such that

$h_n \rightarrow 0$  as  $n \rightarrow \infty$ . Define

$$(2.2) \quad \hat{\psi}_r(x) = (nh_n^d)^{-1} \sum_{j=1}^n Y_j^r K\left(\frac{X_j - x}{h_n}\right),$$

and

$$\hat{\phi}(x) = \hat{\psi}_0(x).$$

Throughout this paper, the notation  $(a)_B$  will stand for  $-B$ ,  $a$  or  $B$  depending on whether  $a < -B$ ,  $|a| \leq B$  or  $a > B$ .

In view of our knowledge of  $\phi$ , nonparametric kernel estimate of  $\mu_r(x)$  is

$$(2.3) \quad \hat{\mu}_r(x) = \frac{\hat{\psi}_r(x)}{\hat{\phi}(x)}$$

For estimation of  $\sigma^2$ , we assume that  $|E(Y|X)| \leq B$  w.p.1 for some  $B < \infty$ .

We estimate  $\sigma^2(x)$  by

$$(2.4) \quad \hat{\sigma}^2(x) = \hat{\mu}_2(x) - (\hat{\mu}_1(x))_B^2, \text{ where } \hat{\mu} = (\hat{\mu}_1)_B.$$

Estimators  $\tilde{\theta}_r(x) = (\hat{\psi}_r(x)/\hat{\phi}(x))$  for  $\mu_r(x)$  and  $\tilde{\sigma}^2(x) = (\tilde{\theta}_2(x) - \tilde{\theta}_1(x))^2$  for  $\sigma^2(x)$  are considered in Singh (1989). Only under some moment conditions on  $Y$  and the existence of  $\phi$ , it has been established there that the asymptotic variance of the regression estimate  $\tilde{\theta}_1(x) = \tilde{\theta}(x)$ , up to the order  $o\left((nh_n^d)^{-1}\right)$ , is given by

$$(2.5) \quad V(\tilde{\theta}(x)) = (nh_n^d)^{-1} \frac{\sigma^2(x) \int K^2}{\phi(x)},$$

$$\text{where } \int K^2 = \int_{\mathbb{R}^d} K^2(x) dx,$$

(see also Singh and Ullah (1985), Singh et al (1987). It follows from the same arguments that the asymptotic variances of the regression estimates  $\tilde{\mu}(x)$  and  $\hat{\mu}(x) = \hat{\mu}_1(x)$  are the same as given in (2.5). Thus we estimate  $V(\hat{\mu}(x))$  by

$$(2.6) \quad \hat{V}(\hat{\mu}(x)) = (nh_n^d)^{-1} \frac{\hat{\sigma}^2(x) \int K^2}{\phi(x)}$$

where  $\hat{\sigma}^2(x)$  is as given in (2.4).

Notice that  $\tilde{\theta}_1(x) = \sum_{j=1}^n Y_j W_{nj}(x)$  and  $\tilde{\theta}_2(x) = \sum_{j=1}^n Y_j^2 W_{nj}(x)$ , where  $W_{nj}(x) = \{K((X_j - x)/h_n)\} \{ \sum_{j=1}^n K((X_j - x)/h_n) \}^{-1}$ , are well known Nadaraya-Watson type kernel estimators of the regressions of  $Y$  and  $Y^2$  on  $X$ , respectively. Therefore  $\tilde{e}_j(x) = (Y_j - \tilde{\theta}_1(x))$  are the nonparametric kernel estimators of the residuals  $e_j(x)$  and  $\tilde{\sigma}^2(x) = (\tilde{\theta}_2(x) - \tilde{\theta}_1(x))^2 = \sum_{j=1}^n (Y_j - \tilde{\theta}_1(x))^2 W_{nj}(x)$  is the Nadaraya-Watson type kernel estimator of the regression of the square of the residuals  $(Y_j - \tilde{\theta}_1(x))$  on  $X$ , which is a natural way to estimate  $\sigma^2(x) = E(e^2 | X=x) = \text{var}(e | X=x)$ .

Throughout this paper, let

$$(2.7) \quad A \subseteq \mathbb{R}^d \text{ such that } \text{ess-inf}_{x \in A} (\text{w.r.t. } \lambda) \phi(x) > 0.$$

Definition:

For a real valued statistic  $\theta^*(x) = \theta^*((Y_1, X_1), \dots, (Y_n, X_n); x)$  defined for  $x \in \mathbb{R}^d$  and for  $p \geq 1$ , we say  $\theta^*$  is  $L_p$ -norm consistent estimator of a real valued function  $\theta(x)$  on  $\mathbb{R}^d$  if

$$(2.8) \quad \|\theta^* - \theta\|^p = \int_A E|\theta^*(x) - \theta(x)|^p d\lambda(x) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

We remark that  $L_p$ -norm convergence of the type (2.8) over a subset  $A$  of  $\mathbb{R}^d$  satisfying condition similar to (2.7), has been used as a standard by several authors for examining a global consistency of a regression estimate, see Section 4, for example.

For a real valued function  $g$  on  $\mathbb{R}^d$  we say  $g \in L_p$  if  $\int_{\mathbb{R}^d} |g(u)|^p du < \infty$ , and for a measure  $\nu$  on  $\mathbb{R}^d$ , we say  $g \in L_p(\nu)$  if  $\int_{\mathbb{R}^d} |g(\cdot)|^p d\nu(\cdot) < \infty$ . For the sake of simplicity in writing, we denote, whenever convenient,  $\int_{\mathbb{R}^d} g(u) du$  by  $\int g$  and  $\int g(\cdot) d\nu(\cdot)$  by  $\int g d\nu$ . Unless stated otherwise, all integrals throughout this paper are taken over the space  $\mathbb{R}^d$ . For  $t \in \mathbb{R}^1$ , let  $K^*(t) = \text{ess-sup}_{\|u\|>t} |K(u)|$ , where the ess-sup is taken with respect to the Lebergue measure on  $\mathbb{R}^d$ . (Note that  $K^* \in L_p$  whenever  $|K(\cdot)|$  is nonincreasing in  $\|\cdot\|$  and belongs to  $L_p$ ). Throughout this paper it is assumed that  $K$  and  $K^*$  belong to  $L_1 \cap L_2$ , and no further indication of it will be made in the paper. Unless stated otherwise, all convergences are w.r.t.  $n \rightarrow \infty$ .

### 3. THE $L_p$ -NORM CONVERGENCES OF ESTIMATORS

In this section we establish three theorems. Theorem 3.1 establishes the  $L_p$ -norm convergence of  $\hat{\mu}_r$  under certain moment conditions on  $Y$ . The statistics  $\hat{\mu}_r$  are the kernel estimators of  $\mu_r$ , the regression of  $Y^r$  on  $X$  for  $r \geq 0$ , though only the particular cases of our interest here are with  $r = 1$  and  $2$ . The  $L_p$ -norm convergences of  $\hat{\sigma}^2$  and  $\hat{V}(\hat{\mu})$  are established in Theorems 3.2 and 3.3 respectively.

Theorem 3.1. *Let  $h_n \rightarrow 0$ ,  $r \geq 0$  and  $1 \leq p < \infty$ . Then*

$$(3.1) \quad \|\hat{\mu}_r - \mu_r\|^p = o(1) \text{ with A, in (2.6), } \equiv R^d \text{ for } p = 1,$$

*provided for  $1 \leq p \leq 2$ ,  $E|Y|^{2r} < \infty$  and  $nh_n^d \rightarrow \infty$ ; and for  $p > 2$ , there exists an  $0 < \eta < 2$  such that with  $w = 2(p-\eta)/(2-\eta)$ ,  $E|Y|^{r w} < \infty$ ,  $K \in L_w$  and  $n^{\eta/2} h_n^{d(p-1)} \rightarrow \infty$ .*

Theorem 3.2. *Let  $1 \leq p < \infty$ , and the conditions of Theorem 3.1 hold for  $r = 1$  and  $2$ . Further let there exist a finite constant  $B$  such that  $|E(Y|X)| \leq B$  w.p.1, and  $\hat{\sigma}^2$  be defined as in (2.4). Then*

$$(3.2) \quad \|\hat{\sigma}^2 - \sigma^2\|^p = o(1) \text{ with A, in (2.6), } \equiv R^d \text{ for } p = 1.$$

Theorem 3.3. *Let  $1 \leq p < \infty$ , and the conditions for Theorem 3.2 hold. Let  $\hat{V}(\hat{\mu}(x))$  be defined as in (2.6) and  $V(\hat{\mu}(x))$  stand for the asymptotic variance of  $\hat{\mu}(x)$ . Then*

$$(3.3) \quad (nh_n^d)^p \|\hat{V}(\hat{\mu}) - V(\hat{\mu})\|^p = o(1).$$

To prove Theorems 3.1, 3.2 and 3.3 we establish first the following lemmas.

Lemma 3.1     Let  $h_n \rightarrow 0$ ,  $1 \leq p < \infty$  and  $r \geq 0$ . If  $E|Y|^r < \infty$ , then

$$(3.4) \quad \int \phi^{-p}(x) |E\hat{\psi}_r(x) - \psi_r(x)|^p d\lambda(x) = o(1)$$

Proof.

Our proof involves the following steps. We first prove that  $|E\hat{\psi}_r(x) - \psi_r(x)| = o(1)$  a.e.  $\lambda(x)$ . Then we show that the sequence (in  $n$ ) of functions  $\phi^{-p+1}(x) |E\hat{\psi}_r(x) - \psi_r(x)|^p \leq g_n(x)$  a.e.  $\lambda(x)$  for some sequence of functions  $g_n(x)$  which converges to  $g(x)$  as  $n \rightarrow \infty$  a.e.  $\lambda(x)$ . We then show that  $g \in L_1$  and then we complete the proof by using generalized dominated convergence theorem.

Since  $\{(Y_1, X_1), \dots, (Y_n, X_n)\}$  is a random sample on  $(Y, X)$ ,

$$(3.5) \quad E\hat{\psi}_r(x) = h_n^{-d} \int K \left( \frac{u-x}{h_n} \right) \psi_r(u) du$$

Next note that  $\int |\psi_r(u)| du = E|Y|^r < \infty$ . Therefore,  $\psi_r \in L_1$  and hence almost all points  $u \in \mathbb{R}^d$  are in the Lebesgue set of  $\psi_r$  (see, Natanson (1955), pp.255-266 or Wheeden and Zygmund (1977), pp.100-109). Hence, since  $K^* \in L_1$ , by Theorem 1.25 of Stein and Weiss (1975), p.13,

$$(3.6) \quad \lim_{n \rightarrow \infty} h_n^{-d} \int K \left( \frac{u-x}{h_n} \right) \psi_r(u) du = \psi_r(x) \int K \quad \text{a.e. in } x \in \mathbb{R}^d$$

Hence since  $\int K = 1$ , we conclude from (3.5) and (3.6) that  $|E\hat{\psi}_r(x) - \psi_r(x)| \rightarrow 0$  a.e.  $\lambda(x)$ .



Next notice that

$$\begin{aligned} \phi^{-p+1}(x) |E\hat{\psi}_r(x) - \psi_r(x)|^p &\leq 2^{p-1} \phi^{-p+1}(x) \left\{ |E\hat{\psi}_r(x)|^p + |\psi_r(x)|^p \right\} \\ &\leq 2^{p-1} \phi^{-p+1}(x) \left\{ |h_n^{-d} \int K\left(\frac{u-x}{h_n}\right) \psi_r(u) du|^p \right. \\ &\quad \left. + |\psi_r(x)|^p \right\} = g_n(x), \text{ say.} \end{aligned}$$

Notice that from (3.6),  $g_n(x) \rightarrow 2^p \phi^{-p+1}(x) |\psi_r(x)|^p = g(x)$ , say, a.e. in  $x \in \mathbb{R}^d$ . But  $g \in L_1$  since  $\int \phi^{-p+1}(x) |\psi_r(x)|^p dx \leq E|Y|^{rP} < \infty$ , which completes the proof of the Lemma.  $\square$

Lemma 3.2. Let  $r \geq 0$  and  $1 \leq p \leq \infty$ . If for  $1 \leq p \leq 2$ ,  $E|Y|^{2r} < \infty$  and  $nh_n^d \rightarrow \infty$ , then

$$(3.7) \quad \int_A \phi^{-p}(x) E|\hat{\psi}_r(x) - E\hat{\psi}_r(x)|^p d\lambda(x) = o(1) \text{ with } A \equiv \mathbb{R}^d \text{ for } p = 1.$$

Further (3.7) holds for  $p > 2$  if there exists an  $0 < \eta < 2$  such that, with  $w = 2(p-\eta)/(2-\eta)$ ,  $E|Y|^{rw} < \infty$ ,  $K \in L_w$  and  $n^{\eta/2} h_n^{d(p-1)} \rightarrow \infty$ .

Proof.

First consider the case  $1 \leq p \leq 2$ . By Hölder's inequality  $E|\hat{\psi}_r(x) - E\hat{\psi}_r(x)|^p \leq (\text{var} \hat{\psi}_r(x))^{p/2}$ . But since  $(Y_1, X_1), \dots, (Y_n, X_n)$  are i.i.d.,

$$\text{var} \hat{\psi}_r(x) \leq \left( nh_n^{2d} \right)^{-1} E \left( Y_1^r K \left( \frac{X_1 - x}{h_n} \right) \right)^2$$

Therefore

$$(3.8) \quad \left( nh_n^d \right)^{p/2} E|\hat{\psi}_r(x) - E\hat{\psi}_r(x)|^p \leq \left( h_n^{-d} \int K^2 \left( \frac{u-x}{h_n} \right) \psi_{2r}(u) du \right)^{p/2}$$

The r.h.s. of (3.8) converges to  $(\psi_{2r}(x) \int K^2)^{p/2}$  a.e.  $\lambda$ , since  $\psi_{2r} \in L_1$  by virtue of the fact that  $E|Y|^{2r} < \infty$  and the arguments used to prove the convergence of the rhs (3.5) to  $\psi_r(x) \int K$  can be applied. Hence, we conclude that if  $nh_n^d \rightarrow \infty$ , then

$$(3.9) \quad \phi^{-p+1}(x) E|\hat{\psi}_r(x) - E\hat{\psi}_r(x)|^p \rightarrow 0$$

Now for  $p = 1$ , the l.h.s. of (3.9) is bounded by  $2(nh_n^d)^{-1} \sum_{j=1}^n E|Y_j^r K((X_j - x)/h_n)| = 2h_n^{-d} \int |K(u-x)/h_n| |\Psi_r(u)| du = g_n(x)$ , say, where  $\Psi_r(x) = E(|Y|^r | X=x) \phi(x)$ . Again, by the arguments given earlier,  $g_n(x) \rightarrow g(x) = 2\Psi_r(x) \int |k|$  which belongs to  $L_1$  since  $E|Y|^r < \infty$ . Hence (3.9) followed by the generalized dominated convergence theorem gives (3.7) for  $p = 1$ .

For  $1 < p \leq 2$ , the l.h.s. of (3.7) from (3.8) is bounded by

$$\begin{aligned} & (nh_n^d)^{-p/2} \int_A \phi^{-p}(x) \left( \int K^2(u) \psi_{2r}(x+h_n u) du \right)^{p/2} d\lambda(x) \\ & \leq \left( nh_n^d \right)^{-p/2} \left( \alpha(A) \right)^{-p/2} \left( \iint K^2(u) \psi_{2r}(x+h_n u) dudx \right)^{p/2} \\ & = \left( nh_n^d \right)^{-p/2} \left( \alpha(A) \right)^{-p/2} \left( \int K^2 \right) E|Y|^{2r} \right)^{p/2} \end{aligned}$$

by Hölder's inequality. The proof of (3.7) for  $1 < p \leq 2$  is now complete since  $nh_n^d \rightarrow \infty$ .

Now we prove (3.11) for  $p > 2$ . Let  $0 < \eta < 2$  and  $w = 2(p-\eta)/(2-\eta)$ .

By Hölder's inequality,

$$(3.10) \quad E|\hat{\psi}_r(x) - E\hat{\psi}_r(x)|^p \leq \left( E|\hat{\psi}_r(x) - E\hat{\psi}_r(x)|^w \right)^{(2-\eta)/2} \left( \text{var}(\hat{\psi}_r(x)) \right)^{\eta/2}$$

Now, since  $w > 1$ ,  $E|\hat{\psi}_r(x) - E\hat{\psi}_r(x)|^w$  is bounded by  $2^w h_r^{-dw} E|Y_1^r K((X_1-x)/h_n)|^w$   
 $= 2^w h_n^{-d(w-1)}$  times  $\int |K(u)|^w \psi_{rw}(x+h_n u) du$  and  $\text{var}(\hat{\psi}_r(x))$  is bounded by  
 $(nh_n^d)^{-1} E|Y_1^r K((X_1-x)/h_n)|^2 = (nh_n^d)^{-1} \int K^2(u) \psi_{2r}(x+h_n u) du$ . Hence, from (3.10)  
and by Hölder inequality, the l.h.s. of (3.7) for  $p > 2$  is bounded by  
 $2^w h_n^{-d(w-1)(2-\eta)/2} (nh_n^d)^{-\eta/2} = 2^w n^{-\eta/2} h_n^{-d(p-1)}$  times

$$\int_A \phi^{-p}(x) \left\{ \int |K(u)|^w \psi_{rw}(x+h_n u) du \right\}^{(2-\eta)/2} \left\{ \int |K(u)|^2 \psi_{2r}(x+h_n u) du \right\}^{\eta/2} d\lambda(x),$$

which, again by Hölder Inequality, is bounded by  $c_n \cdot c_n'$ , where

$$c_n = \left\{ \int_A \phi^{-p+1}(x) \int |K(u)|^w \psi_{rw}(x+h_n u) du dx \right\}^{(2-\eta)/2}$$

$$\leq \left( (\alpha(A))^{-(p-1)} E|Y|^{rw} \int |K|^w \right)^{(2-\eta)/2}$$

and

$$c_n' = \left\{ \int_A \phi^{-p+1}(x) \int |K(u)|^2 \psi_{2r}(x+h_n u) du dx \right\}^{\eta/2}$$

$$\leq \left( (\alpha(A))^{-(p-1)} E|Y|^{2r} \int |K|^2 \right)^{\eta/2}.$$

Hence we conclude that the l.h.s. of (3.7) for  $p > 2$  is  $O(n^{-\eta/2} h_n^{-d(p-1)})$  under the moment condition in  $Y$  stated there. Now the proof of the Lemma is complete.  $\square$

### Proof of Theorem 3.1

It can be easily seen that

$$|\hat{\mu}_r(x) - \mu_r(x)|^p \leq 2^{p-1} (|\hat{\mu}_r(x) - \mu_r(x)|^p + |\hat{\mu}_r(x) - E\hat{\mu}_r(x)|^p)$$

$$= 2^{p-1} \phi^{-p}(x) (|E\hat{\psi}_r(x) - \psi_r(x)|^p + |\hat{\psi}_r(x) - E\hat{\psi}_r(x)|^p)$$

from (2.1) and (2.3). Therefore

$$\begin{aligned} \|\hat{\mu}_r - \mu_r\|^P &\leq 2^{P-1} \left\{ \int_A \phi^{-P}(x) |E\hat{\psi}_r(x) - \psi_r(x)|^P d\lambda(x) \right. \\ &\quad \left. + \int_A \phi^{-P}(x) |E\hat{\psi}_r(x) - E\hat{\psi}_r(x)|^P d\lambda(x) \right\}. \end{aligned}$$

The proof is complete from Lemmas 3.1 and 3.2.  $\square$

### Proof of Theorem 3.2

Note that

$$\begin{aligned} |\hat{\sigma}^2(x) - \sigma^2(x)|^P &\leq |(\hat{\mu}_2(x) - \mu_2(x)) - (\tilde{\mu}^2(x) - \mu^2(x))|^P \\ &\leq 2^{P-1} \left\{ |\hat{\mu}_2(x) - \mu_2(x)|^P + (2B)^P (|\hat{\mu}(x) - \mu(x)| \wedge B)^P \right\} \end{aligned}$$

since  $|E(Y|X)| = |\mu(X)| \leq B$  w.p.1 and by definition,  $\tilde{\mu}$  is the retraction of  $\hat{\mu}_1$  to the interval  $[-B, B]$ . Thus proof for (3.2) follows from (3.1) applied with  $r = 1$  and 2.  $\square$

### Proof of Theorem 3.3

To prove (3.3), recall from Section 2 that the asymptotic variance of  $\hat{\mu}(x) = \hat{\mu}_1(x)$ , up to under  $(nh_n^d)^{-1}$  is

$$V(\hat{\mu}(x)) = (nh_n^d)^{-1} \frac{\sigma^2(x) \int K^2}{\phi(x)}.$$

Therefore, from (2.5)

$$\begin{aligned} (nh_n^d / \int K^2) |\hat{V}(\hat{\mu}(x)) - V(\hat{\mu}(x))| &\leq \phi^{-1}(x) |\hat{\sigma}^2(x) - \sigma^2(x)| \\ &\leq (\alpha(A))^{-1} |\hat{\sigma}^2(x) - \sigma^2(x)| \end{aligned}$$

Hence proof for (3.3) follows from (3.2).  $\square$

#### 4. CONCLUDING REMARKS

Note that for  $p = 1$  the integrals in (3.1) and (3.2) are over the whole space  $\mathbb{R}^d$ , and for  $p > 1$ , the set  $A$  in the definition (2.8) could be any arbitrary subset of the support of  $\phi$  satisfying the restriction (2.7). We remark that Stone (1982), in consideration of "in probability convergence" of  $\int_A |\theta^*(x) - \theta(x)|^p dx$  for a class of regression estimators  $\theta^*$  has used the same restriction (2.7) on  $A$ , and Härdle (1984) in consideration of some results on mean integrated squared error (i.e. the case of  $p=2$ ) of regression estimators  $\hat{\mu} = \hat{\mu}_1$  and  $\mu^* = (\hat{\psi}_1/\hat{\phi})$  have taken  $A = (0,1)^d$  with the same restriction (2.7) on  $A$ . In consideration of uniform (over  $A$ ) weak and strong convergences of  $\mu = (\hat{\psi}_1/\hat{\phi})$ ,  $A$  is invariably taken as a compact subset of the support of  $\phi$  with the restriction (2.7), (e.g. see Nadaraya (1964, 1965, 1970), Schuster and Yakowitz (1979), Mack and Silverman (1981), among others). The condition (2.7) on  $A$  is also imposed in Singh and Ahmad (1987) in consideration of uniform (over  $A$ ) mean square consistency of their regression estimator  $\mu_* = (\hat{\psi}_1/\hat{\phi})_b$ , where  $b$  is an w.p.1 bound for  $|Y|$ .

If  $\phi$  is known, it makes sense to consider Nadaraya-Watson type estimator  $\hat{\mu} = \hat{\mu}_1$ , utilizing the knowledge of  $\phi$ . Greblicki and Krzyzak (1980), Johnston (1982) and Härdle (1986) have considered such estimators as well. Under the assumptions that  $\phi$  and  $\mu$  are continuous at  $x$  and  $E(Y^2) < \infty$ , Greblicki and Kryzyak established weak consistency of  $\hat{\mu}$  at  $x$  and Johnston examined the asymptotic distribution of the maximal deviation of  $\hat{\mu}(x)$  from  $\mu(x)$  while Härdle established  $L_2$ -norm consistency of  $\hat{\mu}$  with  $A = (0,1)$ .

As indicated earlier, estimations of heteroskedasticity or of asymptotic variance of the regression estimates under weaker conditions

have drawn little attention in the literature, and certainly none of the works mentioned in the preceding two paragraphs have discussed such estimations. Recently Singh (1989) considered estimators  $\tilde{\theta}_r(x) = (\hat{\psi}_r(x)/\hat{\phi}(x))$  for  $r \geq 0$  and  $\tilde{\sigma}^2(x) = \{\tilde{\theta}_2^2(x) - (\tilde{\theta}_1(x))^2\}$  respectively for the  $r$ th order regression  $\mu_r(x) = E(Y^r|X=x)$  and heteroskedasticity  $\sigma^2(x)$ , and established weak and strong consistencies, as well as the asymptotic normality of  $\tilde{\theta}_r$  and  $\tilde{\sigma}^2$  only under certain moment conditions on  $Y$ . Weak and strong consistencies of the estimator  $\tilde{V}(\tilde{\theta}_1(x)) = (nh_n^d)^{-1}(\tilde{\sigma}^2(x)/\hat{\phi}(x))JK^2$  of the asymptotic variance  $V(\tilde{\theta}_1(x)) = (nh_n^d)^{-1}(\sigma^2(x)/\phi(x))JK^2$  of the regression estimate  $\tilde{\theta}_1(x)$  follow implicitly from his similar results on  $\tilde{\theta}_r$  and  $\tilde{\sigma}^2$ . We wish to point out that the weak and strong consistencies of  $\hat{\mu}_r$ ,  $\hat{\sigma}^2$  and  $\hat{V}(\hat{\mu})$  also follow by the same arguments. However, we note that weak and strong consistencies do not imply  $L_p$ -norm consistency for any  $p \geq 1$ . In this paper we have established, for  $1 \leq p < \infty$ , the  $L_p$ -norm consistencies of  $\hat{\mu}_r$ ,  $\hat{\sigma}^2$  and  $\hat{V}(\hat{\mu})$ , again only under certain moment conditions on  $Y$ .

Singh and Giles (1991) considered regression estimator  $\mu_* = (\hat{\psi}_1/\hat{\phi})_b$  and heteroskedasticity estimator  $\sigma_*^2 = ((\hat{\psi}_2/\hat{\phi})_b - \mu_*^2)$  under the condition that  $|Y|$  is bounded w.p. 1 by  $b$ , and established the  $L_1$ -norm consistencies (and hence  $L_p$ -norm consistencies for any  $p \geq 1$ , since  $(\mu_* - \mu) \leq b$  and  $|\sigma_*^2 - \sigma^2| \leq b^2$ ). In our estimations of the regression, heteroskedasticity and the asymptotic variance of the regression estimate though  $\phi$  is assumed to be known but  $Y$  need not be bounded.

## REFERENCES

- Box, G.E.P. and Hill, W.J. (1974) Correcting inhomogeneity of variance with power transformation weighting. *Technometrics* 16, 385-389.
- Carroll, R.J. (1982) Adapting for heteroscedasticity in linear models. *Ann. Statist.* 10, 1224-1233.
- Deheuvels, P. (1974). Conditions nécessaires et suffisantes de convergence ponctuelle presque sûre et uniforme presque sûre des estimateurs de al densité. *C.R. Acad. Sci Paris* 278, 1217-1220.
- Froehlich, B.R. (1973). Some estimators for a random coefficient regression model. *J. Amer. Statist. Assoc.* 68, 329-335,
- Fuller, W.A. and Rao, J.N.K. (1978). Estimation for a linear regression model with unknown diagonal covariance matrix. *Ann. Statist.* 6, 1149-1158.
- Greblick, W. and Krzyzak, A. (1980). Asymptotic properties of kernel estimates of a regression function. *J. Statist. Plan. Inf.* 4, 81-90.
- Hall, P. (1981). Laws of the iterated logarithm for nonparametric density estimators. *Z. Wahrscheinlichkeitstheorie verw. Gebiete* 50, 47-61.
- Härdle, W. (1984). Robust regression function estimation. *J. Multi. Anal.* 14, 169-180.
- Härdle, W. and Luckhaus, S. (1984). Uniform consistency of a class of regression function estimates. *Ann. Statist.* 12, 612-623.
- Härdle, W. (1986). Approximations to the mean integrated squared error with applications to optimal bandwidth selection for nonparametric regression function estimators. *J. Multi. Anal.* 18, 150-168.
- Hildreth, C. and Houck, J.P. (1968). Some estimators for a linear model with random coefficients. *J. Amer. Statist. Assoc.* 63, 584-595.
- Jobson, J.D. and Fuller, W.A. (1980). Least squares estimation when the matrix and parameter vector are functionally related. *J. Amer. Statist. Assoc.* 75, 176-181.
- Johnston, G.J. (1982). Probabilities of maximal deviations of nonparametric regression function estimation. *J. Multi. Anal.* 12, 402-414.
- Judge, G.G., Hill, W.J., Griffiths, W.E., Lütkepohl, H. and Lee, T.C. (1988). *The Theory and Practice of Econometrics*. John Wiley.
- Mack, Y.P. and Silverman, B.W. (1982). Weak and strong uniform consistency of kernel regression estimates. *Z. Wahrscheinlichkeitstheorie verw. Gebiete* 61, 405-415.

- MacKinnon, J.G. and White, H. (1985). "Some heteroskedastic-consistent covariance matrix estimators with improved finite sample properties. *J. of Econometrics* 29, 305-327.
- Müller, H.G. and Stadtmüller, U. (1987). Estimation of heteroskedasticity in regression analysis. *Ann. Statist.* 15, 2, 610-625.
- Nadaraya, E.A. (1964). On estimating regression. *Theor. Prob. Appl.* 9, 141-142.
- Nadaraya, E.A. (1965). On nonparametric estimates of a density function and regression curves. *Theor. Prob. Appl.* 10, 186-190.
- Nadaraya, E.A. (1970). Remarks on nonparametric estimates for density functions and regression curves. *Theor. Prob. Appl.* 15, 134-137.
- Natanson, I.P. (1955). *Theory of Functions of a Real Variable*. Frederick Ungar, New York.
- Noda, K. (1976). Estimation of a regression function by the Parzen kernel-type density estimators. *Ann. Inst. Statist. Math.* 28, 221-234.
- Rosenblatt, M. (1969). Conditional probability density and regression estimates. In: *Multivariate Analysis II*, ed. Krishnaiah, P.R. 25-31.
- Schuster, E.F. (1972). Joint asymptotic distribution of the estimated regression function at a finite number of distinct points. *Ann. Math. Statist.* 42, 84-88.
- Schuster, E.F. and Yakowitz, S. (1979). Contributions to the theory of nonparametric regression, with applications to system identification. *Ann. Statist.* 7, 139-149.
- Singh, Radhey S. and Ullah, A. (1985). Nonparametric time-series estimation of joint DGP, conditional DGP, and vector autoregression. *Econometric Th.* 1, 27-52.
- Singh, Radhey S. and Ullah, A. and Carter R.A.L. (1987). Nonparametric inference in econometrics: New applications. *Time Series and Econometric Modeling* eds. I.B. MacNeill and Umphrey G.J., 253-278.
- Singh, Radhey S. and Ahmad, M. (1987). Modified nonparametric kernel estimates of a regression function and their consistencies with rates. *Ann. Inst. Statist. Math.* 39, 549-562.
- Singh, Radhey S. (1989). Nonparametric Estimation of heteroskedasticity in regression with consistency and normality under weaker conditions. University of Guelph mimeo.
- Singh, Radhey S. and Giles, David E.A. (1991). Nonparametric estimation of regression and heteroscedasticity with  $L_1$ -norm consistencies. University of Canterbury mimeo.
- Stein, E.M. and Weiss, G. (1975). *Introduction to Fourier Analysis on Euclidean Spaces*. Princeton, New Jersey.



- Stone, C. (1982). Optimal global rates of convergence for nonparametric regression. *Ann. Statis.* 5, 1595-645.
- Watson, G.S. (1964). Smooth regression analysis, *Sankya Ser. A.* 26, 359-372.
- Wheeden, R.L. and Zygmund, A. (1977). *Measure and Integrals.* Dekker, New York.
- White, H. (1980). A heteroskedasticity-consistent covariance matrix and a direct test for heteroskedasticity. *Econometrica* 48, 817-838.

## LIST OF DISCUSSION PAPERS\*

- No. 8701 Stochastic Simulation of the Reserve Bank's Model of the New Zealand Economy, by J. N. Lye.
- No. 8702 Urban Expenditure Patterns in New Zealand, by Peter Hampton and David E. A. Giles.
- No. 8703 Preliminary-Test Estimation of Mis-Specified Regression Models, by David E. A. Giles.
- No. 8704 Instrumental Variables Regression Without an Intercept, by David E. A. Giles and Robin W. Harrison.
- No. 8705 Household Expenditure in Sri Lanka: An Engel Curve Analysis, by Mallika Dissanayake and David E. A. Giles.
- No. 8706 Preliminary-Test Estimation of the Standard Error of Estimate in Linear Regression, by Judith A. Clarke.
- No. 8707 Invariance Results for FIML Estimation of an Integrated Model of Expenditure and Portfolio Behaviour, by P. Dorian Owen.
- No. 8708 Social Cost and Benefit as a Basis for Industry Regulation with Special Reference to the Tobacco Industry, by Alan E. Woodfield.
- No. 8709 The Estimation of Allocation Models With Autocorrelated Disturbances, by David E. A. Giles.
- No. 8710 Aggregate Demand Curves in General-Equilibrium Macroeconomic Models: Comparisons with Partial-Equilibrium Microeconomic Demand Curves, by P. Dorian Owen.
- No. 8711 Alternative Aggregate Demand Functions in Macro-economics: A Comment, by P. Dorian Owen.
- No. 8712 Evaluation of the Two-Stage Least Squares Distribution Function by Imhof's Procedure by P. Cribbitt, J. N. Lye and A. Ullah.
- No. 8713 The Size of the Underground Economy: Problems and Evidence, by Michael Carter.
- No. 8714 A Computable General Equilibrium Model of a Fisherine Method to Close the Foreign Sector, by Ewen McCann and Keith McLaren.
- No. 8715 Preliminary-Test Estimation of the Scale Parameter in a Mis-Specified Regression Model, by David E. A. Giles and Judith A. Clarke.
- No. 8716 A Simple Graphical Proof of Arrow's Impossibility Theorem, by John Fountain.
- No. 8717 Rational Choice and Implementation of Social Decision Functions, by Manimay Sen.
- No. 8718 Divisia Monetary Aggregates for New Zealand, by Ewen McCann and David E. A. Giles.
- No. 8719 Telecommunications in New Zealand: The Case for Reform, by John Fountain.
- No. 8801 Workers' Compensation Rates and the Demand for Apprentices and Non-Apprentices in Victoria, by Pasquale M. Sgro and David E. A. Giles.
- No. 8802 The Adventures of Sherlock Holmes, the 48% Solution, by Michael Carter.
- No. 8803 The Exact Distribution of a Simple Pre-Test Estimator, by David E. A. Giles.
- No. 8804 Pre-testing for Linear Restrictions in a Regression Model With Student-t Errors, by Judith A. Clarke.
- No. 8805 Divisia Monetary Aggregates and the Real User Cost of Money, by Ewen McCann and David Giles.
- No. 8806 The Management of New Zealand's Lobster Fishery, by Alan Woodfield and Pim Borren.
- No. 8807 Poverty Measurement: A Generalization of Sen's Result, by Prasanta K. Pattanaik and Manimay Sen.
- No. 8808 A Note on Sen's Normalization Axiom for a Poverty Measure, by Prasanta K. Pattanaik and Manimay Sen.
- No. 8809 Budget Deficits and Asset Sales, by Ewen McCann.
- No. 8810 Unorganized Money Markets and 'Unproductive' Assets in the New Structuralist Critique of Financial Liberalization, by P. Dorian Owen and Otton Solis-Fallas.
- No. 8901 Testing for Financial Buffer Stocks in Sectoral Portfolio Models, by P. Dorian Owen.
- No. 8902 Provisional Data and Unbiased Prediction of Economic Time Series by Karen Browning and David Giles.
- No. 8903 Coefficient Sign Changes When Restricting Regression Models Under Instrumental Variables Estimation, by David E. A. Giles.

*(Continued on next page)*

- No. 8904 Economies of Scale in the New Zealand Electricity Distribution Industry, by David E. A. Giles and Nicolas S. Wyatt.
- No. 8905 Some Recent Developments in Econometrics: Lessons for Applied Economists, by David E. A. Giles.
- No. 8906 Asymptotic Properties of the Ordinary Least Squares Estimator in Simultaneous Equations Models, by V. K. Srivastava and D. E. A. Giles.
- No. 8907 Unbiased Estimation of the Mean Squared Error of the Feasible Generalised Ridge Regression Estimator, by V. K. Srivastava and D. E. A. Giles.
- No. 8908 An Unbiased Estimator of the Covariance Matrix of the Mixed Regression Estimator, by D. E. A. Giles and V. K. Srivastava.
- No. 8909 Pre-testing for Linear Restrictions in a Regression Model with Spherically Symmetric Disturbances, by Judith A. Giles.
- No. 9001 The Durbin-Watson Test for Autocorrelation in Nonlinear Models, by Kenneth J. White.
- No. 9002 Determinants of Aggregate Demand for Cigarettes in New Zealand, by Robin Harrison and Jane Chetwyd.
- No. 9003 Unemployment Duration and the Measurement of Unemployment, by Manimay Sengupta.
- No. 9004 Estimation of the Error Variance After a Preliminary-Test of Homogeneity in a Regression Model with Spherically Symmetric Disturbances, by Judith A. Giles.
- No. 9005 An Expository Note on the Composite Commodity Theorem, by Michael Carter.
- No. 9006 The Optimal Size of a Preliminary Test of Linear Restrictions in a Mis-specified Regression Model, by David E. A. Giles, Offer Lieberman, and Judith A. Giles.
- No. 9007 Inflation, Unemployment and Macroeconomic Policy in New Zealand: A Public Choice Analysis, by David J. Smyth and Alan E. Woodfield.
- No. 9008 Inflation — Unemployment Choices in New Zealand and the Median Voter Theorem, by David J. Smyth and Alan E. Woodfield.
- No. 9009 The Power of the Durbin-Watson Test when the Errors are Heteroscedastic, by David E. A. Giles and John P. Small.
- No. 9010 The Exact Distribution of a Least Squares Regression Coefficient Estimator After a Preliminary t-Test, by David E. A. Giles and Virendra K. Srivastava.
- No. 9011 Testing Linear Restrictions on Coefficients in a Linear Regression Model with Proxy variables and Spherically Symmetric Disturbances, by Kazuhiro Ohtani and Judith A. Giles.
- No. 9012 Some Consequences of Applying the Goldfeld-Quandt Test to Mis-Specified Regression Models, by David E. A. Giles and Guy N. Saxton.
- No. 9013 Pre-testing in a Mis-specified Regression Model, by Judith A. Giles.
- No. 9014 Two Results in Balanced-Growth Educational Policy, by Alan E. Woodfield.
- No. 9101 Bounds on the Effect of Heteroscedasticity on the Chow Test for Structural Change, by David Giles and Offer Lieberman.
- No. 9102 The Optimal Size of a Preliminary Test for Linear Restrictions when Estimating the Regression Scale Parameter, by Judith A. Giles and Offer Lieberman.
- No. 9103 Some Properties of the Durbin-Watson Test After a Preliminary t-Test, by David Giles and Offer Lieberman.
- No. 9104 Preliminary-Test Estimation of the Regression Scale Parameter when the Loss Function is Asymmetric, by Judith A. Giles and David E. A. Giles.
- No. 9105 On an Index of Poverty, by Manimay Sengupta and Prasanta K. Pattanaik.
- No. 9106 Cartels May Be Good For You, by Michael Carter and Julian Wright.
- No. 9107  $L_p$ -Norm Consistencies of Nonparametric Estimates of Regression, Heteroskedasticity and Variance of Regression Estimate when Distribution of Regressor is Known, by Radhey S. Singh.

\* Copies of these Discussion Papers may be obtained for \$4 (including postage, price changes occasionally) each by writing to the Secretary, Department of Economics, University of Canterbury, Christchurch, New Zealand. A list of the Discussion Papers prior to 1987 is available on request.