



*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

*No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.*

CANTER

9116✓

Department of Economics  
UNIVERSITY OF CANTERBURY

CHRISTCHURCH, NEW ZEALAND

ISSN 1171-0705



ANNINI FOUNDATION  
RIGULTURAL/ECONOMICS  
LIBRARY

FEB 05 1992

ESTIMATION OF THE SCALE PARAMETER  
AFTER A PRE-TEST FOR HOMOGENEITY IN A  
MIS-SPECIFIED REGRESSION MODEL

Judith A. Giles

*Discussion Paper*

No. 9116



This paper is circulated for discussion and comments. It should not be quoted without the prior approval of the author. It reflects the views of the author who is responsible for the facts and accuracy of the data presented. Responsibility for the application of material to specific cases, however, lies with any user of the paper and no responsibility in such cases will be attributed to the author or to the University of Canterbury.

Department of Economics, University of Canterbury  
Christchurch, New Zealand

***Discussion Paper No. 9116***

December 1991

**ESTIMATION OF THE SCALE PARAMETER  
AFTER A PRE-TEST FOR HOMOGENEITY IN A  
MIS-SPECIFIED REGRESSION MODEL**

**Judith A. Giles**

# ESTIMATION OF THE SCALE PARAMETER AFTER A PRE-TEST FOR HOMOGENEITY IN A MIS-SPECIFIED REGRESSION MODEL

Judith A. Giles

Department of Economics  
University of Canterbury  
Private Bag, Christchurch  
New Zealand

*Key Words and Phrases: pre-test estimation; mis-specified regression models; spherical symmetry; multivariate Student-t.*

## ABSTRACT

The risk properties of estimators of the scale parameter after a pre-test for homogeneity of the error variances in the two sample linear regression model has received quite an amount of attention in the literature. This literature typically assumes normal disturbances and a properly specified model. In this paper we consider that both equations may be mis-specified by the omission of relevant regressors and that the error distributions may belong to a wider class than the normal distribution. We derive and analyse the exact risk (under quadratic loss) of the pre-test estimator of the scale parameter for the first sub-sample.

## 1. INTRODUCTION AND MODEL FRAMEWORK

We consider a simple heteroscedastic linear regression model in which the error variance is constant within each sample but it may be different between the samples,

$$\begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} X_1 & 0 \\ 0 & X_2 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} + \begin{bmatrix} Z_1 & 0 \\ 0 & Z_2 \end{bmatrix} \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} + \begin{bmatrix} e_1 \\ e_2 \end{bmatrix} \quad (1)$$

or

$$y = X\beta + Z\gamma + e$$

where for  $i=1,2$ ,  $y_i$  is a  $(T_i \times 1)$  vector of observations on the dependent variable,  $X_i$  is a known  $(T_i \times k_i)$  non-stochastic design matrix of rank  $k_i$  ( $< T_i$ ),  $Z_i$  is a fixed  $(T_i \times p_i)$  matrix of full rank,  $\gamma_i$  is a

$(p_i \times 1)$  coefficient vector, and  $\beta_i$  is a  $(k_i \times 1)$  vector of unknown parameters. Let  $T = T_1 + T_2$ .

We assume that  $E(e_i) = 0$ , and that  $E(e_i e_i') = \sigma_{e_i}^2 I_{T_i}$ . Let  $\phi = \sigma_{e_1}^2 / \sigma_{e_2}^2$  so that

$$E(ee') = \begin{bmatrix} \sigma_{e_1}^2 I_{T_1} & 0 \\ 0 & \sigma_{e_2}^2 I_{T_2} \end{bmatrix} = \sigma_{e_2}^2 \begin{bmatrix} \phi I_{T_1} & 0 \\ 0 & I_{T_2} \end{bmatrix} = \sigma_{e_2}^2 \Sigma. \quad (2)$$

We suppose that  $e$  has a non-normal distribution of the form  $f(e) = \int_0^\infty f_N(e) f(\tau) d\tau$ , where  $f_N(e)$  is  $f(e)$  when  $e \sim N(0, \tau^2 \Sigma)$ , and  $f(\tau)$  is supported on  $[0, \infty)$ . Then  $\sigma_{e_2}^2 = E(\tau^2)$ ,  $\sigma_{e_1}^2 = \phi E(\tau^2)$ , and  $e$  has an elliptically symmetric distribution ( $ESD_N$ ) when  $\phi \neq 1$  but a spherically symmetric distribution ( $SSD_N$ ) when the error variances are equal.<sup>1,2</sup> This is the commonly called scale mixture of normal distributions family, of which the multivariate Student-t (Mt) distribution is the most well known member. The Mt family arises if  $f(\tau)$  is an inverted gamma density with, say, scale parameter  $\sigma_2^2$  and degrees of freedom parameter  $\nu$ . We then write  $e \sim \text{Mt} \left( 0, (\nu \sigma_2^2 / (\nu - 2)) \Sigma \right)$ . For this member the marginal distributions are univariate Student-t with thicker tails than under a corresponding normality assumption for  $\nu < \infty$ . The kurtosis increases as  $\nu$  decreases and  $\nu = \infty$  corresponds to normality.

Rather than model (1) being estimated we suppose that the proposed model is

$$y = X\beta + u, \quad u \sim N \left( 0, \sigma_{e_2}^2 \Sigma \right) \quad (3)$$

which the researcher assumes is correctly specified. In fact, (3) is

<sup>1</sup> A discussion of this family of distributions is beyond the scope of this paper. See, for example, Kelker (1970), Muirhead (1982), and Dickey and Chen (1985).

<sup>2</sup> It would be relatively straightforward to extend our analysis to the case of different mixing distributions for each sample when we have independent mixing distributions. It is unclear, however, how we would proceed if they are dependent.

mis-specified as  $u \sim \text{ESD}_N(Z\gamma, \sigma_e^2 \Sigma)$ . Note that (3) reflects the fact that  $\sigma_{u_i}^2 = \sigma_{e_i}^2$ ,  $i=1,2$ . The researcher is interested in estimating  $\sigma_{e_1}^2$  but he is uncertain of the homogeneity of the error variances and so conducts a pre-test of

$$H_0 : \phi = 1 \text{ vs } H_1 : \phi < 1, \quad (4)$$

where  $\phi$  is a measure of the hypothesis error and we assume, for simplicity, a one-sided alternative hypothesis though the analysis could be easily extended to the two-sided case.

Assuming the usual least squares estimators of the error variances, the researcher, mistakenly proceeding as if (3) is properly specified, has three options for the estimation of  $\sigma_{e_1}^2$ :

(1) He could assume that the variances are equal and use the so-called always pool estimator of  $\sigma_{e_1}^2$ ,  $s_A^2$ :

$$s_A^2 = (v_1 s_1^2 + v_2 s_2^2) / (v_1 + v_2) \quad (5)$$

where  $s_i^2 = (y_i - X_i b_i)'(y_i - X_i b_i) / v_i$ ,  $v_i = T_i - k_i$ , and  $b_i = (X_i' X_i)^{-1} X_i' y_i$ ,  $i=1,2$ .

(2) He could proceed as if the error variances are unequal, ignore the second sample, and use the so-called never pool estimator of  $\sigma_{e_1}^2$ ,  $s_N^2$ :

$$s_N^2 = s_1^2 \quad (6)$$

(3) He could undertake a preliminary test of the validity of  $H_0$  and use  $s_A^2$  if he accepts  $H_0$  or  $s_N^2$  if he rejects  $H_0$ . As the researcher assumes that model (3) is correctly specified, he uses the usual  $J$  test for homoscedasticity, given by  $J = s_2^2 / s_1^2$ , and proceeds as if  $f(J) = \phi^{-1} f(F_{(v_2, v_1)})$ , where  $F_{(v_2, v_1)}$  is a central  $F$  variate with  $v_2$  and  $v_1$  degrees of freedom. Under this option, he is in fact using the pre-test estimator  $s_P^2$ :

$$s_P^2 = \begin{cases} s_N^2 & \text{if } J > c \\ s_A^2 & \text{if } J \leq c \end{cases} = I_{[0, c)}(J) s_A^2 + I_{(c, \infty)}(J) s_N^2 \quad (7)$$

where  $I_{[...]}(J)$  equals one if  $J$  lies within the subscripted range, zero otherwise, and  $c$  is the critical value of the pre-test corresponding to a nominal test size of  $\alpha$  such that  $\int_c^\infty f(F(v_2, v_1)) = \alpha$ .

Assuming normal disturbances,  $e \sim N(0, \sigma_e^2 \Sigma)$ , and a correctly specified design matrix, the risk of  $s_P^2$ , defined as  $\rho(\sigma_e^2, s_P^2) = E(s_P^2 - \sigma_e^2)^2$ , is considered by Bancroft (1944), Toyoda and Wallace (1975), Ohtani and Toyoda (1978), and Bancroft and Han (1983).<sup>3</sup> Figure 1 presents a typical relative risk function under their assumptions, where the relative risk is  $R(s_P^2) = \rho(\sigma_e^2, s_P^2) / \sigma_e^4$ .<sup>4</sup>

We see from Figure 1 that it is only preferable to pool the samples around the neighbourhood of the null hypothesis and that we should never use the never pool estimator, as there is a family of pre-test estimators with  $c \in (0, 2)$  which strictly dominate this estimator (see Toyoda and Wallace (1975)). Ohtani and Toyoda (1978) prove that the pre-test estimator which uses  $c=1$  has the smallest risk of this dominating family.

Giles (1992) extends these aforementioned studies to the error distribution framework considered in this paper, though she still maintains the (unrealistic) assumption of a correctly specified design matrix. In the case that she examines we know that the test statistic  $J$  maintains the same null and non-null distributions as under normal disturbances from the results of King (1979) and Chmielewski (1981). Figure 2 illustrates typical relative risk functions under Mt disturbances  $\left[ e \sim Mt \left( 0, (\nu \sigma_e^2 / (\nu - 2)) \Sigma \right) \right]$  when  $\nu = 5$ .

<sup>3</sup>A related, though not identical, pre-test problem is considered by Yancey *et al.* (1983) and Ohtani (1987). Both of these studies also assume normal errors.

<sup>4</sup>We lose no loss in generality in considering relative risk; the results could equally be interpreted as the risk functions when  $\sigma_e^2 = 1$ .






Figure 1. Relative risk functions for  $s_N^2$ ,  $s_A^2$ , and  $s_P^2$  when  $e \sim N(0, \sigma_2^2 \Sigma)$ ,  $v_1=16$ ,  $v_2=8$ ,  $k_1=k_2=3$ .




Figure 2. Relative risk functions for  $s_N^2$ ,  $s_A^2$ , and  $s_P^2$  when  $e \sim Mt\left(0, (\nu\sigma_2^2/(\nu-2))\Sigma\right)$ ,  $v_1=16$ ,  $v_2=8$ ,  $k_1=k_2=3$ ,  $\nu=5$ .

We see from Figure 2 that for non-normal disturbances we should sometimes always pre-test, *even if the error variances are equal*. Giles (1992) proves that the optimal critical value to use in these circumstances is  $c=1$ .

In this paper we recognise that in reality departures from the standard regression assumptions are likely to occur simultaneously and we, accordingly, examine the risk properties of  $s_N^2$ ,  $s_A^2$ , and  $s_P^2$  when the disturbances are elliptically symmetric and we have omitted relevant regressors from the design matrix. The impact of excluding regressors on the risk functions of the estimators of the error variance for this particular pre-test problem has yet to be examined in the pre-test literature, even under normality. Other studies, though, have investigated the effect of this mis-specification after other pre-tests (see, for example, Ohtani (1983), Mittelhammer (1984), Giles (1986), Ohtani (1987), Giles and Clarke (1989), and Giles (1991b)).

To undertake this task we need to first examine the distribution of the test statistic  $J$ , which is now no longer a function of a central  $F$  variate under either the null or the alternative hypotheses. We consider this in the next section. We follow in section 3 with the derivation of the exact risk functions, and then in section 4 with some numerical evaluations of these risk functions, assuming for the purpose of this discussion that the variance mixing distribution is inverted gamma so that our errors are  $M_t$ . The final section provides some concluding remarks.

## 2. THE DISTRIBUTION OF $J$ WITH OMITTED REGRESSORS

If the model is mis-specified in the way investigated here then both the null and the non-null distributions of  $J$  depend on  $v_1$ ,  $v_2$ , the degree of mis-specification of the design matrix, and the variance mixing distribution,  $f(\tau)$ . This is shown by Theorem 1 and Corollary 1.

**Theorem 1.** Under the stated assumptions, the density function of  $J = s_2^2/s_1^2$  is

$$f(J) = \phi^{-1} \sum_{r=0}^{\infty} \sum_{s=0}^{\infty} \frac{\theta_1^s \theta_2^r v_1^{v_1/2+s} v_2^{v_2/2+r} J^{v_2/2+r-1}}{r!s!B\left(\frac{v_2}{2}+r; \frac{v_1}{2}+s\right) \left(v_1+v_2J\right)^{(v_1+v_2)/2+r+s}} \times \int_0^{\infty} e^{-(\theta_1+\theta_2)/\tau^2} \tau^2 \left(\tau^2\right)^{-(r+s)} f(\tau) d\tau, \quad (8)$$

where  $\theta_1 = \gamma_1' Z_1' M_1 Z_1 \gamma_1 / (2\phi)$ ,  $\theta_2 = \gamma_2' Z_2' M_2 Z_2 \gamma_2 / 2$ ,  $M_i = I_{T_i} - X_i' (X_i' X_i)^{-1} X_i'$ , and  $B(.,.)$  is the beta function.

**Proof.** See the appendix.

**Corollary 1.** Under the null hypothesis,  $\phi=1$ , and

$$f(J) = \sum_{r=0}^{\infty} \sum_{s=0}^{\infty} \frac{\theta_{10}^s \theta_2^r v_1^{v_1/2+s} v_2^{v_2/2+r} J^{v_2/2+r-1}}{r!s!B\left(\frac{v_2}{2}+r; \frac{v_1}{2}+s\right) \left(v_1+v_2J\right)^{(v_1+v_2)/2+r+s}} \times \int_0^{\infty} e^{-(\theta_{10}+\theta_2)/\tau^2} \tau^2 \left(\tau^2\right)^{-(r+s)} f(\tau) d\tau, \quad (9)$$

where  $\theta_{10} = \gamma_1' Z_1' M_1 Z_1 \gamma_1 / 2$ .

**Proof.** As  $\theta_1 = \theta_{10}$  when  $\phi=1$ , (9) follows from (8).

If  $e \sim N(0, \sigma_e^2 \Sigma)$  and, say,  $\sigma_e^2 = \sigma_2^2$  and  $\sigma_{e_1}^2 = \sigma_1^2$ , then from (8)

$$f_N(J) = \phi^{-1} f\left(F'_{(v_2, v_1; \lambda_2, \lambda_1)}\right) \quad (10)$$

where  $\lambda_i = \theta_i / \sigma_i^2$ ,  $i=1,2$ . Note that (10) does not collapse to a central F density under  $H_0$ ; that is, the standard assumption that J has a central F density under  $H_0$  is invalid if we have omitted relevant regressors, irrespective of our assumption on the distribution of the

disturbances. So, there will be a difference between the nominal and true sizes of the test.

In this section we have established the distribution of the test statistic  $J$  which we use to test for the homogeneity of the error variances. In the next section we derive the exact risk functions of the never pool, always pool, and pre-test estimators of  $\sigma_{e_1}^2$ .

### 3. THE RISK FUNCTIONS

**Theorem 2.** If we use the mis-specified model (3) rather than the true model (1) when  $e \sim \text{ESD}_N(0, \sigma_{e_2}^2 \Sigma)$  and the pre-test is of  $H_0$  in (4) then

$$\rho(\sigma_{e_1}^2, s_N^2) = \phi^2 \left[ v_1(v_1+2)E(\tau^4) - v_1^2 \left( E(\tau^2) \right)^2 + 8\theta_1 E(\tau^2) + 4\theta_1^2 \right] / v_1^2 \quad (11)$$

$$\begin{aligned} \rho(\sigma_{e_1}^2, s_A^2) = & \left\{ \phi^2 \left[ v_1(v_1+2)E(\tau^4) + 4(v_1+2)\theta_1 E(\tau^2) - 2v_1(v_1+v_2) \left( E(\tau^2) \right)^2 \right. \right. \\ & + \left. \left. \left( 2\theta_1 - E(\tau^2)(v_1+v_2) \right)^2 \right] + 2\phi \left[ v_1 v_2 E(\tau^4) - v_2(v_1+v_2) \left( E(\tau^2) \right)^2 \right. \right. \\ & + \left. \left. 2v_2 E(\tau^2)(\theta_1 - \theta_2) + 4\theta_1 \theta_2 \right] + v_2(v_2+2)E(\tau^4) + 4(v_2+2)\theta_2 E(\tau^2) + 4\theta_2^2 \right\} \\ & / (v_1+v_2)^2 \quad (12) \end{aligned}$$

$$\begin{aligned} \rho(\sigma_{e_1}^2, s_P^2) = & \left\{ \phi^2 (v_1+v_2)^2 \left[ v_1(v_1+2)E(\tau^4) - v_1^2 \left( E(\tau^2) \right)^2 + 8\theta_1 E(\tau^2) + 4\theta_1^2 \right] \right. \\ & + \int_0^\infty \left\{ \phi^2 v_2 \left[ -(2v_1+v_2) \left( v_1(v_1+2)\tau^4 Q_{04}^\tau + 4(v_1+2)\theta_1 \tau^2 Q_{06}^\tau + 4\theta_1^2 Q_{08}^\tau \right) \right. \right. \\ & + \left. \left. 2E(\tau^2)v_1(v_1+2) \left( v_1 \tau^2 Q_{02}^\tau + 2\theta_1 Q_{04}^\tau \right) \right] + 2v_1^2 \phi \left[ v_1 v_2 \tau^4 Q_{22}^\tau \right. \right. \\ & - \left. \left. v_2(v_1+v_2)E(\tau^2)\tau^2 Q_{20}^\tau + 2v_1 \theta_2 \tau^2 Q_{42}^\tau - 2(v_1+v_2)\theta_2 E(\tau^2)Q_{40}^\tau \right] \right\} \end{aligned}$$

$$\begin{aligned}
& +2v_2\theta_1\tau^2Q_{24}^\tau+4\theta_1\theta_2\tau^2Q_{44}^\tau\Big]+v_1^2\Big[v_2(v_2+2)\tau^4Q_{40}^\tau \\
& +4(v_2+2)\tau^2\theta_2Q_{60}^\tau+4\theta_2^2Q_{80}^\tau\Big]\Big]f(\tau)d\tau\Big)/\Big(v_1^2(v_1+v_2)^2\Big)
\end{aligned} \tag{13}$$

where

$$Q_{ij}^\tau = \text{Pr.} \left[ F'_{(v_2+i, v_1+j; \lambda_{2\tau}, \lambda_{1\tau})} \leq \left( v_2(v_1+j)c\phi \right) / \left( v_1(v_2+i) \right) \right]$$

$i, j=0, 1, 2, \dots$  and  $\lambda_{i\tau} = \theta_i / \tau^2$ ,  $i=1, 2$ .

**Proof.**

See the appendix.

**Remarks:**

(1) Aside from depending on the arguments of the model  $v_1$  and  $v_2$ , the risk functions depend on first, the true error variances  $\sigma_{e_1}^2$  and  $\sigma_{e_2}^2$  via  $\phi$ ; secondly, they depend on  $f(\tau)$ ; thirdly,  $\rho(\sigma_{e_1}^2, s_P^2)$  depends on the nominal significance level of the pre-test; and finally, the risk functions depend on the degree of mis-specification in each sample, via  $\theta_1$  and  $\theta_2$ .

(2) The data enter the risk functions only through  $\theta_1$  and  $\theta_2$ .

(3) It is straightforward to show that (11)-(13) collapse first to the appropriate expressions derived by Giles (1992) when the design matrix is in fact correctly specified, and secondly, to the risk functions derived by Toyoda and Wallace (1975), for example, when (3) is the valid model specification.

(4) If  $\alpha=0$  then  $c=\infty$  and  $Q_{ij}^\tau=1$  so that we never reject  $H_0$ . Then the risk of the pre-test estimator collapses to that of the always pool estimator. Conversely, if  $\alpha=1$  then  $c=0$  and  $Q_{ij}^\tau=0$  so that we always reject  $H_0$  and the risk of the pre-test estimator equals that of the never pool estimator.

(5)  $\lim_{\phi \rightarrow 0} \left[ \rho(\sigma_{e_1}^2, s_P^2) \right] = \lim_{\phi \rightarrow 0} \left[ \rho(\sigma_{e_1}^2, s_N^2) \right] = 0$  while  $\lim_{\phi \rightarrow 0} \left[ \rho(\sigma_{e_1}^2, s_A^2) \right] = \left( v_2(v_2+2)E(\tau^4) + 4(v_2+2)\theta_2E(\tau^2) + 4\theta_2^2 \right) / (v_1+v_2)^2 > 0$ . Pre-testing leads us to

follow the appropriate strategy of ignoring the prior information when

that information is very false.

(6)  $\rho(\sigma_{e_1}^2, s_N^2)$  is independent of  $\theta_2$  and therefore bounded as  $\theta_2 \rightarrow \infty$ , for a given value of  $\theta_1$ . However, this risk function is bounded as  $\theta_1 \rightarrow \infty$ , for a given value of  $\theta_2$ . Similarly,  $\rho(\sigma_{e_1}^2, s_P^2)$  is unbounded as  $\theta_1 \rightarrow \infty$ , given  $\theta_2$ , but it is bounded (by  $\rho(\sigma_{e_1}^2, s_N^2)$ ) as  $\theta_2 \rightarrow \infty$ , given  $\theta_1$ . Intuitively, if the model for the second sample is badly mis-specified relative to the model for the first sample, then pre-testing will lead us to ignore the second sample, which is the appropriate strategy.  $\rho(\sigma_{e_1}^2, s_A^2)$ , on the other hand, is unbounded as  $\theta_1 \rightarrow \infty$ , given  $\theta_2$ , or as  $\theta_2 \rightarrow \infty$ , given  $\theta_1$ .

Further,  $\left[ \rho(\sigma_{e_1}^2, s_A^2) - \rho(\sigma_{e_1}^2, s_N^2) \right]$  and  $\left[ \rho(\sigma_{e_1}^2, s_A^2) - \rho(\sigma_{e_1}^2, s_P^2) \right]$  are unbounded as  $\theta_1 \rightarrow \infty$ , given  $\theta_2$ , or as  $\theta_2 \rightarrow \infty$ , given  $\theta_1$ , while  $\left[ \rho(\sigma_{e_1}^2, s_N^2) - \rho(\sigma_{e_1}^2, s_P^2) \right]$  is bounded (and equal to zero) as  $\theta_2 \rightarrow \infty$ , given  $\theta_1$ , but it is unbounded as  $\theta_1 \rightarrow \infty$ , given  $\theta_2$ . These results imply, in particular, that the risk of the always pool estimator can be infinitely higher than that of the never pool estimator and the pre-test estimator even if the error variances are equal. That is, there is no guarantee of a reduction in risk by imposing valid prior information if we have omitted relevant regressors. This accords with the results of, for instance, Mittelhammer (1984) and Giles (1991b) in the case of estimating the coefficient vector in the classical linear regression model after a pre-test for exact linear restrictions.

(7) For any given degree of mis-specification through the omitted variables, there exists a family of pre-test estimators which strictly dominate the never pool estimator, and of this family of dominating estimators the pre-test estimator which uses  $c=1$  has the smallest risk.<sup>5</sup> This generalises the result of Ohtani and Toyoda (1978) and it holds for all  $\theta_1$  and  $\theta_2$ , and for all feasible members of

<sup>5</sup>Though not included in this paper, it is straightforward to analytically prove this result using the approach outlined by Giles (1991a,b, 1992). Details are available from Giles (1990) or the author.



the  $ESD_N$  family.

Further, these same pre-test estimators can also strictly dominate the always pool estimator, given  $\theta_1$  and  $\theta_2$ . The condition under which this result occurs depends on  $v_1$ ,  $v_2$ ,  $\theta_1$ ,  $\theta_2$ ,  $\alpha$ , and on the variance mixing distribution  $f(\tau)$ .

(8) The risks of  $s_N^2$  and  $s_A^2$  have two possible  $\phi$  intersections:<sup>6</sup>

$$\begin{aligned} \phi_i = (v_1/v_2) & \left\{ v_1 \left[ v_1 v_2 E(\tau^4) - v_2 (v_1 + v_2) \left( E(\tau^2) \right)^2 + 2v_2 E(\tau^2)(\theta_1 - \theta_2) + 4\theta_1 \theta_2 \right] \right. \\ & \pm \left[ v_1^2 \left[ v_1 v_2 E(\tau^4) - v_2 (v_1 + v_2) \left( E(\tau^2) \right)^2 + 2v_2 E(\tau^2)(\theta_1 - \theta_2) + 4\theta_1 \theta_2 \right]^2 \right. \\ & + v_2 \left[ (2v_1 + v_2) \left( v_1 (v_1 + 2) E(\tau^4) + 4\theta_1 (v_1 + 2) E(\tau^2) + 4\theta_1^2 - 2v_1 (v_1 + v_2) \right. \right. \\ & \times E(\tau^2) \left( v_1 E(\tau^2) + 2\theta_1 \right) \left. \left. \left[ v_2 (v_2 + 2) E(\tau^4) + 4(v_2 + 2)\theta_2 E(\tau^2) - 4\theta_2^2 \right]^{1/2} \right] \right\} \\ & \wedge \left[ (2v_1 + v_2) \left( v_1 (v_1 + 2) E(\tau^4) + 4\theta_1 (v_1 + 2) E(\tau^2) + 4\theta_1^2 \right) \right. \\ & \left. \left. - 2v_1 (v_1 + v_2) E(\tau^2) \left( v_1 E(\tau^2) + 2\theta_1 \right) \right] \right\} \\ & = \omega \pm \kappa, \end{aligned} \quad (14)$$

$i=1,2$ . Let  $\phi_1 = \omega + \kappa$  and  $\phi_2 = \omega - \kappa$ . Giles (1992) shows that if the disturbances are  $ESD_N$  but the design matrix is correctly specified then there always exists two possible events but only one feasible intersection - neither the always pool estimator nor the never pool estimator can strictly dominate each other. However, when relevant regressors have been excluded there are four possible events: (i)  $0 < \phi_1 < 1$ ,  $\phi_2 > 1$ ; (ii)  $\phi_1 > 1$ ,  $\phi_2 < 0$ ; (iii)  $0 < \phi_1 < 1$ ,  $\phi_2 < 0$ ; (iv)  $\phi_1 > 1$ ,  $\phi_2 > 1$ . If  $v_1$ ,  $v_2$ ,  $\theta_1$ , and  $\theta_2$  are such that cases (ii) or (iv) result then the always pool estimator is strictly dominated by the never pool estimator for all  $\phi \in (0,1]$ . We discuss this result further in the following section.

---

<sup>6</sup>We require  $\left[ (2v_1 + v_2) \left( v_1 (v_1 + 2) E(\tau^4) + 4\theta_1 (v_1 + 2) E(\tau^2) + 4\theta_1^2 \right) - 2v_1 (v_1 + v_2) \times E(\tau^2) \left( v_1 E(\tau^2) + 2\theta_1 \right) \right] \neq 0$ .

#### 4. NUMERICAL EVALUATIONS OF THE RISK FUNCTIONS

To illustrate the risk functions we have numerically evaluated them for the special case of Mt disturbances, which arise when  $\tau$  is an inverted gamma variate. Then  $e \sim \text{Mt}\left(0, (\nu\sigma_2^2/(\nu-2))\Sigma\right)$  and

$$\rho_{\text{Mt}}\left(\sigma_{e_1}^2, s_N^2\right) = 2\phi^2\sigma_2^4\left(\nu^2v_1(v_1+\nu-2)+4\lambda_1\nu(\nu-2)(\nu-4)+2\lambda_1^2(\nu-2)^2(\nu-4)\right)/\left(v_1^2(\nu-2)^2(\nu-4)\right) \quad (15)$$

$$\begin{aligned} \rho_{\text{Mt}}\left(\sigma_{e_1}^2, s_A^2\right) = & \sigma_2^4\left\{\phi^2\left[v_2^2\nu^2(\nu-4)+2v_1\nu^2(v_1+\nu-2)-4\lambda_1\nu(\nu-2)(\nu-4)(v_2-2)+4\lambda_1^2(\nu-2)^2(\nu-4)\right]+2\phi\left[v_2\nu^2\left(2v_1-v_2(\nu-4)\right)+2v_2\nu(\nu-2)(\nu-4)(\lambda_1-\lambda_2)+4\lambda_1\lambda_2(\nu-2)^2(\nu-4)\right]+v_2(v_2+2)\nu^2(\nu-2)+4(v_2+2)\lambda_2\nu(\nu-2)(\nu-4)+4\lambda_2^2(\nu-2)^2(\nu-4)\right\}/\left((v_1+v_2)^2(\nu-2)^2(\nu-4)\right) \end{aligned} \quad (16)$$

$$\begin{aligned} \rho_{\text{Mt}}\left(\sigma_{e_1}^2, s_P^2\right) = & \sigma_2^4\left\{2\phi^2(v_1+v_2)^2\left[v_1\nu^2(v_1+\nu-2)+4\lambda_1\nu(\nu-2)(\nu-4)+4\lambda_1^2(\nu-2)^2(\nu-4)\right]+\phi^2v_2\left[-(\nu-2)(2v_1+v_2)\left(v_1(v_1+2)\nu^2Q_{040}+4(v_1+2)\nu(\nu-4)Q_{061}+4\lambda_1^2(\nu-2)(\nu-4)Q_{082}\right)+2\nu(\nu-4)v_1(v_1+v_2)\right.\right. \\ & \times\left.\left(v_1\nu Q_{021}+2\lambda_1(\nu-2)Q_{042}\right)\right]+2v_1^2\phi\left[v_1v_2\nu^2(\nu-2)Q_{220}-v_2(v_1+v_2)\nu^2\right. \\ & \times\left.\left(\nu-4\right)Q_{201}+2v_1\lambda_2\nu(\nu-2)(\nu-4)Q_{421}-2(v_1+v_2)\lambda_2\nu(\nu-2)(\nu-4)Q_{402}+2v_2\lambda_1\nu(\nu-2)(\nu-4)Q_{241}+4\lambda_1\lambda_2(\nu-2)^2(\nu-4)Q_{442}\right] \\ & \left.+v_1^2(\nu-2)\left[v_2(v_2+2)\nu^2Q_{400}+4(v_2+2)\lambda_2\nu(\nu-4)Q_{601}+4\lambda_2^2(\nu-2)(\nu-4)Q_{802}\right]\right\}/\left(v_1^2(v_1+v_2)^2(\nu-2)^2(\nu-4)\right) \end{aligned} \quad (17)$$

where

$$Q_{ijn} = \sum_{r=0}^{\infty} \sum_{s=0}^{\infty} \frac{(2\lambda_2/\nu)^r (2\lambda_1/\nu)^s \Gamma(\nu/2+r+s+n-2)}{r!s! \left(1+2(\lambda_1+\lambda_2)/\nu\right)^{\nu/2+r+s+n-2} \Gamma(\nu/2+n-2)}$$

$$\times I_w \left( (v_2+i)/2+r; (v_1+j)/2+s \right) \quad i, j, n=0, 1, 2, \dots,$$

$\lambda_i = \theta_i / \sigma_2^2$  ( $i=1, 2$ ) and  $I_w(\cdot, \cdot)$  is Pearson's incomplete beta function with  $w = c\phi v_2 / (v_1 + c\phi v_2)$ .

We have evaluated (15)-(17) for a wide range of the arguments :  $v_1=v_2=20$ ;  $v_1=16$ ,  $v_2=8$ ;  $v_1=25$ ,  $v_2=10$ ;  $k_1=k_2=3, 4, 5$ ;  $\alpha=0.01, 0.05, 0.30, 0.50, 0.75$  and those values of  $\alpha$  associated with a critical value of unity;  $\nu=5, 10, 100, \infty$ ;  $\lambda_1 \in [0, 20]$ ;  $\lambda_2 \in [0, 20]$ ; and  $\phi \in [0.05, 1]$ . The FORTRAN computer programs, executed on a VAX 6230 computer, used various subroutines from Press *et al.* (1986) and Davies' (1980) algorithm. Figures 3 to 8 present some typical results; full details of which are available on request. These figures consider  $\nu=5$  and  $\nu=\infty$  (normal errors) for three degrees of mis-specification: first, when regressors are excluded from the model for sample one but not from that for sample two ( $\lambda_1=3, \lambda_2=0$ ); secondly, when the design matrix for sample one is correctly specified but that for sample two is not ( $\lambda_1=0, \lambda_2=3$ ); and thirdly, when both models are mis-specified to the same degree ( $\lambda_1=\lambda_2=3$ ). As in Figures 1 and 2, we consider risk relative to  $\sigma_2^4$  and parameterise with respect to  $\lambda_1$  and  $\lambda_2$  rather than with respect to  $\theta_1$  and  $\theta_2$  to eliminate the scale parameter  $\sigma_2^2$ . Equivalently, the figures represent the risks of the estimators when  $\sigma_2^2=1$ .

The figures illustrate the features discussed in the previous section. In particular, they highlight that the always pool estimator can be strictly dominated by both the never pool estimator and the pre-test estimator. Then, as the never pool estimator is itself always strictly dominated by at least the pre-test estimator which uses  $c=1$ , it is always preferable to pre-test and to use a critical value of unity.

Giles (1992), in the correctly specified design matrix case, shows that the pre-test estimator which uses  $c=1$  can strictly dominate both of its component estimators for relatively small  $\nu$ . Our

numerical evaluations support her findings but they also suggest that the always pool estimator will be strictly dominated by the pre-test estimator which uses  $c=1$  if the models for either sample are, or for both samples is, sufficiently mis-specified, irrespective of the value of  $\nu$ .

Regarding the possibility of the strict dominance of the always pool estimator by the never pool estimator, which is infeasible in the properly specified model, our numerical evaluations suggest that this result depends not only on the degree of mis-specification but also on the other arguments in the problem. Specifically, it will typically be observed if  $\nu_1 \approx \nu_2$ ,  $\lambda_2 \gg \lambda_1$ , and  $\nu$  is relatively large. Intuitively, the gain in information from the second sample in terms of additional degrees of freedom is outweighed by the loss in the 'quality' of the information due to the relatively higher specification error in the second sample. For small values of  $\nu$  we find there is still a small  $\phi$ -range, in the neighbourhood of  $H_0$ , over which it is better to always pool the samples than to never pool them.

If, however, the mis-specification in the first sample is significantly higher than that in the second sample then there is a  $\phi$ -range over which the always pool estimator has smaller risk than the never pool estimator, irrespective of the value of  $\nu$ . *Ceteris paribus*, the width of this range increases with  $\nu$  and with  $\lambda_1$ .

## 5. CONCLUDING REMARKS

In this paper we have considered the risk properties of estimators of the error variance after a pre-test for homogeneity, when the joint distribution of the unobservable errors in each sample is  $SSD_N$  but it is assumed to be normal, and there is simultaneously a possible mis-specification of the design matrix. We showed first that the classical test for homoscedasticity in our model is no longer valid when regressors have been omitted. Then, both the null and non-null distributions of  $J$  depend on  $\nu_1$ ,  $\nu_2$ , the degree of mis-specification of the design matrix, and the variance mixing distribution,  $f(\tau)$ .




Figure 3. Relative risk functions for  $s_N^2$ ,  $s_A^2$ , and  $s_P^2$  when  $e \sim Mt\left(0, (\nu\sigma_2^2/(\nu-2))\Sigma\right)$ ,  $\nu_1=16$ ,  $\nu_2=8$ ,  $k_1=k_2=3$ ,  $\nu=5$ ,  $\lambda_1=3$ ,  $\lambda_2=0$ .




Figure 4. Relative risk functions for  $s_N^2$ ,  $s_A^2$ , and  $s_P^2$  when  $e \sim N(0, \sigma_2^2 \Sigma)$ ,  $\nu_1=16$ ,  $\nu_2=8$ ,  $k_1=k_2=3$ ,  $\lambda_1=3$ ,  $\lambda_2=0$ .




Figure 5. Relative risk functions for  $s_N^2$ ,  $s_A^2$ , and  $s_P^2$  when  $e \sim \text{Mt}\left(0, (\nu\sigma_2^2/(\nu-2))\Sigma\right)$ ,  $v_1=16$ ,  $v_2=8$ ,  $k_1=k_2=3$ ,  $\nu=5$ ,  $\lambda_1=0$ ,  $\lambda_2=3$ .




Figure 6. Relative risk functions for  $s_N^2$ ,  $s_A^2$ , and  $s_P^2$  when  $e \sim N(0, \sigma_2^2\Sigma)$ ,  $v_1=16$ ,  $v_2=8$ ,  $k_1=k_2=3$ ,  $\lambda_1=0$ ,  $\lambda_2=3$ .






Figure 7. Relative risk functions for  $s_N^2$ ,  $s_A^2$ , and  $s_P^2$  when  $e \sim \text{Mt}\left(0, (\nu\sigma_2^2/(\nu-2))\Sigma\right)$ ,  $\nu_1=16$ ,  $\nu_2=8$ ,  $k_1=k_2=3$ ,  $\nu=5$ ,  $\lambda_1=\lambda_2=3$ .




Figure 8. Relative risk functions for  $s_N^2$ ,  $s_A^2$ , and  $s_P^2$  when  $e \sim N(0, \sigma_2^2 \Sigma)$ ,  $\nu_1=16$ ,  $\nu_2=8$ ,  $k_1=k_2=3$ ,  $\lambda_1=\lambda_2=3$ .

Our analysis of the risk functions of the never pool, always pool, and pre-test estimators of  $\sigma_{e_1}^2$  showed first that the dominance of the never pool estimator by a family of pre-test estimators is robust to the mis-specification of the design matrix and of the error distribution. It is straightforward to show that the pre-test estimator which uses  $c=1$  has the smallest risk of this family of dominating pre-test estimators irrespective of the degree of mis-specification of the design matrix or of the form of the variance mixing distribution.

We showed secondly that if we have excluded variables then the never pool estimator can also strictly dominate the always pool estimator. This result is impossible when we have properly specified the model, and it suggests that the risk of the always pool estimator is (qualitatively) less robust to the mis-specification than is the never pool estimator.

Practically, our analysis suggests, given that the degrees of model mis-specification and hypothesis error are unknown, that it is generally preferable to pre-test rather than to impose or ignore the prior information without testing. Then the optimal critical value is unity irrespective of the degrees of freedom of the model. Typically this critical value results in a (nominal) size that is far greater than the usual test sizes of 1% or 5%.

It remains for future research to consider the extension of this analysis to the two-sided alternative hypothesis case. The sensitivity of the results to the particular form of non-normality also requires attention. In particular, it is unclear whether they will extend to the case of iid non-normal disturbances.

#### APPENDIX

Proof of Theorem 1.

$$f(J) = \int_0^\infty f_N(J)f(\tau)d\tau, \quad (A.1)$$

where  $f_N(J)$  is the density function of  $J$  when  $e \sim N(0, \tau^2 \Sigma)$ . Under this normality assumption

$$e^* \equiv \begin{bmatrix} (Z_1 \gamma_1 + e_1) / \sqrt{\phi} \\ (Z_2 \gamma_2 + e_2) \end{bmatrix} \sim N \left( \begin{bmatrix} Z_1 \gamma_1 / \sqrt{\phi} \\ Z_2 \gamma_2 \end{bmatrix}, \tau^2 I_T \right)$$

and so

$$J = \frac{s_2^2}{s_1^2} = \frac{v_1 (Z_2 \gamma_2 + e_2)' M_2 (Z_2 \gamma_2 + e_2)}{v_2 (Z_1 \gamma_1 + e_1)' M_1 (Z_1 \gamma_1 + e_1)} = \frac{v_1 e^{**} M_2^* e^*}{v_2 e^{**} M_1^* e^*}$$

where  $M_1^*$  and  $M_2^*$  are  $(T \times T)$  idempotent matrices partitioned as  $M_i^* =$

$$\begin{bmatrix} M_i & 0 \\ 0 & 0 \end{bmatrix} \text{ and } M_i^* = \begin{bmatrix} 0 & 0 \\ 0 & M_i \end{bmatrix} \text{ with } r(M_i^*) = r(M_i) = v_i, \quad i=1,2. \text{ Under the}$$

normality assumption, it is straightforward to show that the quadratic forms  $(e^{**} M_2^* e^* / \tau^2)$  and  $(e^{**} M_1^* e^* / \tau^2)$  are independent with  $(e^{**} M_i^* e^* / \tau^2) \sim \chi_{v_i; \lambda_i \tau}^2$  and  $\lambda_i \tau = \theta_i / \tau^2$ ,  $i=1,2$ . So,  $f_N(J) = \phi^{-1} f(F' (v_2, v_1; \theta_2 / \tau^2, \theta_1 / \tau^2))$ . Given the density function of a doubly non-central  $F$  variate and using (A.1) equation (8) follows directly. ■

**Proof of Theorem 2.**

$$\rho(\sigma_{e_1}^2, s_N^2) = E(s_N^4) - 2\phi E(\tau^2) E(s_N^2) + \phi^2 (E(\tau^2))^2 \quad (\text{A.2})$$

and  $E(s_N^2) = \int_0^\infty E_N(s_N^2) f(\tau) d\tau$ , where  $E_N(s_N^2) = E(s_N^2)$  when  $e \sim N(0, \tau^2 \Sigma)$ . Then,  $e^* \sim N(0, \tau^2 I_T)$ ,  $e^{**} M_1^* e^* / \tau^2 \sim \chi_{v_1; \lambda_1 \tau}^2$ ,  $E_N(e^{**} M_1^* e^* / \tau^2) = v_1 + 2\lambda_1 \tau$ , and so  $E(s_N^2) = \phi(v_1 E(\tau^2) + 2\theta_1) / v_1$ . Using the same approach,  $E(s_N^4) = \phi^2(v_1(v_1+2)E(\tau^4) + 4(v_1+2)\theta_1 E(\tau^2) + 4\theta_1^2) / v_1^2$ . Substituting these expressions into (A.2) yields  $\rho(\sigma_{e_1}^2, s_N^2)$ .

$$\rho(\sigma_{e_1}^2, s_A^2) = E(s_A^4) - 2\phi E(\tau^2) E(s_A^2) + \phi^2 (E(\tau^2))^2 \quad (\text{A.3})$$

and  $E(s_A^2) = \int_0^\infty E_N(s_A^2) f(\tau) d\tau$ , where  $E_N(s_A^2) = E(s_A^2)$  when  $e \sim N(0, \tau^2 \Sigma)$ . Then,  $e^* \sim N(0, \tau^2 I_T)$ ,  $e^{**} M_i^* e^* / \tau^2 \sim \chi_{v_i; \lambda_i \tau}^2$ ,  $E_N(e^{**} M_i^* e^* / \tau^2) = v_i + 2\lambda_i \tau$  ( $i=1,2$ ), and

so as  $s_A^2 = (\phi e^{**} M_1^* e^{**} + e^{**} M_2^* e^{**}) / (v_1 + v_2)$  we have  $E_N(s_A^2) = \tau^2 \left( \phi(v_1 + 2\lambda_{1\tau}) + v_2 + 2\lambda_{2\tau} \right) / (v_1 + v_2)$ . Integrating this expression with respect to  $\tau$  gives  $E(s_A^2) = \left[ \phi \left( v_1 E(\tau^2) + 2\theta_1 \right) + v_2 E(\tau^2) + 2\theta_2 \right] / (v_1 + v_2)$ . Similarly,  $E(s_A^4) = \left( \phi^2 \left[ v_1(v_1 + 2) E(\tau^4) + 4(v_1 + 2)\theta_1 E(\tau^2) + 4\theta_1^2 \right] + 2\phi \left[ v_1 v_2 E(\tau^4) + 2v_1 \theta_2 E(\tau^2) + 2v_2 \theta_1 E(\tau^2) + 4\theta_1 \theta_2 \right] + v_2(v_2 + 2) E(\tau^4) + 4(v_2 + 2)\theta_2 E(\tau^2) + 4\theta_2^2 \right) / (v_1 + v_2)^2$ . Substituting these expressions into (A.3) completes the derivation of  $\rho \left( \sigma_e^2, s_A^2 \right)$ .

Finally,

$$\rho \left( \sigma_e^2, s_P^2 \right) = E \left( s_P^4 \right) - 2\phi E(\tau^2) E \left( s_P^2 \right) + \phi^2 \left( E(\tau^2) \right)^2. \quad (A.4)$$

Using the aforementioned notation we write  $s_P^2 = \left( \phi(v_1 + v_2)(e^{**} M_1^* e^{**}) + \left[ v_1 e^{**} M_2^* e^{**} - \phi v_2 e^{**} M_1^* e^{**} \right] I_{[0, \phi]} \left( (v_1 e^{**} M_2^* e^{**}) / (v_2 e^{**} M_1^* e^{**}) \right) \right) / \left( v_1(v_1 + v_2) \right)$ . Using Lemma 1 of Clarke et al. (1987)  $E_N \left[ (e^{**} M_2^* e^{**} / \tau^2) \times I_{[0, \phi]} \left( (v_1 e^{**} M_2^* e^{**}) / (v_2 e^{**} M_1^* e^{**}) \right) \right] = v_2 Q_{20}^\tau + 2\lambda_{2\tau} Q_{40}^\tau$  and  $E_N \left[ (e^{**} M_1^* e^{**} / \tau^2) \times I_{[0, \phi]} \left( (v_1 e^{**} M_2^* e^{**}) / (v_2 e^{**} M_1^* e^{**}) \right) \right] = v_1 Q_{02}^\tau + 2\lambda_{1\tau} Q_{04}^\tau$ . So,  $E_N(s_P^2) = \left( \phi(v_1 + v_2)(v_1 \tau^2 + 2\theta_1) + v_1 v_2 \tau^2 (Q_{20}^\tau - \phi Q_{02}^\tau) + 2v_1 \theta_2 Q_{40}^\tau - 2v_2 \theta_1 \phi Q_{04}^\tau \right) / \left( v_1(v_1 + v_2) \right)$  from which we obtain  $E(s_P^2)$  by integrating with respect to  $\tau$ . Likewise,  $E_N(s_P^4) = \left( \phi^2(v_1 + v_2)^2 \left[ v_1(v_1 + 2)\tau^4 + 4(v_1 + 2)\theta_1 \tau^2 + 4\theta_1^2 \right] - \phi^2 v_2(2v_1 + v_2) \left[ v_1(v_1 + 2)\tau^4 Q_{04}^\tau + 4(v_1 + 2)\theta_1 \tau^2 Q_{06}^\tau + 4\theta_1^2 Q_{08}^\tau \right] + v_1^2 \left[ v_2(v_2 + 2)\tau^4 Q_{40}^\tau + 4(v_2 + 2)\theta_2 \tau^2 Q_{60}^\tau + 4\theta_2^2 Q_{80}^\tau \right] + 2\phi v_1^2 \left[ v_1 v_2 \tau^4 Q_{22}^\tau + 2v_1 \theta_2 \tau^2 Q_{42}^\tau + 2v_2 \theta_1 Q_{24}^\tau + 4\theta_1 \theta_2 Q_{44}^\tau \right] \right) / \left( v_1(v_1 + v_2) \right)^2$  which we integrate with respect to  $\tau$  to obtain  $E(s_P^4)$ . Substituting these expressions into (A.4) completes the proof. ■

#### ACKNOWLEDGEMENT

The author thanks David Giles and Mike Veall for helpful comments on the material pertaining to this paper.

## REFERENCES

- Bancroft, T.A., (1944). On biases in estimation due to the use of preliminary tests of significance. *Annals of Mathematical Statistics*, 15, 190-204.
- Bancroft, T.A. and Han, C-P., (1983). A note on pooling variances. *Journal of the American Statistical Association*, 78, 981-983.
- Chmielewski, M.A., (1981). Invariant tests for the equality of K scale parameters under spherical symmetry. *Journal of Statistical Planning and Inference*, 5, 341-346.
- Clarke, J.A., Giles, D.E.A. and Wallace, T.D., (1987). Estimating the error variance in regression after a preliminary test of restrictions on the coefficients. *Journal of Econometrics*, 34, 293-304.
- Davies, R.B., (1980). The distribution of a linear combination of  $\chi^2$  random variables (Algorithm AS 155). *Applied Statistics*, 29, 323-333.
- Dickey, J.M. and Chen, C-H., (1985). Direct subjective-probability modelling using ellipsoidal distributions: in Bernardo, J.M., DeGroot, M.H., Lindley, D.V. and Smith, A.F.M. (eds.), *Bayesian statistics 2*. Amsterdam: North-Holland.
- Giles, D.E.A., (1986). Preliminary-test estimation in mis-specified regressions. *Economics Letters*, 21, 325-328.
- Giles, D.E.A. and Clarke, J.A., (1989). Preliminary-test estimation of the scale parameter in a mis-specified regression model. *Economics Letters*, 30, 201-205.
- Giles, J.A., (1990). Preliminary-test estimation of a mis-specified linear model with spherically symmetric disturbances. Ph.D. thesis, University of Canterbury.
- Giles, J.A., (1991a). Pre-testing for linear restrictions in a regression model with spherically symmetric disturbances. *Journal of Econometrics* 50, 377-398.
- Giles, J.A., (1991b). Pre-testing in a mis-specified regression model. *Communications in Statistics: Theory and Methods* 20, 3221-3238.
- Giles, J.A., (1992). Estimation of the error variance after a preliminary-test of homogeneity in a regression model with spherically symmetric disturbances. *Journal of Econometrics*, forthcoming.

- Kelker, D., (1970). Distribution theory of spherical distributions and a location-scale parameter generalization. *Sankhya A*, 32, 419-430.
- King, M.L., (1979). Some aspects of statistical inference in the linear regression model. Ph.D. thesis, University of Canterbury.
- Mittelhammer, R.C., (1984). Restricted least squares, pre-test, OLS and Stein rule estimators: Risk comparisons under model misspecification. *Journal of Econometrics*, 25, 151-164.
- Muirhead, R.J., (1982). *Aspects of multivariate statistical theory*. New York: John Wiley & Sons.
- Ohtani, K., (1983). Preliminary test predictor in the linear regression model including a proxy variable. *Journal of the Japan Statistical Society*, 13, 11-19.
- Ohtani, K., (1987). Some small sample properties of a pre-test estimator of the disturbance variance in a misspecified linear regression. *Journal of the Japan Statistical Society*, 17, 81-89.
- Ohtani, K. and Toyoda, T., (1978). Minimax regret critical values for a preliminary test in pooling variance. *Journal of the Japan Statistical Society*, 8, 15-20.
- Press, W.H., Flannery, B.P., Teukolsky, S.A., and Vetterling, W.T., (1986). *Numerical recipes: The art of scientific computing*. New York: Cambridge University Press.
- Toyoda, T. and Wallace, T.D., (1975). Estimation of variance after a preliminary test of homogeneity and optimal levels of significance for the pre-test. *Journal of Econometrics*, 3, 395-404.
- Yancey, T.A., Judge, G.G. and Mandy, D.M., (1983). The sampling performance of pre-test estimators of the scale parameter under squared error loss. *Economics Letters*, 12, 181-186.



## LIST OF DISCUSSION PAPERS\*

- No. 8801 Workers' Compensation Rates and the Demand for Apprentices and Non-Apprentices in Victoria, by Pasquale M. Sgro and David E. A. Giles.
- No. 8802 The Adventures of Sherlock Holmes, the 48% Solution, by Michael Carter.
- No. 8803 The Exact Distribution of a Simple Pre-Test Estimator, by David E. A. Giles.
- No. 8804 Pre-testing for Linear Restrictions in a Regression Model With Student-t Errors, by Judith A. Clarke.
- No. 8805 Divisia Monetary Aggregates and the Real User Cost of Money, by Ewen McCann and David Giles.
- No. 8806 The Management of New Zealand's Lobster Fishery, by Alan Woodfield and Pim Borren.
- No. 8807 Poverty Measurement: A Generalization of Sen's Result, by Prasanta K. Pattanaik and Manimay Sen.
- No. 8808 A Note on Sen's Normalization Axiom for a Poverty Measure, by Prasanta K. Pattanaik and Manimay Sen.
- No. 8809 Budget Deficits and Asset Sales, by Ewen McCann.
- No. 8810 Unorganized Money Markets and 'Unproductive' Assets in the New Structuralist Critique of Financial Liberalization, by P. Dorian Owen and Otton Solis-Fallas.
- No. 8901 Testing for Financial Buffer Stocks in Sectoral Portfolio Models, by P. Dorian Owen.
- No. 8902 Provisional Data and Unbiased Prediction of Economic Time Series by Karen Browning and David Giles.
- No. 8903 Coefficient Sign Changes When Restricting Regression Models Under Instrumental Variables Estimation, by David E. A. Giles.
- No. 8904 Economies of Scale in the New Zealand Electricity Distribution Industry, by David E. A. Giles and Nicolas S. Wyatt.
- No. 8905 Some Recent Developments in Econometrics: Lessons for Applied Economists, by David E. A. Giles.
- No. 8906 Asymptotic Properties of the Ordinary Least Squares Estimator in Simultaneous Equations Models, by V. K. Srivastava and D. E. A. Giles.
- No. 8907 Unbiased Estimation of the Mean Squared Error of the Feasible Generalised Ridge Regression Estimator, by V. K. Srivastava and D. E. A. Giles.
- No. 8908 An Unbiased Estimator of the Covariance Matrix of the Mixed Regression Estimator, by D. E. A. Giles and V. K. Srivastava.
- No. 8909 Pre-testing for Linear Restrictions in a Regression Model with Spherically Symmetric Disturbances, by Judith A. Giles.
- No. 9001 The Durbin-Watson Test for Autocorrelation in Nonlinear Models, by Kenneth J. White.
- No. 9002 Determinants of Aggregate Demand for Cigarettes in New Zealand, by Robin Harrison and Jane Chetwyd.
- No. 9003 Unemployment Duration and the Measurement of Unemployment, by Manimay Sengupta.
- No. 9004 Estimation of the Error Variance After a Preliminary-Test of Homogeneity in a Regression Model with Spherically Symmetric Disturbances, by Judith A. Giles.
- No. 9005 An Expository Note on the Composite Commodity Theorem, by Michael Carter.
- No. 9006 The Optimal Size of a Preliminary Test of Linear Restrictions in a Mis-specified Regression Model, by David E. A. Giles, Offer Lieberman, and Judith A. Giles.
- No. 9007 Inflation, Unemployment and Macroeconomic Policy in New Zealand: A Public Choice Analysis, by David J. Smyth and Alan E. Woodfield.
- No. 9008 Inflation — Unemployment Choices in New Zealand and the Median Voter Theorem, by David J. Smyth and Alan E. Woodfield.
- No. 9009 The Power of the Durbin-Watson Test when the Errors are Heteroscedastic, by David E. A. Giles and John P. Small.
- No. 9010 The Exact Distribution of a Least Squares Regression Coefficient Estimator After a Preliminary t-Test, by David E. A. Giles and Virendra K. Srivastava.
- No. 9011 Testing Linear Restrictions on Coefficients in a Linear Regression Model with Proxy variables and Spherically Symmetric Disturbances, by Kazuhiro Ohtani and Judith A. Giles.

*(Continued on next page)*



- No. 9012      Some Consequences of Applying the Goldfeld-Quandt Test to Mis-Specified Regression Models, by David E. A. Giles and Guy N. Saxton.
- No. 9013      Pre-testing in a Mis-specified Regression Model, by Judith A. Giles.
- No. 9014      Two Results in Balanced-Growth Educational Policy, by Alan E. Woodfield.
- No. 9101      Bounds on the Effect of Heteroscedasticity on the Chow Test for Structural Change, by David Giles and Offer Lieberman.
- No. 9102      The Optimal Size of a Preliminary Test for Linear Restrictions when Estimating the Regression Scale Parameter, by Judith A. Giles and Offer Lieberman.
- No. 9103      Some Properties of the Durbin-Watson Test After a Preliminary t-Test, by David Giles and Offer Lieberman.
- No. 9104      Preliminary-Test Estimation of the Regression Scale Parameter when the Loss Function is Asymmetric, by Judith A. Giles and David E. A. Giles.
- No. 9105      On an Index of Poverty, by Manimay Sengupta and Prasanta K. Pattanaik.
- No. 9106      Cartels May Be Good For You, by Michael Carter and Julian Wright.
- No. 9107      Lp-Norm Consistencies of Nonparametric Estimates of Regression, Heteroskedasticity and Variance of Regression Estimate when Distribution of Regression is Known, by Radhey S. Singh.
- No. 9108      Optimal Telecommunications Tariffs and the CCITT, by Michael Carter and Julian Wright.
- No. 9109      Price Indices : Systems Estimation and Tests, by David Giles and Ewen McCann.
- No. 9110      The Limiting Power of Point Optimal Autocorrelation Tests, by John P. Small.
- No. 9111      The Exact Power of Some Autocorrelation Tests When the Disturbances are Heteroscedastic, by John P. Small.
- No. 9112      Some Consequences of Using the Chow Test in the Context of Autocorrelated Disturbances, by David Giles and Murray Scott.
- No. 9113      The Exact Distribution of  $R^2$  when the Disturbances are Autocorrelated, by Mark L. Carrodus and David E. A. Giles.
- No. 9114      Optimal Critical Values of a Preliminary Test for Linear Restrictions in a Regression Model with Multivariate Student-t Disturbances, by Jason K. Wong and Judith A. Giles.
- No. 9115      Pre-Test Estimation in a Regression Model with a Misspecified Error Covariance Matrix, by K. V. Albertson.
- No. 9116      Estimation of the Scale Parameter After a Pre-test for Homogeneity in a Mis-specified Regression Model, by Judith A. Giles.

\* Copies of these Discussion Papers may be obtained for \$4 (including postage, price changes occasionally) each by writing to the Secretary, Department of Economics, University of Canterbury, Christchurch, New Zealand. A list of the Discussion Papers prior to 1988 is available on request.