



**AgEcon** SEARCH  
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search  
<http://ageconsearch.umn.edu>  
[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

## THE MACRO-MINERAL PROFILE OF FOUR TROPICAL GRASSES AT DIFFERENT STAGES OF REGROWTH

F.G. Youssef, R.H. Rastogi, N.K. Persad\* and H. Ramlal\*  
(Department of Livestock Science, The University of the West Indies,  
St. Augustine, Trinidad, W.I.)

### ABSTRACT

Pangola grass (*Digitaria decumbens*), Lucuntu grass (*Ischaemum timorense*), Star grass (*Cynodon niemfuensis*) and Tanner grass (*Brachiaria radicans*) were planted on Piarco fine sand soil at Central Experiment Station, Centeno, Trinidad, using a split plot design with grasses as main plots, and regrowth cutting intervals: 4, 5, 6, 7 and 8 weeks as sub-plots, in order to study their effects on the macro-mineral profile of the grasses. There were four replicates per treatment.

When grasses were compared, it was found that there were significant ( $P < .001$ ) differences in overall means of nitrogen (N), calcium (Ca), phosphorus (P), sodium (Na) and magnesium (Mg) content, but not of potassium (K) content. Star grass had the highest levels (g/kg dry

matter) of N (12.8), P (3.4) and Mg (2.4); while Ca (7.3) and K (10.4) content were highest in Lucuntu grass. The lowest content of N (8.1) and Ca (2.8) were found in Tanner grass, of K (7.3) and Mg (1.6) in Pangola grass, and of P (1.8) and Na (0.4) in Lucuntu grass.

For all minerals studied except calcium, there were significant ( $P < .001$ ) differences at different stages of regrowth, with the highest levels at 4 weeks of age decreasing gradually to reach the lowest levels at 8 weeks.

The results are discussed in consideration of meeting the macro-mineral requirements of ruminants for optimum health and production.

---

\*Central Experiment Station, Centeno, Trinidad, W.I.