



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

Efficacy of Potassium Salts of Fatty Acids in the Management of Thrips and Whitefly on Snap Beans

Geoffrey Ongoya Wafula¹, James W. Muthomi¹, John H. Nderitu¹ & George N. Chemining'wa¹

¹Department of Plant Science and Crop Protection, University of Nairobi. P. O Box 29053-00625 Kangemi, Nairobi, Kenya

Correspondence: James W. Muthomi, Department of Plant Science and Crop Protection, University of Nairobi. P. O. Box 29053-00625 Kangemi, Nairobi, Kenya. Tel: 254-722-984-179. E-mail: james_wanjohi@yahoo.com

Received: June 18, 2017 Accepted: July 31, 2017 Online Published: August 5, 2017

doi:10.5539/sar.v6n4p45

URL: <https://doi.org/10.5539/sar.v6n4p45>

Abstract

Snap bean farmers mainly rely on chemical pesticides to manage insect pests but the introduction of strict maximum residue levels (MRLs) by European markets poses a challenge to the use of pesticides. This has necessitated the search for alternative pest management options that do not result in harmful residues on produce. Therefore, this study was carried out to determine the efficacy of potassium salts of fatty acids as alternative to synthetic chemicals in management of snap bean pests. Field experiments were carried out in farmers' fields in Mwea and Embu for two cropping cycles between July 2013 and January 2014. Potassium salts of fatty acids at 0.5%, 1% and 1.5% spray solution was compared with chemical pesticide confidor® (Imidacloprid) and farmers practice [Thunder® (Imidacloprid + Betacyfluthrin) and Karate® (Lambda Cyhalothrin). Population of thrips, whitefly, and yield of pest damaged pods, in addition to marketable pod yield was determined. Application of potassium salts of fatty acids at 1% and 1.5% significantly ($P<0.05$) reduced white fly and thrips populations by up to 54%. Weight of pest damaged pods was also significantly ($P<0.05$) reduced by up to 76% while the weight of marketable pod yield was significantly ($P<0.05$) increased by up to 112%. The results demonstrated that potassium salts of fatty acids are a viable alternative to synthetic chemical pesticides thereby enabling farmers to produce with acceptable residue levels that meet market requirements.

Keywords: market access, pesticide residues, *Phaseolus vulgaris* L., potassium salts of fatty acids

1. Introduction

Snap bean (*Phaseolus vulgaris* L.) is grown specifically for the immature green pods primarily for export market to European Union and elite local urban markets (Infonet-Biovision, 2014). The production of snap beans, one of Kenya's most important export vegetable crops, is steadily rising (HCDA, 2014). Snap beans from Kenya are exported to United Kingdom, France, Holland, Germany, United Arab Emirates and South Africa (HCDA, 2013). Local consumption of Snap beans has also increased over the last few years, providing a domestic market (HCDA, 2013). In the year 2013, Kenya exported over 31,973 tons of snap beans valued at over US\$ 105 million (HCDA, 2014).

Snap bean production is mainly by small scale farmers and it is estimated that over 50,000 smallholder families are involved in snap bean production in Kenya contributing to the larger agricultural sector (Infonet-Biovision, 2014). The agricultural sector plays an important role in Kenya's economy contributing directly and indirectly to the countries GDP by upto 24% and 27% in the year 2011 (MOA, 2012). Production of snap beans in Kenya is constrained by pests which include thrips (*Frankliniella* spp. and *Megalurotrips sjostedti*), whiteflies (*Bemisia tabaci* and *Trialeurodes vaporariorum*), bean flies (*Ophiomyia* spp) and aphids (*Aphis fabae*) (Monda et al., 2003; Nderitu et al., 2007).

In Kenya, farmers rely on insecticides to manage pests in snap beans but the introduction of maximum residue levels (MRLs) for export vegetables by European retailers poses a challenge to the use of pesticides (EU, 2013; Nderitu et al., 2009). Over-reliance on sythethic chemicals pesticides has often led to non compliance by Kenyan exporters leading losses that are passed over to farmers (Mwangi, 2013; Ndung'u, 2013). Non-compliance to the market standards has also led to enhanced scrutiny of Kenyan produce with 10% of all Kenyan bean produce being sampled (Mwangi, 2013). Farmers and exporters of fresh vegetables are as a result, incurring losses through product rejects and financial bills passed to them after the MRL tests whose average cost per sample is

approximately US\$ 247 (Mwangi, 2013; Ndung'u, 2013). The use of pest management options that are environmentally friendly and reduce reliance on synthetic chemical pesticides would be a major step towards sustainable pest management (Nderitu et al., 2009). Alternative pest management options such as use of potassium salts of fatty acids (insecticidal soaps) can be an option to overcome this challenge (Ciancio et al., 2010). Potassium salts of fatty acid work only on direct contact with the pest by washing away the protective coat on the insect surface and penetrate the cell membrane causing disruption of its permeability causing desiccation of the insect (Dheeraj et al., 2013; Mohamad et al., 2013). Potassium salts of fatty acids work on most soft-bodied insect pests (Koppert, 2013) and have been successfully used in the management of aphids, whiteflies, scales and mealy bugs with very good efficacy (Dheeraj et al., 2013; Hollingsworth, 2005; Mohamad et al., 2013). Unlike synthetic chemical pesticides, potassium salts of fatty acids are fast acting with a quick knock down effect on the pests and break down quickly after application leaving no residues (Dheeraj et al., 2013). They are also easy to handle because of their low toxicity to humans and are, therefore, are user friendly (Dheeraj et al., 2013). As a result, potassium salts of fatty acids have no MRL or PHI requirements (Koppert, 2013). This study was undertaken to determine the efficacy of potassium salts of fatty acids in the management of snap bean pests.

2. Materials and Methods

2.1 Description of Experimental Sites

The experiments were carried out in farmers' fields in the main snap bean-producing regions of Kenya in Mwea, Kirinyaga County and Embu County. Mwea has well established irrigation infrastructure that allows year round horticultural production and falls in lower midland zone 4 (LM4). The area is semiarid with Nitosols soils and average rainfall of about 850 mm, range of 500 - 1250 mm. The temperature ranges from 15.6 °C to 28.6 °C with a mean of about 22 °C (Jaetzold et al., 2009). Snap bean production has recently been introduced in Embu East district of Embu County and its production is spreading fast, therefore likely to have a high uptake of new technologies. Embu East district is located in Upper Midland 2 (UM2) agro-ecological zone with average annual precipitation of 1206 mm and average maximum temperature of 28.8 °C and average minimum temperature 9.6 °C. Snap bean production at both sites is mainly for export and is carried out by small scale farmers organized into self-help groups within the irrigation scheme.

2.2 Experimental Treatments, Design and Layout

The experimental treatments consisted of three concentration of potassium salts of fatty acids (Table 1) compared with chemical pesticide confidor® (Imidacloprid) and farmers practice [Thunder® (Imidacloprid 100 g/L + Betacyfluthrin 45g/L) and Karate® (Lambda Cyhalothrin 25 g/kg)].

Table 1. Description of experimental materials

Product	Active ingredient (AI)	Application rate
Potassium salts	Potassium salts of fatty acids	5ml per litre of water (0.5%), 10 ml per litre (1%) and 15ml per litre of water (1.5%)
Confidor WG	Imidacloprid 700g/Kg	5g per 15 litres of water
Thunder®	Imidacloprid 100 g/L + Betacyfluthrin 45g/L	10 ml in 20 litres of water
Karate®	Lambda Cyhalothrin 25 g/kg	6.5 ml in 20 litres of water

Snap beans variety Serengeti were grown under irrigation over two crop cycles in July to October 2013 and October 2013 to January 2014. The experimental plots were 3 m by 2 m with 1 m alleys between the plots and 1.5 m alleys between the blocks and each treatment was replicated four times. The crop was planted in single rows with spacing of 10 cm x 30 cm. The experiment was laid out in a randomized complete block design. Fertilizer application was done once at planting using di-ammonium phosphate (18%N and 46% P₂O₅) at the rate of 490 kg per ha. Top dressing was done at 21 days after emergence with calcium ammonium nitrate at the rate of 490 kg per ha. The first weeding was done two weeks after emergence (WAE) followed by a second weeding two weeks later. Diseases were controlled using Kocide® (Copper Hydroxide 61.4%) for rust and rots, Oshothane® (Mancozeb) against leaf spots and Ortiva® (Azoxystrobin 250g/L) against mildews. The experimental treatments were applied as follows:

- 0.5 % potassium salts applied weekly from 3 weeks after emergence until early pod formation
- 1 % potassium salts applied weekly from 3 weeks after emergence until early pod formation
- 1.5 % potassium salts applied weekly from 3 weeks after emergence until early pod formation
- Chemical pesticide confidor® (Imidacloprid) applied every two weeks

- v. Farmer practice –application of Thunder[®] (Imidacloprid 100 g/L + Betacyfluthrin 45g/L) and Karate[®] (Lambda Cyhalothrin 25 g/kg) on pest detection
- vi. Control - spray with water only

2.3 Assessment of Whitefly and Thrips Population

Population of whiteflies was assessed using yellow sticky trap counts for adults and leaf counts for nymphs. A yellow sticky trap was placed at the centre of each plot and the number of adult whitefly was counted bi-weekly (Hirano et al., 1995; Hoelmer et al., 1998) at two, four, six and eight weeks after emergence (WAE). The number of whitefly nymphs was counted bi-weekly at two, four, six and eight WAE by sampling ten lower leaves from ten plants in a zig-zag manner from inner rows of each plot (Soto et al., 2002).

Population of thrips was assessed weekly from the start of flowering on ten flowers sampled from ten plants from the inner rows in each plot. Sampling was done three times at six, seven and eight weeks after emergence. The sampled flowers were immediately put in 70% ethanol until the insects were counted in the laboratory. Each flower was placed in a petri dish, dissected and the thrips washed into the ethanol. The thrips were observed under a dissecting microscope and counted using a tally counter. Both adults and larvae thrips stages were counted and identified based on morphological characteristics (Nderitu, et al., 2009).

2.4 Assessment of Pod Yield and Quality

Harvesting and grading of pods was done twice every week for two weeks. The pods were harvested from three inner rows in each plot and graded into marketable and non-marketable yields. The marketable pods were further graded into extra-fine (6 to 7.5 mm in diameter and 8 to 12 cm long) and fine (6.5 to 9 mm in diameter and 10 to 13 cm long) according to USAID-KHCP (2011). The non-marketable pods were further graded into pest damaged pods and other rejects based on pest damage symptoms such as feeding marks, scarring and malformation (Infonet-Biovision, 2013). The weight of the each grade of the harvested pods for each plot was recorded.

2.5 Cost Benefit Analysis of the Pest Management Options

Net return was calculated as the total of the entire marketable yield for each treatment multiply by the average price per kilogram minus the total costs while the cost benefit ratio was calculated by dividing the total cost by the net return as follows:

Total marketable pod yield = Total extra-fine + Total fine

$$\text{Average Price} = \frac{\text{Price for extra-fine} + \text{Price for fine}}{2}$$

Total cost = Land preparation cost + Labour + Cost of inputs

Gross returns = Total marketable x Average price

Net returns = Gross returns – Total cost

$$\text{Cost benefit ratio} = \frac{\text{Total Cost}}{\text{Net Returns}}$$

2.6 Statistical Data Analysis

Analysis of variance (ANOVA) was carried out on the data from the two seasons using GenStat Edition 13 software and tested for significance by F-test at 95% level. The treatment means were then compared using the least significant difference (LSD) test at P=0.05 where the F-test was significant (Mead et al., 2003).

3. Results

Potassium salts of fatty acids at rate of 1% and 1.5% significantly ($P < 0.05$) reduced both adult and nymph whitefly population compared to untreated plots. Similar results were observed for the number of thrips per flower. Application of potassium salts of fatty acids at 1.5% compared well with confidor[®] (Imidacloprid) in reducing white fly and thrips population but the application of application of Thunder[®] (Imidacloprid + Betacyfluthrin) followed by Karate[®] (Lambda Cyhalothrin) was less effective (Figure 1). The application of potassium salts of fatty acids at 1.5% and confidor[®] (Imidacloprid) resulted in the least amount of pest damaged pods.




Figure 1. Population of whitefly and thrips and pod weight of pest damaged snap bean pods after application of potassium salts of fatty acid as compared to pesticides Confidor® (Imidacloprid) and Thunder® (Imidacloprid + Betacyfluthrin) followed by Karate® (Lambda Cyhalothrin)

There was a general reduction in whitefly population with increase in concentration potassium salts of fatty acids in all the treated plots compared to plots without treatment. (Tables 2). Potassium salts of fatty acids at 1 % and 1.5% had the highest reduction of up to 50% and 54% in adult and nymph whitefly population. Application of

confidor® (Imidacloprid) and Thunder® (Imidacloprid + Betacyfluthrin) followed by Karate® (Lambda Cyhalothrin) reduced the whitefly population by up to 47% and 46 %, respectively. No whitefly incidence was recorded in Embu during the first and second season. Application of potassium salts of fatty acids at 1% and 1.5% significantly ($P < 0.05$) reduced the thrips population at both Embu and Mwea site by up to 58 % and their activity was comparable to the application of chemicals Confidor® and Thunder® followed by Karate® (Table 3). The level of thrips population per flower was higher in Mwea than in Embu.

Potassium salts of fatty acids at the rate of 1 % and 1.5% resulted in the highest and significant ($P < 0.05$) increase of extra-fine and fine pods by up to 112% and 46 %, respectively, in Mwea (Table 4). However, in Embu, the application of both potassium salts of fatty acids and chemicals did not have any significant (P -value > 0.05) effect on the yield of both extra-fine and fine pods during the first season and second season (Table 4.0). Potassium salts of fatty acids significantly ($P < 0.05$) reduced pest damaged extra-fine and fine pods (Table 5.0). Potassium salts of fatty acids at the rate 1.5% and chemical treatment with confidor® potassium salts spray solution had the highest reduction of pest damage in both the extra-fine and fine grade of 76% and 69% respectively in Mwea (Table 5). In Embu, potassium salts of fatty acids had significant ($P < 0.05$) effect on the pest damaged pods. Potassium salts of fatty acids at the rate of 1.5% potassium salts and chemical treatment with confidor® had the highest reduction of 70% and 58% reduction respectively in pest damaged fine pods (Table 5). The use of potassium salts at 1% and 1.5% had the least cost-benefit ratio, followed by Confidor® (imidachloprid) (Figure 2). The least profitable pest management options were potassium salts at 0.5 % and in plots with application of Thunder® (Imidacloprid + Betacyfluthrin) followed by Karate® (Lambda Cyhalothrin) which was comparable to control plots.

Table 2. Mean number adult whitefly per sticky card and whitefly nymphs per leaf for different rates of potassium salts per season in snap beans in Mwea

Treatments	Adult whitefly			Whitefly nymphs		
	Season 1	Season 2	Mean	Season 1	Season 2	Mean
0.5 % Potassium salts	92.0c	68.7b	80.3b	32.5bc	34.9b	33.4b
1 % Potassium Salts	68.4ab	33.4a	50.9a	30.5b	17.3a	18.9a
1.5 % Potassium Salts	62.6a	30.7a	46.6a	19.9a	14.9a	17.3a
Confidor®	82.4bc	27.9a	55.1a	21.6a	18.5a	20.1a
Thunder® + Karate®	76.6b	33.4a	54.6a	21.2a	20.0a	20.4a
Control (Water only)	110.9d	78.7b	94.6c	37.1c	38.7b	37.8b
LSD ($p \leq 0.05$)	13.6	16.4	10.1	5.4	9.7	6.1
C.V%	1.7	8.4	3.9	8.4	11.9	10.0

Treatments with different letters in the same column are significantly different at 5% probability

Table 3. Mean number thrips per flower for different rates of potassium salts per season in snap beans in Mwea and Embu

Treatments	Mwea			Embu		
	Season 1	Season 2	Mean	Season 1	Season 2	Mean
0.5 % Potassium salts	2.90b	2.40cd	2.65b	0.33b	1.70bc	1.00bc
1 % Potassium Salts	2.00a	1.70ab	1.85ab	0.27ab	0.93a	0.60ab
1.5 % Potassium Salts	1.90a	1.60a	1.78a	0.18a	0.79a	0.49a
Confidor®	2.10a	1.80ab	1.95ab	0.17a	0.88a	0.53ab
Thunder® + Karate®	2.30ab	2.20bc	2.25ab	0.29b	1.27ab	0.78b
Control (Water only)	3.60c	2.70d	3.15c	0.35b	1.97c	1.16c
LSD ($p \leq 0.05$)	0.60	0.56	0.48	0.11	0.50	0.26
C.V%	15.80	3.20	13.2	14.40	8.80	9.70

Treatments with different letters in the same column are significantly different at 5% probability

Table 4. Total yield (kg/ ha) of marketable snap bean pods in each grade for different rates of potassium salts in snap beans per season plantings in Mwea and Embu

	Extra fine pods			Fine pods		
	Season 1	Season 2	Mean	Season 1	Season 2	Mean
Mwea						
0.5 % Potassium salts	2394ab	1529a	1962a	2654a	5679bc	4166b
1 % Potassium Salts	2760bc	3049c	2905b	2754a	6055c	4404b
1.5 % Potassium Salts	4072cd	3236c	3654b	2754a	5706c	4230b
Confidor®	3138c	3099c	3118b	2974a	5658bc	4316b
Thunder® + Karate®	3748c	2355abc	3050b	3898a	4180ab	4039b
Control (Water only)	1624a	1821ab	1722a	1946a	4092a	3019a
LSD ($p \leq 0.05$)	1052	1203	943	1624	1509	947
C.V%	15	13	11.5	30.1	16.8	5.5
Embu						
0.5 % Potassium salts	5826a	1165a	3496a	6621a	1324a	3973a
1 % Potassium Salts	7090a	1418a	4254a	6068a	1214a	3640a
1.5 % Potassium Salts	7386a	1477a	4432a	5801a	1160a	3481a
Confidor®	5936a	1187a	3562a	6271a	1254a	3763a
Thunder® + Karate®	6085a	1217a	3651a	5976a	1195a	3586a
Control (Water only)	5510a	1102a	3306a	4043a	847a	2445a
LSD ($p \leq 0.05$)	2564	513	1538	1983	421	1201
C.V%	12.7	12.7	12.7	12.4	12.5	12.4

Treatments with different letters in the same column are significantly different at 5% probability

Table 5. Total yield (kg/ ha) of pest damaged snap bean pods in each grade for different rates of potassium salts in snap beans per season plantings in Mwea and Embu

	Extra fine pods			Fine pods		
	Season 1	Season 2	Mean	Season 1	Season 2	Mean
Mwea						
0.5 % Potassium salts	149b	66a	108b	224ab	265b	244b
1 % Potassium Salts	128ab	69a	98ab	187a	114a	150a
1.5 % Potassium Salts	90a	28a	59a	164a	79a	121a
Confidor®	72a	98a	85ab	156a	110a	133a
Thunder® + Karate®	203c	103a	153c	274bc	328b	301b
Control (Water only)	221c	274b	247d	328c	465c	396c
LSD ($p \leq 0.05$)	57	89	41	70	106	63
C.V%	11.6	24.8	13.3	28.4	10.0	16.7
Embu						
0.5 % Potassium salts	234b	91c	163b	389b	78b	233b
1 % Potassium Salts	84a	47b	66a	268a	54a	161a
1.5 % Potassium Salts	115a	23a	69a	178a	36a	107a
Confidor®	39a	8a	24a	246a	49a	148a
Thunder® + Karate®	288b	59b	173b	459b	92b	275b
Control (Water only)	456c	91c	274c	592c	119c	355c
LSD ($p \leq 0.05$)	101	20	61.2	100	20	60.4
C.V%	24.1	24.1	24.1	18.9	18.9	18.9

Treatments with different letters in the same column are significantly different at 5% probability




Figure 2. Cost-benefit ratios for potassium salts of fatty acids and conventional chemicals in management of white fly and thrips in snap beans

4. Discussion

Potassium salts of fatty acids at 1.5% concentration significantly ($P < 0.05$) reduced whitefly population and was effective in the management of whitefly causing up to 65% reduction in whitefly population. However, potassium salts at 0.5% had no significant ($P > 0.05$) effect compared to the untreated control plots. These findings are consistent with reports by Mohamad et al., (2013) and Dheeraj et al., (2013) who reported effect of potassium salts on aphids and whiteflies, respectively. Similar results have been reported by Liu et al., (2000), Vavrina et al., (1995) and Ciancio et al., (2010). Hollingsworth, (2005) also reported the control of scales and mealybugs using potassium salts of fatty acids.

The effectiveness of potassium salts of fatty acids in this study increased with the increase in their concentration in the spray solution. The results agree with the findings by Liu et al., (2000) where increase in the concentration of potassium salts resulted in higher mortalities in whitefly. These results demonstrated that potassium salts are just as effective as synthetic chemical pesticides in the management of whitefly (Dheeraj et al., 2013).

Potassium salts of fatty acids work only on direct contact with the pest (Liu et al., 2000) and work on most soft-bodied insect pests (Koppert, 2013). They have been successfully used in the management of aphids, whiteflies, scales and mealy bugs (Mohamad et al., 2013; Dheeraj et al., 2013; Hollingsworth, 2005).

The results of this study showed that potassium salts of fatty acids at 1.5% concentration were also effective in management of thrips causing up to 68% reduction in their population. The results are consistent with the findings by Clinton et al., (2011) who reported the management of thrips in onions and by Heidi and Cullen (2008), Ciancio et al. (2010), Mohamad et al. (2013) and Dheeraj et al. (2013) on aphids and other arthropod pests. Potassium salts of fatty acids wash away the protective coating on the insect surface and penetrate the cell membrane causing disruption of the cell membrane permeability which leads to desiccation of the insects (Mohamad et al., 2013; Dheeraj et al., 2013; Ciancio et al., 2010).

Snap bean pod yield was significantly ($P > 0.05$) increased by up to 151% by application of 1% and 1.5% potassium salts of fatty acids and the yield increase was comparable to that obtained from the plots treated with conventional chemicals confidor® (Imidacloprid) and Thunder® (Imidacloprid + Betacyfluthrin) followed by Karate® (Lambda Cyhalothrin). The results concur with findings by Clinton et al., (2011) who reported an increase in yield with decrease in pest population after application of potassium salts of fatty acids and by Nderitu et al., (2009), Delkhoshi et al., (2012) and El-Mohamedy et al., (2008) as a result of application of pest control measures to reduce pest populations. However, the results of the current study contradicts the results by Vavrina et al., (1995) who reported that application of higher rates of potassium salts of fatty acids reduced yield of tomato fruits due to reduced biomass as a result of phytotoxicity at higher concentrations of potassium salts of

fatty acids.

The study showed that potassium salts of fatty acids are as effective as synthetic chemicals in managing pests in snap bean production resulting in increased yields and produce quality as reported by Dheeraj et al. (2013) and Mohamad et al. (2013). However, unlike synthetic chemical pesticides, potassium salts of fatty acids are fast acting with a quick knock down effect on the pests and break down quickly after application leaving no residues on the crop (Dheeraj et al., 2013; Hollingsworth, 2005). The cost benefit analysis showed that the use of potassium salts for management of thrips and whitefly was just as profitable as using the chemical pesticides. Potassium salts at 1.5% concentration was most effective and more profitable compared to concentrations of 0.5% and 1.5%. Though the potassium salts were as equally effective as synthetic chemicals, they pose no risk of residues on the crop and therefore offer the best alternative to chemical pesticides.

The results above demonstrate that potassium salts of fatty acids at the concentration of 1.5% potassium salts in spray solution are effective in the management of thrips and whiteflies on snap beans and can be used as alternative to synthetic chemical pesticides in snap bean production. Snap bean farmers depend on continuous and excessive use of synthetic pesticides often leading to accumulation of hazardous residues, development of resistant pest strains and loss of biodiversity due to toxicity to non-target organisms, in addition to the direct negative effect on health of the producers (Cardona et al., 2012; Das et al., 2010; Gorman et al., 2007; Kumar et al., 2012). High residue levels in fresh vegetables is the common cause of non-compliance to European market standards thus leading to enhanced scrutiny of Kenyan produce at the market with 10% of all Kenyan bean produce being sampled (Mwangi, 2013). Potassium salts of fatty acids are fast acting, break down quickly after application leaving no residues and are easy to handle because of their low toxicity to humans (Dheeraj et al., 2013). Therefore, they have no MRL or PHI requirements (Koppert, 2013) thus they would enable snap bean produce meet the stringent European market thereby reducing the losses currently experienced by fresh vegetable farmers and exporters (Mwangi, 2013; Ndung'u, 2013).

Acknowledgement

The authors wish to thank the Kenya Agricultural Productivity and Agribusiness Project (KAPAP) for funding this study under the Grant Number KAPAP/comp.2/06/Vol.II/39 and Koppert Biological Systems (K) for providing logistical support.

References

- Cardona, C., Frei, A., Bueno, J. M., Diaz, J., Gu, H., & Dorn, S. (2012). Resistance to *Thrips palmi* (Thysanoptera: Thripidae) in Bean. *Journal of Economic Entomology*, 95(5), 1066-1073. <https://doi.org/10.1603/0022-0493-95.5.1066>.
- Ciancio, A., & Mukerji, K. G. (2010). *Integrated management of arthropod pests and insect borne diseases*. Springer science + Business BV, London New York. <https://doi.org/10.1007/978-90-481-8606-8>
- Clinton, C. S., Erik, B. G. F., Kenzie, J. B., & Lamont, D. S. (2011). *Insecticide rotations for control of thrips in onions*. Ontario Malheur Experiment Station, Oregon State University.
- Das, S., Roy, S., & Mukhopadhyay, A. (2010). Diversity of arthropod natural enemies in the tea plantations of North Bengal with emphasis on their association with tea pests. *Current Science*, 99, 1457-1463.
- Delkhoshi, H., & Jalilian, J. (2012). Effect of pre-sowing seed treatment and spraying of bio-organic nutrient on yield and yield components of maize (*Zea mays* L.). *International Research Journal of Applied and Basic Sciences*, 3(9), 1867-1873.
- Dheeraj, J., Susan, V. P., & Nisha, S. (2013). Fatty acid metallic salts and pyrethroids - environmental friendly pesticides. *International Journal of Scientific Research and Reviews*, 2(1), 43- 51.
- El-Mohamedy, R. S. R., & Abd El-Baky, M. M. H. (2008). Evaluation of different types of Seed Treatment on Control of Root Rot Disease, Improvement Growth and Yield Quality of Pea Plant in Nobaria Province. *Research Journal of Agriculture and Biological Sciences*, 4(6), 611-622.
- European Union (EU) (2013). Maximum residue levels (MRLs) for pesticides in food. www.cbi.eu. Accessed 20th March 2014.
- Fritz, L. L., Heinrichs, E. A., Machado, V., Andreis, T. F., Pandolfo, M., de Salles, S. M., de Oliveira, J. V., & Fiuza, L. M. (2013). Impact of lambdacyhalothrin on arthropod natural enemy populations in irrigated rice fields in southern Brazil. *International Journal of Tropical Insect Science*, 33(3), 178-187. <https://doi.org/10.1017/S1742758413000192>.

- Gorman, K., Devine, D., Bennison, J., Coussons, P., Punchard, N., & Denholm, I. (2007). Report of resistance to the neonicotinoid insecticide imidacloprid in *Trialeurodes vaporariorum* (Hemiptera: Aleyrodidae). *Pest Management Science*, 63, 555-558. <https://doi.org/10.1002/ps.1364>
- Heidi, K., & Cullen, E. M. (2008). Efficacy and nontarget effects of reduced-risk insecticides on *Aphis glycines* Hemiptera: Aphididae) and its biological control agent *Harmonia axyridis* (Coleoptera: Coccinellidae). *Journal of Economic Entomology*, 101, 2391- 398.
- Hirano, K., Budiyo, E., Swastika, N., & Fujii, K. (1995). Population dynamics of the whitefly, *Bemisia tabaci* (Gennadius) (Homoptera: Aleyrodidae), in Java, Indonesia, with special reference to spatio-temporal changes in the quantity of food resources. *Ecological Research*, 1(1), 75-85. <https://doi.org/10.1007/BF02347657>.
- Hoelmer, K. A., Roltsch, V. J., Chu, E. C., & Hekneberry, T. J. (1998). Selectivity of Whitefly Traps in Cotton for *Eretmocerus eremicus* (Hymenoptera: Aphelinidae), a Native Parasitoid of *Bemisia argentifolii* (Homoptera: Aleyrodidae). *Environmental Entomology*, 27(4), 1039-1044. <https://doi.org/10.1093/ee/27.4.1039>
- Hollingsworth, R. G. (2005). Limonene, a citrus Extract, for control of mealybugs and scale insects. *Journal of Economic Entomology*, 98(3), 772-779. <https://doi.org/10.1603/0022-0493-98.3.772>
- Horticultural Crop Development Authority (HCDA) (2013). Export statistics for horticultural crops in 2012. Horticultural Crop Development Authority, Nairobi, Kenya.
- Horticultural Crop Development Authority (HCDA) (2014). Vegetables- French Beans. http://www.hcda.or.ke/?page_id=10541. Accessed 16th May 2014.
- Infonet-Biovision (2013). Infonet-Biovision. <http://www.infonet-biovision.org/default/ct/118/crops>. Accessed 6th July 2013
- Infonet-Biovision (2014). Infonet-Biovision. <http://www.infonet-biovision.org/default/ct/118/crops>. Accessed 16th May 2014.
- Jaetzold, R., Schmidt, H., Hornetz, B., & Shisanya, C. (2009). Farm Management Handbook of Kenya VOL. II— Natural Conditions and Farm Management Information –2nd Edition. Nairobi, Kenya.
- Koppert (2013). www.koppert.com: <http://www.koppert.com/products/products-pests-diseases/products/detail/savona-3/>. Accessed 6th July 2013.
- Kumar, R., Kranthi, S., Nitharwal, M., Jat, S. L., & Monga, D. (2012). Influence of pesticides and application methods on pest and predatory arthropods associated with cotton. *Phytoparasitica*, 10(10), 12-241. <https://doi.org/10.1007/s12600-012-0241-5>
- Liu, T., & Stansly, P. A. (2000). Insecticidal activity of surfactants and oils against silverleaf whitefly (*Bemisia argentifolii*) nymphs (Homoptera: Aleyrodidae) on collards and tomato. *Pest Management Science*, 56, 861-866. [https://doi.org/10.1002/1526-4998\(200010\)56:10<861::AID-PS217>3.3.CO;2-P](https://doi.org/10.1002/1526-4998(200010)56:10<861::AID-PS217>3.3.CO;2-P)
- Mead, R., Curnow, R. N., & Hasted, A. M. (2003). Statistical methods in agriculture and experimental biology. Third Edition. p 122. Chapman & Hall / CRC.
- Ministry of Agriculture (2012). National Horticulture Policy. http://www.kilimo.go.ke/kilimo_docs/pdf/National_Horticulture_Policy.pdf. Accessed 16th May 2014.
- Mohamad, S. F. S., Mohamad, S., & Aziz, A. A. (2013). The susceptibility of aphids, *Aphis gossypii* Glover to lauric acid based natural pesticide. *Procedia Engineering*, 53, 20-28. <https://doi.org/10.1016/j.proeng.2013.02.004>
- Monda, E. O., Munene, S., & Ndegwa, A. (2003). Snap bean production constraints in Kenya. *African Crop Science Conference Proceedings*, 6, 683-687.
- Mwangi, T. (2013). Threat to market access for Kenya's beans and snowpeas to EU. *Hortfresh Journal*, 5-6. http://hortfreshjournal.com/old/articles/threat_%20to_Mmarket_access_for%20kenyas_beans.pdf
- Nderitu, J. H., Wambua, E. M., Olubayo, F., Kasina, J. M., & Waturu, C. N. (2007). Evaluation of snap bean (*Phaseolus vulgaris* L.) cultivars and breeding lines for resistance to thrips (Thysanoptera: Thripidae) pests in Kenya. *Journal of Entomology*, 4, 202-208. <https://doi.org/10.3923/je.2007.202.209>
- Nderitu, J., Mwangi, F., Nyamasyo, G., & Kasina, M. (2009). Evaluation of cropping systems as a strategy for managing snap bean flower thrips in Kenya. *International Journal of Sustainable Crop Production*, 4(6), 22-25.

- Ndung'u, A. (2013). Kenya spills the beans. *Horticultural News*, 2, 36-37.
- Soto, A., Ohlenschläger, F., & García-Marín F. (2002). Distribution and Sampling of the whiteflies *Aleurothrixus floccosus*, *Dialeurodes citri*, and *Parabemisia myricae* (Homoptera: Aleyrodidae) in citrus in Spain. *Journal of Economic Entomology*, 95(1), 167-173. <https://doi.org/10.1603/0022-0493-95.1.167>
- United States Agency for International Development-Kenya Horticulture Competitiveness Project (USAID-KHCP) (2011). The European Market for Green Beans Market Survey #04. United States Agency for International Development-Kenya Horticulture Competitiveness Project Monthly Bulletin.
- Vavrina, C. S., Stansly, P. A., & Liu, T. X. (1995). Household Detergent on Tomato: Phytotoxicity and Toxicity to Silverleaf Whitefly. *Hortscience*, 30(7), 1406-1409.

Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.

This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).