

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search http://ageconsearch.umn.edu aesearch@umn.edu

Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.



Evolution in Well-being and Happiness after Increases in Consumption of Fruit and Vegetables

Redzo Mujcica and Andrew J. Oswald

July 2016

No: 1128

Warwick Economics Research Papers

ISSN 2059-4283 (online) ISSN 0083-7350 (print)

1	Evolution in Well-being and Happiness after Increases in
2	Consumption of Fruit and Vegetables
3 4	
5	2016
6	2010
7	
8	
9	Redzo Mujcic ^a and Andrew J. Oswald ^{b*}
10	
11	
12	
13	
14	^a Faculty of Health and Behavioural Sciences, University of Queensland, Brisbane QLD 4072,
15	Australia.
16	^b Department of Economics and CAGE Centre, University of Warwick, Coventry, CV4 7AL,
17	UK.
18	
19	
20	
21	
22	
23 24	2016 <i>ajph</i>
24 25	2010 ajpn
23 26	
20 27	*To whom correspondence should be addressed: phone: +44 (0)2476 523510. Email:
28	andrew.oswald@warwick.ac.uk. Full address: University of Warwick, Coventry CV4 7AL,
29	United Kingdom.
30	
31	
32	
33	WORD COUNT: 3100 words approx. (excluding supplemental material).

Abstract

- 36 OBJECTIVES
- 37 To explore whether improvements in psychological well-being occur after increases in fruit
- 38 and vegetable consumption.
- 39 METHODS
- 40 Longitudinal food diaries were examined on 12,000 randomly-sampled Australian adults over
- 41 2007, 2009, and 2013. The study estimated fixed-effects regression equations on individuals'
- 42 happiness and life satisfaction. It adjusted for a large set of other influences, including people's
- 43 changing incomes and personal circumstances. Prospective analysis, Granger-causality tests,
- 44 and instrumental-variable estimation were also done.
- 45 RESULTS
- 46 Increases in fruit and vegetable intake were predictive of increases in happiness and life
- 47 satisfaction. Well-being improvements were of up to 0.24 life-satisfaction points (for an
- 48 increase of 8 portions a day), which is equal in size to the psychological gain of moving from
- 49 unemployment to employment. Improvements occurred within 24 months.
- 50 CONCLUSIONS
- 51 People's motivation to eat healthy food is weakened by the fact that physical-health benefits
- 52 accrue decades later. This study offers a new possibility. Public-health policy could emphasize
- 53 immediate well-being improvement from healthy eating.
- 54 POLICY IMPLICATIONS
- 55 Citizens could be shown longitudinal evidence that 'happiness' gains from healthy eating can
- 56 occur quickly and many years before enhanced physical health.

57 Fruit and vegetables are known to provide important health benefits (*1*, *2*). Yet in Western 58 society the typical citizen eats an unhealthy diet (US data are available at <u>www.cdc.gov/brfss</u> 59 and European data at <u>www.eufic.org</u>). The difficulty of persuading people to consume more 60 fruit and vegetables remains a serious one (*3-7*).

This study explores a new approach to the problem. The paper is designed partly for the scientific researcher and partly for the public-health practitioner. It uncovers evidence consistent with a longitudinal connection between the consumption of certain foods (especially fruit and vegetables) and later subjective well-being, and a channel that appears to be independent of long-run health.

In disciplines beyond public-health research, the study of happiness and well-being has 66 67 generated a large modern literature. It is summarized in sources such as 8, 9. The potential influence of food has been virtually ignored. Traditional research on well-being has focused 68 69 upon the role of economic, personal, and political influences (see e.g. 9, 10, 11) and in character 70 has been steadily moving in emphasis from cross-sectional to longitudinal analysis (e.g. 12). 71 That the scholarly literature has developed in the way described is perhaps unsurprising. First, 72 most data sets do not record information on the foods eaten by individuals; second, the main 73 contributors to the happiness and well-being literature have been researchers from the classic 74 social-science disciplines. Hence it is perhaps understandable that the role of food in the list 75 of determinants of well-being has so far been given little attention, even though, in an important 76 line of work, researchers (such as 13) have, within a different literature, drawn attention to the 77 potential social significance of diet.

The present study uses a representative panel of 12,000 individuals to trace the potential linkages running from diet to later life satisfaction and happiness. It is intended as a complement to the aforementioned literature on socio-economic influences. In its style, the study fits within an emerging panel-data literature on human well-being. The analysis is first done by following individuals between 2007 and 2009. Just as the project was completed, however, new data were released, which makes it possible to check the calculations also for the period 2009 to 2013 (these replication findings are reported in supplemental tables S9-S11).

There are precursors to this paper. Innovative research by Tamlin Conner and collaborators (*14*) has found -- using data on daily food diaries on 281 students tracked over a three-week period -- that a high level of fruit and vegetable consumption appears to be predictive of greater emotional well-being on the following day. Various cross-sectional papers have also pointed to the possible existence of a statistical connection between psychological well-being and the amount of fruit and vegetables eaten, and have shown that 91 this correlation survives the inclusion of a large number of covariates (15).

92 There is also a small longitudinal literature which suggests there may be positive benefits 93 from a high intake of fruit and vegetables, although, crucially, that literature has not been able 94 to control for some of the key confounders such as individuals' levels of income (16-19). There 95 have also been three important small randomized controlled trials: on nutritional counseling 96 and on the provision of healthy food and snacks (20, 21, 22), which find some evidence that a 97 higher intake of fruit and vegetables may be associated with improved psychological health (as 98 well as physical health). Another set of writings has tried to understand obesity and its links 99 to subjective well-being (e.g. 23, 24). These suggest that there is an inverse -- although 100 relatively small -- correlation between body mass index BMI and mental well-being.

101 This paper documents not cross-sectional patterns but rather the longitudinal (the so-called 102 'change-on-change') linkages between fruit and vegetable consumption and mental well-being; 103 such an approach helps ensure that any observed relationship is not merely a spurious cross-104 sectional pattern caused by omitted confounding factors such as personality, background 105 wealth, or prior family upbringing. Cognizant of the work of others (25), this paper examines 106 whether the level of fruit and vegetable consumption today is predictive of the level of later 107 well-being, while inquiring into reverse-causality concerns hitherto unaddressed in the 108 happiness literature.

109

110 METHODS

111

112 The main data in this study come from Waves 7 and 9 (years 2007 and 2009) of the Household, 113 Income and Labour Dynamics in Australia (HILDA) Survey, a nationally representative panel 114 survey that began in 2001. The HILDA Survey collects annual longitudinal information from 115 members of Australian households who are at least 15 years of age. It provides information on 116 a total of 13,969 individuals from 7,682 different households interviewed since the first wave. 117 Data are collected each year by face-to-face interviews and self-completion questionnaires. 118 The former technique is mainly used to gather the demographic and socio-economic 119 information, while the latter is adopted to measure health and lifestyle choices.

After excluding respondents with missing information on the key outcome and control variables, the total sample available for this study consists of 12,389 individuals (aged 15 to 93) and 20,136 person-year observations. No observations are deliberately dropped. As would be expected, however, the sample sizes vary slightly across the different well-being measures. Two questions relating to fruit and vegetable consumption are available in Waves 7 and 9.
The corresponding questionnaires ask:

126 - Including tinned, frozen, dried and fresh fruit, on how many days in a usual
127 week do you eat fruit?

128 - Including tinned, frozen and fresh vegetables, on how many days in a usual
129 week do you eat vegetables?

with possible responses ranging from 0 ("do not eat any fruit or vegetables in a usual week")
to 7 days per week. For individuals who respond with some positive frequency to the questions
above, the following is also asked:

- On a day when you eat fruit, how many serves of fruit do you usually eat?
- On a day when you eat vegetables, how many serves of vegetables do you usually eat?

135 The survey respondents are shown flashcards to visually define a serving size or portion 136 (photographs of these are given as Figures S3-S4 in the Supplemental Material), with possible 137 answers ranging from '1' to '6 or more' portions. This visual approach is for simplicity and 138 clarity (see, e.g., 26). We multiply the responses to the above paired (frequency and quantity) 139 questions to form a weekly consumption amount of fruit and vegetables, respectively. We then 140 divide each resulting product by seven to arrive at the average daily amount. The average 141 intake of fruit by each survey respondent is then added to their average intake of vegetables to 142 compute the combined average daily consumption of fruit and vegetables. The mean value is 143 3.84 serves per day with a standard deviation of 2.01. Some respondents said they did not 144 consume any fruit or vegetables in a typical week. This group forms the 'none' or 'zero' 145 consumption category. Approximately 85% of respondents have fewer than 3 daily servings 146 of fruit; 60% consume fewer than 3 daily servings of vegetables. A small fraction of people 147 consume, on average, both more than 5 servings of fruit (1.83%) or vegetables (7.75%) each 148 day. Table S8 contains more detailed summary statistics on the separate fruit and vegetable 149 intake measures.

The first dependent variable examined is self-reported life satisfaction, derived from the question: "All things considered, how satisfied are you with your life?" Respondents are told to: "Pick a number between 0 and 10 to indicate how satisfied you are", and that "the more satisfied you are the higher number you should pick". Overall, the mean score for the sampled individuals in Australia is 7.91 with a standard deviation of 1.41. About two-thirds of 155 respondents report a life satisfaction score of more than 7 out of 10. As an additional check, a 156 second measure is used. A generic health variable available in the HILDA data set is the 157 Medical Outcomes Short Form (SF-36) Questionnaire. The SF-36 is a one of the most widely 158 used and validated self-completion measures of health status available, consisting of 36 159 items/questions; 35 of them are used to derive eight health subscales/indices. The respondent 160 is asked '*how much of the time in the past four weeks..*' did he/she experience particular types 161 of feelings/symptoms, including '... been a happy person'. The resulting response distribution 162 for the latter question is as follows: 1% (None of the time); 4.8% (A little of the time); 13.9% 163 (Some of the time); 19.5% (A good bit of the time); 51.9% (Most of the time); 8.9% (All of the 164 time). The individuals' responses are coded as from 1 (None of the time) to 6 (All of the time), 165 with a mean happiness score of 4.43 out of 6.

166

167 **RESULTS**

168

Figure 1 is a simple graphical illustration of the study's key result for life satisfaction. A similar histogram holds also for happiness data. The plot in Figure 1 is based on a so-called fixed-effect regression equation. It depicts the (uncorrected) longitudinal relationship -- the change-on-change relationship -- between people's subjective well-being and nine different levels of fruit and vegetable consumption. Further descriptive information is provided in the Supplemental Material. Alternative kinds of scatter plots are given as Figures S1 and S2 in that material.

The regression analyses reported in Table 1 provide formal evidence. These correct for other influences following sources such as (8) and (27). The key coefficient in the first column of Table 1 is 0.03 ($\beta = 0.03$, 95% confidence interval, or CI = [0.01, 0.04], p = .002). This implies that a change from the lowest levels to the highest levels of fruit and vegetable consumption would, on average, be associated with a rise in life satisfaction of approximately 0.24 life-satisfaction points.

The implied effect-size is substantial. At first glance, the number 0.24 might be thought to indicate that the consequences of fruit and vegetable intake are minor. That interpretation is mistaken; it stems from a blurring of the distinction between inter-person variance and intraperson variance. As in much of the longitudinal public-health research, this study tries to understand not the (inevitably high) cross-sectional variation in human well-being but instead intra-person changes that might be capable of being influenced by public interventions. In column 1 of Table 1, this requires that a number such as 0.24 (which is 8 times the coefficient of 0.03) has to be added to the number 7.81. As can be seen from the later right-hand-side columns of Table 1, the effect is the equivalent in absolute size to (in the negative direction) that of becoming unemployed or approximately half the size of the emotional consequence of marital separation. Such an effect-size is large.

193 If Model 1 of Table 1 were the only regression result available, it would be plausible to 194 believe that the relationship is spurious. It might be being driven by omitted variables -- for 195 example, someone, say, becoming richer through time and becoming happier and 196 simultaneously eating in a healthier way because they could now afford it, or, say, divorcing a 197 spouse and becoming less happy and also eating in a less healthy way. However, the later 198 columns of Table 1 imply that such interpretations would be incorrect. The analyses here 199 include extra covariates: the natural logarithm of household income, age, education, whether 200 working, marital status, health, children, alcohol and food patterns, Body Mass Index, and 201 exercise (for a detailed specification of these variables see Tables S7-S8 in the Supplemental 202 Material). In Table 1, there is no detectable influence from BMI. A later table, Table S3 in 203 the Supplemental Material, however, is consistent with the existence of an inverse relationship 204 between current BMI and future well-being.

Figure 1 uses coefficients from longitudinal estimates. Fixed-effect estimation is equivalent here to a first-difference estimator, as discussed in 28, so they emerge, in effect, from regressing the change in well-being between 2007 and 2009 on the change over that period in variables such as food consumption, income, marital status, and so on. This is why, in Table 1, attributes such as gender and ethnicity are omitted; they are unchanging and thus have automatically been differenced out. Table 2 repeats the calculations for the alternative dependent variable of feeling happy. Results are similar.

212 An open scientific issue is whether diet might have slow-acting effects on mental well-213 being. The analyses reported in Table 3 explore this. They treat the data as if from a 214 prospective setting. Here the regression equations reveal that fruit and vegetable consumption 215 in the current year is predictive of higher well-being -- measured either as life satisfaction or 216 as happiness -- in the future even after controlling for current well-being (as well as controlling 217 for the list of covariates in the tables). Hence, in the life-satisfaction equation in Table 3, for 218 example, where the dependent variable is life satisfaction measured in period t+1, a variable 219 for fruit and vegetable consumption in period t is statistically significant at the 99.9% 220 confidence level ($\beta = 0.03$, 95% CI = [0.01, 0.04], p < .001), while holding constant life 221 satisfaction in period t, which itself enters, as would be expected, with a large positive 222 coefficient. Similar results are found for happiness in Table 3. The Supplemental Material

223 provides the equation specifications.

224 Such prospective analysis is subject to a potential objection. It is that some form of 225 correlation might run in both directions simultaneously. To check for this, a form of Granger 226 causality test was done, and is given in the Supplemental Material. Tables S4 and S5 test 227 whether fruit and vegetable consumption in the future can be predicted from the level of life 228 satisfaction or happiness in the current period. In neither case is there evidence for such 229 reverse-causality; the effect does not achieve statistical significance in either of the tables. In 230 Table S4, in fact, the variable has the wrong point-estimate sign ($\beta = -0.003$, 95% CI = [-0.03, 231 0.02], p > .250).

We checked whether the findings can be reproduced on a new round of the panel data set, which was released, towards the end of our project, for the year of 2013. The paper's key results can be replicated; the findings are presented in supplemental tables S9-S11. It can be seen in the extra tables that the coefficients remain essentially identical to those presented in the main body of the paper.

We also did a test for whether fruits and vegetables should be separated into two independent variables – rather than combined into the number of daily F&V portions variable that has been traditional in research on physical health. The results (not reported) suggested that for happiness and life-satisfaction equations it was appropriate to combine them into a single F&V variable. The null hypothesis of an identical well-being gradient for fruit intake and vegetables intake could not be rejected.

243 Last, we made another effort, in addition to the Granger causality tests, to tackle the 244 inevitably complex issue of causality. To do so we exploit a public campaign that was designed 245 to encourage healthy eating in Australia. Scientifically, the advantage of such a campaign is 246 that, from a researcher's point of view, an advertising campaign of this kind could be seen as 247 an exogenous positive 'shock' to people's motivation to eat a greater number of portions of 248 fruit and vegetables. Hence it offers the possibility of a form of natural experiment: as the 249 campaign came in, with different timings in different states, it might be expected that it would 250 shift people's consumption decisions at these particular points in time. Any consequences, for 251 mental well-being and physical well-being, might then go on to be detectable.

Known as the "Go For 2&5 Campaign", this initiative began in the state of Western Australia in the year 2004. It spread, at different speeds, into most of the other Australian states. Two-stage least squares estimation can then be done (as described in reference *29*). The instrumental-variable estimates are provided in supplemental tables S12-S15.

In this form of inquiry, we exploit the fact that different Australian states had different

257 number of years over which they systematically promoted the consumption of fruit and 258 vegetables. Victoria did so for zero years; New South Wales for 2 years; Tasmania for 4 years; 259 South Australia for 4 years; Queensland for 5 years; the Northern Territories for 7 years; ACT 260 for 7 years; and Western Australia for 10 years. Thus we create a variable for Intensity of 261 Campaign. This adds up the length (i.e. number of years) that a state had previously had a 262 campaign. All states in our analysis are thereby given an integer-valued entry, from 0 for 263 Victoria to 10 for Western Australia, as a measure of the different intensities of the public fruitand-vegetable campaign in the different states. In plainer English, the citizens of each region 264 265 can be thought of as having a different level of 'publicly-sponsored push' to eat in a healthy 266 way. That policy variable can be viewed as an extraneous influence upon later state levels of 267 consumption of fruit and vegetables.

268 Analytically, we then take two steps. The first is to estimate a Consumption of Fruit+Veg 269 equation (not a well-being equation) for the year 2013. We then test whether a variable for 270 Campaign Intensity comes in positively in that equation. We find that it does, with a 271 statistically significant coefficient. Hence there is evidence that the Australian healthy-eating 272 campaign had an effect on fruit and vegetable intake. Then, in the second stage of our two-273 stage least-squares estimation, a set of instrumented well-being regression equations for the 274 year 2013 are estimated. The purpose is to correct for simultaneity bias and the possibility of 275 reverse causality. After doing so, an instrumented variable for fruit and vegetable consumption 276 is found to enter positively in a well-being equation (as in Table S12). Hence there is some 277 evidence that the Australian healthy-eating campaign may have improved people's levels of 278 life satisfaction and happiness. Nevertheless, it is not possible statistically to be certain of that 279 conclusion. As is often found in the statistical literature on two-stage least-squares estimation, 280 the level of statistical power here is insufficient for us to obtain truly small standard errors in 281 the second-stage equations. The paper's confidence levels do not exceed 75% when using this 282 final form of statistical method.

283

284 **DISCUSSION**

285

This study is a longitudinal examination of the links between food and people's psychological well-being. It examines data on the lives of a nationally representative sample of approximately 12,000 individuals between 2007 and 2009, and is able to check, and replicate, its main findings for additional newly-released data over the period 2009 to 2013. Prospective analysis and Granger-causality tests are also done. By using information on the Australian "Go for 2&5 Campaign" it also attempts to offer instrumental-variable estimation.

This study's findings are consistent with the idea that eating certain foods is a form of investment in future happiness. The implications of fruit and vegetable consumption are estimated to be substantial and to operate within the space of two years -- too quickly to be a reflection of the physical advantages of diet for outcomes such as cardiovascular disease documented by earlier researchers (2). Moreover, as shown in Table S6 of the Supplemental Material, the fruit-and-vegetables effect still operates if the regression equation includes an extra covariate for self-reported health.

In a sense, the paper offers a new possibility for future public-policy programs to encourage healthy eating – the possibility that citizens in western society could be given evidence that 'happiness' gains from healthy eating may show up much more quickly than any long-distant improvement to their physical health. If individuals weigh up the likely benefits of fruit and vegetables in their diet, and set that against any perceived costs, both pecuniary and non-pecuniary, of doing so, scientific evidence of extra gains from a healthy diet may help persuade people to raise their intake of fruit and vegetables.

306 Two main issues remain to be tackled. First, although at the end of this study we 307 attempted to address the causality problem by using instrumental-variable methods, a huge 308 randomized trial would lead to a natural form of scientific evidence. The well-being research 309 literature is, however, far from such a point; a randomized trial would have its own inherent 310 difficulties, because a double-blind procedure would not be feasible, so placebo effects would 311 be hard to disentangle; and large-scale longitudinal studies, of the sort described in this study, 312 would still be required as part of a body of persuasive evidence. Second, the channels from 313 eating certain food types to subjective well-being remain to be properly understood. For 314 example, (18, 30) discuss a variety of intriguing possibilities. These include a potential 315 influence from vitamin B12 upon the eventual production of human serotonin, as well as the 316 idea of a role for folate deficiency (see also 31). A further potential channel (32) is that 317 microbiota may modulate brain chemistry. Lastly, it may be possible eventually to link the 318 current research to a new literature on antioxidants that is suggestive of a connection between 319 human optimism and carotenoid in the blood (33). Further connections between the biology 320 and practical public-health policy of healthy eating (34) remain to be forged. These issues 321 demand attention.

323	References
324 325	1. Patterson BH, Block G (1988). Food choices and the cancer guidelines. American Journal
325	of Public Health 78, 282-286.
320	 2. Ames BN, Shigenaga MK, Hagen TM (1993). Oxidants, antioxidants, and the degenerative
328	diseases of aging. <i>Proceedings of the National Academy of Sciences of the USA</i> 90, 7915-
328 329	7922.
330	3. Serdula MK, Gillespie C, Kettel-Khan L, et al. (2004). Trends in fruit and vegetable
331	consumption among adults in the United States: Behavioural Risk Factor Surveillance
332	System, 1994-2000. American Journal of Public Health 94, 1014-1018.
333	 4. Lucan SC, Barg FK, Long JA (2010). Promoters and barriers to fruit, vegetable, and fast-
333 334	food consumption among urban, low-income African Americans – A qualitative approach.
335	American Journal of Public Health 100, 631-635.
336	5. Kwon SC, Wyatt LC, Kranick JA, et al. (2015). Physical activity, fruit and vegetable intake,
337	
	and health-related quality of life among older Chinese, Hispanics, and Blacks in New York
338	City. American Journal of Public Health 105, S544-S552.
339 240	6. Sacks R, Yi SS, Nonas C (2015). Increasing access to fruits and vegetables: Perspectives
340	from the New York City experience. <i>American Journal of Public Health</i> 105, E29-E37.
341	7. Kesse-Guyot E, Andreeva VA, Lassale C, et al. (2014). Clustering of midlife lifestyle
342	behaviors and subsequent cognitive function: A longitudinal study. American Journal of
343	Public Health 104, E170-E177.
344	8. Dolan P, Peasgood T, White M (2008). Do we really know what makes us happy? A review
345	of the economic literature on the factors associated with subjective well-being. <i>Journal of</i>
346	Economic Psychology 29, 94-122.
347	9. Diener E (2013). The remarkable changes in the science of subjective well-being.
348	Perspectives on Psychological Science 8, 663-666.
349	10. Easterlin RA (2003). Explaining happiness. Proceedings of the National Academy of
350	Sciences of the USA 100(19), 11176-11183.
351	11. Oswald AJ, Wu S (2010). Objective confirmation of subjective measures of human well-
352	being: Evidence from the USA. Science 327, 576-579.
353	12. Lucas RE (2007). Long-term disability is associated with lasting changes in subjective well-
354	being: Evidence from two nationally representative longitudinal studies. Journal of
355	Personality and Social Psychology 92, 717-730.

- 356 13. Otero G, Pechlaner G, Liberman G, Gurcan E (2015). The neoliberal diet and inequality in
 357 the United States. *Social Science & Medicine* 142, 47-55.
- 14. White BA, Horwath CC, Conner T S (2013). Many apples a day keep the blues away–Daily
 experiences of negative and positive affect and food consumption in young adults. *British Journal of Health Psychology* 18(4), 782-798.
- 15. Blanchflower DG, Oswald AJ, Stewart-Brown S (2011). Is psychological well-being linked
 to the consumption of fruit and vegetables? *Social Indicators Research* 114, 785-801.
- 363 16. Jacka FN, Kremer PJ, Berk M, et al. (2011). A prospective study of diet quality and mental
 364 health in adolescents. *PLOS ONE* 6, e24805.
- 365 17. Blank L, Grimsley M, Goyder E, et al. (2007). Community-based lifestyle interventions:
 366 changing behaviour and improving health. *Journal of Public Health* 29, 236-245.
- 367 18. Rooney C, McKinley MC, Woodside JV (2013). The potential role of fruit and vegetables
 368 in aspects of psychological well-being: A review of the literature and future directions.
 369 *Proceedings of the Nutrition Society* 72, 420-432.
- Henriquez Sanchez P, Ruano C, de Irala J, Ruiz-Canela M, Martinez-Gonzalez MA,
 Sanchez-Villegas A (2012). Adherence to the Mediterranean diet and quality of life in the
 SUN project. *European Journal of Clinical Nutrition* 66, 360-368.
- 20. Steptoe A, Perkins-Porras L, Rink E, Capuccio FP (2004). Quality of life and self-rated
 health in relation to changes in fruit and vegetable intake and in plasma vitamins C and E
 in a randomized trial of behavioural and nutritional counselling. *British Journal of Nutrition*92, 177–184.
- 377 21. Salah M, Grimble G (2013). The effect of a fresh fruit and vegetable delivery on the health
 378 and independence of seniors. *International Journal of Food, Nutrition and Public Health*379 6, 123-141.
- Smith AP, Rogers R (2014). Positive effects of a healthy snack (fruit) versus an unhealthy
 snack (chocolate/crisps) on subjective reports of mental and physical health: A preliminary
 intervention study. *Frontiers in Nutrition* <u>10.3389/fnut.2014.00010</u>.
- 383 23. Oswald AJ, Powdthavee N (2007). Obesity, unhappiness and the challenge of affluence:
 384 Theory and evidence. *Economic Journal* 117, F441-F454.
- 385 24. Graham C (2008). Happiness and health: Lessons and questions for public policy.
 386 *Health Affairs* 27, 72-87.
- 387 25. Gardner MP, Wansink B, Kim J, Park SB (2011). Better moods for better eating? How
 388 mood influences food choice. *Journal of Consumer Psychology* 24, 320-335.

- 389 26. Tourangeau R, Rips LJ, Rasinski K (2000). *The psychology of survey response*. Cambridge
 390 University Press.
- 391 27. Blanchflower DG, Oswald AJ (2008). Is well-being U-shaped over the life cycle? *Social*392 *Science & Medicine* 66, 1733-1749.
- 28. Liker JK, Augustyniak S, Duncan GJ (1985). Panel data and models of change: A
 comparison of first difference and conventional two-wave models. *Social Science Research*14, 80-101.
- 396 29. Angrist JD, Imbens GW, Rubin DB (1996). Identification and causal effects using
 397 instrumental variables. *Journal of the American Statistical Society* 91(434), 444-455.
- 30. Young SN (2007). How to increase serotonin in the human brain without drugs. *Journal of Psychiatry and Neuroscience* 32, 394-399.
- 31. Botez MI, Young SN, Bachevalier, J, et al. (1979) Folate-deficiency and decreased 5hyrdoxytryptamine synthesis in man and rat. *Nature* 278, 182-183.
- 402 32. Dinan TG, Cryan JF (2013). Melancholic microbes: a link between gut microbiota and
 403 depression? *Neurogastroentorolgy & Motility* 25, 713-719.
- 33. Boehm JK, Williams DR, Rimm EB, Ryff C, Kubzansky LD (2013). Association between
 optimism and serum antioxidants in the Midlife in the United States Study. *Psychosomatic Medicine* 75, 2-10.
- 407 34. Pomerleau J, Lock K, Knai C, et al. (2005). Interventions designed to increase adult fruit
 408 and vegetable intake can be effective: A systematic review of the literature. *Journal of*409 *Nutrition* 135, 2486-2495.
- 410 35. Watson N, Wooden M (2002). *The household, income and labour dynamics in Australia*411 (*Hilda*) survey: Wave 1 survey methodology, HILDA Project Technical Paper Series, No
- 412 1/02, Melbourne Institute of Applied Economics and Social Research.
- 413

414	Author Contributions
415	Author contributions: RM had the idea for the study; RM and AJO designed the research; RM
416	led the study and wrote up the first results; AJO made suggestions for changes; RM and AJO
417	analyzed the data; both authors revised the draft. AJO wishes to record that the main credit for
418	this work is due to RM.
419	
420	Declaration of Conflicting Interests
421	The authors declare no conflict of interest.
422	
423	Acknowledgments
424	This paper replaces an earlier draft by RM alone; the estimation methods, and results, in this
425	paper differ from those in that older version. Helpful comments were received in a seminar at
426	the London School of Economics. We also thank Kylie Ball, M Lynne Cooper, Paul Frijters,
427	George Grimble, David W. Johnston, James Oswald, Ciara Rooney, Fiona Scott, Daniel Sgroi,
428	and Simon Young for helpful ideas. All errors are our own. Financial support from the ESRC
429	through the CAGE Centre at Warwick University is gratefully acknowledged. This paper uses
430	unit record data from the Household, Income and Labour Dynamics in Australia (HILDA)
431	Survey. The HILDA Project was initiated and is funded by the Australian Government
432	Department of Social Services (DSS) and is managed by the Melbourne Institute of Applied
433	Economic and Social Research (Melbourne Institute). The findings and views reported in this
434	paper, however, are those of the authors and should not be attributed to either DSS or the
435	Melbourne Institute.

436 437 438 439 Table 1. Life Satisfaction Equations: Fixed-effects Regression Models of Changes in Life Satisfaction on Changes in Fruit and Vegetable Consumption and Covariates. Longitudinal Survey Data on 12,000 Adults,

HILDA Survey 2007 and 2009.

	Model 1 (no covariates)		Model 2 (partial set of covari			Model 3 (full set of covariates)	
Independent variable	β	р	β	p	β	<i>p</i>	
Fruit and vegetable portions/day	0.03 [0.01, 0.04]	.002	0.03 [0.01, 0.04]	.003	0.02 [0.01, 0.04]	.010	
Log of household income			0.02 [-0.03, 0.06]	.452	0.02 [-0.03, 0.06]	.476	
Age			-0.01 [-0.05, 0.04]	.837	-0.01 [-0.06, 0.04]	.758	
Age ²			0.01 [-0.04, 0.05]	.766	0.01 [-0.04, 0.06]	.720	
Masters or doctorate			-0.31 [-0.86, 0.24]	.271	-0.32 [-0.87, 0.23]	.256	
Bachelor or honors			-0.07 [-0.48, 0.35]	.755	-0.05 [-0.46, 0.36]	.812	
Graduate diploma or certificate			-0.18 [-0.51, 0.16]	.304	-0.17 [-0.51, 0.17]	.315	
Advanced diploma			-0.09 [-0.46, 0.27]	.618	-0.10 [-0.47, 0.27]	.609	
Professional qualification			-0.01 [-0.30, 0.28]	.944	-0.02 [-0.31, 0.27]	.894	
Year 12 high school			-0.21 [-0.41,-0.01]	.036	-0.20 [-0.40, 0.00]	.045	
Full-time student			-0.01 [-0.15, 0.13]	.894	0.00 [-0.15, 0.14]	.965	
Unemployed			-0.21 [-0.43, 0.01]	.058	-0.22 [-0.44, 0.00]	.050	
Not in the labor force			-0.02 [-0.13, 0.09]	.695	-0.04 [-0.15, 0.07]	.508	
Married			-0.01 [-0.18, 0.16]	.917	-0.01 [-0.18, 0.16]	.895	
Separated			-0.57 [-0.89, -0.26]	.000	-0.58 [-0.89, -0.26]	.000	
Divorced			-0.32 [-0.63, -0.01]	.042	-0.33 [-0.64, -0.02]	.036	
Widowed			-0.45 [-0.99, 0.09]	.099	-0.46 [-1.00, 0.08]	.097	
Long-term health condition			-0.14 [-0.22, -0.07]	.000	-0.14 [-0.22, -0.07]	.000	
# children under the age of 4			-0.01 [-0.10, 0.08]	.838	-0.01 [-0.09, 0.08]	.881	
# children aged 5-14			0.06 [-0.02, 0.14]	.121	0.06 [-0.01, 0.14]	.108	
Drink alcohol 1 or 2 days/week					0.02 [-0.09, 0.14]	.697	
Drink alcohol 2 or 3 days/week					-0.01 [-0.11, 0.09]	.889	
Drink alcohol 3 or 4 days/week					-0.03 [-0.17, 0.10]	.619	
Drink alcohol 5 or 6 days/week					-0.04 [-0.20, 0.12]	.638	
Drink alcohol everyday					-0.14 [-0.34, 0.06]	.159	
Non-smoker					0.04 [-0.09, 0.17]	.532	
Never eat red meat					0.20 [-0.16, 0.55]	.273	
Never eat fish					-0.09 [-0.20, 0.02]	.107	
Eat breakfast regularly					0.11 [0.03, 0.18]	.004	
Drink low fat or skinny milk					-0.04 [-0.12, 0.04]	.316	
Avoid fatty foods					-0.05 [-0.12, 0.01]	.105	
BMI					0.01 [0.00, 0.01]	.115	
Exercise regularly					0.09 [0.03, 0.14]	.002	
Constant	7.81 [7.74, 7.88]	.000	7.90 [6.80, 9.00]	.000	7.75 [6.65, 8.85]	.000	
Overall <i>R</i> ²	.02		.03		.03		
Number of individuals	12,385		12,385		12,385		
Number of observations	20,127		20,127		20,127		

 $\begin{array}{c} 440\\ 441\\ 442 \end{array}$

Note: Values in parentheses are 95% confidence intervals. Dependent variable is Life Satisfaction [range: 0-10]. HILDA Survey data: Australia. Further details of the data set are available in reference (35). With two waves of data, a fixed-effects estimator is equivalent to a first-difference estimator; see, for example, reference 28.

Table 2. Happiness Equations: Fixed-effects Regression Models of Changes in 'Been a Happy Person'

443 444 on Changes in Fruit and Vegetable Consumption and Covariates. Longitudinal Survey Data on 12,000 445 Adults, HILDA Survey 2007 and 2009.

446

	Model 1 (no covariates)		Model 2 (partial set of covariates)		Model 3 (full set of covariates)	
Independent variable	β	р	β	р	β	р
Fruit and vegetable portions/day	0.02 [0.01, 0.03]	.003	0.02 [0.01, 0.04]	.002	0.02 [0.003, 0.03]	.017
Log of household income			0.02 [-0.02, 0.05]	.369	0.02 [-0.02, 0.05]	.320
Constant	4.35 [4.30, 4.40]	.000	4.29 [3.40, 5.17]	.000	4.31 [3.42, 5.20]	.000
Other covariates included	No		Yes (a partial set)		Yes (a full set)	
Overall <i>R</i> ²	.02		.01		.03	
Number of individuals	12,360		12,360		12,360	
Number of observations	20,054		20,054		20,054	

447 448 Note: Values in parentheses are 95% confidence intervals. Dependent variable is Been a Happy Person [range: 1-6]. HILDA Survey

data: Australia. 'Partial set' and 'Full set' are as defined in columns 2 and 3, respectively, of Table 1. The full estimation results (with

449 a complete set of control variable coefficient estimates) are available in Table S1 in the Supplemental Material.

450 Table 3. Prospective Analyses of Life Satisfaction and Happiness on Lagged Fruit and Vegetable 451

Consumption. HILDA Survey 2007 (period *t*) and 2009 (period t+1)

1	5	2
4	J	L

	Life satisfaction t+1	Been a happy person t+1
Independent variable	β	β
Fruit and vegetable portions/day t	0.03 [0.01, 0.04]	0.02 [0.01, 0.03]
Life satisfaction t	0.49 [0.47, 0.50]	
Been a happy person t		0.45 [0.43, 0.47]
Log of household income t	0.03 [0.00, 0.07]	0.03 [0.00, 0.05]
Constant	3.98 [3.55, 4.41]	2.36 [2.04, 2.68]
Full set of other covariates:	Yes	Yes
Adjusted R ²	.31	.26
Number of observations	7,742	7,694

Note: Values in parentheses are 95% confidence intervals. First dependent variable is Life Satisfaction [range: 0-10] in period t+1 (year 2009). Second dependent variable is Been a Happy Person [range: 1-6] in period t+1 (year 2009). Period t denotes the year 2007. The full estimation results (with a complete set of control variable coefficient estimates) are available in Tables S2 and S3 in the Supplemental Material. The table's title uses the term 'prospective' for simplicity; it would be possible to object to this on strict semantic grounds; we obtained the data after the Wave 2 information, on year 2009, had been collected.




Fig. 1. Longitudinal changes in fruit and vegetable (F&V) consumption are positively correlated with longitudinal changes in satisfaction with life. The vertical axis here measures life satisfaction; the horizontal axis measures daily F&V portions. The 0 on the horizontal axis denotes less than one portion of fruit and vegetables per day, 1 denotes higher than one portion but less than two portions per day, ... and 8 denotes eight-and-above portions a day. The sample size is 12,385 Australian individuals measured in years 2007 and 2009.

An equivalent diagram would hold symmetrically for reductions in F&V consumption (not drawn above).

This figure is not cross-sectional. It is derived from a fixed-effects regression equation with nine banded dummy variables for the above nine different levels of fruit and vegetable (F&V) daily consumption. Formal test statistics are presented in Table 1, which treats F&V as a continuous variable.

485 END OF MANUSCRIPT......
486
487
488

SUPPLEMENTAL FILES: LATER MATERIAL IS FOR REFEREES ONLY AND/OR ONLINE PUBLICATION AS EXTRA INFORMATION FOR READERS.

491	
492	Supplemental Files (For Referees or Online Publication Only)
493	
494	
495	
496	
497	
498	
499	
500	Tables S1-S8
501	Figures \$1-\$4

 501
 Figures S1-S4

 502
 Tables S9-S15

504 Table S1 (Full Estimation Results for Table 2). Happiness Equations: Fixed-effects Regression Models of

Changes in 'Been a Happy Person' on Changes in Fruit and Vegetable Consumption and Covariates, HILDA Survey 2007 and 2009

	Model 1	-)	Model 2)	Model 3)
	(no covariates)		(partial set of covariates)		(full set of covariates)	
Independent variable	β	р	β	р	β	р
Fruit and vegetable portions/day	0.02 [0.01, 0.03]	.003	0.02 [0.01, 0.04]	.002	0.02 [0.003, 0.03]	.017
Log of household income			0.02 [-0.02, 0.05]	.369	0.02 [-0.02, 0.05]	.320
Age			-0.01 [-0.05, 0.03]	.736	0.00 [-0.04, 0.04]	.832
Age squared			0.01 [-0.03, 0.05]	.525	0.01 [-0.03, 0.05]	.571
Masters or doctorate			0.10 [-0.41, 0.61]	.697	0.15 [-0.35, 0.65]	.560
Bachelor or honors			-0.13 [-0.53, 0.26]	.507	-0.10 [-0.49, 0.29]	.614
Graduate diploma or certificate			-0.05 [-0.36, 0.27]	.778	-0.01 [-0.32, 0.30]	.942
Advanced diploma			-0.30 [-0.73, 0.13]	.173	-0.30 [-0.74, 0.15]	.190
Professional qualification			0.08 [-0.16, 0.32]	.493	0.08 [-0.16, 0.32]	.511
Year 12 high school			-0.04 [-0.23, 0.15]	.706	-0.01 [-0.20, 0.18]	.891
Full-time student			-0.03 [-0.16, 0.09]	.620	-0.03 [-0.15, 0.10]	.653
Unemployed			0.05 [-0.10, 0.19]	.528	0.05 [-0.10, 0.19]	.519
Not in the labor force			-0.10 [-0.19, -0.02]	.015	-0.11 [-0.19, -0.03]	.010
Married			-0.02 [-0.18, 0.14]	.808	-0.02 [-0.18, 0.14]	.80
Separated			-0.23 [-0.48, 0.03]	.083	-0.24 [-0.49, 0.02]	.069
Divorced			0.01 [-0.27, 0.29]	.942	-0.01 [-0.29, 0.27]	.958
Widowed			-0.14 [-0.47, 0.19]	.405	-0.15 [-0.48, 0.17]	.358
Long-term health condition			-0.07 [-0.13, -0.01]	.024	-0.06 [-0.12, 0.00]	.040
# children under the age of 4			0.03 [-0.03, 0.10]	.321	0.04 [-0.03, 0.11]	.233
# children aged 5-14			0.02 [-0.04, 0.08]	.460	0.03 [-0.03, 0.09]	.339
Drink alcohol 1 or 2 days/week					-0.05 [-0.14, 0.04]	.244
Drink alcohol 2 or 3 days/week					-0.02 [-0.10, 0.05]	.570
Drink alcohol 3 or 4 days/week					-0.07 [-0.17, 0.04]	.209
Drink alcohol 5 or 6 days/week					-0.04 [-0.16, 0.08]	.516
Drink alcohol everyday					0.03 [-0.12, 0.18]	.673
Non-smoker					0.01 [-0.09, 0.12]	.776
Never eat red meat					-0.02 [-0.27, 0.24]	.907
Never eat fish					0.05 [-0.04, 0.14]	.250
Eat breakfast regularly					0.12 [0.05, 0.18]	.000
Drink low fat or skinny milk					-0.01 [-0.07, 0.05]	.776
Avoid fatty foods					0.00 [-0.05, 0.05]	.935
BMI					-0.01 [-0.02, 0.00]	.009
Exercise regularly					0.14 [0.10, 0.19]	.000
Constant	4.35 [4.30, 4.40]	.000	4.29 [3.40, 5.17]	.000	4.31 [3.42, 5.20]	.000
Overall <i>R</i> ²	.02		.01		.03	
Number of individuals	12,360		12,360		12,360	
Number of observations	20,054		20,054		20,054	

 Note: Values in parentheses are 95% confidence intervals. Dependent variable is Been a Happy Person [range: 1-6].

514Table S2 (Full Estimation Results for First Part of Table 3). Prospective Analysis of Life Satisfaction:515Linear Regression Model of Life Satisfaction on Lagged Fruit and Vegetable Consumption and Covariates,516HILDA Survey 2007 (period t) and 2009 (period t+1)

Life satisfaction : 0.48 [0.47, 0.50] 49.31 0.00 Log of household income : 0.03 [0.00, 0.07] 1.78 0.07 Age : 0.02 [0.01, 0.03] 3.87 0.00 Male : 0.02 [0.01, 0.03] 3.87 0.00 Male : 0.01 [-0.05, 0.06] 0.20 8.44 Masters or doctorate : 0.01 [-0.07, 0.01] 1.77 0.77 Bachelor or honors : 0.06 [-0.14, 0.03] 1.21 2.22 Advanced diploma or certificate : 0.06 [-0.14, 0.03] 1.21 2.22 Advanced diploma : Professional qualification : 0.06 [-0.13, 0.02] 1.47 1.44 Year 12 high school : 1.75 0.88 Unemployed : 0.03 [-0.15, 0.21] 0.32 7.44 Not in the labor force : 0.03 [-0.15, 0.21] 0.32 7.44 Not in the labor force : 0.03 [-0.15, 0.21] 0.32 7.44 Not in the labor force : 0.03 [-0.15, 0.21] 0.32 7.44 Not in the labor force : 0.03 [-0.16, 0.05] 1.36 .077 Divorced : 0.03 [-0.17, 0.05] 1.36 .077 Divorced : 0.03 [-0.17, 0.05] 1.36 .077 Divorced : 0.01 [-0.22, 0.10] 0.15 .88 Widowed : 0.02 [-0.06, 0.09] 0.43 .666 Drink alcohol 1 or 2 days' week : 0.02 [-0.06, 0.09] 0.43 .666 Drink alcohol 3 or 4 days' week : 0.02 [-0.06, 0.09] 0.43 .666 Drink alcohol 5 or 6 days' week : 0.02 [-0.06, 0.09] 0.43 .666 Drink alcohol 5 or 6 days' week : 0.03 [-0.07, 0.11] 0.43 .666 Eat breakfast regularly : 0.04 [-0.04, 0.12] 0.96 .333 Drink alcohol 5 or 6 days' week : 0.03 [-0.07, 0.11] 0.43 .666 Eat breakfast regularly : 0.04 [-0.04, 0.19] 0.75 .098 Avoid fatty foods : BML : 0.00 [-0.01, 0.10] 1.75 .088 Avoid fatty foods : 0.03 [-0.07, 0.11] 1.65 .099 Drink alcohol 5 or 6 days' week ; 0.03 [-0.07, 0.11] 0.43 .666 Drink alcohol 5 or 6 days' week ; 0.03 [-0.07, 0.11] 0.43 .666 Drink alcohol 5 or 6 days' week ; 0.03 [-0.07, 0.11] 0.43 .666 Drink alcohol 5 or 6 days' week ; 0.04 [-0.00, 0.14] 0.76 .444 Nor-smoker : 0.08 [0.01, 0.15] 2.19 .022 Drink alcohol 5 or 6 days' week ; 0.06 [-0.01, 0.11] 2.29 .022 Drink alcohol 5 or 6 days' wee	Independent variable	β	t	р	
$\begin{aligned} & \text{Deg of household income}_1 & 0.03 [0.00, 0.07] & 1.78 & 0.77 \\ & \text{Age}_1 & 0.02 [-0.03, 0.01] & 3.16 & 0.00 \\ & \text{Age squared}_1 & 0.02 [-0.03, 0.01] & 3.16 & 0.00 \\ & \text{Age squared}_1 & 0.02 [-0.03, 0.01] & 3.16 & 0.00 \\ & \text{Male}_1 & 0.01 [-0.05, 0.06] & 0.20 & 8.43 \\ & \text{Masters of octorate}_1 & 0.13 [-0.27, 0.01] & 1.77 & 0.77 \\ & \text{Bachelor or honors}_1 & -0.08 [-0.20, 0.04] & 1.38 & 1.66 \\ & \text{Graduate diploma or certificate}_1 & -0.06 [-0.14, 0.03] & 1.21 & 222 \\ & \text{Advanced diploma}_1 & -0.06 [-0.14, 0.03] & 1.21 & 222 \\ & \text{Advanced diploma}_1 & -0.06 [-0.13, 0.02] & 1.47 & 1.44 \\ & \text{Year 12 high school}_1 & -0.06 [-0.13, 0.02] & 1.47 & 1.44 \\ & \text{Year 12 high school}_1 & -0.08 [-0.16, 0.00] & 1.86 & 0.66 \\ & \text{Full-time student}_1 & 0.12 [-0.01, 0.25] & 1.75 & 0.88 \\ & \text{Unemployed}_1 & 0.03 [-0.15, 0.21] & 0.32 & .748 \\ & \text{Married}_1 & 0.13 [0.05, 0.21] & 0.31 & 0.00 \\ & \text{Separated}_1 & -0.13 [0.05, 0.21] & 0.31 & 0.00 \\ & \text{Separated}_1 & -0.11 [-0.27, 0.05] & 1.36 & 1.75 \\ & \text{Divorced}_1 & 0.21 [-0.28, 0.15] & 6.28 & 0.001 \\ & \text{widowed}_1 & 0.22 [-0.08, 0.16] & 0.23 & .002 \\ & \text{Unemployed}_1 & 0.02 [-0.06, 0.09] & 0.43 & .666 \\ & \text{Drink alcohol 2 or 3 days/ week_1 & 0.02 [-0.03, 0.14] & 1.29 & .199 \\ & \text{Drink alcohol 2 or 3 days/ week_1 & 0.03 [-0.07, 0.13] & 0.63 & .522 \\ & \text{Drink alcohol 2 or 5 days/ week_1 & 0.03 [-0.07, 0.13] & 0.63 & .522 \\ & \text{Drink alcohol 2 or 6 days/ week_1 & 0.04 [-0.06, 0.14] & 0.76 & .444 \\ & \text{Non-smoker}_1 & 0.02 [-0.07, 0.13] & 0.63 & .522 \\ & Drink alcohol 2 or 7 days/ week_1 & 0.03 [-0.07, 0.13] & 0.63 & .522 \\ & \text{Drink alcohol 2 or 7 days/ week_1 & 0.03 [-0.07, 0.13] & 0.63 & .522 \\ & \text{Drink alcohol 2 or 7 days/ week_1 & 0.03 [-0.07, 0.13] & 0.63 & .522 \\ & \text{Drink alcohol 2 or 7 days/ week_1 & 0.02 [-0.07, 0.13] & 0.63 & .522 \\ & \text{Drink alcohol 2 or 7 days/ week_1 & 0.02 [-0.07, 0.11] & 0.43 & .666 \\ & \text{Et breakfast regularly_1 & 0.03 [-0.04, 0.09] & 0.85 & .397 \\ & \text{Drink alcohol 2 or 7 days/ week_1 & 0.02 [-0.07, 0.11] & 0.43 & .666 \\ $	Fruit and vegetable portions/day t	0.03 [0.01, 0.04]	3.82	.000	
Age : -0.02 [-0.03, -0.01] 3.16 .000 Age squared : 0.02 [0.01, 0.03] 3.87 .000 Male : 0.01 [-0.05, 0.06] 0.20 .844 Maters or doctorate : -0.13 [-0.27, 0.01] 1.77 .077 Bachelor or honors : -0.06 [-0.14, 0.03] 1.21 .222 Advanced diploma or certificate : -0.06 [-0.13, 0.02] 1.47 .144 Year 12 high school : -0.08 [-0.16, 0.00] 1.86 .066 Full-time student : 0.12 [-0.01, 0.25] 1.75 .084 Unemployed : 0.03 [-0.15, 0.21] 0.32 .744 Not in the labor force : -0.03 [-0.11, 0.05] 0.83 .400 Maried : 0.13 [0.05, 0.21] 3.11 .000 Separated : -0.01 [-0.27, 0.05] 1.36 .172 Divorced : -0	Life satisfaction t	0.48 [0.47, 0.50]	49.31	.000	
Age squared : $0.02 [0.01, 0.03]$ 3.87 0.00 Male : $0.01 [-0.05, 0.06]$ 0.20 8.44 Masters or doctorate : $-0.13 [-0.27, 0.01]$ 1.77 0.77 Bachelor or honrs : $-0.08 [-0.14, 0.03]$ 1.21 2.22 Advanced diploma or certificate : $-0.06 [-0.14, 0.03]$ 1.21 2.22 Advanced diploma i $-0.12 [-0.21, -0.02]$ 2.36 0.018 Professional qualification : $-0.06 [-0.13, 0.02]$ 1.47 1.44 Year 12 high school : $-0.08 [-0.16, 0.00]$ 1.86 0.66 Unemployed : $0.03 [-0.15, 0.21]$ 0.32 7.48 Not in the labor force : $-0.03 [-0.11, 0.05]$ 0.83 400 Married : $0.13 [0.05, 0.21]$ 3.11 0.001 Separated : $-0.01 [-0.25, 0.07]$ 0.34 7.33 Widowed : $0.02 [0.00, 0.41]$ 3.22 0.001 Long-term health condition : $-0.21 [-0.28, -0.15]$ 6.28 0.000 ℓ rikidren under the age of 4 : $0.01 [-0.05, 0.07]$ 0.34 7.33 Trink alcohol 2 or 3 days/ week : $0.03 [-0.07, 0.13]$ 0.65 0.99 Drink alcohol 1 or 2 days/ week : $0.03 [-0.07, 0.13]$ 0.63 5.29 Nort- art erd meat : $-0.03 [-0.07, 0.13]$ 0.63 5.29 Drink alcohol 2 or 3 days/ week : $0.03 [-0.07, 0.13]$ 0.63 5.29 Drink alcohol 2 or 3 days/ week : $0.03 [-0.07, 0.13]$ 0.63 5.29 Drink alcohol 2 or 3 days/ week :<	Log of household income t	0.03 [0.00, 0.07]	1.78	.075	
Male : 0.01 [-0.05, 0.06] 0.20 844 Masters or doctorate : -0.13 [-0.27, 0.01] 1.77 0.77 Bachelor or honors : -0.08 [-0.20, 0.04] 1.88 1.66 Graduate diploma or cerificate : -0.06 [-0.14, 0.03] 1.21 223 Advanced diploma : -0.06 [-0.14, 0.03] 1.21 223 Advanced diploma : -0.06 [-0.13, 0.02] 1.47 1.44 Year 12 high school : -0.08 [-0.16, 0.00] 1.86 0.66 Full-time student : 0.12 [-0.01, 0.25] 1.75 0.88 Mot in the labor force : -0.03 [-0.11, 0.05] 0.83 .400 Married : 0.13 [0.05, 0.21] 3.11 0.002 Separated : -0.01 [-0.12, 0.10] 0.15 88 Divorced : -0.01 [-0.0, 0.41] 3.22 0.00 Long-term health condition : -0.21 [-0.28, -0.15] 6.28 0.00 Unikalcohol 2 or 3 days/ week : 0.02 [-0.06, 0.09] 0.43 6.66 Drink alcohol 2 or 4 days/ week : 0.02 [-0.07, 0.13] 0.63 5.24 Drink alcohol 3 or 4 days/ week : 0.02 [-0.07, 0.13]	Age t	-0.02 [-0.03, -0.01]	3.16	.002	
Masters or doctorate , $-0.13 [-0.27, 0.01]$ 1.77 0.77 Bachelor or honors , $-0.08 [-0.27, 0.01]$ 1.77 0.77 Bachelor or honors , $-0.08 [-0.27, 0.01]$ 1.38 1.69 Graduate diploma , $-0.06 [-0.14, 0.03]$ 1.21 2.22 Advanced diploma , $-0.06 [-0.13, 0.02]$ 1.47 1.44 Year 12 high school , $-0.06 [-0.15, 0.00]$ 1.86 0.66 Full-time student , $0.12 [-0.01, 0.25]$ 1.75 0.88 Unemployed , $0.03 [-0.15, 0.21]$ 0.32 $.744$ Not in the labor force , $-0.03 [-0.11, 0.05]$ 0.36 $.177$ Divorced , $-0.11 [-0.27, 0.05]$ 1.36 $.177$ Divorced , $-0.01 [-0.12, 0.10]$ 0.15 883 Widowed , $0.26 [0.10, 0.41]$ 3.22 $.000$ Long-term health condition , $-0.21 [-0.28, 0.015]$ 6.28 $.000$ Dirik alcohol 1 or 2 days/ week , $0.02 [-0.07, 0.01]$ 1.65 $.099$ Dirik alcohol 2 or 3 days/ week , $0.02 [-0.07, 0.01]$ 0.63 $.524$ $.$	Age squared t	0.02 [0.01, 0.03]	3.87	.000	
Bachelor or homors : $-0.08 [-0.20, 0.04]$ 1.38 1.66 Graduate diploma or certificate : $-0.06 [-0.14, 0.03]$ 1.21 2.22 Advanced diploma : $-0.12 [-0.21, -0.02]$ 2.36 0.018 Professional qualification : $-0.06 [-0.13, 0.02]$ 1.47 1.44 Year 12 high school : $-0.08 [-0.16, 0.00]$ 1.86 0.66 Unemployed : $0.03 [-0.15, 0.21]$ 0.32 7.44 Not in the labor force : $-0.03 [-0.11, 0.05]$ 0.83 409 Married : $0.13 [0.05, 0.21]$ 3.11 0.00 Separated : $-0.01 [-0.12, 0.10]$ 1.56 1.75 Divorced : $-0.01 [-0.12, 0.10]$ 1.56 1.88 Widowed , $0.26 [0.10, 0.41]$ 3.22 0.00 Long-term health condition : $-0.21 [-0.28, -0.15]$ 6.28 0.00 # children under the age of 4 : $0.03 [-0.07, 0.01]$ 1.65 0.99 Drink alcohol 1 or 2 days/ week : $0.02 [-0.06, 0.09]$ 0.43 6.66 Drink alcohol 3 or 4 days/ week : $0.03 [-0.07, 0.13]$ 0.63 5.22 Drink alcohol 5 or 6	Male t	0.01 [-0.05, 0.06]	0.20	.845	
Graduate diploma or certificate : -0.06 [-0.14, 0.03] 1.21 .223 Advanced diploma : -0.12 [-0.21, -0.02] 2.36 .014 Professional qualification : -0.06 [-0.13, 0.02] 1.47 .144 Year 12 high school : -0.08 [-0.16, 0.00] 1.86 .066 Full-time student : 0.12 [-0.01, 0.25] 1.75 .086 Unemployed : 0.03 [-0.15, 0.21] 0.32 .744 Not in the labor force : -0.03 [-0.11, 0.05] 0.83 .409 Married : 0.13 [0.05, 0.21] 3.11 .002 Separated : -0.01 [-0.27, 0.05] 1.36 1.77 Divorced : 0.02 [0.10, 0.41] 3.22 .001 Long-term health condition : -0.21 [-0.28, -0.15] 6.28 .000 Long-term health condition : -0.21 [-0.28, -0.15] 6.28 .000 Drink alcohol 1 or 2 days/ week : 0.001 [-0.07, 0.01] 1.65 .099 Drink alcohol 2 or 3 days/ week : 0.04 [-0.04, 0.12] 0.96 .333 Drink alcohol 3 or 4 days/ week : 0.03 [-0.07, 0.13] 0.63 .525 <td< td=""><td>Masters or doctorate t</td><td>-0.13 [-0.27, 0.01]</td><td>1.77</td><td>.077</td></td<>	Masters or doctorate t	-0.13 [-0.27, 0.01]	1.77	.077	
Advanced diploma : $-0.12 [-0.21, -0.02]$ 2.36 0.14 Professional qualification : $-0.06 [-0.13, 0.02]$ 1.47 1.44 Year 12 high school : $-0.08 [-0.16, 0.00]$ 1.86 0.66 Full-time student : 0.12 [-0.01, 0.25] 1.75 0.88 Unemployed : $0.03 [-0.15, 0.21]$ 0.32 7.49 Married : 0.13 [0.05, 0.21] 3.11 0.00 Separated : $-0.01 [-0.12, 0.10]$ 0.15 88 Widowed : 0.26 [0.10, 0.41] 3.22 0.00 Long-term health condition : $-0.21 [-0.28, -0.15]$ 6.28 0.00 # children under the age of 4 : $0.01 [-0.05, 0.07]$ 0.34 7.33 # children under the age of 4 : $0.00 [-0.03, 0.14]$ 1.29 1.99 Dirik alcohol 1 or 2 days/ week : $0.00 [-0.03, 0.14]$ 1.29 1.99 Dirik alcohol 1 or 2 days/ week : $0.03 [-0.07, 0.13]$ 0.63 5.22 Drink alcohol 2 or 3 days/ week : $0.04 [-0.06, 0.14]$ 0.76 4.44 Non-smoker : $0.03 [-0.07, 0.11]$ 0.43 6.66 Drink alcohol 2 or 3 day	Bachelor or honors t	-0.08 [-0.20, 0.04]	1.38	.169	
Professional qualification : -0.06 [-0.13, 0.02] 1.47 1.44 Year 12 high school : -0.08 [-0.16, 0.00] 1.86 0.66 Full-time student : 0.12 [-0.01, 0.25] 1.75 0.88 Unemployed : 0.03 [-0.15, 0.21] 0.32 7.44 Not in the labor force : -0.03 [-0.11, 0.05] 0.83 409 Married : 0.13 [0.05, 0.21] 3.11 0.00 Separated : -0.01 [-0.27, 0.05] 1.36 1.75 Divorced : -0.01 [-0.12, 0.10] 0.15 883 Widowed : 0.26 [0.10, 0.41] 3.22 0.00 Long-term health condition : -0.21 [-0.28, -0.15] 6.28 0.00 # children under the age of 4 : 0.01 [-0.05, 0.07] 0.34 .733 # children aged 5-14 : -0.03 [-0.07, 0.01] 1.65 0.99 Drink alcohol 1 or 2 days/ week : 0.04 [-0.04, 0.12] 0.96 3.30 Drink alcohol 2 or 3 days/ week : 0.03 [-0.07, 0.13] 0.63 5.22 Drink alcohol 5 or 6 days/ week : 0.03 [-0.07, 0.13] 0.63 5.22 Drink alcohol everyday : 0.03 [-0.07,	Graduate diploma or certificate t	-0.06 [-0.14, 0.03]	1.21	.225	
Year 12 high school t -0.08 [-0.16, 0.00] 1.86 0.66 Full-time student t 0.12 [-0.01, 0.25] 1.75 0.80 Unemployed t 0.03 [-0.15, 0.21] 0.32 7.44 Not in the labor force t -0.03 [-0.11, 0.05] 0.83 4.00 Married t 0.13 [0.05, 0.21] 3.11 0.00 Separated t -0.01 [-0.27, 0.05] 1.36 1.75 Divorced t -0.01 [-0.12, 0.10] 0.15 8.83 Widowed t 0.26 [0.10, 0.41] 3.22 0.00 Long-term health condition t -0.21 [-0.28, -0.15] 6.28 0.00 # children under the age of 4 t 0.01 [-0.05, 0.07] 0.34 7.33 # children aged 5-14 t -0.03 [-0.07, 0.01] 1.65 0.99 Drink alcohol 1 or 2 days/ week t 0.02 [-0.06, 0.09] 0.43 665 Drink alcohol 2 or 3 days/ week t 0.03 [-0.07, 0.1] 0.63 5.25 Drink alcohol 3 or 4 days/ week t 0.03 [-0.07, 0.13] 0.63 5.25 Drink alcohol 5 or 6 days/ week t 0.03 [-0.07, 0.13] 0.63 5.29 Non-smoker t 0.03 [-0.07, 0.11] </td <td>Advanced diploma t</td> <td>-0.12 [-0.21, -0.02]</td> <td>2.36</td> <td>.018</td>	Advanced diploma t	-0.12 [-0.21, -0.02]	2.36	.018	
Full-time student i 0.12 [-0.01, 0.25] 1.75 0.88 Unemployed i 0.03 [-0.15, 0.21] 0.32 749 Not in the labor force i -0.03 [-0.11, 0.05] 0.83 409 Married i 0.13 [0.05, 0.21] 3.11 0.00 Separated i -0.01 [-0.27, 0.05] 1.36 1.75 Divorced i -0.01 [-0.12, 0.10] 0.15 883 Widowed i 0.26 [0.10, 0.41] 3.22 0.00 Long-term health condition i -0.21 [-0.28, 0.15] 6.28 0.00 # children under the age of 4_i 0.01 [-0.05, 0.07] 0.34 7.33 # children under the age of 4_i 0.02 [-0.06, 0.09] 0.43 663 Drink alcohol 1 or 2 days/ week i 0.04 [-0.04, 0.12] 0.96 339 Drink alcohol 2 or 3 days/ week i 0.03 [-0.07, 0.13] 0.63 529 Drink alcohol 5 or 6 days/ week i 0.04 [-0.04, 0.12] 0.96 339 Drink alcohol 2 veryday i 0.03 [-0.07, 0.11] 0.43 663 Drink alcohol 5 or 6 days/ week i 0.03 [-0.07, 0.11] 0.43 663 Drink alcohol e	Professional qualification t	-0.06 [-0.13, 0.02]	1.47	.142	
Unemployed $_1$ 0.03 [-0.15, 0.21] 0.32 .744 Not in the labor force $_1$ -0.03 [-0.11, 0.05] 0.83 .409 Married $_1$ 0.13 [0.05, 0.21] 3.11 .000 Separated $_1$ -0.01 [-0.27, 0.05] 1.36 .173 Divorced $_1$ -0.01 [-0.12, 0.10] 0.15 .883 Widowed $_1$ 0.26 [0.10, 0.41] 3.22 .000 Long-term health condition $_1$ -0.21 [-0.28, -0.15] 6.28 .000 # children under the age of 4 $_1$ -0.03 [-0.07, 0.01] 1.65 .099 Drink alcohol 1 or 2 days/ week $_1$ -0.03 [-0.07, 0.01] 1.65 .099 Drink alcohol 2 or 3 days/ week $_1$ 0.06 [-0.03, 0.14] 1.29 .190 Drink alcohol 3 or 4 days/ week $_1$ 0.03 [-0.07, 0.13] 0.63 .522 Drink alcohol 2 or 3 days/ week $_1$ 0.04 [-0.06, 0.14] 0.76 .444 Non-smoker $_1$ 0.04 [-0.06, 0.14] 0.76 .444 Non-smoker $_1$ 0.03 [-0.07, 0.11] 0.43 .665 Drink alcohol 2 or 3 days/ week $_1$ 0.03 [-0.07, 0.11] 0.43 .666 Drink	Year 12 high school t	-0.08 [-0.16, 0.00]	1.86	.063	
Not in the labor force $_1$ -0.03 [-0.11, 0.05] 0.83 .409 Married $_1$ 0.13 [0.05, 0.21] 3.11 .000 Separated $_1$ -0.01 [-0.27, 0.05] 1.36 .172 Divorced $_1$ -0.01 [-0.12, 0.10] 0.15 .881 Widowed $_1$ 0.26 [0.10, 0.41] 3.22 .001 Long-term health condition $_1$ -0.21 [-0.28, -0.15] 6.28 .000 # children under the age of 4 $_1$ 0.01 [-0.05, 0.07] 0.34 .732 # children under the age of 4 $_1$ -0.03 [-0.07, 0.01] 1.65 .099 Drink alcohol 1 or 2 days/ week $_1$ 0.02 [-0.06, 0.09] 0.43 .662 Drink alcohol 2 or 3 days/ week $_1$ 0.04 [-0.04, 0.12] 0.96 .330 Drink alcohol 3 or 4 days/ week $_1$ 0.03 [-0.07, 0.13] 0.63 .529 Drink alcohol 2 veryda $_1$ 0.04 [-0.04, 0.12] 0.96 .330 Drink alcohol 2 veryda $_1$ 0.04 [-0.07, 0.11] 0.63 .529 Drink alcohol 2 veryda $_1$ 0.03 [-0.07, 0.11] 0.63 .529 Drink alcohol 2 veryda $_1$ 0.03 [-0.07, 0.11] 0.66 .529	Full-time student t	0.12 [-0.01, 0.25]	1.75	.080	
Married ι 0.13 [0.05, 0.21] 3.11 .000 Separated ι -0.11 [-0.27, 0.05] 1.36 .172 Divorced ι -0.01 [-0.12, 0.10] 0.15 .883 Widowed ι 0.26 [0.10, 0.41] 3.22 .001 Long-term health condition ι -0.21 [-0.28, -0.15] 6.28 .000 # children under the age of 4 ι 0.01 [-0.05, 0.07] 0.34 .732 # children aged 5-14 ι -0.03 [-0.07, 0.01] 1.65 .099 Drink alcohol 1 or 2 days/ week ι 0.02 [-0.66, 0.09] 0.43 .662 Drink alcohol 2 or 3 days/ week ι 0.04 [-0.04, 0.12] 0.96 .333 Drink alcohol 5 or 6 days/ week ι 0.03 [-0.07, 0.13] 0.63 .529 Drink alcohol everyday ι 0.04 [-0.06, 0.14] 0.76 .444 Non-smoker ι 0.08 [0.01, 0.15] 2.19 .029 Never eat red meat ι -0.13 [-0.28, 0.03] 1.54 .122 Never eat fish ι 0.03 [-0.04, 0.09] 0.85 .397 Drink low fat or skinny milk ι 0.05 [-0.01, 0.11] 1.65 .098 BMI ι -0	Unemployed t	0.03 [-0.15, 0.21]	0.32	.749	
Separated i -0.11 [-0.27, 0.05] 1.36 .175 Divorced i -0.01 [-0.12, 0.10] 0.15 8.88 Widowed i 0.26 [0.10, 0.41] 3.22 0.00 Long-term health condition i -0.21 [-0.28, -0.15] 6.28 0.00 # children under the age of 4 i 0.01 [-0.05, 0.07] 0.34 .733 # children aged 5-14 i -0.03 [-0.07, 0.01] 1.65 0.99 Drink alcohol 1 or 2 days/ week i 0.02 [-0.06, 0.09] 0.43 .663 Drink alcohol 2 or 3 days/ week i 0.04 [-0.04, 0.12] 0.96 .336 Drink alcohol 5 or 6 days/ week i 0.03 [-0.07, 0.13] 0.63 .522 Drink alcohol everyday i 0.04 [-0.06, 0.14] 0.76 .444 Non-smoker i 0.08 [0.01, 0.15] 2.19 .022 Never eat red meat i -0.13 [-0.28, 0.03] 1.54 .122 Never eat fish i 0.02 [-0.07, 0.11] 0.43 .665 Eat breakfast regularly i 0.03 [-0.04, 0.09] 0.85 .397 Drink low fat or skinny milk i 0.05 [-0.01, 0.10] 1.75 .086 Avoid fa	Not in the labor force t	-0.03 [-0.11, 0.05]	0.83	.409	
Divorced ι -0.01 [-0.12, 0.10]0.15.883Widowed ι 0.26 [0.10, 0.41]3.22.000Long-term health condition ι -0.21 [-0.28, -0.15]6.28.000# children under the age of 4 ι 0.01 [-0.05, 0.07]0.34.733# children aged 5-14 ι -0.03 [-0.07, 0.01]1.65.099Drink alcohol 1 or 2 days/ week ι 0.02 [-0.06, 0.09]0.43.663Drink alcohol 2 or 3 days/ week ι 0.04 [-0.04, 0.12]0.96.336Drink alcohol 3 or 4 days/ week ι 0.03 [-0.07, 0.13]0.63.529Drink alcohol 5 or 6 days/ week ι 0.03 [-0.07, 0.13]0.63.529Drink alcohol everyday ι 0.04 [-0.06, 0.14]0.76.444Non-smoker ι 0.08 [0.01, 0.15]2.19.029Never eat red meat ι -0.13 [-0.28, 0.03]1.54.122Never eat fish ι 0.05 [-0.01, 0.10]1.75.086Avoid fatty foods ι 0.05 [-0.01, 0.10]1.75.086BMI ι -0.01 [-0.01, 0.00]2.79.002Constant3.98 [3.55, 4.41]18.34.000	Married t	0.13 [0.05, 0.21]	3.11	.002	
Widowed ι 0.26 [0.10, 0.41] 3.22 .00 Long-term health condition ι -0.21 [-0.28, -0.15] 6.28 .000 # children under the age of 4 ι 0.01 [-0.05, 0.07] 0.34 .732 # children aged 5-14 ι -0.03 [-0.07, 0.01] 1.65 .099 Drink alcohol 1 or 2 days/ week ι 0.02 [-0.06, 0.09] 0.43 .663 Drink alcohol 2 or 3 days/ week ι 0.04 [-0.04, 0.12] 0.96 .336 Drink alcohol 3 or 4 days/ week ι 0.03 [-0.07, 0.13] 0.63 .529 Drink alcohol 5 or 6 days/ week ι 0.03 [-0.07, 0.13] 0.63 .529 Drink alcohol 5 or 6 days/ week ι 0.03 [-0.07, 0.13] 0.63 .529 Drink alcohol 9 or 4 days/ week ι 0.03 [-0.07, 0.13] 0.63 .529 Drink alcohol 9 or 6 days/ week ι 0.03 [-0.07, 0.13] 0.63 .529 Drink alcohol 9 veryday ι 0.04 [-0.06, 0.14] 0.76 .444 Non-smoker ι 0.03 [-0.07, 0.11] 0.43 .665 Never eat red meat ι 0.03 [-0.04, 0.09] 0.85 .397 Drink low fat or skinny milk ι 0.05 [-0.01, 0.10] <td< td=""><td>Separated t</td><td>-0.11 [-0.27, 0.05]</td><td>1.36</td><td>.175</td></td<>	Separated t	-0.11 [-0.27, 0.05]	1.36	.175	
Long-term health condition t -0.21 [-0.28, -0.15] 6.28 .000 # children under the age of 4 t 0.01 [-0.05, 0.07] 0.34 .732 # children aged 5-14 t -0.03 [-0.07, 0.01] 1.65 .099 Drink alcohol 1 or 2 days/ week t 0.02 [-0.06, 0.09] 0.43 .663 Drink alcohol 2 or 3 days/ week t 0.04 [-0.04, 0.12] 0.96 .336 Drink alcohol 5 or 6 days/ week t 0.03 [-0.07, 0.13] 0.63 .529 Drink alcohol 5 or 6 days/ week t 0.04 [-0.06, 0.14] 0.76 .448 Non-smoker t 0.08 [0.01, 0.15] 2.19 .029 Never eat red meat t 0.03 [-0.07, 0.11] 0.43 .665 Eat breakfast regularly t 0.04 [-0.06, 0.14] 0.76 .448 Non-smoker t 0.08 [0.01, 0.15] 2.19 .029 Never eat red meat t 0.03 [-0.07, 0.11] 0.43 .665 Eat breakfast regularly t 0.03 [-0.04, 0.09] 0.85 .397 Drink low fat or skinny milk t 0.05 [-0.01, 0.10] 1.75 .088 Avoid fatty foods t 0.05 [-0.01, 0.01] 2.29 .022	Divorced t	-0.01 [-0.12, 0.10]	0.15	.881	
# children under the age of 4 t $0.01 [-0.05, 0.07]$ 0.34 $.732$ # children aged 5-14 t $-0.03 [-0.07, 0.01]$ 1.65 $.099$ Drink alcohol 1 or 2 days/ week t $0.02 [-0.06, 0.09]$ 0.43 $.663$ Drink alcohol 2 or 3 days/ week t $0.06 [-0.03, 0.14]$ 1.29 $.196$ Drink alcohol 3 or 4 days/ week t $0.04 [-0.04, 0.12]$ 0.96 $.336$ Drink alcohol 5 or 6 days/ week t $0.03 [-0.07, 0.13]$ 0.63 $.529$ Drink alcohol 2 or 3 days/ week t $0.04 [-0.04, 0.12]$ 0.96 $.336$ Drink alcohol 5 or 6 days/ week t $0.03 [-0.07, 0.13]$ 0.63 $.529$ Drink alcohol everyday t $0.04 [-0.06, 0.14]$ 0.76 $.448$ Non-smoker t $0.08 [0.01, 0.15]$ 2.19 0.029 Never eat red meat t $-0.13 [-0.28, 0.03]$ 1.54 $.122$ Never eat fish t $0.02 [-0.07, 0.11]$ 0.43 $.666$ Eat breakfast regularly t $0.03 [-0.04, 0.09]$ 0.85 $.397$ Drink low fat or skinny milk t $0.05 [-0.01, 0.10]$ 1.75 $.086$ Avoid fatty foods t	Widowed t	0.26 [0.10, 0.41]	3.22	.001	
# children aged 5-14 t $-0.03 [-0.07, 0.01]$ 1.65 .099 Drink alcohol 1 or 2 days/ week t $0.02 [-0.06, 0.09]$ 0.43 .663 Drink alcohol 2 or 3 days/ week t $0.06 [-0.03, 0.14]$ 1.29 .196 Drink alcohol 3 or 4 days/ week t $0.04 [-0.04, 0.12]$ 0.96 .336 Drink alcohol 5 or 6 days/ week t $0.03 [-0.07, 0.13]$ 0.63 .529 Drink alcohol everyday t $0.04 [-0.06, 0.14]$ 0.76 .444 Non-smoker t $0.08 [0.01, 0.15]$ 2.19 .029 Never eat red meat t $-0.13 [-0.28, 0.03]$ 1.54 .123 Never eat fish t $0.02 [-0.07, 0.11]$ 0.43 .666 Eat breakfast regularly t $0.03 [-0.04, 0.09]$ 0.85 .397 Drink low fat or skinny milk t $0.05 [-0.01, 0.10]$ 1.75 .086 Avoid fatty foods t $0.05 [-0.01, 0.10]$ 1.75 .086 BMI t $-0.01 [-0.01, 0.00]$ 2.79 .007 Exercise regularly t $0.06 [0.01, 0.11]$ 2.29 .027 Constant $3.98 [3.55, 4.41]$ 18.34 .000	Long-term health condition t	-0.21 [-0.28, -0.15]	6.28	.000	
Drink alcohol 1 or 2 days/ week t 0.02 [-0.06, 0.09] 0.43 .663Drink alcohol 2 or 3 days/ week t 0.06 [-0.03, 0.14] 1.29 .196Drink alcohol 3 or 4 days/ week t 0.04 [-0.04, 0.12] 0.96 .336Drink alcohol 5 or 6 days/ week t 0.03 [-0.07, 0.13] 0.63 .529Drink alcohol everyday t 0.04 [-0.06, 0.14] 0.76 .448Non-smoker t 0.08 [0.01, 0.15] 2.19 0.029 Never eat red meat t -0.13 [-0.28, 0.03] 1.54 .123Never eat fish t 0.02 [-0.07, 0.11] 0.43 .666Eat breakfast regularly t 0.03 [-0.04, 0.09] 0.85 .397Drink low fat or skinny milk t 0.05 [-0.01, 0.10] 1.75 .086Avoid fatty foods t 0.05 [-0.01, 0.11] 1.65 .098BMI t -0.01 [-0.01, 0.00] 2.79 .002Exercise regularly t 0.06 [0.01, 0.11] 2.29 .022Adjusted R^2 .31.31.31	# children under the age of 4 $_{\rm t}$	0.01 [-0.05, 0.07]	0.34	.732	
Drink alcohol 2 or 3 days/ week t $0.06 [-0.03, 0.14]$ 1.29 $.190$ Drink alcohol 3 or 4 days/ week t $0.04 [-0.04, 0.12]$ 0.96 $.336$ Drink alcohol 5 or 6 days/ week t $0.03 [-0.07, 0.13]$ 0.63 $.529$ Drink alcohol everyday t $0.04 [-0.06, 0.14]$ 0.76 $.448$ Non-smoker t $0.08 [0.01, 0.15]$ 2.19 $.029$ Never eat red meat t $-0.13 [-0.28, 0.03]$ 1.54 $.122$ Never eat fish t $0.02 [-0.07, 0.11]$ 0.43 $.662$ Eat breakfast regularly t $0.03 [-0.04, 0.09]$ 0.85 $.397$ Drink low fat or skinny milk t $0.05 [-0.01, 0.10]$ 1.75 $.086$ Avoid fatty foods t $0.05 [-0.01, 0.11]$ 1.65 $.098$ BMI t $-0.01 [-0.01, 0.00]$ 2.79 $.002$ Constant $3.98 [3.55, 4.41]$ 18.34 $.000$	# children aged 5-14 t	-0.03 [-0.07, 0.01]	1.65	.099	
Drink alcohol 3 or 4 days/ week t $0.04 [-0.04, 0.12]$ 0.96 $.336$ Drink alcohol 5 or 6 days/ week t $0.03 [-0.07, 0.13]$ 0.63 $.529$ Drink alcohol everyday t $0.04 [-0.06, 0.14]$ 0.76 $.448$ Non-smoker t $0.08 [0.01, 0.15]$ 2.19 0.029 Never eat red meat t $-0.13 [-0.28, 0.03]$ 1.54 $.123$ Never eat fish t $0.02 [-0.07, 0.11]$ 0.43 $.663$ Eat breakfast regularly t $0.03 [-0.04, 0.09]$ 0.85 $.397$ Drink low fat or skinny milk t $0.05 [-0.01, 0.10]$ 1.75 $.086$ Avoid fatty foods t $0.05 [-0.01, 0.10]$ 1.75 $.098$ BMI t $-0.01 [-0.01, 0.00]$ 2.79 $.002$ Constant $3.98 [3.55, 4.41]$ 18.34 $.000$	Drink alcohol 1 or 2 days/ week t	0.02 [-0.06, 0.09]	0.43	.665	
Drink alcohol 5 or 6 days/ week t $0.03 [-0.07, 0.13]$ 0.63 $.529$ Drink alcohol everyday t $0.04 [-0.06, 0.14]$ 0.76 $.448$ Non-smoker t $0.08 [0.01, 0.15]$ 2.19 $.029$ Never eat red meat t $-0.13 [-0.28, 0.03]$ 1.54 $.123$ Never eat fish t $0.02 [-0.07, 0.11]$ 0.43 $.669$ Eat breakfast regularly t $0.03 [-0.04, 0.09]$ 0.85 $.397$ Drink low fat or skinny milk t $0.05 [-0.01, 0.10]$ 1.75 $.086$ Avoid fatty foods t $0.05 [-0.01, 0.10]$ 1.65 $.098$ BMI t $-0.01 [-0.01, 0.00]$ 2.79 $.002$ Constant $3.98 [3.55, 4.41]$ 18.34 $.000$	Drink alcohol 2 or 3 days/ week t	0.06 [-0.03, 0.14]	1.29	.196	
Drink alcohol everyday t 0.04 [- 0.06 , 0.14] 0.76 $.448$ Non-smoker t 0.08 [0.01 , 0.15] 2.19 $.029$ Never eat red meat t -0.13 [-0.28 , 0.03] 1.54 $.123$ Never eat fish t 0.02 [-0.07 , 0.11] 0.43 $.665$ Eat breakfast regularly t 0.03 [-0.04 , 0.09] 0.85 $.397$ Drink low fat or skinny milk t 0.05 [-0.01 , 0.10] 1.75 $.086$ Avoid fatty foods t 0.05 [-0.01 , 0.01] 1.65 $.098$ BMI t -0.01 [-0.01 , 0.00] 2.79 $.002$ Constant 3.98 [3.55 , 4.41] 18.34 $.000$	Drink alcohol 3 or 4 days/ week t	0.04 [-0.04, 0.12]	0.96	.336	
Non-smoker t $0.08 [0.01, 0.15]$ 2.19 $.029$ Never eat red meat t $-0.13 [-0.28, 0.03]$ 1.54 $.123$ Never eat fish t $0.02 [-0.07, 0.11]$ 0.43 $.663$ Eat breakfast regularly t $0.03 [-0.04, 0.09]$ 0.85 $.397$ Drink low fat or skinny milk t $0.05 [-0.01, 0.10]$ 1.75 $.086$ Avoid fatty foods t $0.05 [-0.01, 0.10]$ 1.75 $.098$ BMI t $-0.01 [-0.01, 0.00]$ 2.79 $.002$ Exercise regularly t $0.06 [0.01, 0.11]$ 2.29 $.022$ Constant $3.98 [3.55, 4.41]$ 18.34 $.000$	Drink alcohol 5 or 6 days/ week t	0.03 [-0.07, 0.13]	0.63	.529	
Never eat red meat t -0.13 [-0.28, 0.03] 1.54 .123 Never eat fish t 0.02 [-0.07, 0.11] 0.43 .665 Eat breakfast regularly t 0.03 [-0.04, 0.09] 0.85 .397 Drink low fat or skinny milk t 0.05 [-0.01, 0.10] 1.75 .086 Avoid fatty foods t 0.05 [-0.01, 0.10] 1.65 .098 BMI t -0.01 [-0.01, 0.00] 2.79 .005 Exercise regularly t 0.06 [0.01, 0.11] 2.29 .022 Constant 3.98 [3.55, 4.41] 18.34 .000	Drink alcohol everyday t	0.04 [-0.06, 0.14]	0.76	.448	
Never eat fish t $0.02 [-0.07, 0.11]$ 0.43 .665 Eat breakfast regularly t $0.03 [-0.04, 0.09]$ 0.85 .397 Drink low fat or skinny milk t $0.05 [-0.01, 0.10]$ 1.75 .080 Avoid fatty foods t $0.05 [-0.01, 0.10]$ 1.65 .098 BMI t $-0.01 [-0.01, 0.00]$ 2.79 .005 Exercise regularly t $0.06 [0.01, 0.11]$ 2.29 .022 Constant $3.98 [3.55, 4.41]$ 18.34 .000	Non-smoker t	0.08 [0.01, 0.15]	2.19	.029	
Eat breakfast regularly t $0.03 [-0.04, 0.09]$ $0.85 .392$ Drink low fat or skinny milk t $0.05 [-0.01, 0.10]$ $1.75 .080$ Avoid fatty foods t $0.05 [-0.01, 0.11]$ $1.65 .098$ BMI t $-0.01 [-0.01, 0.00]$ $2.79 .005$ Exercise regularly t $0.06 [0.01, 0.11]$ $2.29 .022$ Constant $3.98 [3.55, 4.41]$ $18.34 .000$	Never eat red meat t	-0.13 [-0.28, 0.03]	1.54	.123	
Drink low fat or skinny milk t $0.05 [-0.01, 0.10]$ 1.75 .080 Avoid fatty foods t $0.05 [-0.01, 0.10]$ 1.65 .098 BMI t $-0.01 [-0.01, 0.00]$ 2.79 .009 Exercise regularly t $0.06 [0.01, 0.11]$ 2.29 .022 Constant $3.98 [3.55, 4.41]$ 18.34 .000	Never eat fish t	0.02 [-0.07, 0.11]	0.43	.665	
Drink low fat or skinny milk t $0.05 [-0.01, 0.10]$ 1.75 .080 Avoid fatty foods t $0.05 [-0.01, 0.11]$ 1.65 .098 BMI t $-0.01 [-0.01, 0.00]$ 2.79 .005 Exercise regularly t $0.06 [0.01, 0.11]$ 2.29 .022 Constant $3.98 [3.55, 4.41]$ 18.34 .000	Eat breakfast regularly t		0.85	.397	
BMI t $-0.01 [-0.01, 0.00]$ 2.79 $.005$ Exercise regularly t $0.06 [0.01, 0.11]$ 2.29 $.022$ Constant $3.98 [3.55, 4.41]$ 18.34 $.000$ Adjusted R^2 $.31$	Drink low fat or skinny milk t		1.75	.080	
BMI t -0.01 [-0.01, 0.00] 2.79 .005 Exercise regularly t 0.06 [0.01, 0.11] 2.29 .022 Constant 3.98 [3.55 , 4.41] 18.34 .000 Adjusted R^2 .31	Avoid fatty foods t	0.05 [-0.01, 0.11]	1.65	.098	
Exercise regularly t $0.06 [0.01, 0.11]$ 2.29 $.022$ Constant $3.98 [3.55, 4.41]$ 18.34 $.000$ Adjusted R^2 $.31$	BMI t		2.79	.005	
Constant 3.98 [3.55, 4.41] 18.34 .000 Adjusted R ² .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 .31 <t< td=""><td>Exercise regularly t</td><td></td><td>2.29</td><td>.022</td></t<>	Exercise regularly t		2.29	.022	
				.000	
	Adjusted <i>R</i> ²	31			
	Number of observations	7,742			

Note: Values in parentheses are 95% confidence intervals. Dependent variable is *Life Satisfaction* [range: 0-10] in period t+1 (year 2009).

524 525 526 Table S3 (Full Estimation Results for Second Part of Table 3). Prospective Analysis of Happiness: Linear Regression Model of 'Been a Happy Person' on Lagged Fruit and Vegetable Consumption and Covariates, HILDA Survey 2007 (period t) and 2009 (period t+1)

527

Independent variable	β	t	р
Fruit and vegetable portions/day t	0.02 [0.01, 0.03]	3.97	.000
Been a happy person t	0.45 [0.43, 0.47]	44.26	.000
Log of household income t	0.03 [0.00, 0.05]	1.84	.066
Age t	-0.01 [-0.02, 0.00]	2.93	.003
Age squared t	0.01 [0.00, 0.02]	2.87	.004
Male t	0.01 [-0.04, 0.05]	0.23	.822
Masters or doctorate t	-0.01 [-0.12, 0.10]	0.21	.833
Bachelor or honors t	-0.03 [-0.13, 0.06]	0.65	.514
Graduate diploma or certificate t	-0.02 [-0.09, 0.05]	0.57	.569
Advanced diploma t	0.00 [-0.08, 0.07]	0.08	.936
Professional qualification t	-0.04 [-0.10, 0.02]	1.35	.176
Year 12 high school t	-0.02 [-0.09, 0.05]	0.58	.560
Full-time student t	-0.02 [-0.12, 0.08]	0.32	.745
Unemployed t	-0.21 [-0.35, -0.07]	2.87	.004
Not in the labor force t	-0.02 [-0.08, 0.04]	0.56	.572
Married t	0.10 [0.03, 0.16]	2.87	.004
Separated t	0.09 [-0.03, 0.22]	1.41	.157
Divorced t	0.09 [0.00, 0.18]	1.96	.050
Widowed t	0.31 [0.19, 0.43]	4.97	.000
Long-term health condition t	-0.24 [-0.29, -0.19]	8.96	.000
# children under the age of 4 $_{\rm t}$	-0.03 [-0.08, 0.01]	1.39	.165
# children aged 5-14 t	-0.03 [-0.06, 0.00]	1.93	.054
Drink alcohol 1 or 2 days/ week t	0.11 [0.05, 0.17]	3.71	.000
Drink alcohol 2 or 3 days/ week t	0.04 [-0.02, 0.11]	1.27	.206
Drink alcohol 3 or 4 days/ week t	0.06 [0.00, 0.12]	1.83	.067
Drink alcohol 5 or 6 days/ week t	0.13 [0.05, 0.20]	3.25	.00
Drink alcohol everyday t	0.03 [-0.05, 0.11]	0.72	.473
Non-smoker t	0.03 [-0.02, 0.09]	1.23	.217
Never eat red meat t	-0.01 [-0.13, 0.12]	0.13	.899
Never eat fish t	-0.03 [-0.10, 0.04]	0.77	.44
Eat breakfast regularly t	0.02 [-0.03, 0.07]	0.65	.510
Drink low fat or skinny milk t	-0.01 [-0.05, 0.03]	0.52	.604
Avoid fatty foods t	0.03 [-0.02, 0.08]	1.08	.279
BMI t	0.00 [-0.01, 0.00]	1.78	.074
Exercise regularly t	0.06 [0.01, 0.10]	2.66	.008
Constant	2.36 [2.04, 2.68]	14.40	.000
Adjusted R ²	.26		
Number of observations	7,694		

Note: Values in parentheses are 95% confidence intervals. Dependent variable is Been a Happy Person [range: 1-6] in period t+1 (year 2009).

531	Table S4. Granger Causality Test: Linear Regression Model of Fruit and Vegetable Consumption on
532	Lagged Life Satisfaction and Covariates, HILDA Survey 2007 (period t) and 2009 (period t+1)

Dependent variable: Fruit and vegetable consumption t+1			
Independent variable	β	t	р
Life satisfaction t	-0.003 [-0.03, 0.02]	0.22	.827
Fruit and vegetable portions/day t	0.55 [0.53, 0.57]	57.23	.000
Log of household income t	0.01 [-0.04, 0.06]	0.33	.739
Age t	0.02 [0.01, 0.04]	3.11	.002
Age squared t	-0.01 [-0.03, 0.00]	1.37	.170
Male t	-0.16 [-0.24, -0.09]	4.16	.000
Masters or doctorate t	0.20 [0.01, 0.39]	2.07	.038
Bachelor or honors t	0.29 [0.13, 0.46]	3.54	.000
Graduate diploma or certificate t	0.19 [0.07, 0.31]	3.02	.003
Advanced diploma t	0.19 [0.06, 0.32]	2.79	.005
Professional qualification t	0.15 [0.05, 0.25]	2.87	.004
Year 12 high school t	0.12 [0.00, 0.23]	2.02	.043
Full-time student t	0.27 [0.10, 0.45]	3.06	.002
Unemployed t	0.01 [-0.23, 0.26]	0.08	.934
Not in the labor force t	-0.03 [-0.13, 0.08]	0.54	.591
Married t	0.04 [-0.07, 0.15]	0.67	.500
Separated t	-0.18 [-0.40, 0.04]	1.61	.107
Divorced t	-0.10 [-0.25, 0.06]	1.24	.216
Widowed t	-0.12 [-0.33, 0.09]	1.13	.259
Long-term health condition t	0.02 [-0.07, 0.11]	0.52	.605
# children under the age of 4 t	-0.04 [-0.11, 0.04]	0.92	.360
# children aged 5-14 t	-0.02 [-0.07, 0.04]	0.61	.541
Drink alcohol 1 or 2 days/week t	-0.02 [-0.12, 0.09]	0.29	.769
Drink alcohol 2 or 3 days/week t	-0.02 [-0.14, 0.10]	0.29	.772
Drink alcohol 3 or 4 days/week t	0.01 [-0.10, 0.12]	0.25	.806
Drink alcohol 5 or 6 days/week t	0.05 [-0.08, 0.18]	0.70	.484
Drink alcohol everyday t	0.03 [-0.11, 0.17]	0.42	.678
Non-smoker t	0.25 [0.16, 0.35]	5.20	.000
Never eat red meat t	0.09 [-0.13, 0.30]	0.81	.419
Never eat fish t	-0.19 [-0.31, -0.07]	3.10	.002
Eat breakfast regularly t	0.19 [0.10, 0.27]	4.31	.000
Drink low fat or skinny milk t	0.01 [-0.06, 0.08]	0.31	.758
Avoid fatty foods t	0.15 [0.06, 0.23]	3.48	.001
BMI t	0.00 [-0.01, 0.01]	0.06	.951
Exercise regularly t	0.21 [0.13, 0.28]	5.63	.000
Constant	0.40 [-0.18, 0.97]	1.35	.177
Adjusted R ²	.4	2	
Number of observations	7,7		

Note: Values in parentheses are 95% confidence intervals. Dependent variable is *Fruit and Vegetable Consumption* (portions per day) in period t+1 (year 2009). It should be noted that Granger causality examines how an outcome variable of interest is correlated with lagged values of the same variable (from previous periods) as well as lagged values of other explanatory variables. This method is analogous to prospective analysis, but is not equivalent to identifying the true causal effect of one variable on another (where, for example, a change in the variable *X* strictly leads to a change in the variable *Y*).

		2

Dependent variable: Fruit and vegetable consumption t+1 β Independent variable t p0.03 [-0.01, 0.06] 1.63 .104 Been a happy person t Fruit and vegetable portions/day t 56.72 .000 0.55 [0.53, 0.57] Log of household income t 0.22 .826 0.01 [-0.04, 0.05] 3.29 .001 Age t 0.02 [0.01, 0.04] 1.61 .108 Age squared t -0.01 [-0.03, 0.00] Male t .000 -0.17 [-0.24, -0.09] 4.20 Masters or doctorate t 2.17 .030 0.21 [0.02, 0.40] Bachelor or honors t 3.66 .000 0.30 [0.14, 0.47] Graduate diploma or certificate t 3.18 .001 0.20 [0.08, 0.32] Advanced diploma t 2.81 .005 0.19 [0.06, 0.32] Professional qualification t 2.99 .003 0.16 [0.05, 0.26] Year 12 high school t 2.07 .039 0.12 [0.01, 0.24] 3.06 .002 Full-time student t 0.27 [0.10, 0.45] Unemployed t 0.03 [-0.22, 0.27] 0.23 .822 Not in the labor force t -0.02 [-0.13, 0.08] 0.46 .644 Married t 0.05 [-0.06, 0.16] 0.88 .379 Separated t -0.17 [-0.39, 0.05] 1.53 .126 Divorced t -0.09 [-0.24, 0.06] 1.17 .242 Widowed t -0.10 [-0.31, 0.11] 0.95 .344 Long-term health condition t 0.04 [-0.05, 0.14] 0.97 .333 # children under the age of 4 t -0.04 [-0.12, 0.04] 0.98 .325 # children aged 5-14 t -0.02 [-0.07, 0.04] 0.62 .534 Drink alcohol 1 or 2 days/ week t -0.01 [-0.12, 0.09] 0.27 .787 Drink alcohol 2 or 3 days/week t 0.21 .831 -0.01 [-0.13, 0.11] Drink alcohol 3 or 4 days/ week t 0.19 .849 0.01 [-0.10, 0.12] Drink alcohol 5 or 6 days/ week t 0.62 .533 0.04 [-0.09, 0.17] Drink alcohol everyday t 0.33 .740 0.02 [-0.11, 0.16] 4.97 .000 Non-smoker t 0.24 [0.15, 0.34] Never eat red meat t 0.10 [-0.12, 0.31] 0.87 .386 Never eat fish t 3.27 .001 -0.20 [-0.32, -0.08] Eat breakfast regularly t 4.19 .000 0.18 [0.10, 0.27] Drink low fat or skinny milk t 0.25 .801 0.01 [-0.06, 0.08] Avoid fatty foods t 3.49 .000 0.15 [0.07, 0.23] BMI t 0.05 .957 0.00 [-0.01, 0.01] 5.50 .000 Exercise regularly t 0.20 [0.13, 0.28] 0.27 [-0.29, 0.82] 0.93 Constant .350 Adjusted R^2 .42 7,694 Number of observations

540Table S5. Granger Causality Test: Linear Regression Model of Fruit and Vegetable Consumption on541Lagged 'Been a Happy Person' and Covariates, HILDA Survey 2007 (period t) and 2009 (period t+1)

Note: Values in parentheses are 95% confidence intervals. Dependent variable is *Fruit and Vegetable Consumption* (portions per day) in period t+1 (year 2009). It should be noted that Granger causality examines how an outcome variable of interest is correlated with lagged values of the same variable (from previous periods) as well as lagged values of other explanatory variables. This method is analogous to prospective analysis, but is not equivalent to identifying the true causal effect of one variable on another (where, for example, a change in the variable *X* strictly leads to a change in the variable *Y*).

Health), HILDA Survey 2007 and 2009

550

Dependent variable: Life Satisfaction β Independent variable t р Fruit and vegetable portions/day 0.02 [0.01, 0.03] 1.99 .047 Self-reported health 0.29 [0.25, 0.34] 12.22 .000 Log of household income 0.02 [-0.03, 0.06] 0.72 .468 -1.01Age -0.03 [-0.07, 0.02] .314 Age squared 0.02 [-0.02, 0.07] 1.00 .318 Masters or doctorate -0.22 [-0.78, 0.33] -0.79.428 Bachelor or honors 0.10 [-0.32, 0.52] 0.49 .627 Graduate diploma or certificate -0.05 [-0.39, 0.29] -0.29 .770 Advanced diploma -0.01 [-0.40, 0.37] -0.08 .939 Professional qualification 0.02 [-0.26, 0.30] 0.13 .896 Year 12 high school -1.19 .236 -0.12 [-0.31, 0.08] Full-time student -0.01 [-0.16, 0.13] -0.16 .872 Unemployed -0.24 [-0.46, -0.02] -2.11.035 Not in the labor force -0.03 [-0.14, 0.08] -0.48.632 Married 0.01 [-0.16, 0.17] 0.09 .930 Separated -0.55 [-0.86, -0.23 -3.40 .001 Divorced -0.35 [-0.66, -0.04] -2.22 .026 Widowed -1.88 -0.54 [-1.09, 0.02] .060 Long-term health condition -0.09 [-0.16, -0.02] -2.38 .017 # children under the age of 4 0.01 [-0.08, 0.09] 0.13 .895 # children aged 5-14 0.07 [-0.01, 0.15] 1.61 .107 Drink alcohol 1 or 2 days/week -0.06 .953 0.00 [-0.12, 0.11] Drink alcohol 2 or 3 days/week -0.01 [-0.11, 0.08] -0.26 .794 Drink alcohol 3 or 4 days/week -0.05 [-0.18, 0.09] -0.67 .500 Drink alcohol 5 or 6 days/week -0.06 [-0.22, 0.10] -0.76 .450 Drink alcohol everyday -0.17 [-0.37, 0.03] -1.67 .095 Non-smoker 0.04 [-0.09, 0.16] 0.61 .541 Never eat red meat 0.17 [-0.18, 0.52] 0.94 .346 Never eat fish -0.08 [-0.18, 0.03] -1.37 .171 Eat breakfast regularly 0.10 [0.02, 0.17] 2.54 .011 Drink low fat or skinny milk -0.04 [-0.11, 0.04] -0.97 .332 Avoid fatty foods -0.05 [-0.12, 0.01] -1.54 .124 BMI 0.01 [0.00, 0.02] 2.12 .034 Exercise regularly 0.05 [-0.01, 0.10] 1.72 .086 7.09 [5.99, 8.20] 12.57 .000 Constant Overall R² .09 Number of individuals 12,288 19,778 Number of observations

55

Note: Values in parentheses are 95% confidence intervals. Dependent variable is *Life Satisfaction* [range: 0-10]. For the self-reported health measure (covariate), individuals in the HILDA Survey were asked: "*In general, would you say your health is: Excellent, Very Good, Good, Fair, or Poor*". The resulting response distribution was as follows: 3% (Poor); 12.8% (Fair); 35.2% (Good); 36.8% (Very Good); 12.1% (Excellent). In the analysis above, these individual responses are coded from 1 (Poor) to 5 (Excellent), with the average reported score being 3.42 out of 5.

 Table S7. Description of Demographic and Socioeconomic Covariates

Variable	Description	Mean	SD	Min	Max
Age	Years of age	45.16	17.89	15	93
Age squared	Years of age squared, divided by 100	23.59	17.37	2.25	86.49
Income	Log of equivalized household income	10.15	1.02	0	13.01
Male		0.47	0.50	0	1
Full-time student		0.07	0.26	0	1
Education dummy 1	Masters or doctorate	0.04	0.19	0	1
Education dummy 2	Bachelor or honors	0.14	0.34	0	1
Education dummy 3	Grad diploma, grad certificate	0.06	0.23	0	1
Education dummy 4	Advanced diploma, diploma	0.09	0.29	0	1
Education dummy 5	Professional qualification (any certificate I, II, III, IV)	0.22	0.41	0	1
Education dummy 6	Year 12	0.15	0.36	0	1
Education dummy 7	Year 11 and below (baseline category)	0.30	0.46	0	1
Employment status 1	Unemployed	0.03	0.16	0	1
Employment status 2	Not in the labor force	0.30	0.46	0	1
Employment status 3	Employed (baseline category)	0.68	0.47	0	1
Married		0.51	0.50	0	1
Separated		0.03	0.18	0	1
Divorced		0.10	0.29	0	1
Widowed		0.05	0.22	0	1
Long-term health issues	Have a long-term health condition, disability or impairment	0.23	0.42	0	1
Number of children 1	Number of children under the age of 4	0.16	0.48	0	4
Number of children 2	Number of children aged 5-14	0.31	0.71	0	6

561 Table S8. Description of Dietary and Lifestyle Covariates


•	~	-
5	1	\mathbf{a}
	h	2
•	~	_

Variable	Description	Mean	SD	Min	Max
Daily fruit intake	Average number of fruit serves based on weekly intake	1.42	1.15	0	≥5
Daily vegetable intake	Average number of vegetable serves based on weekly intake	2.43	1.34	0	≥5
Weekly fruit intake frequency	Number of days in a usual week that fruit is eaten	5.31	2.17	0	7
Weekly vegetable intake frequency	Number of days in a usual week that vegetables are eaten	5.75	1.55	0	7
Usual fruit intake quantity	On those days, number of fruit serves eaten	1.79	1.07	0	≥5
Usual vegetable intake quantity	On those days, number of vegetable serves eaten	2.89	1.28	0	≥5
Alcohol intake 1	Drink alcohol: never, no longer, or rarely	0.38	0.48	0	1
Alcohol intake 2	Drink alcohol 1 or 2 days per week	0.20	0.40	0	1
Alcohol intake 3	Drink alcohol 2 or 3 days per week	0.12	0.32	0	1
Alcohol intake 4	Drink alcohol 3 or 4 days per week	0.14	0.35	0	1
Alcohol intake 5	Drink alcohol 5 or 6 days per week	0.09	0.29	0	1
Alcohol intake 6	Drink alcohol everyday	0.08	0.27	0	1
Non-smoker	Do not smoke cigarettes at all	0.80	0.40	0	1
Eat breakfast regularly	Eat breakfast seven times a week	0.70	0.46	0	1
Low fat/skim milk	Drink low fat or skinny milk	0.49	0.50	0	1
Avoid fatty foods	Eat fried potatoes, French fires, hot chips or wedges less than once a month	0.26	0.44	0	1
No fish intake	Never eat fresh, frozen, tinned fish, or shellfish	0.11	0.31	0	1
No meat intake	Never eat red meat (beef, veal, lamb, pork)	0.03	0.17	0	1
Regular physical exercise	Exercise at least three times a week per week; moderately to intensively	0.51	0.50	0	1
BMI	Body Mass Index	26.59	5.66	9.6	85.3

Note: Average Daily fruit intake = (Weekly fruit intake frequency \times Usual fruit intake quantity) divided by 7 days. Similarly, average Daily vegetable intake = (Weekly vegetable intake frequency \times Usual vegetable intake quantity) divided by 7 days. The Weekly intake frequency and Usual intake quantity variables correspond to the fruit and vegetable intake 'frequency' and 'quantity' survey questions presented in the Methods section. A standard serve (or portion) of fruit is 150 grams. A standard serve of vegetables is 75 grams.

569 Supplemental Figures:


- 570 These are derived from change-on-change regression equations.
- 571
- 572
- 573
- 574



575 576

577 Fig. S1. Scatter plot of change in fruit-and-vegetable consumption and change in satisfaction with life. To derive 578 this scatter plot, we generate fractional changes (such as -0.3 change in fruit and vegetable consumption) in 579 portions consumed between the two periods due to the fruit-and-vegetable intake variable being the average daily 580 amount of fruit and vegetables consumed (derived from the total weekly amount - this is how the questions in the 581 HILDA Survey are asked). Hence, the F+V variable (and its changes between the two periods) does not take on 582 a whole number (portion) for some individuals due to the averaging performed to get from the weekly to average 583 daily amounts. To get rid of the ensuing tens of thousands data points, we used the Stata command (called 584 'binscatter') that bands the points and produces a line of best fit.





587 588

Fig. S2. Scatter plot of change in fruit-and-vegetable consumption and change in happiness. To derive this scatter plot, we generate fractional changes (such as -0.3 change in fruit and vegetable consumption) in portions consumed between the two periods due to the fruit-and-vegetable intake variable being the average daily amount of fruit and vegetables consumed (derived from the total weekly amount - this is how the questions in the HILDA Survey are asked). Hence, the F+V variable (and its changes between the two periods) does not take on a whole number (portion) for some individuals due to the averaging performed to get from the weekly to average daily amounts. To get rid of the ensuing tens of thousands data points, we used the Stata command (called 'binscatter') that bands the points and produces a line of best fit.




Fig. S3. Vegetable servings size (Showcard K25, HILDA Survey, Waves 7 and 9)




Fig. S4. Fruit servings size (Showcard K27, HILDA Survey, Waves 7 and 9)

Further Supplemental Appendix: The Results Re-Estimated On New Data From 2009-2013, and Instrumental-Variable Estimation.

Table S9. (Table 1 Redone on Further Data). Life Satisfaction Equations: Fixed-effects RegressionModels of Changes in Life Satisfaction on Changes in Fruit and Vegetable Consumption and Covariates,HILDA Survey 2009 and 2013

	Model 1		Model 2		Model 3	
	(no covariate	s)	(partial set of covari	ates)	(full set of covaria	ites)
Independent variable	β	р	β	р	β	р
Fruit and vegetable portions/day	0.04 [0.02, 0.06]	.000	0.04 [0.03, 0.06]	.000	0.04 [0.02, 0.05]	.000
Log of household income			0.01 [-0.03, 0.05]	.682	0.01 [-0.03, 0.05]	.596
Age			-0.02 [-0.05, 0.00]	.082	-0.03 [-0.05, 0.00]	.050
Age ²			0.02 [-0.01, 0.04]	.165	0.02 [0.00, 0.05]	.108
Masters or doctorate			0.04 [-0.30, 0.38]	.810	0.04 [-0.30, 0.38]	.816
Bachelor or honors			-0.14 [-0.49, 0.22]	.444	-0.15 [-0.50, 0.20]	.411
Graduate diploma or certificate			-0.06 [-0.31, 0.19]	.647	-0.06 [-0.31, 0.19]	.643
Advanced diploma			0.10 [-0.14, 0.35]	.421	0.11 [-0.13, 0.35]	.372
Professional qualification			-0.13 [-0.34, 0.08]	.237	-0.12 [-0.33, 0.09]	.254
Year 12 high school			-0.09 [-0.26, 0.07]	.268	-0.07 [-0.24, 0.09]	.396
Full-time student			0.01 [-0.12, 0.14]	.876	0.01 [-0.12, 0.14]	.836
Unemployed			-0.22 [-0.40, -0.04]	.018	-0.22 [-0.40, -0.05]	.014
Not in the labor force			-0.05 [-0.13, 0.04]	.318	-0.05 [-0.14, 0.04]	.276
Married			0.05 [-0.09, 0.19]	.452	0.04 [-0.10, 0.18]	.568
Separated			-0.48 [-0.74, -0.23]	.000	-0.51 [-0.76, -0.26]	.000
Divorced			0.14 [-0.12, 0.40]	.293	0.12 [-0.14, 0.38]	.359
Widowed			0.04 [-0.26, 0.33]	.808	0.02 [-0.27, 0.32]	.875
Long-term health condition			-0.22 [-0.30, -0.15]	.000	-0.22 [-0.30, -0.15]	.000
# children under the age of 4			-0.08 [-0.14, -0.02]	.008	-0.07 [-0.13, -0.02]	.013
# children aged 5-14			-0.04 [-0.09, 0.01]	.142	-0.04 [-0.09, 0.01]	.138
Drink alcohol 1 or 2 days/week					-0.05 [-0.14, 0.05]	.321
Drink alcohol 2 or 3 days/week					-0.03 [-0.12, 0.07]	.579
Drink alcohol 3 or 4 days/week					-0.04 [-0.16, 0.07]	.478
Drink alcohol 5 or 6 days/week					-0.12 [-0.26, 0.02]	.097
Drink alcohol everyday					-0.11 [-0.27, 0.05]	.190
Non-smoker					0.03 [-0.08, 0.14]	.639
Never eat red meat					0.03 [-0.25, 0.32]	.829
Never eat fish					-0.08 [-0.19, 0.02]	.130
Eat breakfast regularly					0.13 [0.06, 0.20]	.001
Drink low fat or skinny milk					0.03 [-0.03, 0.09]	.332
Avoid fatty foods					0.01 [-0.05, 0.07]	.817
BMI					0.00 [-0.01, 0.01]	.773
Exercise regularly					0.15 [0.09, 0.20]	.000
Constant	7.76 [7.70, 7.83]	.000	8.41 [7.79, 9.04]	.000	8.38 [7.72, 9.04]	.000
Overall R^2	.02		.03		.04	
Number of individuals	16,242		16,242		16,242	
Number of observations	23,985		23,985		23,985	

Note: Values in parentheses are 95% confidence intervals. Dependent variable is Life Satisfaction [range: 0-10].

	Model 1 (no covariates)		Model 2 (partial set of covariates)		Model 3 (full set of covariates)	
Independent variable	β	р	β	р	β	р
Fruit and vegetable portions/day	0.02 [0.01, 0.04]	.000	0.02 [0.01, 0.04]	.001	0.02 [0.01, 0.03]	.006
Log of household income			-0.01 [-0.03, 0.03]	.790	-0.01 [-0.03, 0.03]	.823
Other covariates included	No		Partial		Full	
Constant	4.33 [4.28, 4.38]	.000	5.04 [4.53, 5.56]	.000	5.06 [4.51, 5.60]	.000
Overall R ²	.01		.01		.02	
Number of individuals	16,206		16,206		16,206	
Number of observations	23,917		23,917		23,917	

Table S10. (Table 2 Redone on Further Data) Happiness Equations: Fixed-effects RegressionModels of Changes in 'Been a Happy Person' on Changes in Fruit and Vegetable Consumption andCovariates, HILDA Survey Data 2009 and 2013

Note: Values in parentheses are 95% confidence intervals. Dependent variable is Been a Happy Person [range: 1-6].

Table S11. (Full Version of Table S10). Happiness Equations: Fixed-effects Regression Models of Changes in 'Been a Happy Person' on Changes in Fruit and Vegetable Consumption and Covariates, HILDA Survey 2009 and 2013

	Model 1 (no covariates	s)	Model 2 (partial set of covari	ates)	Model 3 (full set of covaria	ites)
Independent variable	β	p	β	р	β	p
Fruit and vegetable portions/day	0.02 [0.01, 0.04]	.000	0.02 [0.01, 0.04]	.001	0.02 [0.01, 0.03]	.006
Log of household income			-0.01 [-0.03, 0.03]	.790	-0.01 [-0.03, 0.03]	.823
Age			-0.02 [-0.04, 0.00]	.068	-0.02 [-0.04, 0.00]	.039
Age squared			0.01 [-0.01, 0.03]	.318	0.01 [-0.01, 0.03]	.205
Masters or doctorate			0.13 [-0.16, 0.42]	.374	0.14 [-0.15, 0.43]	.356
Bachelor or honors			0.00 [-0.27, 0.28]	.985	0.00 [-0.28, 0.28]	.994
Graduate diploma or certificate			0.14 [-0.08, 0.35]	.212	0.14 [-0.08, 0.36]	.219
Advanced diploma			0.13 [-0.10, 0.37]	.262	0.14 [-0.10, 0.37]	.253
Professional qualification			0.07 [-0.10, 0.25]	.422	0.07 [-0.11, 0.25]	.441
Year 12 high school			0.01 [-0.14, 0.15]	.912	0.02 [-0.13, 0.17]	.789
Full-time student			0.09 [-0.03, 0.20]	.154	0.09 [-0.03, 0.21]	.141
Unemployed			0.01 [-0.13, 0.16]	.854	0.01 [-0.14, 0.15]	.899
Not in the labor force			-0.10 [-0.17, -0.03]	.004	-0.11 [-0.18, -0.04]	.003
Married			0.00 [-0.12, 0.11]	.970	0.00 [-0.12, 0.11]	.951
Separated			-0.09 [-0.28, 0.11]	.393	-0.10 [-0.30, 0.09]	.300
Divorced			0.14 [-0.05, 0.34]	.155	0.14 [-0.06, 0.33]	.172
Widowed			0.00 [-0.28, 0.29]	.973	0.00 [-0.28, 0.29]	.989
Long-term health condition			-0.17 [-0.23, -0.11]	.000	-0.17 [-0.23, -0.11]	.000
# children under the age of 4			0.00 [-0.05, 0.05]	.969	0.00 [-0.04, 0.05]	.860
# children aged 5-14			-0.01 [-0.06, 0.03]	.647	-0.01 [-0.06, 0.03]	.648
Drink alcohol 1 or 2 days/week					-0.02 [-0.10, 0.06]	.607
Drink alcohol 2 or 3 days/week					0.02 [-0.05, 0.09]	.598
Drink alcohol 3 or 4 days/week					-0.06 [-0.15, 0.04]	.231
Drink alcohol 5 or 6 days/week					-0.08 [-0.19, 0.03]	.135
Drink alcohol everyday					-0.04 [-0.17, 0.09]	.549
Non-smoker					-0.04 [-0.13, 0.06]	.415
Never eat red meat					0.14 [-0.09, 0.38]	.232
Never eat fish					-0.03 [-0.12, 0.05]	.427
Eat breakfast regularly					0.04 [-0.02, 0.10]	.156
Drink low fat or skinny milk					0.00 [-0.05, 0.05]	.921
Avoid fatty foods					0.03 [-0.02, 0.09]	.190
BMI					0.00 [-0.01, 0.01]	.991
Exercise regularly					0.14 [0.09, 0.18]	.000
Constant	4.33 [4.28, 4.38]	.000	5.04 [4.53, 5.56]	.000	5.06 [4.51, 5.60]	.000
Overall R^2	.01		.01		.02	
Number of individuals	16,206		16,206		16,206	
Number of observations	23,917		23,917		23,917	

Note: Values in parentheses are 95% confidence intervals. Dependent variable is Been a Happy Person [range: 1-6].

Table S.12. Additional Life Satisfaction Equations: Instrumental-Variables Regression Models of 'Life Satisfaction' using 'Intensity of Go for 2&5 Campaign' as an Instrument for 'Fruit and Vegetable Consumption', HILDA Survey 2013

	Model 1 (no covariates)		Model 2 (partial set of covariates)		Model 3 (full set of covariates)	
Independent variable	β	р	β	р	β	р
Fruit and vegetable portions/day	0.10 [-0.93, 1.13]	.852	0.33 [-0.26, 0.92]	.276	0.31 [-0.24, 0.85]	.270
Log of household income			0.02 [-0.01, 0.05]	.165	0.02 [-0.01, 0.05]	.248
Other covariates included	No		Partial		Full	
Constant	7.56 [3.73, 11.39]	.000	7.83 [5.76, 9.90]	.000	7.88 [6.46, 9.30]	.000
Number of observations	13,788		13,788		13,788	

Note: Values in parentheses are 95% confidence intervals. Dependent variable is *Life Satisfaction* [range: 0-10]. The first-stage equations can be found in Table S.13.

Table S.13. First-Stage Regressions for Instrumented Life Satisfaction Equations in Table S.12: Regression Model of 'Fruit and Vegetable Consumption' on 'Intensity of Go for 2&5 Campaign', HILDA Survey 2013

	Model 1 (no covariates)		Model 2 (partial set of covariates)		Model 3 (full set of covariates)	
Independent variable	β	р	β	р	β	р
Intensity of campaign	0.01 [-0.003, 0.02]	.176	0.01 [0.003, 0.02]	.012	0.02 [0.01, 0.03]	.005
Log of household income			-0.01 [-0.05, 0.03]	.562	-0.03 [-0.07, 0.01]	.084
Other covariates included	No		Partial		Full	
Constant	3.68 [3.64, 3.73]	.000	3.40 [2.98, 3.83]	.000	2.51 [2.09, 2.94]	.000
First-stage F-statistic	1.83		6.31		8.05	
Number of observations	13,788		13,788		13,788	

Note: Values in parentheses are 95% confidence intervals. Dependent variable is *Fruit and Vegetable Consumption* (portions per day). First-stage *F*-statistic relates to a test of weak instruments, with a commonly suggested cutoff point of 10 for a strong instrument.

Table S.14. Additional Happiness Equations: Instrumental-Variables Regression Models of 'Been a Happy Person' using 'Intensity of Go for 2&5 Campaign' as an Instrument for 'Fruit and Vegetable Consumption', HILDA Survey 2013

	Model 1 (no covariates)		Model 2 (partial set of covariates)		Model 3 (full set of covariates)	
Independent variable	β	р	β	р	β	р
Fruit and vegetable portions/day	-0.38 [-1.39, 0.62]	.453	-0.00 [-0.43, 0.43]	.999	0.02 [-0.37, 0.42]	.907
Log of household income			0.01 [-0.01, 0.03]	.317	0.01 [-0.01, 0.03]	.439
Other covariates included	No		Partial		Full	
Constant	5.83 [2.12, 9.55]	.002	5.02 [3.52, 6.52]	.000	4.73 [3.70, 5.77]	.000
Number of observations	13,748		13,748		13,748	

Note: Values in parentheses are 95% confidence intervals. Dependent variable is *Been a happy person* [range: 1-6]. The first-stage equations can be found in Table S.15.

Table S.15. First-Stage Regressions for Instrumented Happiness Equations in Table S.14: Regression Model of 'Fruit and Vegetable Consumption' on 'Intensity of Go for 2&5 Campaign', HILDA Survey 2013

	Model 1 (no covariates)		Model 2 (partial set of covariates)		Model 3 (full set of covariates)	
Independent variable	β	р	β	р	β	р
Intensity of campaign	0.01 [-0.003, 0.02]	.169	0.01 [0.003, 0.02]	.011	0.02 [0.01, 0.03]	.004
Log of household income			-0.01 [-0.05, 0.03]	.541	-0.03 [-0.07, 0.01]	.078
Other covariates included	No		Partial		Full	
Constant	3.68 [3.64, 3.73]	.000	3.40 [2.98, 3.82]	.000	2.51 [2.08, 2.93]	.000
First-stage F-statistic	1.89		6.50		8.27	
Number of observations	13,748		13,748		13,748	

Note: Values in parentheses are 95% confidence intervals. Dependent variable is *Fruit and Vegetable Consumption* (portions per day). First-stage *F*-statistic relates to a test of weak instruments, with a commonly suggested cutoff point of 10 for a strong instrument.