



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

Statistics
Netherlands school of economics
ECONOMETRIC INSTITUTE

LOCAL LIMIT THEOREMS FOR SAMPLE EXTREMES

GIANNINI FOUNDATION OF
AGRICULTURAL ECONOMICS

MAR 4 1980
WITHDRAWN

L. de HAAN and S.I. RESNICK

Erasmus

REPORT 7922/S

Local Limit Theorems for Sample Extremes

L. de Haan* and S. I. Resnick**

Erasmus University, Rotterdam and Colorado State University

Abstract

A local limit theorem for maxima of i.i.d. random variables is proved. Also it is shown that under the so-called von Mises' conditions the density of the normalized maximum converges to the limit density in L_p ($0 < p \leq \infty$) provided both the original density and the limit density are in L_p . Finally an occupation time result is proved. The methods of proof are different from those used for the corresponding results concerning partial sums.

* Grateful acknowledgment is made to Colorado State University, Department of Statistics, for hospitality and support.

**Supported by NSF Grant No. MCS 78-00915

American Mathematical Society 1980 subject classifications. Primary 60F05; Secondary 60F25.

Key words and phrases: local limit theorem, extreme values, regular variation, occupation time, density convergence.

1. Introduction. In extreme value theory not much is known about the quality of convergence in the case of weak convergence of the normalized maximum of a sample to one of the limit distributions. Attempts to establish uniform rates of convergence have been sporadic and not completely satisfactory. There is an unpublished manuscript by A. A. Balkema dealing with the general case and specific cases have been considered by P. Hall (1978) and W. J. Hall and J. Wellner (1978). Large deviation results can be found in de Haan and Hordijk (1972) and C. W. Anderson (1978). Our focus in this paper is on local limit results.

As usual, the results for maxima parallel those for sums but the methods are completely different. Also, in the theory of partial sums of iid random variables, local limit results are frequently related to occupation time theorems (Breiman, 1968, p. 229; Darling and Kac, 1957). In extreme value theory this relation seems to be rather weak.

In Section 2 we give some preliminaries. Section 3 discusses the limit behavior of the probability the normalized sample maximum is in a certain interval. Section 4 treats density convergence in the uniform and L_p metrics. Finally in Section 5 we derive an occupation time result using the structure of extremal processes.

2. Preliminaries. We first recall some of the properties of distribution functions in the domain (D) of one of the extreme value distributions Φ_α , Ψ_α , and Λ (cf. Gnedenko, 1943). The properties are formulated in a way slightly different from de Haan, 1970.

If $F \in D(\Phi_\alpha)$, then the function $-\log F$ is regularly varying with exponent $-\alpha$ and

$$-\log F(x) = c(x) \exp\left\{-\int_1^x \frac{a(t)}{t} dt\right\}$$

$$\text{with } a(x) = c(x) \left\{ \int_1^\infty s^{-1} (-\log F(s)) ds \right\}^{-1} =$$

$$= \left[\int_x^\infty t^{-1} \{-\log F(t)\} \right]^{-1} \{-\log F(x)\}; \text{ then } \lim_{x \rightarrow \infty} a(x) = \alpha.$$

If the von Mises condition $\lim_{x \rightarrow \infty} \{-\log F(x)\}^{-1} x F'(x) = \alpha$ holds, then $F \in D(\Phi_\alpha)$ and

$$-\log F(x) = \{-\log F(1)\} \exp\left\{-\int_1^x \frac{a(t)}{t} dt\right\}$$

$$\text{with } a(x) = \{-F(x) \log F(x)\}^{-1} x F'(x).$$

We will further use the property that the positive norming constants $\{a_n\}$ in the limit relation $\lim_{n \rightarrow \infty} F^n(a_n x) = \Phi_\alpha(x)$ are regularly varying with exponent α^{-1} (being defined as the inverse function of $-1/\log F$ at the point n).

If $F \in D(\Lambda)$ and $F(x) < 1$ for all x then

$$-\log F(x) = c(x) \exp\left\{-\int_1^x \frac{a(t)}{f_1(t)} dt\right\}$$

$$\text{with } a(x) + 1 = c(x) \quad c(x) = 2 \left\{ \int_x^\infty (-\log F(t)) dt \right\}^{-2} (-\log F(x)) \left\{ \int_x^\infty \int_y^\infty (-\log F(t)) dt dy \right\}$$

and $f_1(x) = \left\{ \int_x^\infty \int_y^\infty (-\log F(t)) dt dy \right\} \left\{ \int_x^\infty (-\log F(t)) dt \right\}^{-1}$; then $\lim_{x \rightarrow \infty} a(x) = 1$

and $\lim_{x \rightarrow \infty} f_1'(x) = 0$. We also need $f_2(x) = \int_x^\infty (-\log F(t)) dt / (-\log F(x))$, which

is asymptotic to f_1 . If the von Mises type condition

$$\lim_{x \rightarrow \infty} \frac{F'(x) \left(\int_x^\infty (-\log F(t)) dt \right)}{\{-\log F(x)\}^2} = 1$$

holds, the $F \in D(\Lambda)$ and

$$-\log F(x) = (-\log F(1)) \exp \left\{ - \int_1^x \frac{a(t)}{f_0(t)} dt \right\}$$

with $a(x) = \frac{1}{F(x)}$ and $f_0(x) = \frac{-\log F(x)}{F'(x)}$. (The von Mises type condition given here is implied by the well-known von Mises condition given on p. 112 of deHaan, 1970; see also von Mises, 1936.) Any of the functions $f(f_0, f_1,$

or $f_2)$ is called an auxiliary function and satisfies $\lim_{t \rightarrow \infty} \frac{f(t + xf(t))}{f(t)} = 1$

uniformly on finite x -intervals. In the limit relation $\lim_{n \rightarrow \infty} F^n(a_n x + b_n) = \Lambda(x)$

we may then take b_n as the inverse function of $1/(-\log F)$ at the point n and $a_n = f(b_n)$. We will also need the property that $\{a_n\}$ is slowly varying.

The following lemma on regularly varying functions is well-known.

Lemma 1. Suppose U is regularly varying with exponent α . Take $\varepsilon > 0$. Then there exists x_0 such that for $x, y \geq x_0$

$$(1-\varepsilon) \left(\frac{x}{y} \right)^{\alpha-\varepsilon} < \frac{U(x)}{U(y)} < (1+\varepsilon) \left(\frac{x}{y} \right)^{\alpha+\varepsilon}$$

Proof. The inequalities follow easily from the representation for regularly varying functions (de Haan, 1970. Theorem 1.2.2; Feller, 1971, VIII. 9).

Remark. Obviously one can also prove that for any $\varepsilon_1, \varepsilon_2 > 0$ eventually

$$(1-\varepsilon_1)\left(\frac{x}{y}\right)^{\alpha-\varepsilon_2} < \frac{U(x)}{U(y)} < (1+\varepsilon_1)\left(\frac{x}{y}\right)^{\alpha+\varepsilon_2}.$$

Next we prove an inequality which is crucial in our attack.

Lemma 2. Suppose $F \in D(\Lambda)$ with auxiliary function f and $\varepsilon > 0$. There exists a t_0 such that for all t and x with $t, t + x f(t) \geq t_0$

$$(1-\varepsilon)\left\{\frac{-\log F(t)}{-\log F(t + x f(t))}\right\}^{-\varepsilon} \leq \frac{f(t + x f(t))}{f(t)} \leq (1+\varepsilon)\left\{\frac{-\log F(t)}{-\log F(t + x f(t))}\right\}^{\varepsilon}.$$

Proof. It is known (Balkema and de Haan, 1972) that there exists F_1 such that $-\log F_1$ has an increasing (negative) density and $-\log F(t) \sim \log F_1(t)(t \rightarrow \infty)$. Since F and F_1 have the same auxiliary function it is sufficient to prove the result for F_1 . Let U be the inverse function of $1/\{-\log F_1(x)\}$, then $f \circ U$ is slowly varying (de Haan, 1974). We apply lemma 1 for this function $f \circ U$ and replace x by $1/\{-\log F_1(U(t) + z f(U(t)))\}$ and y by t in the statement of lemma 1. Then for $U(t), U(t) + z f(U(t)) \geq t_0$

$$\begin{aligned} (1-\varepsilon)\{-t \log F_1(U(t) + z f(U(t)))\}^{+\varepsilon} &\leq \frac{f(U(t) + z f(U(t)))}{f(U(t))} \\ &\leq (1+\varepsilon)\{-t \log F_1(U(t) + z f(U(t)))\}^{-\varepsilon}. \end{aligned}$$

Introduce a new variable $s = U(t)$ to get the statement of the lemma.

Remark. Here again we may take the epsilons in the exponents different from the other ones.

We now use lemma 1 to prove a needed variant. This lemma can be found with a different proof in Pickands (1968).

Lemma 3. Suppose $V: \mathbb{R}^+ \rightarrow \mathbb{R}$ satisfies

$$\lim_{t \rightarrow \infty} \frac{V(tx) - V(t)}{L(t)} = \log x$$

for some positive (necessarily slowly varying function) L and all $x > 0$.

Take $\varepsilon > 0$. Then for $x, y \geq x_0$

$$(1-\varepsilon)^2 \frac{1 - (x/y)^{-\varepsilon}}{\varepsilon} - \varepsilon < \frac{V(x) - V(y)}{L(y)} < (1+\varepsilon)^2 \frac{(x/y)^{\varepsilon} - 1}{\varepsilon} + \varepsilon.$$

Proof. We may assume L is such that $V(x) = \int_1^x \frac{L(t)}{t} dt + L(x)$ (de Haan, 1970, p. 34). Then

$$\frac{V(x) - V(y)}{L(y)} = \int_y^x \frac{L(t)}{L(y)} \frac{dt}{t} + \frac{L(x)}{L(y)} - 1.$$

Apply lemma 1 for $U = L$; then for $x, y \geq x_0$

$$\begin{aligned} (1-\varepsilon) \frac{(1-(x/y)^{-\varepsilon})}{\varepsilon} + (1-\varepsilon) \left(\frac{x}{y}\right)^{-\varepsilon} - 1 &< \frac{V(x) - V(y)}{L(y)} \\ &< (1+\varepsilon) \frac{(x/y)^{\varepsilon} - 1}{\varepsilon} + (1+\varepsilon) \left(\frac{x}{y}\right)^{\varepsilon} - 1 \end{aligned}$$

and this is the statement of lemma 3.

Remark. Here also one can prove the refinement

$$(1-\varepsilon_1) \frac{1 - (\frac{x}{y})^{-\varepsilon_2}}{\varepsilon_2} - \varepsilon_1 < \frac{V(x) - V(y)}{L(y)} < (1+\varepsilon_1) \frac{(\frac{x}{y})^{\varepsilon_2} - 1}{\varepsilon_2} + \varepsilon_1.$$

3. Local limit theorem. We prove statements of the following form. Suppose X_1, X_2, \dots are i.i.d. random variables with common df F . Set $M_n = \sum_{i=1}^n X_i$

for $n = 1, 2, \dots$. If $\lim_{n \rightarrow \infty} F^n(a_n x + b_n) = G(x)$ for all x then under an additional smoothness condition $\lim_{n \rightarrow \infty} a_n P\left\{a_n \left| \frac{M_n - b_n}{a_n} - x \right| \leq h\right\} = 2hG'(x)$ uniformly for all x .

Theorem a) If $\lim_{n \rightarrow \infty} F^n(a_n x + b_n) = \Lambda(x) := \exp\{-e^{-x}\}$ for all x , $a_n \rightarrow \infty$ and

for all x

$$\lim_{t \rightarrow \infty} f_2(t+x) - f_2(t)$$

$$\lim_{t \rightarrow \infty} \left\{ \frac{\int_{t+x}^{\infty} (-\log F(x)) ds}{-\log F(t+x)} - \frac{\int_t^{\infty} (-\log F(x)) ds}{-\log F(t)} \right\} = 0,$$

then for all $h > 0$

$$\lim_{n \rightarrow \infty} a_n P\{a_n^{-1} (M_n - b_n) \in (x, x + a_n^{-1} h]\} = \lim_{n \rightarrow \infty} a_n \{F^n(a_n x + b_n + h) - F^n(a_n x + b_n)\}$$

$= h\Lambda'(x)$ uniformly for all x .

b) If $\lim_{n \rightarrow \infty} F^n(a_n x) = \Phi_\alpha(x)$ and for all x

$$\lim_{t \rightarrow \infty} t \left\{ \frac{-\log F(t+x)}{\int_{t+x}^{\infty} s^{-1} (-\log F(s)) ds} - \frac{-\log F(t)}{\int_t^{\infty} s^{-1} (-\log F(s)) ds} \right\} = 0,$$

then for all $h > 0$

$$\lim_{n \rightarrow \infty} a_n P\{a_n^{-1} M_n \in (x, x + a_n^{-1} h]\} = \lim_{n \rightarrow \infty} a_n \{F^n(a_n x + h) - F^n(a_n x)\} =$$

$= h \Phi'_\alpha(x)$ uniformly for all x .

c) If $\lim_{n \rightarrow \infty} F^n(a_n x + b_n) = \Lambda(x)$ for all x , $a_n \rightarrow 0$, $F(x) < 1$ for all x

and for all x

$$\lim_{t \rightarrow \infty} \frac{1}{f_1(t)} \left\{ \frac{f_1(t+x)}{f_2(t+x)} - \frac{f_1(t)}{f_2(t)} \right\} = 0$$

where f_1 and f_2 are defined in Section 2, then for all $h > 0$

$$\begin{aligned} \lim_{n \rightarrow \infty} a_n^{-1} P\{a_n^{-1}(M_n - b_n) \in (x, x + a_n h]\} &= \\ &= \lim_{n \rightarrow \infty} a_n^{-1} \{F^n(a_n x + b_n + h) - F^n(a_n x + b_n)\} = h \Lambda'(x) \end{aligned}$$

uniformly for all x .

Remark. We only treat the cases where $F(x) < 1$ for all x . We do not provide a result when $F^n(a_n x + b_n) \rightarrow \Lambda(x)$ and a_n tends to a finite positive constant (e.g. F exponential).

Proof. a) We have to prove

$$a_n [F^n(a_n x_n + b_n + h) - F^n(a_n x_n + b_n)] - h \Lambda'(x_n) \rightarrow 0$$

for any sequence $\{x_n\} (n \rightarrow \infty)$. Considering subsequences we may assume that both $\{x_n\}$ and $\{a_n x_n + b_n\}$ converge where $\{n\}$ now represents a subsequence of the integers. First consider the case $a_n x_n + b_n \rightarrow c < \infty$ (hence $x_n \rightarrow -\infty$). Then both $F^n(a_n x_n + b_n + h)$ and $F^n(a_n x_n + b_n)$ converge geometrically fast to zero; since a_n is slowly varying, the result follows.

Next consider the case $x_n \rightarrow -\infty$ and $a_n x_n + b_n \rightarrow \infty (n \rightarrow \infty)$.

Using the simple inequality $u < -\log(1-u)$ for $0 < u < 1$ twice we get

$$\begin{aligned}
 0 &\leq F^n(a_n x_n + b_n + h) - F^n(a_n x_n + b_n) \leq \\
 &\leq F^n(a_n x_n + b_n + h) \{ \log F^n(a_n x_n + b_n + h) - \log F^n(a_n x_n + b_n) \} = \\
 &= F^n(a_n x_n + b_n + h) (-\log F^n(a_n x_n + b_n)) \left\{ 1 - \frac{-\log F(a_n x_n + b_n + h)}{-\log F(a_n x_n + b_n)} \right\} \\
 &\leq F^n(a_n x_n + b_n + h) \{-\log F^n(a_n x_n + b_n)\} \{ \log(-\log F(a_n x_n + b_n + h)) + \\
 &\quad + \log(-\log F(a_n x_n + b_n)) \}.
 \end{aligned}$$

First we want to change the second factor. According to the representation of section 2

$$\frac{-\log F^n(a_n x_n + b_n)}{-\log F^n(a_n x_n + b_n + h)} = \frac{c(a_n x_n + b_n)}{c(a_n x_n + b_n + h)} \exp \left\{ \int_0^h \left(\frac{a(a_n x_n + b_n + s)}{f_1(a_n x_n + b_n + s)} \right) ds \right\},$$

which tends to 1 as $n \rightarrow \infty$ since $f_1 \rightarrow \infty$. So $-\log F^n(a_n x_n + b_n) < (1+\epsilon) \log F^n(a_n x_n + b_n + h)$ for sufficiently large n .

The third factor multiplied by a_n can be written as

$$(*) \quad a_n [\log c(b_n + a_n x_n + h) - \log c(b_n + a_n x_n)] + a_n \int_{x_n}^{x_n + \frac{h}{a_n}} \left(\frac{a(b_n + s a_n) f(b_n)}{f(b_n + s a_n)} \right) ds.$$

The first term of (*) is asymptotic to

$c f(b_n) \{c(b_n + a_n x_n + h) - c(b_n + a_n x_n)\}$ which by lemma 2 is at most

$$f(b_n + a_n x_n) \{c(b_n + a_n x_n + h) - c(b_n + a_n x_n)\} \left\{ \frac{-\log F(b_n)}{-\log F(b_n + a_n x_n)} \right\}^\epsilon (1-\epsilon)^{-1} \quad \text{for}$$

sufficiently large n .

Now $f(t)(c(t+h) - c(t)) = \frac{f(t)}{c} \left(\frac{f_1(t+h)}{f(t+h)} - \frac{f_1(t)}{f(t)} \right) =$
 $= c^{-1} \frac{f_1(t+h)}{f(t+h)} (f(t) - f(t+h)) + c^{-1} (f_1(t+h) - f_1(t))$. The first term of this
 expression tends to zero as $t \rightarrow \infty$ by assumption, the second one by the
 conditions for the domain of attraction ($\lim_{t \rightarrow \infty} f_1'(t) = 0$).

For the second term of (*) we use the lemma 2 again and get

$$\begin{aligned} a_n \int_{x_n}^{x_n+h/a_n} \left(\frac{a(b_n + s a_n) f(b_n)}{f(b_n + s a_n)} \right) ds &\leq \frac{1+\varepsilon}{1-\varepsilon} a_n \int_{x_n}^{x_n+h/a_n} \left(\frac{-\log F(b_n)}{-\log F(b_n + s a_n)} \right)^\varepsilon ds \\ &\leq \left(\frac{1+\varepsilon}{1-\varepsilon} \right) h \left(\frac{-\log F(b_n)}{-\log F(b_n + x_n a_n + h)} \right)^\varepsilon. \end{aligned}$$

Collecting the inequalities we get

$$\begin{aligned} 0 &\leq a_n [F^n(a_n x_n + b_n + h) - F^n(a_n x_n + b_n)] \leq \\ &\leq \frac{(1+\varepsilon)^2}{1-\varepsilon} (h+\varepsilon) \left\{ \frac{-\log F(a_n x_n + b_n)}{-\log F(b_n)} \right\}^{1-\varepsilon} \exp \left\{ - \frac{-\log F(a_n x_n + b_n)}{-\log F(b_n)} \right\} \end{aligned}$$

which tends to zero as $n \rightarrow \infty$ since $n(-\log F(a_n x_n + b_n)) = \{-\log F(b_n)\}^{-1}$
 $\{-\log F(a_n x_n + b_n)\} \rightarrow \infty$.

For the case $x_n \rightarrow c > -\infty$ (hence $a_n x_n + b_n \rightarrow \infty$) one quickly sees (since
 $a_n \rightarrow \infty$ implies $F^n(a_n x_n + b_n + h) = F^n(a_n (x_n + h a_n^{-1}) + b_n) \rightarrow \Lambda(c)$) that

in fact $a_n [F^n(a_n x_n + b_n + h) - F^n(a_n x_n + b_n)] \sim F^n(a_n x_n + b_n + h) \{-\log F^n(a_n x_n + b_n)\}$.

$[a_n \{-\log(-\log F(a_n x_n + b_n + h)) + \log(-\log F(a_n x_n + b_n))\}]$ as $n \rightarrow \infty$.

Now the product $F^n(a_n x_n + b_n + h) \{-\log F^n(a_n x_n + b_n)\}$ tends to the density $\Lambda'(c)$. For the third factor proceed as before, using the representation of Section 2 and the fact that $f(t + x f(t))/f(t) \rightarrow 1$ uniformly on finite intervals. It follows that the third factor tends to h .

b) First consider the case $a_n x_n \rightarrow c < \infty$ (hence $x_n \rightarrow c \leq 0$), then from the convergence of $F^n(a_n x_n + h)$ and $F^n(a_n x_n)$ to zero at a geometric rate and regular variation of $\{a_n\}$, the result follows. Next consider the case $x_n \downarrow 0$ and $a_n x_n \rightarrow \infty$ ($n \rightarrow \infty$). As under a) we get

$$0 \leq a_n [F^n(a_n x_n + h) - F^n(a_n x_n)] \leq (1+\epsilon) F^n(a_n x_n + h) \{-\log F^n(a_n x_n + h)\} \\ \{-a_n \log(-\log F(a_n x_n + h)) + a_n \log(-\log F(a_n x_n))\}.$$

The last factor multiplied by x_n can be written as

$$a_n x_n [\log c(a_n x_n + h) - \log c(a_n x_n)] + a_n x_n \int_{a_n x_n}^{a_n x_n + h} \frac{a(s)}{s} ds.$$

Since $\lim_{s \rightarrow \infty} a(s) = \alpha$, the last term is easily seen to converge to αh . The first term is asymptotic to (const.) $a_n x_n \{c(a_n x_n + h) - c(a_n x_n)\}$ and hence converges to zero by the condition of the theorem. Finally we use lemma 1 and the fact that $a_n x_n > M$ for sufficiently large n to get

$$x_n^{-1} F^n(a_n x_n + h) \{-\log F^n(a_n x_n + h)\} = \\ = x_n^{-1} \left(\frac{-\log F(a_n x_n + h)}{-\log F(a_n)} \right) \exp \left\{ - \left(\frac{-\log F(a_n x_n + h)}{-\log F(a_n)} \right) \right\} < \\ < (1+\epsilon) x_n^{-1} \left(x_n + \frac{h}{a_n} \right)^{-\alpha+\epsilon} \exp \{ -(1-\epsilon) \left(x_n + \frac{h}{a_n} \right)^{-\alpha-\epsilon} \} < \\ < (1+\epsilon) x_n^{-1-\alpha+\epsilon} \exp \{ -(1-\epsilon) x_n^{-\alpha-\epsilon} \left(1 + \frac{h}{M} \right)^{-\alpha-\epsilon} \} \rightarrow 0$$

as $n \rightarrow \infty$ since $x_n \downarrow 0$.

Finally if $x_n \rightarrow c > 0$ as under a) we get

$$a_n \{F^n(a_n x_n + h) - F^n(a_n x_n)\} \sim F^n(a_n x_n + h) \{-\log F^n(a_n x_n)\}$$

$\cdot [a_n \{-\log (-\log F(a_n x_n + h)) + \log (-\log F(a_n x_n))\}]$. The first two factors tend to $c\alpha^{-1}\phi'_\alpha(c)$ and for the third factor use the representation of Section 2 to see that it tends to αhc^{-1} .

c) This part of the proof is completely analogous to part a).

Corollary. Under the conditions of part a one has

$$\lim_{n \rightarrow \infty} a_n P\{x < M_n - b_n \leq x + h\} = h e^{-1}$$

and under the conditions of part b

$$\lim_{n \rightarrow \infty} a_n P\{x < M_n - a_n \leq x + h\} = \alpha h e^{-1}$$

We conclude this section by showing that theorem 1 is not true for any F in the domain of attraction of Λ with $|\log a_n| \rightarrow \infty$.

Take any positive differentiable function f with $\lim_{t \rightarrow \infty} f'(t) = 0$,

$\lim_{t \rightarrow \infty} f(t) = \infty$. Define b_n by $n = \exp \left\{ - \int_0^{b_n} \frac{ds}{f(s)} \right\}$ for $n \geq 1$. We may

construct a continuously differentiable function c_0 with the following properties:

$$c_0(b_1) = 0, c_0(b_1 + 1) = \{f(b_1)\}^{-1}$$

and increasing in between. Furthermore for $t > b_1 + 1$ let $c'_0(t) = -\{f(t)\}^{-1}$.

Since $\int_0^\infty \frac{dt}{f(t)} = \infty$, there exists b' such that $c_0(b') = 0$. Take n such

that $b_{n-1} < b' \leq b_n$ and define $c_0(t) = 0$ for $b' \leq t < b_n$ (with a small adaptation to keep c_0 continuously differentiable). Again we take

$c_0(b_n + 1) = \{f(b_n)\}^{-1}$ and increasing in between, etc. This way we get:

- 1) $c_0(t) + \int_0^t \frac{ds}{f(s)}$ is non-decreasing since $f(t) c_0'(t) \geq -1$,
- 2) $\lim_{t \rightarrow \infty} c_0(t) = 0$ since $\lim_{t \rightarrow \infty} f(t) = \infty$ and
- 3) $f(b_n)\{c_0(b_n + 1) - c_0(b_n)\} = 1$ for infinitely many n .

Define now $F(x) = \exp\left\{-\exp\left[-c_0(x) - \int_0^x \frac{ds}{f(s)}\right]\right\}$ then by the representation

in de Haan, 1970, p. 92 $\lim_{n \rightarrow \infty} F^n(a_n x + b_n) = \Lambda(x)$ for all x , where $a_n \rightarrow \infty$.

But (choose $x = 0$ and $h = 1$ in the corollary)

$$a_n [F^n(b_n + 1) - F^n(b_n)] \sim \Lambda'(0) f(b_n) [c_0(b_n + 1) - c_0(b_n)] +$$

$$\Lambda'(0) a_n \int_0^{1/a_n} \frac{f(b_n) ds}{f(b_n + s f(b_n))}$$

and this converges to $2\Lambda'(0)$ if n goes through the subsequences obtained above. If the corollary would be true, the limit should be just $\Lambda'(0)$.

4. Density convergence.

If we assume von Mises type conditions, we can prove the density of the normalized maximum converges to the density of the appropriate extreme value distribution in the L_p metric, $p \leq \infty$ provided both F' and the limit extreme value density are in the space L_p .

Theorem 2: Let $\{X_n, n \geq 1\}$ be i.i.d. with df $F(x)$ which is absolutely continuous with bounded density F' . Let $M_n = \bigvee_{i=1}^n X_i$ and let g_n be the density of M_n normalized as described below.

a) If $F'(x) > 0$ for all x in a neighborhood of ∞ and

$$(4.1) \quad \lim_{x \rightarrow \infty} \frac{x F'(x)}{-\log F(x)} = \alpha$$

then $g_n(x) = n F^{n-1}(a_n x) F'(a_n x) a_n \rightarrow \phi'_\alpha(x)$ uniformly in x .

b) If $F'(x) > 0$ for all x in a neighborhood of ∞ and

$$(4.2) \quad \lim_{x \rightarrow \infty} \frac{F'(x) \int_x^\infty (-\log F(t)) dt}{(\log F(x))^2} = 1$$

then

$$g_n(x) = n F^{n-1}(a_n x + b_n) F'(a_n x + b_n) a_n \rightarrow \Lambda'(x)$$

uniformly in x .

Proof. We suppose the normalizing constants are chosen as described in Section 2.

a) We need show $g_n(x_n) \rightarrow \phi'_\alpha(x_0)$ when $x_n \rightarrow x_0$ and $n \rightarrow \infty$ through some subsequence of the integers for the cases: (i) $x_n \rightarrow x_0 \in (0, \infty)$, (ii) $x_n \rightarrow \infty$, (iii) $x_n \rightarrow 0$, $a_n x_n \rightarrow \infty$, (iv) $x_n \rightarrow x_0 \leq 0$, $a_n x_n \leq K < \infty$.

Case (i): Write $g_n(x_n) = F^{n-1}(a_n x_n) \left(\frac{a_n x_n F'(a_n x_n)}{-\log F(a_n x_n)} \right) \cdot n(-\log F(a_n x_n))/x_n$.

Now use the fact that both $F^n(a_n x) \rightarrow \Phi_\alpha(x)$ (which follows from the von Mises condition (4.1) - see deHaan, 1979, p. 109) and $n(-\log F(a_n x)) \rightarrow x^{-\alpha}$ uniformly in neighborhoods of $+\infty$ (since we have monotone functions converging to continuous limits). So $g_n(x_n) \sim \Phi_\alpha(x_0) \alpha x_0^{-\alpha} / x_0 = \Phi'_\alpha(x_0)$.

Case (ii): Since $n(-\log F(a_n x_n)) \rightarrow 0$, we see by writing g_n as in (i) that $g_n(x_n) \rightarrow 0$.

Case (iii): As above $g_n(x_n) \sim \alpha e^{-(n-1)(-\log F(a_n x_n))} n(-\log F(a_n x_n)) / x_n$. Since $n(-\log F(a_n x_n)) = -\log F(a_n x_n) / (-\log F(a_n))$ we have by lemma 1 for sufficiently large n : $(1-\epsilon)x_n^{-(\alpha+\epsilon)} \leq n(-\log F(a_n x_n)) \leq (1+\epsilon)x_n^{-(\alpha-\epsilon)}$. Therefore

$$\limsup_{n \rightarrow \infty} g_n(x_n) \leq \limsup_{n \rightarrow \infty} \alpha e^{-\frac{(n-1)}{n}(1-\epsilon)x_n^{-(\alpha+\epsilon)}} (1+\epsilon)x_n^{-(\alpha-\epsilon+1)} = 0$$

Case (iv): Since F' is assumed bounded, $g_n(x_n) = 0$ ($nF^{n-1}(K) a_n$). Since na_n is regularly varying and $F^{n-1}(K) \rightarrow 0$ geometrically fast we have $nF^{n-1}(a_n x_n) a_n \rightarrow 0$.

b) Again we show $g_n(x_n) \rightarrow \Lambda'(x_0)$ when $x_n \rightarrow x_0$ and consider cases (i) $x_0 \in (-\infty, \infty)$, (ii) $x_0 = +\infty$, (iii) $x_0 = -\infty$, $a_n x_n + b_n \rightarrow +\infty$, (iv) $x_0 = -\infty$, $a_n x_n + b_n \leq K$. Write

$$g_n(x_n) = F^{n-1}(a_n x_n + b_n) \left(\frac{f_0(b_n)}{f_0(a_n x_n + b_n)} \right) n(-\log F(a_n x_n + b_n))$$

where as in Section 2 $a_n = f_0(b_n)$, $f_0(x) = (-\log F(x)) / F'(x)$.

Case (i): The result follows immediately since $F^{n-1}(a_n x + b_n) \rightarrow \Lambda(x)$ uniformly on $(-\infty, \infty)$, $f_0(b_n) / f_0(b_n + x f(b_n)) \rightarrow 1$ uniformly on finite intervals and $n(-\log F(a_n x + b_n)) \rightarrow e^{-x}$

uniformly on intervals bounded away from $-\infty$.

Case (ii): $F^n(a_n x_n + b_n) \rightarrow 1$, $n(-\log F(a_n x_n + b_n)) \rightarrow 0$ and from lemma 2 for n large

$$(4.3) \quad \frac{f_o(b_n)}{f_o(b_n + x_n f(b_n))} \leq \left(\frac{1}{1-\epsilon} \right) \left(\frac{-\log F(b_n + x_n f_o(b_n))}{-\log F(b_n)} \right)^{-\epsilon}$$

so $g_n(x_n) \leq \left(\frac{1}{1-\epsilon} \right) (n(-\log F(a_n x_n + b_n)))^{1-\epsilon} \rightarrow 0$.

Case (iii): In this case $n(-\log F(a_n x_n + b_n)) \rightarrow \infty$. Using (4.3) we have for large n

$$g_n(x_n) \leq e^{-\left(\frac{n-1}{n}\right)n(-\log F(a_n x_n + b_n))} \left(\frac{1}{1-\epsilon} \right) (n(-\log F(a_n x_n + b_n)))^{1-\epsilon} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Case (iv): As in part (a) since F' is bounded

$$g_n(x) = O(na_n F^{n-1}(K)).$$

Since na_n is regularly varying and $F^{n-1}(K) \rightarrow 0$ at a geometric rate, the result follows.

Remark: If we suppose F' is ultimately non-increasing and $g_n(x)$ converges pointwise to the appropriate extreme value density then (4.1) or (4.2) is true (recall the pointwise density convergence implies weak convergence).

(Cf. de Haan, 1970, Theorem 2.7.1b and Theorem 2.7.3b).

We now show that under the conditions of Theorem 2 a more general version of Theorem 1 holds.

Corollary.

a) Suppose (4.1). For any sequence $d_n \rightarrow \infty$ ($n \rightarrow \infty$)

$$\lim_{n \rightarrow \infty} d_n P\left\{x < \frac{M_n}{a_n} \leq x + d_n^{-1}h\right\} = h\phi'_\alpha(x)$$

uniformly for all x .

b) Suppose (4.2). For any sequence $d_n \rightarrow \infty$ ($n \rightarrow \infty$)

$$\lim_{n \rightarrow \infty} d_n P\left\{x < \frac{M_n - b_n}{a_n} \leq x + d_n^{-1}h\right\} = h \Lambda'(x)$$

uniformly for all x .

Proof.

a)

$$\lim_{n \rightarrow \infty} h^{-1} d_n \int_x^{x+d_n^{-1}h} g_n(t) dt = \phi'_\alpha(x)$$

uniformly in x .

b)

$$\lim_{n \rightarrow \infty} h^{-1} d_n \int_x^{x+d_n^{-1}h} g_n(t) dt = \Lambda'(x)$$

uniformly in x .

We next consider density convergence in the L_p metric:

Theorem 3: Let $\{X_n, n \geq 1\}$ be i.i.d. with df F which is absolutely continuous with density F' . Set $M_n = \bigvee_{i=1}^n X_i$ and let g_n be the density of the normalized M_n .

a) Suppose (4.1) holds and $g_n(x) = nF^{n-1}(a_n x)F'(a_n x)a_n$. If $\int_{-\infty}^{\infty} |F'(x)|^p dx < \infty$ and $p > (1+\alpha)^{-1}$ then

$$\int_{-\infty}^{\infty} |g_n(x) - \phi'_\alpha(x)|^p dx \rightarrow 0$$

b) Suppose (4.2) holds and $g_n(x) = nF^{n-1}(a_n x + b_n)F'(a_n x + b_n)a_n$. If $\int_{-\infty}^{\infty} |F'(x)|^p dx < \infty$ then

$$\int_{-\infty}^{\infty} |g_n(x) - \Lambda'(x)|^p dx \rightarrow 0.$$

Remark: Under the von Mises type conditions (4.1) or (4.2) we thus get L_p convergence when F' and the limit density are in L_p .

Proof: Since $(\phi'_\alpha(x))^p = \alpha^p x^{-p(\alpha+1)} \exp\{-px^{-\alpha}\}$ we have $\int_0^\infty (\phi'_\alpha(x))^p dx < \infty$

if and only if $p > (1+\alpha)^{-1}$. Hence

$$(4.4) \quad \lim_{M \rightarrow \infty} \int_{[0, M^{-1}] \cup [M, \infty)} (\phi'_\alpha(x))^p dx = 0.$$

Next we prove that the right tail of $\int g_n^p$ is eventually small. For $M > 0$

$$\begin{aligned} \int_M^\infty (g_n(x))^p dx &= \int_M^\infty (nF^{n-1}(a_n x) F'(a_n x) a_n)^p dx \\ &\leq \int_M^\infty (n a_n F'(a_n x))^p dx = n^p a_n^{p-1} \int_{a_n M}^\infty (F'(y))^p dy. \end{aligned}$$

Now using (4.1) and Karamata's Theorem (eg. de Haan, 1970, p. 15, Theorem 1.2.9)

we get since $-\log F$ is regularly varying

$$\begin{aligned} n^p a_n^{p-1} \int_{a_n M}^\infty (F'(y))^p dy &\sim n^p a_n^{p-1} \alpha^p \int_{a_n M}^\infty (-\log F(y))^p y^{-p} dy \\ &\sim n^p a_n^{p-1} \alpha^p (p(1+\alpha)-1)^{-1} (-\log F(a_n M))^p (a_n M)^{-p+1} \\ &\sim n^p a_n^{p-1} \alpha^p (p(1+\alpha)-1)^{-1} (-\log F(a_n))^p M^{-p\alpha} (a_n M)^{-p+1} \\ &= \alpha^p (p(1+\alpha)-1)^{-1} M^{1-p(\alpha+1)} \end{aligned}$$

(as $n \rightarrow \infty$) where we used $-\log F(a_n) = n^{-1}$. We conclude

$$(4.5) \quad \lim_{M \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_M^\infty (g_n(x))^p dx = 0.$$

We now consider the region $(-\infty, M^{-1}]$. It is convenient to define

δ_n by $-\log F(\delta_n) \sim n^{-1/2}$ so that $\delta_n \rightarrow \infty$ but $\delta_n/a_n \rightarrow 0$. (Otherwise if along a subsequence $\delta_n/a_n \rightarrow c > 0$ then

$$\begin{aligned} n(-\log F(\delta_n)) &\sim n^{1/2} \sim n(-\log F(a_n \cdot (\delta_n/a_n))) \\ &\rightarrow c^{-\alpha}.) \end{aligned}$$

We have

$$\int_{-\infty}^{M^{-1}} (g_n(x))^p dx = \int_{-\infty}^{\delta_n/a_n} (g_n(x))^p dx + \int_{\delta_n/a_n}^{M^{-1}} (g_n(x))^p dx$$

$$= (1) + (2).$$

$$\text{Now } (1) = n^p a_n^{p-1} \int_{-\infty}^{\delta_n} (F^{n-1}(t)) F'(t))^p dt$$

$$\leq n^p a_n^{p-1} F^{p(n-1)}(\delta_n) \int_{-\infty}^{\infty} (F'(t))^p dt$$

and since $n^p a_n^{p-1}$ is a regularly varying function of n and $F^{p(n-1)}(\delta_n) \leq \exp\{-\frac{1}{2}pn^{\frac{1}{2}}\}$ the above goes to zero.

For (2) we write

$$(2) = \int_{\delta_n/a_n}^{M^{-1}} n^p a_n^{p-1} (F^{n-1}(t) F'(t))^p dt.$$

For (2) we have that for given ϵ , $F'(t) \leq (\alpha+\epsilon)t^{-1}(-\log F(t))$ ultimately. Noting that on the region of integration of (2) $x > \delta_n/a_n$ so that $a_n x > \delta_n \rightarrow \infty$, we have for sufficiently large n that

$$(2) \leq \int_{\delta_n/a_n}^{M^{-1}} (\exp\{-\frac{n-1}{n} \cdot n(-\log F(a_n x))\} (\alpha+\epsilon)n(-\log F(a_n x) \cdot x^{-1})^p dx.$$

Applying lemma 1 we get

$$(2) \leq \int_{\delta_n/a_n}^{M^{-1}} (\exp\{-(\frac{n-1}{n})(1-\epsilon)x^{-\alpha-\epsilon}\} (\alpha+\epsilon)(1+\epsilon)x^{-\alpha+\epsilon} x^{-1})^p dx.$$

and for appropriate positive constants c_1, c_2 this is

$$\leq c_1 \int_0^{c_2 M^{-1}} \exp\{-s^{-\alpha-\epsilon}\} s^{-p(\alpha-\epsilon)-p} ds < \infty.$$

We conclude $\lim_{M \rightarrow \infty} \limsup_{n \rightarrow \infty} (2) = 0$. Combining this with $\lim_{n \rightarrow \infty} (1) = 0$ gives

$$(4.6) \quad \lim_{M \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{-\infty}^{M^{-1}} (g_n(x))^p dx = 0.$$

Next fix $M > 1$ and consider $\int_{M^{-1}}^M |g_n(x) - \phi'_\alpha(x)|^p dx$.

$$\text{Write } g_n(x) = F^{n-1}(a_n x) \frac{a_n x F(a_n x)}{-\log F(a_n x)} \frac{n(-\log F(a_n x))}{x}.$$

Observe on $[M^{-1}, M]$

$$\begin{aligned} F^{n-1}(a_n x)/\phi_\alpha(x) &\rightarrow 1 \text{ uniformly,} \\ a_n x F'(a_n x)/(-\log F(a_n x)) &\rightarrow \alpha \text{ uniformly} \end{aligned}$$

and $n(-\log F(a_n x))x^\alpha \rightarrow 1$ uniformly.

So on $[M^{-1}, M]$

$$g_n(x) = \phi_\alpha(x) \alpha x^{-\alpha-1}(1+\zeta_n(x)) = (1+\zeta_n(x))\phi'_\alpha(x)$$

where $\zeta_n(x) \rightarrow 0$ uniformly in $x \in [M^{-1}, M]$. Therefore,

$$\int_{M^{-1}}^M |g_n(x) - \phi'_\alpha(x)|^p dx = \int_{M^{-1}}^M |\phi'_\alpha(x)|^p (\zeta_n(x))^p dx$$

and since $\phi'_\alpha \in L_p$ and $\zeta_n \rightarrow 0$ uniformly on

$[M^{-1}, M]$, we get

$$(4.7) \quad \lim_{n \rightarrow \infty} \int_{M^{-1}}^M |g_n(x) - \phi'_\alpha(x)|^p dx = 0$$

Finally write

$$\begin{aligned} \int_{-\infty}^{\infty} |g_n(x) - \phi'_\alpha(x)|^p dx &= \int_{-\infty}^{M^{-1}} + \int_{M^{-1}}^M + \int_M^{\infty} \\ &\leq 2^p \{ \int_{-\infty}^{M^{-1}} |g_n(x)|^p dx + \int_{-\infty}^{M^{-1}} |\phi'_\alpha(x)|^p dx \\ &\quad + \int_M^{\infty} (g_n(x))^p dx + \int_M^{\infty} (\phi'_\alpha(x))^p dx \} + \int_{M^{-1}}^M |g_n(x) - \phi'_\alpha(x)|^p dx. \end{aligned}$$

Now let $n \rightarrow \infty$ and then $M \rightarrow \infty$ and use (4.4), (4.5), (4.6), (4.7) to get the desired L_p convergence.

(b) As in part (a), we begin by noting

$$(4.8) \quad \lim_{M \rightarrow \infty} \int_{(-\infty, -M] \cup [M, \infty)} |\Lambda'(x)|^p dx = 0.$$

This is because $\Lambda' \in L_p$ for $p > 0$. Now we replace Λ' by g_n : We have

$$\begin{aligned} \int_M^{\infty} (g_n(x))^p dx &= \int_M^{\infty} (n F^{n-1}(a_n x + b_n) F'(a_n x + b_n) a_n)^p dx \\ &= \int_M^{\infty} (F^{n-1}(a_n x + b_n) \frac{f_o(b_n)}{f_o(a_n x + b_n)} n(-\log F(a_n x + b_n)))^p dx. \end{aligned}$$

Note $F^{n-1}(a_n x + b_n) \leq 1$. Also, given $\epsilon > 0$, for n, M large we have by lemma 2 $\frac{f_o(b_n)}{f_o(a_n x + b_n)} \leq (\frac{1}{1-\epsilon}) (n(-\log F(a_n x + b_n)))^{-\epsilon}$

Therefore,

$$\begin{aligned} \int_M^\infty |g_n(x)|^p dx &\leq (\frac{1}{1-\epsilon})^p \int_M^\infty (n(-\log F(a_n x + b_n)))^{p(1-\epsilon)} dx \\ &= n^{p(1-\epsilon)} a_n^{-1} (\frac{1}{1-\epsilon})^p \int_{a_n M + b_n}^\infty (-\log F(s))^{p(1-\epsilon)} ds, \end{aligned}$$

and by Theorem 2.8.1, p. 113 of de Haan, 1970, this is asymptotic to

$$\begin{aligned} &\sim n^{p(1-\epsilon)} a_n^{-1} (\frac{1}{1-\epsilon})^p \frac{1}{p(1-\epsilon)} (-\log F(a_n M + b_n))^{p(1-\epsilon)-1} \int_{a_n M + b_n}^\infty (-\log F(s)) ds \\ &= (\text{const}) (n(-\log F(a_n M + b_n)))^{p(1-\epsilon)-1} \frac{\int_{a_n M + b_n}^\infty (-\log F(s)) ds}{\int_{b_n}^\infty (-\log F(s)) ds} \cdot \frac{a_n^{-1} \int_{b_n}^\infty (-\log F(s)) ds}{-\log F(b_n)} \\ &\quad \cdot n(-\log F(b_n)). \end{aligned}$$

Now since $F \in D(\Lambda)$ (from 4.2) we have $n(-\log F(a_n M + b_n)) \rightarrow e^{-M}$. Also by de Haan, 1970, p. 90, Lemma 2.5.1

$$\frac{\int_{a_n M + b_n}^\infty (-\log F(s)) ds}{\int_{b_n}^\infty (-\log F(s)) ds} \rightarrow e^{-M}$$

Recall $n(-\log F(b_n)) = 1$, $\frac{\int_{b_n}^\infty (-\log F(s)) ds}{(-\log F(b_n))}$

$\sim f_o(b_n) \sim a_n$ and we see

$$\lim_{n \rightarrow \infty} \sup_M \int_M^\infty |g_n(x)|^p dx \leq (\text{const}) e^{-Mp(1-\epsilon)}.$$

Therefore,

$$(4.9) \quad \lim_{M \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_M^\infty |g_n(x)|^p dx = 0.$$

For the region $(-\infty, -M]$, it is convenient to choose δ_n satisfying $-\log F(\delta_n) \sim n^{-1/2}$ so that $\delta_n \rightarrow \infty$ and $\frac{\delta_n - b_n}{a_n} \rightarrow -\infty$ (since otherwise if along a subsequence $\frac{\delta_n - b_n}{a_n} \rightarrow c > -\infty$ we would have

$$\infty \leftarrow n^{\frac{1}{2}} \sim n(-\log F(\delta_n)) = n(-\log F(a_n(\frac{\delta_n - b_n}{a_n} + b_n))) \rightarrow e^{-c}.$$

We now decompose the integral

$$\int_{-\infty}^{-M} |g_n(x)|^p dx = \int_{-\infty}^{(\delta_n - b_n)/a_n} + \int_{(\delta_n - b_n)/a_n}^{-M} = A + B.$$

For A we have

$$\begin{aligned} A &= \int_{-\infty}^{(\delta_n - b_n)/a_n} |nF^{n-1}(a_n x + b_n)F'(a_n x + b_n)a_n|^p dx \\ &= n^p a_n^{p-1} \int_{-\infty}^{\delta_n} |F^{n-1}(y)F'(y)|^p dy \\ &\leq n^p a_n^{p-1} F^{n-1}(\delta_n) \int_{-\infty}^{\infty} |F'(y)|^p dy \end{aligned}$$

and since a_n is slowly varying and $F^{n-1}(\delta_n) =$

$$\exp\{-(-\log F(\delta_n))(n-1)\} \leq \exp\{-\frac{1}{2}n^{\frac{1}{2}}\} \text{ for } n \text{ large, we have } A \rightarrow 0 \text{ as}$$

$n \rightarrow \infty$. For B we write

$$B = \int_{(\delta_n - b_n)/a_n}^{-M} \left| F^{n-1}(a_n x + b_n) \frac{f_o(b_n)}{f_o(a_n x + b_n)} n(-\log F(a_n x + b_n)) \right|^p dx$$

which for given $\epsilon > 0$ and n sufficiently large is bounded according to lemma 2 by

$$\leq \int_{(\delta_n - b_n)/a_n}^{-M} |F^{n-1}(a_n x + b_n) (\frac{1}{1-\epsilon}) (n(-\log F(a_n x + b_n)))^{1-\epsilon}|^p dx.$$

Note lemma 2 is applicable since if $x > (\delta_n - b_n)/a_n$ then $b_n + a_n x \geq b_n + a_n((\delta_n - b_n)/a_n) = \delta_n \rightarrow \infty$. In the above integral, make the change of variable $y = (n(-\log F(a_n x + b_n)))^{-1}$ and the integral becomes

$$\int_{(n(-\log F(\delta_n)))^{-1}}^{(n(-\log F(a_n(-M) + b_n)))^{-1}} e^{-(\frac{n-1}{n})p} y^{-1} y^{-(1+\epsilon)p} H_n(dy).$$

where $H_n(y) = \frac{1}{(-\log F)^{\leftarrow(ny) - b_n} a_n}$ where the arrow denotes the inverse

function. Since $F \in D(\Lambda)$, $(\frac{1}{-\log F})^{\leftarrow}$ is in the class Π (de Haan, 1970, 1974) and satisfies $H_n(y) \rightarrow \log y$, $y > 0$, $n \rightarrow \infty$. The endpoints of the interval of integration both converge:

$$\theta_1(n) := (n(-\log F(\delta_n)))^{-1} \rightarrow 0$$

$$\theta_2(n) := (n(-\log F(a_n(-M) + b_n)))^{-1} \rightarrow e^{-M}$$

The above integral is bounded by

$$\int_{\theta_1(n)}^{\theta_2(n)} g(y) H_n(dy)$$

(where $g(y) = e^{-(p/2)y^{-1}} y^{-(1+\epsilon)p}$) and using partial integration this becomes

$$g(\theta_2(n)) H_n(\theta_2(n)) - g(\theta_1(n)) H_n(\theta_1(n))$$

$$- \int_{\theta_1(n)}^{\theta_2(n)} H_n(y) g'(y) dy = (i) + (ij) + (iij)$$

Now (i) $\rightarrow g(e^{-M}) \log e^{-M} = g(e^{-M})/M$ as $n \rightarrow \infty$ and note $\lim_{M \rightarrow \infty} g(e^{-M})/M =$

$$\lim_{s \downarrow 0} \frac{g(s)}{-\log s} = 0.$$

For (ij) and (iij) we need to use lemma 3: For (iij), on the region of integration $y > \theta_1(n) = \frac{1}{n(-\log F(\delta_n))}$ so $ny > n^{\frac{1}{2}}$. Therefore, for given

ϵ and n sufficiently large (remember $b_n = (\frac{1}{-\log F})^{\leftarrow}(n)$ and suppose $\theta_2(n) \ll 1$)

$$\int_{\theta_1(n)}^{\theta_2(n)} g'(y) |H_n(y)| dy \leq \int_{\theta_1(n)}^{\theta_2(n)} g'(y) \left[\frac{(1-\epsilon)^2}{\epsilon} (1-y^{-\epsilon}) + \epsilon \right] dy.$$

It is readily seen $\lim_{M \rightarrow \infty} \limsup_{n \rightarrow \infty}$ of this bound is 0. We can handle (ij)

similarly by using lemma 3 and in conclusion we find

$$(4.10) \quad \lim_{M \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{-\infty}^{-M} |g_n(x)|^p dx = 0.$$

Finally on $[-M, M]$ we have

$$\begin{aligned} F^{n-1}(a_n x + b_n) \frac{f_o(b_n)}{f_o(a_n x + b_n)} n(-\log F(a_n x + b_n)) \\ = (1 + \zeta_n(x)) \Lambda'(x) \end{aligned}$$

where $\zeta_n \rightarrow 0$ uniformly on $[-M, M]$ (recall $f_0(t + xf(t))/f_0(t) \rightarrow 1$ as $t \rightarrow \infty$ locally uniformly) so that

$$(4.11) \quad \int_{-M}^M |g_n(x) - \Lambda'(x)|^p dx = \int_{-M}^M \zeta_n(x)^p |\Lambda'(x)|^p dx$$

$\rightarrow 0$ as $n \rightarrow \infty$.

We finish by decomposing $\int_{-\infty}^{\infty} |g_n(x) - \Lambda'(x)|^p dx$ as done at the end of part (a) and then using 4.8, 4.9, 4.10, 4.11. The proof is complete.

5. Occupation Times

Occupation time theorems for sums are frequently related to local limit theorems. See Breiman (1968), Darling and Kac (1957). For the case of maxima we have not found a direct connection but only one of analytic methodology which we discuss below.

In this section it is convenient to have a slightly different representation for $F\epsilon D(\Lambda)$ and we proceed as in de Haan, 1970, Theorem 2.4.2. We restrict attention to the case where $F(x) < 1$ for all x . According to this result $F\epsilon D(\Lambda)$, iff

$$\lim_{x \rightarrow \infty} \frac{(1 - F(x)) \int_x^\infty \int_s^\infty (1 - F(u)) du ds}{\left(\int_x^\infty (1 - F(s)) ds \right)^2} = 1$$

and in this case

$$1 - F(x) = c(x) \exp \left\{ - \int_1^x \frac{a(t)}{f(t)} dt \right\}$$

and we may set

$$f^*(x) = \frac{\int_x^\infty \int_s^\infty (1 - F(u)) du ds}{\int_x^\infty (1 - F(s)) ds}$$

and

$$a(x) = -1 + 2 \frac{(1 - F(x)) \int_x^\infty \int_s^\infty (1 - F(u)) du ds}{\left(\int_x^\infty (1 - F(s)) ds \right)^2}$$

$$c(x) = \frac{(1 - F(x)) \int_x^\infty \int_s^\infty (1 - F(u)) du}{\left(\int_x^\infty (1 - F(s)) ds \right)^2} \bigg/ \frac{\int_1^\infty \int_s^\infty (1 - F(u)) du}{\left(\int_1^\infty (1 - F(s)) ds \right)^2}$$

so that $a(x) \rightarrow 1$, $c(x) \rightarrow c$, $f'(x) \rightarrow 0$ and hence $f^*(t + xf^*(t)) / f^*(t) \rightarrow 1$ locally uniformly. In this case $F^n(a_n x + b_n) \rightarrow \Lambda(x)$ where $b_n = \left(\frac{1}{1-F} \right)^{\leftarrow}(n)$ and $a_n = f^*(b_n)$.

Lemma 4: Suppose $F\epsilon D(\Lambda)$ and $a_n \rightarrow 0$ (i.e. $f^*(x) \rightarrow 0$, $x \rightarrow \infty$). Set

$R(x) = -\log(1 - F(x))$ and suppose $\lim_{t \rightarrow \infty} f^*(t+x)/f^*(t) = 1$ locally uniformly in x . Then for all x

$$a_n(R(x+b_n) - \log n) \rightarrow x$$

or in terms of measures

$$a_n R(\cdot + b_n) \xrightarrow{v} m$$

where m is Lebesgue measure on $(-\infty, \infty)$ and \xrightarrow{v} denotes vague convergence.

Proof: Let $c_0(x) = -\log c(x)$ in the previous representation so that

$$R(x) = c_0(x) + \int_1^x \frac{a(t)}{f^*(t)} dt$$

and $c_0(x) \rightarrow c_0$.

Now

$$a_n(R(x + b_n) - \log n) = dt.$$

$$a_n(c_0(x + b_n) - c_0(b_n)) + a_n \int_{b_n}^{x+b_n} \frac{a(t)}{f^*(t)} dt$$

Since $a_n = f^*(b_n) \rightarrow 0$, the above is

$$o(1) + \int_0^x a(t + b_n) \frac{f^*(b_n)}{f^*(t + b_n)} dt \rightarrow \int_0^x 1 dt = x$$

as $n \rightarrow \infty$ since $a(t + b_n) \rightarrow 1$ and $f^*(b_n)/f^*(t + b_n) \rightarrow 1$ locally uniformly.

Remark: The condition $f^*(t+x)/f^*(t) \rightarrow 1$ locally uniformly is satisfied

by $1 - F(x) = e^{-x^\alpha}$, $x \geq 0$ for $\alpha > 1$ since in this case we may take

$$f^*(x) \sim (1 - F(x))/F'(x) = \frac{1}{\alpha} x^{-\alpha+1}$$

Also in the case $F = N(0,1)$ the condition is satisfied since

$$f^*(x) \sim (1 - F(x))/F'(x) = (1 - N(0,1,x))/n(0,1,x)$$

$$\sim \frac{n(0,1,x)/x}{n(0,1,x)} \sim x^{-1} \text{ as } x \rightarrow \infty$$

(by Mill's ratio). These two examples illustrate that a helpful sufficient

condition is that $\lim_{t \rightarrow \infty} \frac{d}{dt} \log f^*(t) = 0$. This is easy to verify: Set

$g(t) = \log f^*(t)$ and we need to check $g(t+x) - g(t) \rightarrow 0$, $t \rightarrow \infty$, locally

uniformly when $g'(t) \rightarrow 0$, $t \rightarrow \infty$, which is clear.

We now consider an occupation time theorem for maxima and phrase it in terms of random measures. For $x \in \mathbb{R}$ define the point measure $\varepsilon_x(A) = 1$ if $x \in A$, 0 if $x \notin A$. The occupation time measure for maxima is then

$$O(\cdot) = \sum_{k=1}^{\infty} \varepsilon_{M_k}(\cdot).$$

It is more convenient to express O in terms of record values: Say X_j is a record of the sequence $\{X_n, n \geq 1\}$ if $M_j > M_{j-1}$ and let the record value indices be $L(j)$, $j \geq 0$; i.e. $L = 1$ and $L(j) = \inf \{k > L(j-1) | M_k > M_{k-1}\}$. Let $\Delta_j = L(j) - L(j-1)$, $j \geq 0$. Then

$$O(\cdot) = \sum_{k=0}^{\infty} \Delta_{k+1} \varepsilon_{X_{L(k)}}.$$

We first compute the Laplace functional (cf. Neveu, 1977, p. 258) of O :

Proposition: Suppose F is continuous. Then for g continuous with compact support we have

$$\begin{aligned} E \exp \left\{ - \int g(x) O(dx) \right\} &= \\ &= \exp \left\{ - \int \frac{1 - e^{-g(x)}}{1 - e^{-g(x)} F(x)} R(dx) \right\}. \end{aligned}$$

Proof We have

$$\begin{aligned} E \exp \left\{ - \int g(x) O(dx) \right\} &= E \exp \left\{ - \sum_{k=0}^{\infty} \Delta_{k+1} g(X_{L(k)}) \right\} \\ &= E(E(\exp \left\{ - \sum_{k=0}^{\infty} \Delta_{k+1} g(X_{L(k)}) \right\} | X_{L(i)}, i \geq 0)). \end{aligned}$$

Conditional on $\{X_{L(i)}, i \geq 0\}$, the interrecord times are independent

geometrically distributed random variables (Shorrock, 1972) so the above is

$$\begin{aligned} &= E \left(\prod_{k=0}^{\infty} E(e^{-g(X_{L(k)}) \Delta_{k+1}} | X_{L(i)}, i \geq 0) \right) \\ &= E \prod_{k=0}^{\infty} \sum_{n=1}^{\infty} e^{-g(X_{L(k)}) n} F(X_{L(k)})^{n-1} (1 - F(X_{L(k)})) \\ &= E \prod_{k=0}^{\infty} h(X_{L(k)}) \end{aligned}$$

where
$$h(x) = \frac{(1 - F(x))e^{-g(x)}}{1 - e^{-g(x)}F(x)} \quad . \quad \text{So}$$

$$\begin{aligned} E \prod_{k=0}^{\infty} h(X_{L(k)}) &= E \exp \left\{ - \sum_{k=0}^{\infty} (-\log h(X_{L(k)})) \right\} \\ &= E \exp \left\{ - \int (-\log h(x)) \sum_{k=0}^{\infty} \varepsilon_{X_{L(k)}}(dx) \right\} \end{aligned}$$

which is the Laplace functional of the point process with points $\{X_{L(k)}\}$ at the function $-\log h$. We now make use of the well known fact that $\{X_{L(k)}\}$ is Poisson with mean measure R . The Laplace functional of such a process is well known to be

$$\begin{aligned} &\exp \left\{ - \int (1 - e^{-(\log h(x))}) R(dx) \right\} \\ &= \exp \left\{ - \int (1 - h(x)) R(dx) \right\} \end{aligned}$$

and substituting the form of h gives the result.

Theorem 4: Suppose $F \in D(\Lambda)$ is continuous with auxiliary function f satisfying

$$\lim_{t \rightarrow \infty} f(t+x)/f(t) = 1 \text{ locally uniformly and } \lim_{t \rightarrow \infty} f(t) = 0.$$

Set
$$\mu(A) = \int_A (1 - F(y))^{-1} dy \text{ for } A \text{ a Borel subset of } R. \text{ Then}$$

$$a_n 0(\cdot + b_n) = a_n \sum_{k=1}^{\infty} \varepsilon_{M_k} - b_n(\cdot) \Rightarrow \mu$$

in the sense of weak convergence of stochastic point processes.

For a discussion of weak convergence of point processes see Neveu, 1977, p. 282.

Proof: It suffices to show the Laplace functional of $a_n 0(\cdot + b_n)$ converges to that of μ . For g continuous with compact support we have

$$\begin{aligned} &E \exp \left\{ - \int g(x) a_n 0(dx + b_n) \right\} \\ &= E \exp \left\{ - \int a_n g(x - b_n) 0(dx) \right\} \end{aligned}$$

and from the proposition this equals

$$\begin{aligned} & \exp\left\{-\int \frac{1 - e^{-a_n g(x - b_n)}}{1 - e^{-a_n g(x - b_n)} F(x - b_n)} R(dx)\right\} \\ &= \exp\left\{-\int \frac{a_n^{-1}(1 - e^{-a_n g(y)})}{1 - e^{-a_n g(y)} F(y)} a_n R(dy + b_n)\right\}. \end{aligned}$$

We now use the fact that $a_n \rightarrow 0$, $a_n R(\cdot + b_n) \xrightarrow{y} m$ and obtain the convergence of the Laplace functionals to

$$\exp\left\{-\int g(y) \frac{dy}{1 - F(y)}\right\}$$

so the proof is complete.

Remark: It is easy to formulate a completely analogous version of this theorem in continuous time using the structure of extremal processes.

Cf. Resnick (1974).

Remark: An occupation time result for $a_n^{-1}(M_k - b_n)$, $k \geq 1$ is implied by the following:

$$\begin{aligned} E \exp\left\{-\int g(x) \sum_{k=1}^{\infty} \varepsilon_{(M_k - b_n)/a_n}(dx)\right\} &\rightarrow \\ \exp\left\{-\int \frac{1 - e^{-g(s)}}{1 - e^{-g(s)} F(s)} ds\right\}. \end{aligned}$$

This limit Laplace functional corresponds to the point process $\sum_{k=-\infty}^{\infty} \underline{n}_{k+1} \varepsilon_{\underline{t}_k}$ where $\{\underline{t}_k\}$ are the points of a homogeneous Poisson process on \mathbb{R} and (\underline{n}_k) are conditionally independent given $\{\underline{t}_k\}$ with

$$P[\underline{n}_{k+1} = n | \underline{t}_k] = F^{n-1}(\underline{t}_k)(1 - F(\underline{t}_k))$$

for $n \geq 1$. So we have

$$\sum_{k=1}^{\infty} \varepsilon_{(M_k - b_n)/a_n} \Rightarrow \sum_{k=-\infty}^{\infty} \underline{n}_{k+1} \varepsilon_{\underline{t}_k}.$$

An analogous result holds when $F \in D(\Phi_\alpha)$. Both results depend on the proposition and the fact that

$$F^n(a_n x + b_n) \rightarrow \Lambda(x) \quad (F^n(a_n x) \rightarrow \Phi_\alpha(x))$$

iff

$$R(a_n x + b_n) - \log n \rightarrow x \quad (R(a_n x) - \log n \rightarrow \alpha \log x),$$

References

1. Anderson, C. W. (1978). Super-slowly varying functions in extreme value theory. J. R. Statist. Soc. B. 40, 197-202.
2. Balkema, A. A. (1973). Unpublished manuscript.
3. Balkema, A. A. and Haan, L. de (1972). On R. Von Mises' condition for the domain of attraction of $\exp(-e^{-x})$. Ann. Math. Statist. 43, 1352-1354.
4. Breiman, L. (1968). Probability. Addison-Wesley, Menlo Park, California.
5. Darling, D. A. and Kac, M. (1957). On occupation times for Markoff processes. Trans. Amer. Math. Soc. 84, 444-458.
6. Feller, W. (1971). An Introduction to Probability Theory and its Applications. Wiley, New York.
7. Gnedenko, B. V. (1943). Sur la distributio limite du terme maximum d'une série aléatoire. Annals of Math. 44, 423-453.
8. Haan, L. de (1970). On Regular Variation and its Application to the Weak Convergence of Sample Extremes. Mathematical Centre Tract 32, Amsterdam.
9. Haan, L. de (1974). Equivalence classes of regularly varying functions. J. Stochastic Processes and their Applications. 2, 243-259.
10. Haan, L. de and Hordijk, A. (1972). The rate of growth of sample maxima. Ann. Math. Statist. 43, 1185-1196.
11. Hall, W. J. and Wellner, J. A. (1978). The rate of convergence in law of the maximum of an exponential sample. Preprint, University of Rochester.
12. Hall, P. (1978). J. Applied Probability, submitted.
13. Neveu, J. (1977). Processus Ponctuels. Ecole d'été de Probabilités de Saint-Flour VI-1976. Lecture Notes in Math. 598, Springer, Heidelberg.
14. Pickands, J. (1968). Moment convergence of sample extremes. Ann. Math. Statist. 39, 881-889.
15. Resnick, S. I. (1974) Inverses of extremal processes. Advances in Applied Probability 6, 392-406.
16. Shorrock, R. (1972). A limit theorem for inter-record times. J. Applied Probability, 9, 219-223.
17. Von Mises, R. (1936). La distribution de la plus grande de n valeurs. In: Selected Papers II, Amer. Math. Soc., 271-294.

LIST OF REPORTS 1979

- 7900 "List of Reprints, nos 220-230; Abstracts of Reports Second Half 1978.
- 7901/S "Motorists and Accidents (An Empirical Study)", by B.S. van der Laan.
- 7902/S "Estimation of the Minimum of a Function Using Order Statistics", by L. de Haan.
- 7903/S "An Abel and Tauber Theorem Related to Stochastic Compactness", by L. de Haan.
- 7904/E "Effects of Relative Price Changes on Input-Output Ratios - An Empirical Study for the Netherlands", by P.M.C. de Boer.
- 7905/O "Preemptive Scheduling of Uniform Machines Subject to Release Dates", by J. Labetoulle, E.L. Lawler, J.K. Lenstra and A.H.G. Rinnooy Kan.
- 7906/S "II-Regular Variation", by J.L. Geluk.
- 7907/O "Complexity Results for Scheduling Chains on a Single Machine", by J.K. Lenstra and A.H.G. Rinnooy Kan.
- 7908/E "Input-Output and Macroeconomic Forecasting Through the Generalized Inverse", by K.P. Vishwakarma.
- 7909/E "An Application of the Generalized Inverse in Input-Output and Macroeconomic Analysis", by K.P. Vishwakarma.
- 7910/O "A Numerical Comparison of Self Scaling Variable Metric Algorithms", by G. van der Hoek and M.W. Dijkshoorn.
- 7911/E "A Quadratic Engel Curve Demand Model (squaring with the representative consumer)", by J. van Daal and A.S. Louter.
- 7912/E "On Generalized Linear Demand Systems", by A.C.F. Vorst and J. van Daal.
- 7913/M "A Partial Survey of the Use of Algebraic Geometry in Systems and Control Theory", by M. Hazewinkel.
- 7914/M "(Fine) Moduli (Spaces) for Linear Systems: What are they and what are they good for", by M. Hazewinkel.
- 7915/S "A Simple Asymptotic Estimate for the Index of a Stable Distribution", by L. de Haan and S.I. Resnick.
- 7916/O "A Machine Maintenance Model with Application to Linear Programming", by A.H.G. Rinnooy Kan and J. Telgen.
- 7917/E "Inferential Procedures in Stable Distributions for Class Frequency Data on Incomes", by H.K. van Dijk and T. Kloek.
- 7918/M "On Families of Linear Systems: Degeneration Phenomena", by M. Hazewinkel.
- 7919/O "Complexity of Vehicle Routing and Scheduling Problems", by J.K. Lenstra and A.H.G. Rinnooy Kan.
- 7920 "List of Reprints, nos 231-240; Abstracts of Reports First Half 1979".
- 7921/E "Minimization by the Method of Conjugate Gradients", by V. Stern.
- 7922/S "Local Limit Theorems for Sample Extremes", by L. de Haan and S.I. Resnick.

