



*The World's Largest Open Access Agricultural & Applied Economics Digital Library*

**This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.**

**Help ensure our sustainability.**

Give to AgEcon Search

AgEcon Search

<http://ageconsearch.umn.edu>

[aesearch@umn.edu](mailto:aesearch@umn.edu)

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*

*No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.*

~~Netherlands school of economics~~  
ECONOMETRIC INSTITUTE

ON THE SNAPPER, LIEBLER - VITALE,  
LAM THEOREM ON PERMUTATION  
REPRESENTATIONS OF THE SYMMETRIC GROUP

M. HAZEWINKEL - T. VORST

Erasmus

GIANNINI FOUNDATION OF  
AGRICULTURAL ECONOMICS  
LIBRARY

MAY 7 1981  
WITHDRAWN

REPORT 8022/M

# ON THE SNAPPER, LIEBLER-VITALE, LAM THEOREM ON PERMUTATION REPRESENTATIONS OF THE SYMMETRIC GROUP.

Michiel Hazewinkel and Ton Vorst

ABSTRACT. Let  $\kappa = (\kappa_1, \dots, \kappa_m)$  be a descending partition of  $n$  and  $S_\kappa = S_{\kappa_1} \times S_{\kappa_2} \times \dots \times S_{\kappa_m}$  be the corresponding Young subgroup of  $S_n$ . Denote by  $\rho(\kappa)$  the representation of  $S_n$  which one gets by inducing the trivial representation of  $S_\kappa$ . If  $\lambda = (\lambda_1, \dots, \lambda_m)$  is another partition of  $n$  with  $\sum_{i=1}^r \kappa_i \geq \sum_{i=1}^r \lambda_i$  ( $\forall 1 \leq r \leq m$ ) then  $\rho(\kappa)$  is a subrepresentation of  $\rho(\lambda)$ . In this note we give an elementary complete direct proof of this fact.

## Contents

1. Introduction
  2. The Snapper, Liebler-Vitale, Lam Theorem
- References



ON THE SNAPPER, LIEBLER-VITALE, LAM THEOREM ON  
PERMUTATION REPRESENTATIONS OF THE SYMMETRIC GROUP

Michiel Hazewinkel and Ton Vorst

1. INTRODUCTION.

1. Let  $\kappa = (\kappa_1, \dots, \kappa_m)$ ,  $\kappa_i \in \mathbb{N} \cup \{0\}$ ,  $\kappa_1 \geq \kappa_2 \geq \dots \geq \kappa_m \geq 0$  be a descending partition of  $n$ . We identify partitions which differ only by the addition of some additional zero's. An ordering, which we call the specialization order, is defined on the set of all partitions by

$$(1.1) \quad \kappa > \lambda \iff \sum_{i=1}^r \kappa_i < \sum_{i=1}^r \lambda_i, \quad r = 1, 2, \dots$$

The reverse order has been called the dominance order. It occurs in many, seemingly unrelated parts of mathematics [1,2,3], and one of the central occurrences is in the representation theory of the symmetric groups in characteristic zero.

Let  $S_\kappa = S_{\kappa_1} \times \dots \times S_{\kappa_m}$  be the Young subgroup of  $S_n$  ( $S_{\kappa_i}$  is viewed as the permutation subgroup of  $S_n$  permuting the letters  $\kappa_1 + \dots + \kappa_{i-1} + 1, \dots, \kappa_1 + \dots + \kappa_i$ ) corresponding to the partition  $\kappa$  and let  $\rho(\kappa)$  be the representation of  $S_n$  obtained by inducing the trivial representation of  $S_\kappa$  up to  $S_n$ . Also let  $[\kappa]$  be the irreducible representation of  $S_n$  (in characteristic zero) associated to the partition  $\kappa$ . Snapper [5] proved that  $[\kappa]$  occurs in  $\rho(\lambda)$  implies  $\kappa < \lambda$  (this also follows readily from Young's rule) and conjectured the reverse, which he proved for  $m = 2$ . Proofs of the conjecture were given by Liebler-Vitale [4] and Lam [3]. Liebler and Vitale proved more precisely that  $\kappa < \lambda$  implies that  $\rho(\kappa)$  is a subrepresentation of  $\rho(\lambda)$  (which obviously implies, the conjecture because  $[\kappa]$  occurs in  $\rho(\kappa)$ ).

In this note we give a completely elementary direct proof of the Liebler-Vitale result which requires no representation theory at all (beyond the definition of the permutation representations  $\rho(\kappa)$ ) by constructing explicit homomorphisms of representations.

## 2. THE SNAPPER, LIEBLER-VITALE, LAM THEOREM.

### 2.1. Description of the permutation representation $\rho(\kappa)$ .

Let  $W(\kappa)$  be the set of all words of length  $n$  in the symbols  $a_1, \dots, a_m$  such that each  $a_i$  occurs exactly  $\kappa_i$  times. The group  $S_n$  acts in the obvious way on  $W(\kappa)$  ( $\sigma(b_1 \dots b_n) = b_{\sigma(1)} \dots b_{\sigma(n)}$ ,  $\sigma \in S_n$ ) and the vector-space  $V(\kappa)$  with the elements of  $W(\kappa)$  as basis and the action extended linearly is the representation  $\rho(\kappa)$ . We shall denote the elements of  $W(\kappa)$  and the corresponding basis elements of  $V(\kappa)$  with the same symbols.

2.2. "Reduction to the case  $m = 2$ ". It obviously suffices to prove the statement " $\kappa < \lambda \rightarrow \rho(\kappa)$  is a subrepresentation of  $\rho(\lambda)$ " in the case that  $\kappa < \lambda$  and  $\kappa < \mu < \lambda \Rightarrow \kappa = \mu$  or  $\lambda = \mu$ . In this case one easily shows that there exist  $i$  and  $j$ ,  $i > j$  such that  $\lambda_i = \kappa_i + 1$ ,  $\lambda_j = \kappa_j - 1$  and  $\lambda_r = \kappa_r$  for  $r \neq i, j$ . In this case we define a linear map

$$(2.3) \quad \beta_{\lambda, \kappa}: V(\lambda) \rightarrow V(\kappa)$$

by the formula

$$(2.4) \quad \beta_{\lambda, \kappa}(b_1 \dots b_n) = \sum b'_1 \dots b'_n$$

where the sum extends over all words  $b'_1 \dots b'_n$  such that  $b'_t = b_t$  for all but one  $t$ . And for that one  $t$  we have  $b_t = a_i$  and  $b'_t = a_j$ . I.e. the words in the sum on the right are obtained by replacing precisely one occurrence of  $a_i$  by  $a_j$ . This is obviously an  $S_n$ -equivariant map.

We shall prove that  $\beta_{\lambda, \kappa}$  is surjective if (and only if)  $\lambda > \kappa$ . This proves the theorem because the category of  $S_n$ -modules (in characteristic

zero) is semisimple. Alternatively observe that if  $\alpha_{\kappa, \lambda}: V(\kappa) \rightarrow V(\lambda)$  is defined as  $\beta_{\lambda, \kappa}$  with the letters  $a_j$  and  $a_i$  interchanged then  $\alpha_{\kappa, \lambda}$  and  $\beta_{\lambda, \kappa}$  are adjoint to each other in the sense that

$$(2.5) \quad \langle \alpha_{\kappa, \lambda} v, \omega \rangle = \langle v, \beta_{\lambda, \kappa} \omega \rangle, \quad v \in V(\kappa), \omega \in V(\lambda)$$

where the inner products on  $V(\lambda)$  and  $V(\kappa)$  are the ones for which  $W(\lambda)$  and  $W(\kappa)$  form orthonormal bases. This  $\alpha_{\kappa, \lambda}$  is an  $S_n$ -equivariant injection iff  $\beta_{\lambda, \kappa}$  is surjective and it remains to prove that  $\beta_{\lambda, \kappa}$  is surjective if  $\kappa < \lambda$ .

To do this observe that as a vectorspace  $V(\lambda)$  is the direct sum of  $\binom{n}{\lambda} \binom{\lambda_i + \lambda_j - 1}{\lambda_i}$  copies of  $V(\lambda_j, \lambda_i)$  indexed by all words in the symbols  $a_1, \dots, \hat{a}_j, \dots, \hat{a}_i, \dots, a_m, c$  ( $\hat{\phantom{a}}$  denotes deletion) such that  $a_t$  occurs  $\lambda_t$  times and  $c$  occurs  $\lambda_i + \lambda_j$  times. Similarly  $V(\kappa)$  is the direct sum of  $\binom{n}{\kappa} \binom{\kappa_i + \kappa_j - 1}{\kappa_i} = \binom{n}{\lambda} \binom{\lambda_i + \lambda_j - 1}{\lambda_i}$  copies of  $V(\kappa_j, \kappa_i)$  and the homomorphism

(2.4) maps the copies of  $V(\lambda_j, \lambda_i)$  and  $V(\kappa_j, \kappa_i)$ , labelled by the same word in  $a_1, \dots, \hat{a}_j, \dots, \hat{a}_i, \dots, a_m, c$ , into each other and is in fact the direct sum of these induced maps. Hence it is sufficient to prove the surjectivity of  $\beta_{\lambda, \kappa}$  in the case  $m = 2$ .

2.6. Proof of the surjectivity of  $\beta_{\lambda, \kappa}$  in the case  $m = 2$ . Let

$\lambda = (r-1, s+1)$ ,  $\kappa = (r, s)$ ,  $r + s = n$  and write  $x$  for  $a_j$  and  $y$  for  $a_i$ .

Then  $W(r-1, s+1)$  consists of words of length  $n$  in  $(r-1)$   $x$ 's and  $(s+1)$   $y$ 's and  $\beta = \beta_{\lambda, \kappa}$  changes such a word into the sum of all words which can be obtained from this word by changing precisely one  $y$  into an  $x$ . E.g.

$$(2.7) \quad \beta(xxyyyy) = xxxxyy + xxxxyx + xxxyyx$$

We shall now show that  $\beta$  is surjective if  $r \geq s+1$  (We only need the case  $r \geq s+2$ ). Let  $W = W(r-1, s+1) \cup W(r, s)$ . For each pair  $\omega_1 = b_1 \dots b_n$ ,

$\omega_2 = b_1' \dots b_n'$  in  $W$  we define the distance  $d(\omega_1, \omega_2)$  by

$$(2.8) \quad d(\omega_1, \omega_2) = \# \{t | b_t \neq b_t'\}.$$

(This distance is called Hamming distance in coding theory).

Now for  $\omega_0 = x \dots xy \dots y \in W(r, s)$  let

$$(2.9) \quad E_t = \{\omega \in W | d(\omega_0, \omega) = t\}$$

Then  $E_t \subset W(r, s)$  if  $t$  is even and  $E_t \subset W(r-1, s+1)$  if  $t$  is odd.

Note that  $\omega \in E_{2t}$  iff there are precisely  $t$   $y$ 's among the first  $r$  letters and  $t$   $x$ 's among the second  $s$  letters and similarly  $\omega \in E_{2t+1}$  iff there are precisely  $t+1$   $y$ 's among the first  $r$  letters of  $\omega$  and  $t$   $x$ 's among the last  $s$  letters.

Now let

$$(2.10) \quad f = r^{-1} (c_0 \sum_{\omega \in E_1} \omega + c_1 \sum_{\omega \in E_3} \omega + \dots + c_s \sum_{\omega \in E_{2s+1}} \omega)$$

where

$$(2.11) \quad c_t = (-1)^t \binom{r-1}{t}^{-1}$$

We claim that  $\beta(f) = \omega_0$ . To see this observe that since  $\omega \in W(r, s)$  and  $r \geq s+1$  the maximum distance of a  $\omega \in W$  to  $\omega_0$  is  $2s+1$ . Observe that if  $\omega' \in E_{2t+1}$  then  $\beta(\omega')$  is a sum of elements in  $E_{2t}$  and  $E_{2t+2}$  (except when  $t = s$ , then only elements of  $E_{2s}$  can occur by the maximum distance observation).

Now let  $\omega'' = b_1 \dots b_n \in E_{2t}$  ( $t \geq 1$ ) then the coefficient of  $\omega'' \in \beta(f)$  is equal to

$$(2.12) \quad r^{-1} c_t (\# \{i \in \{1, \dots, r\} | b_i = x\}) + r^{-1} c_{t-1} (\# \{i \in \{r+1, \dots, r+s\} | b_i = x\}) =$$

$$r^{-1} c_t (r-t) + r^{-1} c_{t-1} t.$$

(The first contribution comes from the elements in  $E_{2t+1}$  whose  $i$ -th element was  $y$  and is transformed to  $x$  to decrease the distance to  $\omega_0$ ; the second contribution comes from elements of  $E_{2t-1}$  whose  $i$ -th element was  $y$  and is transformed to  $x$  to increase the distance).

By definition of  $c_t$  the right-hand side of (2.12) is zero.

The coefficient of  $\omega_0 \in \beta(f)$  is equal to

$$(2.13) \quad r^{-1} c_0 (\#\{i \in \{1, \dots, r\} | b_i = x\}) = r^{-1} \cdot 1 \cdot r = 1.$$

This proves that  $\omega_0 = \beta(f) \in \text{Im} \beta$  and hence  $\omega \in \text{Im} \beta$  for all  $\omega \in W(r, s)$  because  $\beta$  is  $S_n$ -equivariant and  $S_n$  acts transitively on  $W(r, s)$ . This concludes the proof.

#### REFERENCES

1. T. Brylawsky, The Lattice of Integer Partitions, Discrete Math. 6(1973), 201-219.
2. M. Hazewinkel and C. Martin, Representations of the Symmetric Groups, the Specialization Order, Systems and Grassmann Manifolds, In preparation.
3. T.Y. Lam, Young Diagrams, Schur Functions, the Gale Ryser Theorem and a Conjecture of Snapper, J. Pure Appl. Algebra 10(1977) 81-94.
4. R.A. Liebler and M.R. Vitale, Ordering the Partition Characters of the Symmetric Group, J. Algebra 25(1973) 487-489.
5. E. Snapper, Group Characters and Nonnegative Integral Matrices, J. Algebra 19(1971) 520-535.

Dept. of Mathematics  
Erasmus University Rotterdam  
P.O. Box 1738,  
3000 DR ROTTERDAM. The Netherlands



# LIST OF REPORTS 1980

- 8000 "List of Reprints, nos 241-260, Abstracts of Reports Second Half 1979".
- 8001/O "A Stochastic Method for Global Optimization", by C.G.E. Boender, A.H.G. Rinnooy Kan, L. Stougie and G.T. Timmer.
- 8002/M "The General Linear Group of Polynomial Rings over Regular Rings", by A.C.F. Vorst.
- 8003/O "A Recursive Approach to the Implementation of Enumerative Methods", by J.K. Lenstra and A.H.G. Rinnooy Kan.
- 8004/E "Linearization and Estimation of the Add -Log Budget Allocation Model", by P.M.C. de Boer and J. van Daal.
- 8005/O "The Complexity of the Constrained Gradient Method for Linear Programming", by J. Telgen.
- 8006/S "On Functions with Small Differences", by J.L. Geluk and L. de Haan.
- 8007/O "Analytical Evaluation of Hierarchical Planning Systems", by M.A.H. Dempster, M.L. Fisher, L. Jansen, B.J. Lageweg, J.K. Lenstra and A.H.G. Rinnooy Kan.
- 8008/S "Looking at Multiway Tables (continued)", by A.P.J. Abrahamse and W.M. Lammerts van Bueren.
- 8009/O "An Introduction to Multiprocessor Scheduling", by J.K. Lenstra and A.H.G. Rinnooy Kan.
- 8010/M "On Families of Systems: Pointwise-Local-Global Isomorphism Problems, by M. Hazewinkel and A.M. Perdon.
- 8011/M "Proceedings Filter-Day Rotterdam 1980 (New Trends in Filtering and Identification of Stochastic Systems, 23 jan. 1980), by M. Hazewinkel (ed).
- 8012/E "Further Experience in Bayesian Analysis Using Monte Carlo Integration" by H.K. van Dijk and T. Kloek.
- 8013/M "Note on the Eigenvalues of the Covariance Matrix of Disturbances in the General Linear Model, II", by R.J. Stroeker.
- 8014/M "Factorization Methods for Solving Diophantine Equations", by R.J. Stroeker.
- 8015 "List of Reprints, nos. 261-270, Abstracts of Reports First Half 1980".
- 8016/S "Renewal Functions and Regular Variation", by J.B.G. Frenk.



- 8017/S "Nominal Variables: Dependence and Predictive Power", by  
A.P.J. Abrahamse and W.M. Lammerts van Bueren.
- 8018/O "Scheduling Subject to Resource Constraints: Classification and  
Complexity", by J. Blazewics, J.K. Lenstra and A.H.G. Rinnooy Kan.
- 8019/M "On Lie Algebras and Finite Dimensional Filtering", by M. Hazewinkel.
- 8020/E "Optimal Household Behaviour under Taxation - A Two-Stage Dynamic  
Programming Approach", by V. Stern.
- 8021/ESM "The Certainty Equivalence Property in Stochastic Control Theory",  
by H. v.d.Water and J.C. Willems.
- 8022/M "On the Snapper, Liebler-Vitale, Lam Theorem on Permutation  
Representations of the Symmetric Group", by M. Hazewinkel and  
A.C.F. Vorst.