



The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

Papers downloaded from AgEcon Search may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.

No endorsement of AgEcon Search or its fundraising activities by the author(s) of the following work or their employer(s) is intended or implied.

Erasmus University Rotterdam. Econometric Institute

A Note on a Stochastic Location Problem

==

J.B.G. Frenk

M. Labb 

S. Zhang

Econometric Institute, Erasmus University Rotterdam

Abstract

In this note we give a short and easy proof of the equivalence of Hakimi's one-median problem and the K -server-single-facility-loss median problem as discussed by Chiu and Larson (cf. [1]). The proof makes only use of a stochastic monotonicity result for birth and death processes and the insensitivity of the $M|G|K$ loss model.

WITHDRAWN
GIANNINI FOUNDATION OF
AGRICULTURAL ECONOMICS
LIBRARY

JUN 5 1991

1. Model formulation and results

Consider the K -server-single-facility-loss median model (K -SFLM) as discussed by Chiu and Larson (cf. [1],[2]) and define

$\underline{n}_t^{(K)}(\mathbf{x}) :=$ number of customers being served at time t by one of the servers if the facility is located at $\mathbf{x} \in F$

where the set $F \subseteq \mathbb{R}^2$ denotes some feasible region. As argued in [1] the queueing process underlying the K -SFLM location problem is a $M|G|K$ loss-model. For this queueing process it is well-known (cf. [3]) that for fixed $\mathbf{x} \in F$ $\underline{n}_t^{(K)}(\mathbf{x})$ converges in distribution to some random variable $\underline{n}_\infty^{(K)}(\mathbf{x})$ representing the number of customers being served in steady state if the location is $\mathbf{x} \in F$. Moreover, the distribution of $\underline{n}_\infty^{(K)}(\mathbf{x})$ only depends on the first moment $m_1(\mathbf{x})$ of the service time (Erlang's Loss Formula). If the facility is located at $\mathbf{x} \in F$ the costfunction $Z(\mathbf{x})$ takes the following form (cf. [1])

$$Z(\mathbf{x}) = P_K(\mathbf{x})Q + (1 - P_K(\mathbf{x}))m_1(\mathbf{x}) \quad (1)$$

with

$P_K(\mathbf{x}) := P\{\text{customer "arriving" in steady state at location } \mathbf{x} \text{ is lost}\}$

$m_1(\mathbf{x}) :=$ expected service time of arbitrary customer whenever facility is located at \mathbf{x}

and

$Q :=$ cost per lost customer, $Q \geq 0$.

The main result proved in [1] using lengthy calculations states that the costfunction $Z(\mathbf{x})$ is increasing in $m_1(\mathbf{x})$. This implies that the K -SFLM location problem is solved by determining that location $\mathbf{x} \in F$ which minimizes the expected service time $m_1(\mathbf{x})$. Hence, in the special case where F denotes some network N this boils down to finding the so-called Hakimi median (cf. [5]) at one of the nodes of N .

The above result can be verified easily without any calculations by using a well-known stochastic monotonicity result for birth and death processes. Before proving this we need the following observations. By Little's formula $(1 - P_K(x)m_1(x))$ equals $\frac{1}{\lambda} \mathcal{E}(\underline{n}_\infty^{(K)}(x))$ with λ the arrival rate of the Poisson process. Moreover, by the PASTA property (Poisson Arrivals See Time Averages)

$$P_K(x) = P\{\underline{n}_\infty^{(K)}(x) = K\}.$$

Hence by (1)

$$Z(x) = QP\{\underline{n}_\infty^{(K)}(x) = K\} + \frac{1}{\lambda} \mathcal{E}(\underline{n}_\infty^{(K)}(x)) \quad (2)$$

Lemma 1.

Let $x, x^* \in F$. If $m_1(x) \leq m_1(x^*)$ we obtain $Z(x) \leq Z(x^*)$.

Proof.

In order to prove $Z(x)$ is increasing in $m_1(x)$ it is sufficient to verify that $\mathcal{E}(\underline{n}_\infty^{(K)}(x)) \leq \mathcal{E}(\underline{n}_\infty^{(K)}(x^*))$ and $P\{\underline{n}_\infty^{(K)}(x) = K\} \leq P\{\underline{n}_\infty^{(K)}(x^*) = K\}$ whenever $m_1(x) \leq m_1(x^*)$.

By the sensitivity of the $M|G|K$ loss system with respect to only the first moment $m_1(x)$ this breaks down to prove the above inequalities for the Markovian loss models $\Sigma_1 = M|M_1|K$ and $\Sigma_2 = M|M_2|K$, i.e. loss models with the same Poisson arrival process and negative exponential service times with parameters $\mu_1(x) = (m_1(x))^{-1}$ for Σ_1 and $\mu_1(x^*) = (m_1(x^*))^{-1}$ for Σ_2 . Denote by $\underline{n}_t^{(K)}(\Sigma_i)$ the number of customers in system Σ_i , $i = 1, 2$, at time t , $0 < t < \infty$ and by $\underline{n}_\infty^{(K)}(\Sigma_i)$, $i = 1, 2$ the corresponding limiting random variables.

Clearly these stochastic processes are birth and death processes on the finite state space $\{0, 1, \dots, K\}$ with transition rates $q_{ii+1} = \lambda$, $i = 0, \dots, K-1$, $q_{ii-1} = i\mu_1(x)$, $i = 1, \dots, K$ for Σ_1 and $q_{ii+1} = \lambda$, $i = 0, \dots, K-1$, $q_{ii-1} = i\mu_1(x^*)$, $i = 1, \dots, K$ for Σ_2 .

Since $\mu_1(x) \geq \mu_1(x^*)$ we obtain by a well-known stochastic monotonicity result for birth and death processes (cf. [4, prop.4.2.10]) that for every $t > 0$

$$\underline{n}_t^{(K)}(\Sigma_1) \leq \underline{n}_t^{(K)}(\Sigma_2) \quad d$$

This implies by a limit procedure $\mathcal{E}_{\underline{n}_{\infty}^{(K)}}(\Sigma_1) \leq \mathcal{E}_{\underline{n}_{\infty}^{(K)}}(\Sigma_2)$ and $P\{\underline{n}_{\infty}^{(K)}(\Sigma_1) = K\} \leq P\{\underline{n}_{\infty}^{(K)}(\Sigma_2) = K\}$. Hence the desired result is proved. □

References

- [1] S.S. Chiu, R.C. Larson, *Locating an n-server facility in a stochastic environment*, Comput. & Ops. Res., Vol. 12, No. 6, 509-516, 1985.
- [2] P. Mirchandani, R. Francis, *Discrete location theory*, Wiley, New York,
- [3] J.W. Cohen, *On regenerative processes in Queueing theory*, Lect. Notes Econ. Math. Syst. 121, Springer Verlag, Berlin, 1976.
- [4] D. Stoyan, *Comparison methods for queues and other stochastic models*, Wiley, New York, 1983.
- [5] S.L. Hakimi, *Optimum location of switching centers and the absolute centers and medians of a graph*, Operations Research 12, 450-459, 1964.

LIST OF REPORTS 1989

8900 Publications of the Econometric Institute Second Half 1988:
List of Reprints 515-525.

8901/C A.G. Oerlemans, H.W. van den Meerendonk en G.W. de Wit, "Een onderzoek naar verschillende premiestructuren in de ziektekostenverzekering", 35 pagina's.

8902/A J. van Dalen, J. Koerts and A.R. Thurik, "An analysis of labour productivity wholesaling", 29 pages.

8903/A O.E. Flippo and M. Minoux, "Continuity of the value-function in mathematical programming: a unifying framework with new results", 39 pages.

8904/A N.L. Van der Sar and B.M.S. Van Praag, "Social stratification: individual importance and social welfare", 21 pages.

8905/A L. Pintelon and L.N. Van Wassenhove, "A maintenance management tool", 26 pages.

8906/A C.N. Potts and L.N. Van Wassenhove, "Single tardiness sequencing heuristics", 25 pages.

8907/A C.N. Potts and L.N. Van Wassenhove, "Approximation algorithms for scheduling a single machine to minimize total late work", 13 pages.

8908/A P. Billington, J. Blackburn, J. Maes, R. Millen and L.N. Van Wassenhove, "Multi-product scheduling in capacitated multi-stage serial systems", 30 pages.

8909/A S.L. Hakimi, M. Labb  and E.F. Schmeichel, "The Voronoi partition of a network and its implications in location theory", 17 pages.

8910/A J.L. Geluk, "On conjugate II-variation and the coefficients of power series", 9 pages.

8911/B H. Bart, G.Ph.A. Thijssse, "Similarity invariants for pairs of upper triangular Toeplitz matrices", 40 pages.

8912/A A.R. Thurik and W.H.M. van der Hoeven, "Manufacturing margins: differences between small and large firms", 10 pages.

8913/A R. de Zwart and B.M.S. van Praag, "Earnings of Dutch adolescents", 22 pages.

8914/C R. de Zwart en I. Sevinga, "Schonings- en wegingsprocedures van een scholierenonderzoek", 41 pagina's.

8915 Publications of the Econometric Institute first half 1989.

8916/A Shuzhong Zhang, "On vertices adjacency characterization and optimization of polymatroids", 16 pages.

8917/A Shuzhong Zhang, "On the convergence property of Iri-Imai's method for linear programming", 18 pages.

8918/C M. Warnaar en R. de Zwart, "Uitspraken van middelbare scholieren", 57 pagina's.

8919/A J.B.G. Frenk, M. Labb  and S. Zhang, "A note on a stochastic location problem", 4 pages.

A. Economics, Econometrics and Operations Research
B. Mathematics
C. Miscellaneous

