



AgEcon SEARCH
RESEARCH IN AGRICULTURAL & APPLIED ECONOMICS

The World's Largest Open Access Agricultural & Applied Economics Digital Library

This document is discoverable and free to researchers across the globe due to the work of AgEcon Search.

Help ensure our sustainability.

Give to AgEcon Search

AgEcon Search
<http://ageconsearch.umn.edu>
aesearch@umn.edu

*Papers downloaded from **AgEcon Search** may be used for non-commercial purposes and personal study only. No other use, including posting to another Internet site, is permitted without permission from the copyright owner (not AgEcon Search), or as allowed under the provisions of Fair Use, U.S. Copyright Act, Title 17 U.S.C.*



The Role of Forestry in Carbon Sequestration in General Equilibrium Models *

by

Brent Sohngen¹, Alla Golub² and Thomas W. Hertel³

GTAP Working Paper No. 49
2008

¹Department of Agricultural, Environmental, and Development Economics, Ohio State University, 2120 Fyffe Rd., Columbus, OH 43210, Contact: Sohngen.1@osu.edu

²Center for Global Trade Analysis, Purdue University, West Lafayette, IN 47907: golub@purdue.edu

³ Center for Global Trade Analysis, Purdue University, West Lafayette, IN 47907: Hertel@purdue.edu

*Chapter 11 of the forthcoming book *Economic Analysis of Land Use in Global Climate Change Policy*, edited by Thomas W. Hertel, Steven Rose, and Richard S.J. Tol

Table of Contents

1. Introduction	3
2. The Economics of Forest Management Under Carbon Policies	5
3. Examples From a Global Timber Model.....	16
4. Implications for CGE Modeling.....	22
5. Evaluation of Current Approaches to Forestry Modeling in Global CGE Analysis ..	27
6. Conclusion.....	30
7. References	33

Table 1. Carbon intensity in temperate zone forests (t C/hectare), expressed relative to the baseline carbon intensity (1.0 = no change)	35
Table 2. Annual equivalent carbon gains (losses) and production impacts over 20 years at three different carbon prices	36

Figure 1. Proportion of carbon sequestration in temperate and tropical forests resulting from forestry actions for several different carbon prices over a 20 and 100 year period	37
Figure 2. Biomass and Merchantable timber yield functions for upland hardwoods in the U.S. south	38
Figure 3. Growth rate and average annual flow of timber from upland hardwood type in the Southern U.S.....	39
Figure 4. Present value of upland hardwoods in U.S. South under baseline (no carbon incentive) and three alternative carbon prices (derived from equation 5), assuming management inputs remain \$56/hectare.....	40
Figure 5. Relationship between \$/hectare management inputs, "n", and stocking density (delta, or δ).....	41
Figure 6. Age class distribution of upland hardwood stands in the Southern U.S. in the baseline and climate mitigation scenario (high damage) from Sohngen and Mendelsohn (2006).....	42
Figure 7. Aboveground carbon stored in natural southern pine stands in the United States in the baseline and the high damage carbon mitigation scenario	43
Figure 8. Timber price index for baseline and high damage carbon policy scenario, derived from Sohngen and Mendelsohn (2007).....	44
Figure 9. Annual timber harvests in southern upland hardwood stands in the United States in the baseline and the high damage carbon mitigation scenario.....	45

THE ROLE OF FORESTRY IN CARBON SEQUESTRATION IN GENERAL EQUILIBRIUM MODELS

Brent Sohngen, Alla Golub and Thomas W. Hertel

1. INTRODUCTION

The role of forestry in the global carbon balance is widely recognized, as described in chapter 3¹. A number of models and methods have now been developed to assess the potential role of forestry in climate mitigation policy (e.g., Adams et al., 1999; Stavins, 1999; Sohngen and Mendelsohn, 2003; Richards and Stokes, 2004). These studies suggest that forestry could provide 1.0 – 2.0 Pg C/yr in sequestration services or avoided emissions through reductions in deforestation for \$150 - \$200/ t C, amounting to potentially one-third of efficient global carbon abatement (Sohngen and Mendelsohn, 2003). Given that these prices are well within the range of expected prices associated with fairly stringent carbon abatement policies (see Weyant et al., 2006), current economic analysis suggests that forestry can compete effectively with other mitigation options. However, most of these studies of forestry's role in climate change policy have been partial equilibrium analyses that have ignored the broader implications of climate policy on economic activity. As this book suggests, it is useful to develop general equilibrium modeling frameworks that capture the full range of implications of climate policies on carbon markets and consequently forest carbon sequestration activity.

The chapters in this book show that the state of the art in modeling forestry activities within a general equilibrium context has advanced significantly in recent years. Modeling carbon sequestration in forests within a general equilibrium context, however,

¹ GTAP Working Paper No. 41

remains a formidable task because of the complex dynamic considerations inherent in the economics of managing and modeling forests. While other land uses, such as agriculture, may be modeled within the context of a comparative static, single representative time period, or perhaps in the context of a recursive, dynamic model, forest management involves investments in one time period based on expectations about what will happen many years ahead. In terms of carbon sequestration, the carbon or timber benefits of expenditures on forestry may not accrue for a number of years, presenting difficulties for conducting static (single period) analysis in computable general equilibrium (CGE) models. If CGE modelers instead utilize recursive models, forestry remains difficult to incorporate because of difficulties in capturing the intertemporal management of forest carbon.

In order to provide an economic basis for incorporating forestry management and carbon sequestration into CGE models, this chapter explores the intertemporal forestry management problem in some depth, thereupon providing information and insights that may aid CGE modelers in developing improved representations of forestry in the context of global climate policy analysis. The chapter begins with a description of the forestry management problem, and illustrates how carbon sequestration incentives influence forest management. The results of a global forestry model are then used to show the importance of capturing the management effects in particular regions. The chapter then briefly discusses the influence of other management options on forest carbon. Since one of the most important effects of carbon policy on forestry may be changes in optimal rotation lengths, one section of the chapter describes how CGE modelers may incorporate

short and longer term adjustments in age classes in their models. A brief evaluation of how the models in this book capture forestry concludes the chapter.

2. THE ECONOMICS OF FOREST MANAGEMENT UNDER CARBON POLICIES

Modeling carbon sequestration in forestry requires the assessment of changes in management, rotation ages, set-asides, forest products, land use. Each of these concepts requires some further clarification. Changes in forestry management include actions like thinning, fertilizing, herbicide treatments, pruning, planting, and replanting. Changes in rotation ages are also a form of management change, but are usually considered separately from other management actions. Set-asides refer to removing timberland from the active forest management base and holding the land as a carbon reserve. Forest product storage refers to the storage of carbon in marketed products. For example, many sawnwood products will ultimately be used in construction (e.g., to build houses), or in furniture. These products have long-lives, and during their lives the carbon in the wood products is with-held from the atmosphere. Finally, changes in land use, including afforestation (planting lands that were previously in agriculture) and reductions in deforestation (keeping lands in forests that would otherwise convert to agriculture) are two potential options for carbon sequestration that have been widely discussed in the policy arena.

The economic efficiency of choosing among these sequestration options in any particular region of the world will depend on the relative costs of implementing the actions versus the benefits (e.g., carbon sequestered). Both costs and carbon intensities

vary substantially across regions. Chapter 3² provides information on both the biological growth and the economic parameters that can be used to assess forestry actions in different regions. To get a sense for the relative size of carbon storage available from different types of activities, estimates of the proportion of carbon sequestered or emissions reduced across different activities in temperate and tropical forests for several carbon prices are shown in Figure 1.³ These estimates are derived from the model used in Sohngen and Mendelsohn (2007), which determines an "optimal" allocation of carbon sequestration in different activities and time periods given a constant carbon price over a 20 year, and 100 year period, respectively.

As can be seen in Figure 1 (panels A & B), in the short run (i.e., over the next 20 years), most carbon sequestered in temperate forests will be obtained through extending rotation ages, changing management, and setting aside forests. By contrast, in tropical forests, nearly all the short run carbon sequestration is obtained from reduced deforestation. In the longer-run (Figure 1, panels C & D), afforestation plays a more important role in temperate zones, although in terms of annual equivalent amounts, it remains less than 30% of total temperate forest sequestration for the range of carbon prices considered. Afforestation, and in particular reductions in deforestation, remain dominant components of overall carbon sequestration activities in tropical forests, even over the long run.

The example in Figure 1 illustrates the importance of understanding how carbon is sequestered in forests and through forestry activities when implementing forest carbon

² GTAP Working Paper No. 41

³ Reductions in carbon emissions from avoided deforestation are described as "emissions reductions" in this paper, while increased carbon stored in forests through rotation changes, management changes, afforestation, etc. is described as "carbon sequestration" in this paper.

sequestration in CGE models. Given the relative importance of age class management, the results suggest that it is as important to model what happens to the distribution of age classes (e.g., the rotation age) as it is to model what happens to the total area of forest land. To assess the implications of carbon prices on forest rotation ages and forest management, this section considers these two activities in the context of a simple, single hectare, forestry management model. Later in this chapter, we use a global timber model to expand these results to the regional level.

Modeling forestry rotation ages and management begins with modeling the yield of biomass and merchantable forestry products. Forest yield functions have the typical logistical shape identified with most biological species. Equation (1) presents the biomass growth function for upland hardwoods in the Southern U.S. region, which translates the age of the stand into the cubic meters of biomass on the site:

$$(1) V_a^C \text{ (m}^3\text{/ha)} = \delta * \exp[5.2 - 30/(\text{age})]$$

The parameter δ captures stocking density, which will typically range from 0.6 to 1.4, depending on average management inputs in a region. Stocking density is a unitless parameter that adjusts the yield function for average conditions of specific sites or regions. Stocking density can be assumed to be given exogenously, or it can be assumed to be an endogenous function of management inputs. Initially, we assume that stocking density, δ , is exogenous.

As forests age, the value of the merchantable component of the stock increases. Bigger stems improve stand quality because more material can be utilized for sawtimber,

which is a higher value use of the material. To account for this improvement in value over time, we utilize an additional function to depict the proportion of merchantable timber on a given site (once the forest reaches the commercially viable vintage):

$$(2) \gamma(\text{age}) = 0.85 * [1 - \exp(1 - 0.1 * (\text{age} - 30))]^2$$

The merchantable yield for this southern upland hardwood forest type becomes:

$$(3) \quad \text{if age} \leq 30 \Rightarrow 0$$

$$\text{If age} > 30 \Rightarrow V_a^M (\text{m}^3/\text{ha}) = \gamma(\text{age}) * \delta * \exp[5.2 - 30/(\text{age})]$$

Figure 2 shows biomass yield and merchantable yield for particular forest type. As can be seen in equation (2), merchantable yield can be at most 85% of the biomass in growing stock on the site. Furthermore, with equation (3), we have assumed that there is no merchantable yield before age 30. These are both strong assumptions which may be violated at particular sites where timber operations occur, however, when aggregated across a large number of sites (i.e., a region), the assumption is more realistic.

In a static world, the harvest age occurs where the net present value of the stand is maximized. The net present value of a single rotation when prices are held constant is:

$$(4a) \quad NPV^T = (P^{QA}) * (V_a^M) (1 + r)^{-a} - C,$$

and the net present value of an infinite series of rotations when prices are held constant is:

$$(4b) \quad NPV^\infty = \frac{(P^{QA}) * (V_a^M)(1+r)^{-a} - C}{(1 - (1+r)^{-a})}$$

In equations (4a) and (4b), P^{QA} is the quality adjusted stumpage price for timber, r is the interest rate, C is the planting or regeneration cost, and " a " is the rotation age. Planting costs (investments) occur at the beginning of the rotation, and the benefits of harvests occur when trees are harvested at time " a ". The quality adjustment factor accounts for any quality differences associated with particular timber types that are found in local or national price data. The rotation age can be determined endogenously by maximizing (4) with respect to " a ". If timber prices are \$100/m³, replanting costs are \$56/hectare, $\delta=1.15$, and $r=0.05$, the optimal rotation age in this case will be $a = 48$ years, the biomass yield will be 112 m³/ha and the merchantable yield will be 66 m³ per hectare. The net present value of the stand will be \$642/hectare. For simplicity, assume that the rotation age is rounded to 50 years old. If there are 34 million hectares of this type of land, then on average 0.68 million hectares will be cut each year, or 45 million m³ per year (0.68*66).

A number of important relationships can be analyzed with the yield function and net present value formula shown in equations (1) – (4). First, the optimal rotation age occurs approximately at the point in time when the value of the stand, $(P^{QA})*(V_a^M)$, is growing at the rate of interest, which translates (with constant prices) into the time when the stand itself is growing at the rate of interest. Since the interest rate in this case is 5%, the stand is growing at approximately 5% per year at 50 years, which is the optimal rotation length. Note that this optimal rotation period is not the same time interval as that

which maximizes the flow of timber from the land. That age, often called "maximum sustainable yield," occurs later, as shown in figure 3, which displays the annual percentage growth rate and the annual average flow of timber from this stand of forest. The annual average flow of timber is calculated as the merchantable yield divided by the age. Maximum sustainable yield in this type of forest occurs at age 60, or 10 years later than the optimal rotation age. The profit maximizing rotation period is shorter due to the opportunity cost of forestry investment, as reflected in the discount rate.

For further insight, let us consider the case wherein this type of forest is harvested every 60 years. The merchantable yield of a 60 year old stand, by equation (3) above is 97 m^3 per hectare. If rotations are 60 years in length, then 0.57 million hectares of the 34 million hectares will be harvested each year. This is fewer hectares than with the 50 year rotations, but more timber is harvested on each hectare (97 m^3 for the 60 year old stand versus 66 m^3 for the 50 year old stand). By waiting 10 years, one has obtained 53% more merchantable timber, and the area harvested each year has only gone down by 16% (0.68 million hectares/yr down to 0.57 million hectares/yr). The annual flow of timber could be 55 million m^3 per year (0.57×97) if harvested at the maximum sustainable yield age of 60 years. But commercial forests that are not harvested at maximum sustained yield because the additional 10 years of waiting impose substantial opportunity costs. Once these opportunity costs are taken into account, one will move to shorter rotations.

There a number of reasons why managers might move towards harvesting older age classes. Some governments, for instance, impose minimum rotation ages that are above the economically optimal rotation ages. There may be several explanations for this. Countries that have extensive government holdings may be interested in

maximizing their annual revenues rather than their land value, so the contracts they sign with harvesters are likely to emphasize maximizing annual harvests. Alternatively, countries may choose to impose environmental constraints that lead to longer rotation ages on government or private lands. From an economic perspective, harvesting stands in older rotation ages can make sense if there are non-market benefits associated with older aged stands (see, for example, Hartman, 1979). A different rationale for lengthening the rotation period relates to interest rates. Binkley (1987) has shown that harvests move towards maximum sustained yield when interest rates fall. So if the government believes that the discount rate used by the forest managers is too high, they might wish to lengthen the rotation period. Carbon sequestration offers another reason for lengthening rotations, since the sequestration benefits increase in direct proportion to the yield of timber (Van Kooten et al., 1995). Indeed, this is the dimension of optimal rotation length that is of greatest interest in the context of the present chapter.

When considering the value of forests used for timber and carbon, the present value formula in equation (4) above is adjusted as follows:

$$(5) \quad NPV^C = \frac{(P^{QA}) * (V_a^M)(1+r)^{-a} - C + rP^C \sum_0^a \alpha V_a^C (1+r)^{-a} + 0.3P^C \alpha V_a^M (1+r)^{-a}}{(1 - (1+r)^{-a})}$$

The numerator in equation (5) is the present value of a single rotation of forests where carbon is valued and the harvest age is given by "a"). The denominator accounts for repeating this rotation over an infinite horizon. The first component of the numerator,

$(P^{QA}) * (V_a^M)(1+r)^{-a}$, is the present value of timber benefits where P^{QA} is the quality adjusted price of timber stumpage, V_a^M is the merchantable timber harvested at age "a", and $(1+r)^{-a}$ is the discount factor, where "r" is the discount rate. The second term, C, is the cost of replanting. The third term, $rP^C \sum_0^a \alpha V_a^C (1+r)^{-a}$, is the rental value of holding carbon during the rotation. P^C is the carbon price, and rP^C is the carbon rental value. Note that we assume carbon prices are constant in this case. The term α is in Mg C per m^3 of wood. V_a^C is the biomass yield function described above. The final term is the value of permanent storage of carbon in harvested timber products. For this analysis, we have assumed that 30% of wood is permanently stored. This present value formula will allow modelers to determine the value of land in which produces jointly commercial forest products as well as carbon storage. If this value outweighs the opportunity costs of using the land for agricultural products, then in principle, the land should shift to forestry.

Carbon incentives have a number of impacts on forestry activity. First, they affect the value of land in forests. Figure 4 shows the net present value (bare land value) of forestland in the baseline (no carbon prices) and three carbon prices: $P^C = \$50, \$100,$ and $\$200$ per t C.⁴ The value of land in forests rises from $\$642/\text{ha}$ in the baseline, to $\$4859/\text{ha}$ when carbon prices are $\$200/\text{t C}$. At $\$4859/\text{ha}$, forests can compete strongly with other uses, such as crops. Of course, these values apply only to upland hardwoods, which have fairly high carbon intensity per hectare (Sohngen and Brown, 2006), but nonetheless, high carbon values would have strong impacts on how much land is maintained under forest cover. Second, carbon prices extend rotation ages. The rotation

⁴ Note that for this analysis we have assumed that management inputs are held constant. As shown below, management inputs would change with carbon policy, and so one would expect optimal responses to differ from the examples with carbon prices in this section.

age is increased from 50 to 60 years of age with the \$200/t carbon incentive. This can effect is also evident in figure 4 wherein the maximum land value shifts to the right as carbon prices rise.

In summary, carbon incentives have strong implications for the value of land in forests and non-negligible implications for rotation ages. Each timber type (and landowner or manager) will react differently to carbon prices, depending on the shape of the growth function, interest rates, timber prices, costs of regenerating, and carbon prices. Higher carbon prices increase land values in forestry and extend rotations. Longer rotations increase the supply of timber from forests in the long-run because they move forests closer to maximum sustained yield. At some carbon price, however, it may be optimal to simply set-aside forests because they are more valuable as carbon reserves than as harvested timber forests. There are thus two caveats to the implications of rotation extensions for commercial timber supplies. First, in order to extend rotations, one must choose not to harvest forests. Therefore, in the short-term, the effect of the carbon subsidy is to *reduce* timber supply. A market model would respond to this short-term reduction in supply by increasing timber prices. Second, in the long-run, if carbon prices rise substantially, some forests may be converted to carbon reserves and never harvested. Creation of carbon reserves will serve to reduce timber supply in the long-run, which will tend to counteract the effect of moving towards the maximum sustainable yield. In summary, the impact of carbon prices on commercial forest supplies is complex. In fact, it is even more complex than this, since we have not yet considered the impact of carbon prices on management intensity. We now turn to this issue.

The Effect of Carbon Prices on Timber Management: As shown in figure 1, forest management activities beyond changes in rotation ages potentially provide 20 – 30% of forest carbon sequestration across carbon prices ranging from \$10 - \$100/t C. The analysis above, however, focused solely on age class management. Other types of forest management activities include replanting more intensively, thinning, fertilizing, using herbicide treatments, and other actions. Incorporating management intensity adjustments into CGE models will impose additional demands on CGE modelers. In order to inform researchers seeking to capture this dimension of the intensification response to higher carbon prices, this section describes how management intensity is captured in the global timber supply model of Sohngen and Mendelsohn (2007).⁵

Sohngen and Mendelsohn (2007) handle management actions via the stocking density parameter, δ . Instead of treating it as exogenous, as was done above, here we assume that it is a function of management inputs. Stocking density is written as:

$$(6) \quad \delta = 0.8*(1 + C)^n .$$

In equation (6), C is the planting cost, 0.8 is the value of δ without management inputs (the un-managed natural yield), and "n" is an elasticity parameter of stocking density with respect to management input. In the Southern upland hardwood example above, "n" is assumed to be 0.09. If planting costs are \$56 (as assumed above), then $\delta = 0.8*(1 + 56)^{0.09} = 1.15$.

One can see the influence of management inputs (C, or \$/hectare) and the parameter "n" in Figure 5. Increases in management inputs increase the stocking density

⁵ Obviously there are other options for incorporating management intensity decisions into such models.

of forestland, although the marginal cost of boosting yields is increasing in C (Figure 5). We expect the shape of this marginal cost function to vary by species of tree. Fast-growing plantations, for example, are often more responsive to adjustments in management, and thus may be assigned a higher value for " n ". In Figure 5, we plotted three different yield functions for three levels for " n " used in the global timber model. The highest level, $n = 0.13$, suggests relatively large potential to increase management. Species that are less responsive to management inputs are assigned a lower level for " n ", with the least responsive species being given the value of 0.05.

For simplicity, equations (4) and (5) were not solved for C when they were introduced above, however, since δ is a function of C , equations (4) or (5) can be optimized for C at the same time they are optimized for rotation age. The \$56/hectare management input is the optimal management input under baseline timber prices with no carbon values. In the baseline, with this management effort, the net present value of the stand is \$642/hectare. If management inputs are increased by \$10/hectare to \$66/hectare, the present value of the stand falls to \$641/hectare. The additional costs increased output at harvest time, but the value of these improvements were less than \$10/hectare.

While it would not be profitable to expand management inputs beyond \$56/hectare in the baseline, management inputs would increase if a price is placed on carbon sequestration. With a carbon price of \$100/t C , the value of the stand is \$2702/hectare under a \$56/hectare management level. However, under the \$100/t C carbon price, the optimal management input is \$264/hectare, and the value of the stand is \$2887. Biomass in 48 year old stands initially will be 112 m³/hectare, but at \$100/t C , with management inputs of \$264/hectare, biomass in a 48 year old stand will be 128

m³/hectare, or an increase of 15%. Thus carbon sequestration subsidies can have an important impact on the intensity with which forests are managed, thereby inducing an increase in forest product output.

3. EXAMPLES FROM A GLOBAL TIMBER MODEL

This section illustrates the single-stand responses discussed above by using the results from the global timber model analysis conducted by Sohngen and Mendelsohn (2007). That study developed a baseline case, and then explored a set of carbon policy scenarios. One of the scenarios assumes relatively high damages from climate change. In their "high damage" scenario, carbon prices start at \$22/t C in 2005, and rise to \$187/t C by 2105. This path of carbon prices reflects increasingly stringent emissions restrictions. Because the model is global in scope, carbon prices influence forests and timber prices globally. In this section, we focus on specific management activities in a single region of the model. However, the results are cast in a global context.

Forest Carbon Response to Adjustments in Age Classes: Carbon sequestration is a function of the inventory of forests, which in turn is a function of the age class distribution. One of the most important responses landowners may make to carbon incentives is to adjust the rotation age, particularly in the short-run when the area of forests cannot be expanded. (Even if it could be expanded, new plantings would not provide large quantities of carbon services for a number of years.) Extending rotations would not appear to provide substantial carbon when considered on an incremental, per hectare basis, but widespread adjustment in rotation ages across all commercially harvested hectares in response to carbon incentives, can have important effects on the

total stock of carbon stored in forests for an entire region. Furthermore, because this is just a marginal adjustment, it can be undertaken immediately by forest managers.

Therefore, as we have seen in Figure 1, changes in the age structure of the forest could well account for the majority of the carbon sequestered in temperate forests in the near term (e.g., 20 years), particularly at low carbon prices.

To examine the importance of accounting for changes in rotation ages on carbon sequestration, consider the implications of carbon incentives on upland hardwood stands of the Southern United States. Forests of the same type introduced in the previous section are included in the global timber model of Sohngen and Mendelsohn (2007). That model takes a dynamic optimization approach to managing stands by maximizing the net present value of consumer and producer surplus in timber markets. In the baseline case, the model ignores incentives for carbon sequestration. In the carbon policy scenario, carbon sequestration is modeled by renting carbon in the standing stock of forests. The model rents carbon along the high damage scenario price path introduced above.

Under the baseline, upland hardwood stands in the Southern U.S. region of the model are "optimally" harvested in 50 year rotations. Panel A of Figure 6 shows that the initial stock of Southern upland hardwood stands have a maximum age of 60 years (note that we are only considering the timber forests in this example, not less intensively managed forests that would already be at older rotation ages). In the baseline simulation, these 60 year old stands are harvested in the first time period, and the stock is then converted to a 50 year old stock in years 20 and 30 (Panels B and C, Figure 6) in order to maximize net present value of the forestry activity on this land.

The carbon prices for the high damage scenario induce landowners with upland hardwood stands in the U.S. South to hold timber longer than in the baseline (Figure 6). Stands are held at 60 years of age through the 30th year of the simulation, in the presence of carbon prices, whereas they decline to 50 years of age in the baseline. The total area of forestland also increases by 2.1 million hectares by year 30 in the carbon policy scenario, but the increase in forestland does not result primarily from additional planting of forests. Indeed, in year 20 of the simulation, there is *less* stock in the initial age class under the carbon policy scenario than in the baseline. The increase in forest land area is due to the fact that the stands that would have been harvested at 50 years in the baseline are now maintained in 60 year rotations in years 20 and 30, thus giving rise to an increase in total forestland compared to baseline.

Figure 7 shows that increasing rotation ages by 10 years (from 50 to 60 years) increases total carbon sequestered in natural southern pine forests over the 30 year time-frame (assuming that the first decade starts in 2005). Total storage increases by 436 million t C by the 30th year of the simulation. This increase is attributed to an increase in the rotation age, although the total forestland area has also increased by 2.1 million hectares. Expanding rotations means that each of these additional hectares adds carbon storage of 79 t C per hectare, far exceeding the average storage of carbon of around 49 t C per hectare in these types of stands. This example illustrates why adjusting rotation length is such a powerful vehicle for increasing forest carbon in the near term. Indeed, as the intertemporal optimization model shows, in the first 30 years, it is far more efficient than simply planting new stands and waiting for them to grow.

These results illustrate the importance of accounting for management of timberland when measuring adjustments in carbon sequestration triggered by carbon policy. Most computable general equilibrium (CGE) models that model forestry do not optimize age class management over time. Therefore, they are unable to account for the many adjustments that landowners can make to carbon sequestration incentives, as captured in a dynamic model like Sohngen and Mendelsohn (2003, 2007). One exception is the chapter⁶ by Sands and Kim in this volume, wherein the CGE model endogenizes optimal rotation length via incorporation of a steady-state formula. This will be discussed further below.

Of course, the results shown here also suggest the importance of accounting for age class distributions – particularly if one wishes to track the path of adjustment from one steady state to another. This is necessarily ignored in any formulation that deals only with steady state outcomes.

Forest Product Response: Changing rotation ages will alter the flow of products from the forest sector. Changes in timber supply, in turn, will affect prices, and hence the efficiency of the carbon actions themselves. It is thus useful to consider the response of timber markets to carbon incentives. This is particularly important in the case where some regions do not participate in global policy (e.g., they have a \$0/t C price). The extent of leakage to non-participating regions now depends on what happens to global forest product prices. And these price impacts in turn depend on supply changes from the regions facing carbon prices. As seen above, product supply will depend on total hectares forested, rotation ages and management intensity.

⁶ GTAP Working Paper No. 45

Forest product responses are likely to be quite large when carbon policies are introduced. Figure 8 shows timber prices in the baseline and the high damage carbon policy described above. In the carbon policy simulation, timber prices initially rise around 1% compared to the baseline as global supplies are reduced. By 2045 they fall by around 3% relative to the baseline. The carbon policy induces landowners to hold timber initially, and thus capture carbon rents. In the longer run, additional land supply, higher rotation ages, higher management, etc., increase the total timber supply, causing prices to fall.

One can see the effects of the carbon policy more directly by analyzing the effects on the Southern hardwood type described above. In year 20 of the baseline in the example above, there are 4.9 million hectares of land at age 50. When age = 50 years (the baseline rotation age in the model), the yield of merchantable timber would be 72 m³/ha. All of these hectares would be harvested, so over the 4.9 million hectares harvested, timber supply that decade (data is provided in 10-year age classes) would be 351 million m³/decade, or 35.1 million m³/yr over the decade. In contrast, under the climate policy, there are 5.0 million hectares in age class 60 in the 20th year. The yield of timber for 60 year old stands is 96.7 m³/ha, so total timber harvests are 486 million m³/decade, or 48.6 million m³/yr. Raising rotation ages increases total timber harvests by moving forests towards their maximum sustainable yield.

This quantity, however, applies only to the 20th year of that simulation. A dynamic forestry model will adjust harvest ages in response to present and future timber prices, carbon prices, and opportunity costs. In order to extend rotation ages by the 20th year, some timber must be withheld from markets in initial periods. This is illustrated in

figure 9, where harvests in this forest type are presented for the baseline and for the high damage carbon policy scenario. Initially, harvests are lower in carbon policy scenario than in the baseline because some timber is with-held from markets in order to extend rotation ages with the carbon price. Over the longer run, rotations are extended and forest areas expand, so harvests increase above the baseline scenario.

Response of Forest Management: Management inputs are determined endogenously with timber prices and carbon prices in the global forestry model. As carbon prices rise, timber management inputs also increase in order to increase the carbon density on forested sites. These enhancements to management, as shown in Figure 1, can provide substantial benefits in terms of carbon sequestration.

For the Southern hardwood forests discussed above, management inputs in the baseline are approximately \$56/hectare over the first 50 years of the simulation. With the high damage carbon policy, however, these inputs triple in size in this type, to around \$170/hectare. Southern upland hardwoods currently experience a relatively low level of management intensity overall, and these results indicate that there are efficient opportunities to increase management and enhance carbon sequestration. This is a relatively strong response. Species that are more intensively managed in the baseline, such as Southern pine plantations, experience only 30% increases in management inputs with the high damage carbon policy. This makes sense given that more intensively managed species will already contain fairly high carbon stocks, and thus they are likely to have limited opportunities to improve stocking conditions with positive carbon prices.

Combined Effects of Rotation and Management Changes: One way to gauge the potential effects of both extensions in rotation ages and increases in the management

inputs is to assess the change in carbon on each hectare of land. These two activities have their most pronounced and important effects globally in the temperate zones, so we focus on that region for our discussion. Table 1 shows the ratio of average carbon intensity (t C/hectare) for temperate forests for different constant carbon prices in the shorter term (by 2025) and in the longer term (2055), *relative to baseline*. Thus an entry of 1.00 in table 1 indicates no change under the carbon policy regime compared to the baseline. These results suggest that carbon intensity rises on average as carbon prices rises. Indeed, within the next 20 years or so, the global timber model suggests that, for a price of \$100/t C, carbon intensity in forests could be increased by up to 14% in China and 16% in Japan. And, if the same price were maintained in the future, over the next 50 years, carbon intensity could be increased by up to 47% in Europe and 45% in China, with more modest rises in other regions. Of course these are the most responsive regions. Oceania and Russia show very small responses on this intensive margin. 8667820900

4. IMPLICATIONS FOR CGE MODELING

Accounting for this potentially complex dynamic behavior in forestry can be difficult in static or recursive dynamic CGE models. Fully intertemporal optimization routines require substantial computing resources, which is particularly burdensome if the interest of the modeler lies in the general equilibrium effects, since all sectors must then engage in forward-looking investment behavior. For comparative static analyses, the main issue becomes: Comparative static over what period of time? If the analysts are primarily interested in long-run phenomenon, then they can potentially utilize steady-state, static adjustments using a simple Faustmann model of optimal rotation. If, on the

other hand, analysts are interested in “near-term” phenomena – say the first 30 years -- it will be fairly difficult to capture the dynamic effects described above. (We explore some potential alternatives below.)

Assuming analysts are interested primarily in long-run effects, one way to handle the rotation age adjustments that occur in forestry is to simply assume a uniform age distribution and calculate the steady state effects based on comparing steady states (one for the baseline and one for the carbon policy). With a uniform age distribution, the output of interest is the average annual flow of forest product. Calculating average annual flow is relatively straightforward. The modeler needs to obtain total hectares in the forest type (summed over all age classes) and the optimal rotation age. In the example above, there are 34 million hectares initially. If optimal rotations are 50 years, then 0.68 million hectares will be cut each year on average ($34/50$). The yield at year 50 is $72.4 \text{ m}^3/\text{ha}$, so the average timber supply is $49.2 \text{ million m}^3/\text{yr}$.

When carbon policies are introduced, two things must be considered: the change in land area and the change in rotation ages. Changes in land area are best handled within the CGE framework through a land market whereby forestry competes with other land uses and land rents adjust to equate supply and demand for land of a particular type/location. With carbon incentives, the value of forestland increases because the standing stock of carbon has value. Since the forest land could otherwise be used for other purposes, adding timberland has an opportunity cost. If carbon incentives draw more land into forestry, modelers will need to adjust the average flow of timber from the landscape. The value of land in forests when both timber and carbon are valued can be estimated with equation (5) above. That equation provides the value of land in forests

under the carbon policy, and can be used as the relevant land value to be compared to alternatives, such as agriculture.

Equation (5) can also be used to determine the rotation age of forests under a carbon policy. As noted in the previous section, an increase in rotation ages increases the long run supply of forest product. Under the carbon policy scenario described above, the new optimal rotation age is 60 years. For the same 34 million hectares, average annual flow under the 60 year rotation becomes 54.7 million m^3/yr , which is larger than under the baseline rotation of 50 years. This calculation of a change in timber production assumes that the forest area has remained the same even under the climate policy. Obviously, total forest area may change if the value of forests relative to other uses rises due to the carbon incentives. In the example from the timber model described in the previous section, over 100 years, forest area in Southern upland hardwoods increases to 46 million hectares. With 46 million hectares, the average annual flow of timber would be 74.1 million m^3/yr .

Finally, there is the question of stocking density. This is one area where CGE models are relatively well equipped to deal with the forestry issue. The application of additional inputs to a given unit of land to increase output per unit of land is analogous to non-land/land substitution in the agriculture sectors of these models. Authors can either employ equation (6) above, or they can calibrate the elasticity of substitution in a CES production function involving land and non-land inputs to capture the same intensification possibilities.

In summary, CGE modelers conducting static or recursively dynamic analyses of long run policies must consider multiple factors if they hope to capture the key features of forestry in climate mitigation analysis:

- (A) Calculation of land values using equations (4) in the baseline or (5) under the carbon scenario.
- (B) Calculation of optimal rotation ages using equations (4) in the baseline or (5) under the carbon scenario.
- (C) Calculation of the optimal use of non-land inputs.
- (D) Calculation of land areas using values obtained through (A) and comparing those values to relevant values for other activities, such as agriculture.
- (E) Calculation of average annual flow of timber.

Of course, by altering the supply of forest products, the change in annual timber flow will have an impact on forest product prices as well. The extent of this change will depend on the derived demand elasticity for output from the forestry sector. In a general equilibrium model, this depends on the consumer demand elasticity for products like furniture, housing, paper, etc., as well as the ability of the sectors producing these final goods to substitute between other inputs and forest products.

As shown above, near term analyses create additional complexities for incorporating forestry in CGE models. Carbon sequestration and timber production tend to be complements in the long-run (expanding forest area, increasing forest carbon through management, and increasing rotations all increase production); however, in the short-run they can be substitutes. In regions where increasing the rotation age is the efficient response to carbon prices, increases in rotation ages can only be accomplished

by with-holding timber from markets. While increasing management in forests or increasing the area of forests both lead to additional carbon sequestered, it takes many years for these activities to have an impact on the overall carbon balance. Owing to the forestry growth function, much of the carbon potential in the short-run in temperate regions is likely to result from changes in rotation ages and some changes in management, with subsequent implications for output flows.

The short run production impacts of carbon policy are potentially substantial, and they increase as the carbon price offered to the forestry sector rises. Table 2 presents estimates of carbon sequestration and timber production from the global timber model in Sohngen and Mendelsohn (2007). Carbon sequestration and reduced emissions from management plus changes in rotation ages and from afforestation plus reduced deforestation are shown for different regions of the world and different carbon prices. Changes in timber production are also shown. All values are annual equivalent amounts over the first 20 years period.

In the short-run (here defined as the first 20 years of the policy) output declines as carbon sequestration occurs. These production impacts get steadily larger as carbon prices rise. Globally, carbon sequestration and timber output are substitutes in the short-run. Most carbon sequestered in temperate forests in the short run arises from changes in management and changes in rotation ages. This contrasts with the tropics where most carbon sequestration results from reductions in deforestation. Reductions in deforestation also have strong production effects in the tropics because forestry outputs that are by-products of the deforestation process are reduced. On average, in the temperate zone, each 1 million t C increase in carbon sequestered reduces timber output by 0.3% on

average. In tropical zones, each 1 million t C increase in avoided emissions from deforestation reduces timber output by 1.4% in the short-run.

5. EVALUATION OF CURRENT APPROACHES TO FORESTRY MODELING IN GLOBAL CGE ANALYSIS

While a number of partial equilibrium analyses of forestry and land use have been developed to date (e.g., Murray et al., 2005; Sohngen and Mendelsohn, 2003; Sathaye et al., 2006), few CGE models have fully incorporated land use, and none of these have come to grips with forestry in a comprehensive fashion. The FARM model described in Darwin et al. (1995, 1996) and Darwin (1999) is one example where land use has been incorporated into CGE modeling, but the primary emphasis in that work was on agricultural uses. One of the most important issues in modeling forestry is the disaggregation of forest stocks by age class. This has a big impact on the potential for carbon sequestration. The efficiency of forestry actions for sequestration is determined by the rental value of carbon, but the value of carbon depends on future prices for timber and carbon, as well as land rents for forestry. It has proven difficult thus far to account for these dynamic stock effects when modeling forestry in a CGE context.

Despite the difficulty of incorporating forestry into CGE models, this book highlights several important efforts in that direction. Hertel et al. (2007), for example, incorporate reduced form marginal abatement cost functions derived from a partial equilibrium model into the CGE framework. Their model is static, and considers only near term abatement options. However, it is attractive in that it decomposes the total sequestration response into the intensive (management and aging) and extensive margins,

as identified in Table 2. The problem with this comparative static analysis is that the marginal abatement cost curves are dependent on the time horizon assumed. A longer time horizon will generate flatter cost curves and hence different results. Operationalizing this approach in a recursive dynamic CGE model would thus require that the authors shift the abatement cost schedules over time.

Ahammed and Mi (2007) take a different approach using the GTEM model (Global Trade and Environment Model). GTEM is a recursive dynamic CGE model. Within the model, they explicitly account for the age class distribution of forests, and shift forests from one age class to another over time. Their model does not "optimize" harvest ages as would occur in a fully intertemporal model, but instead, they harvest a fixed proportion of all age classes above 10 years old. Forestry competes with agriculture uses of land via a constant elasticity of transformation (CET) function. And, if land shifts into forests from agriculture, it shifts in during the youngest age class (i.e. new plantings). If land shifts out of forestry, it shifts out proportionally from all age classes in the region. So these authors begin to capture the differential carbon intensity of different age stands, but they fail to capture the potential for changes in the optimal rotation length.

Golub et al. (2007) also utilize a recursive dynamic CGE model, however, they take a different approach from Ahammed and Mi (2007). They focus on how the demand for agricultural and forest products influences the allocation of land across different uses, and how these demands affect decisions to access new land for agriculture and commercial forestry. To capture the dynamics of forestry extraction, they link their model with the dynamic optimization model of the forestry sector presented in Sohngen and Mendelsohn (2007). The forestry model provides results on the value of extracting

timber resources over the period 2005 – 2025. Price changes for the forestry sector in the recursive CGE model are then guided by prices from the forestry model. With technical change in forestry processing sectors allowed to adjust to hit these price targets, the CGE model is solved to determine forestry land use and output. Golub *et al.* (2007) focus on changes in forest lands in the no-carbon price baseline, but their approach could be adapted to the examination of a carbon policy by targeting the associated changes in land rents over time. The strength of this approach is that it draws directly on the intertemporal partial equilibrium model of Sohngen and Mendelsohn (2006). But this may also be considered the main weakness, since the model does not incorporate the forestry sector's responses changing market conditions directly.

Sands and Kim (2007) utilize a recursive CGE model to analyze land use. They account for the age class distribution of forests, but because of the difficulties associated with solving the dynamic optimization problem within their modeling framework, they simplify the problem by assuming that forests are harvested from fixed age class over time. They utilize the underlying yield functions for forest production to determine steady state rotation ages for forests with and without carbon incentives. They also utilize these yield functions to determine the land rents for forestry. Land is converted from one use to another depending on relative land rents in much the same way as the other CGE studies, but using a supply function in which the elasticity of supply depends on the inherent heterogeneity of yields within a watershed. This permits the authors to capture the important impact of carbon policies on optimal rotations. However, in doing so, it ignores the near term market impacts of climate policy (up to 50 years) which may be quite different from the steady state adjustments, as has been shown above.

The studies in this book suggest that the range of approaches for incorporating forestry stocks into CGE models is fairly wide. Ahammad and Mi (2007) and Sands and Kim (2007) take an approach that accounts for forestry stocks directly. They have different methods for harvesting stocks, for example, Ahammad and Mi (2007) harvest proportionally from different age classes (similar to approaches used in static equilibrium modeling in the U.S., see Adams and Haynes, 1980 and Sohngen and Sedjo, 1998). Alternatively, Sands and Kim (2007) harvest from a fixed age class, where this age class is determined by static net present value calculations. Hertel et al. (2007) take an indirect approach and incorporate marginal abatement cost functions from a forestry sector model directly into their static CGE model to determine near-term abatement strategies. Golub et al. (2007) focus on land allocation and access questions. They calibrate their model based on forestry prices, deforestation rates and sensitivity of management input to changes in forest land rents found in Sohngen and Mendelsohn (2007). It is not clear that any one approach is "better" than the others. Rather, each set of authors has focused on different aspects of the problem. Future research should be aimed at merging some of these different approaches in order to permit CGE modelers to better capture the role of forestry in climate change scenarios.

6. CONCLUSION

This chapter evaluates the optimal response of the forestry sector to changing economic circumstances over time – particularly in the context of a carbon policy in which the forestry sector receives rental payments for carbon sequestered. These responses are then considered in light of current approaches to modeling the forestry

sector in CGE models with an eye to evaluating how well these capture the relevant range of economic responses.

The chapter begins with a description of the forestry management problem and shows the importance of age-class management in forestry. Age class management has strong implications for carbon sequestration in forestry, allowing foresters to rapidly obtain additional carbon at relatively low carbon prices. In contrast, afforestation efforts may take many years to begin and because the forests take many years to grow, the bulk of the carbon associated with afforestation occurs in future periods. So capturing the broad outlines of this option is an important, but complex task for CGE models; some compromises are inevitable.

Based on the analysis in this chapter, we find that, in longer-run studies (e.g., 50 years or more), modelers may focus on steady state adjustments in forestry and calculate timber flows and carbon balances based on the rotation ages implied by these steady state formulae. However, for those interested in shorter-run effects (e.g., 20 years), the age class and production effects may be quite different than those implied by steady-state relationships. Over this shorter time frame, we find that forest output is likely to be sacrificed in order to boost forest carbon stocks. This is the opposite response to that which is observed in the long run, over which higher carbon prices also boost steady-state forest output.

In addition to age class adjustments, the paper also examines changes in other types of management activities which influence the stocking density of the forest -- and through that density also carbon stock and output flows. We estimate that other management activities can provide around 20% of total carbon sequestered in the

temperate zone for a given price, and thus are important to consider. The paper presents the methods used to incorporate management in the global timber model of Sohngen and Mendelsohn (2007). CGE modelers could either incorporate this stocking density parameter directly, or calibrate the elasticity of substitution between other inputs and forest land to give an equivalent response in output per hectare of forest land (see Hertel et al. and Golub et al., this volume).

In summary, the forest sector poses one of the most difficult challenges facing CGE modelers interested in capturing the full range of potential responses to climate change policy. Because the forests grow relatively slowly, the forest capital stock can only be adjusted over a period of decades – as opposed to months or years in the case of other sectors (e.g., wearing apparel, autos or wheat). This very slow adjustment means that it is important to account for the age profile of the forest stock, which also adjusts only slowly to a new policy environment. Keeping track of this age profile, and managing the forest optimally, requires use of a long-run, forward-looking model. Yet such models remain relatively rare in the large scale, global CGE modeling literature due to problems of computation and sheer manageability. Until these models become routine, authors will need to continue to make various compromises in their research. Which compromise is least damaging will depend on the objective of the research being undertaken.

7. REFERENCES

- Ahammed H. and R. Mi. 2007. Modeling land use changes and greenhouse gas emissions in GTEM. Chapter X in *Economic Analysis of Land Use in Global Climate Change Policy*. Edited by T. Hertel, S. Rose, and R. Tol. Routledge.
- Adams, D.M., R.J. Alig, B.A. McCarl, J.M. Callaway, and S.M. Winnett. 1999. "Minimum Cost Strategies for Sequestering Carbon in Forests." *Land Economics*. 75:360-74.
- Darwin, R., Tsigas, M., Lewandrowski, J., Ranases, A. 1995. "World agriculture and climate change: economic adaptations." Agricultural Economic Report No. 703, US Department of Agriculture, Economic Research Service, U.S. Government Printing Office, Washington, DC. <http://www.ers.usda.gov/publications/aer703/aer703.pdf>.
- Darwin, R., Tsigas, M., Lewandrowski, J., Ranases, A., 1996. "Land use and cover in ecological economics." *Ecological Economics*. 17: 157–181.
- Darwin, R.F., 1999. "A FARMer's view of the Richardian approach to measuring agricultural effects of climatic change." *Climate Change* 41(3-4): 371–411.
- Golub, A., T. Hertel, and B. Sohngen. 2007. "Modeling the Long Run Supply and Demand for Land." Chapter X in *Economic Analysis of Land Use in Global Climate Change Policy*. Edited by T. Hertel, S. Rose, and R. Tol. Routledge.
- Hertel, T., H-L Lee, S. Rose, and B. Sohngen. 2007. "Analysis of Global Land Use and the Potential for Greenhouse Gas Mitigation in Agriculture and Forestry. Chapter X in *Economic Analysis of Land Use in Global Climate Change Policy*. Edited by T. Hertel, S. Rose, and R. Tol. Routledge.
- Murray, B.C., B.L. Sohngen, A.J. Sommer, B.M. Depro, K.M. Jones, B.A. McCarl, D. Gillig, B. DeAngelo, and K. Andrasko. 2005. EPA-R-05-006. "Greenhouse Gas Mitigation Potential in U.S. Forestry and Agriculture." Washington, D.C: U.S. Environmental Protection Agency, Office of Atmospheric Programs.
- Richards, K.R., and C. Stokes. 2004. A Review of Forest Carbon Sequestration Cost Studies: A Dozen Years of Research. *Climatic Change*. 63: 1–48.
- Sands R. and M-K Kim. 2007. Modeling the Competition for Land: Methods and Application to Climate Policy. Chapter X in *Economic Analysis of Land Use in Global Climate Change Policy*. Edited by T. Hertel, S. Rose, and R. Tol. Routledge.
- Sathaye, J., W. Makundi, L. Dale, and P. Chan, and K. Andrasko. 2007. GHG Mitigation Potential, Costs and Benefits in Global Forests: A Dynamic Partial Equilibrium Approach. *Energy Journal*. 27: 127-162.

Sohngen, B. and R. Mendelsohn. 2003. "An Optimal Control Model of Forest Carbon Sequestration." *American Journal of Agricultural Economics*. 85(2): 448-457.

Sohngen, B. and R. Mendelsohn. 2007. "A Sensitivity Analysis of Carbon Sequestration." Chapter 19 in *Human-Induced Climate Change: An Interdisciplinary Assessment*. Edited by M. Schlesinger, et al. Cambridge: Cambridge University Press.

Stavins, R. 1999. "The Costs of Carbon Sequestration: A Revealed Preference Approach." *American Economic Review*. 89: 994-1009.

Weyant, J.P, F.C. de la Chesnaye, G.J. Blanford. 2006. "Overview of EMF-21: Multigas Mitigation and Climate Policy." *Energy Journal*. 27: 1-32.


Table 1. Carbon intensity in temperate zone forests (t C/hectare), expressed relative to the baseline carbon intensity (1.0 = no change).

	By 2025			By 2055		
	\$10/t	\$50/t	\$100/t	\$10/t	\$50/t	\$100/t
	Proportional Change in t C/ha					
US	1.00	1.06	1.04	1.03	1.11	1.20
Canada	0.99	1.08	1.10	1.06	1.20	1.25
Europe	1.00	1.03	1.05	1.01	1.19	1.47
Russia	1.00	1.00	0.99	1.00	1.05	1.10
China	1.03	1.15	1.14	1.04	1.40	1.45
Oceania	1.00	1.00	1.00	1.00	1.00	1.01
Japan	0.98	0.98	1.16	1.01	1.03	1.14

Table 2. Annual equivalent carbon gains (losses) and production impacts over 20 years at three different carbon prices,

	Carbon due to Changes in Management and Rotation Ages.			Carbon Due to Afforestation/Reduced Deforestation			Change in Timber Production Relative to Baseline		
	\$10/t	\$50/t	\$100/t	\$10/t	\$50/t	\$100/t	\$10/t	\$50/t	\$100/t
	Million t Ceq./yr			Million t Ceq./yr			Million m ³ /yr		
Temperate									
US	6.8	73.5	102.7	3.5	17.8	43.1	-0.4%	-2.3%	-0.9%
Canada	-2.0	15.2	33.5	4.4	13.4	25.6	0.6%	-4.8%	-6.2%
Europe	7.0	52.5	102.4	-0.4	2.1	2.6	-1.2%	-1.4%	-4.6%
Russia	22.7	74.0	111.3	-1.5	-0.9	2.3	-2.9%	-15.7%	-29.1%
China	14.9	73.9	98.5	0.6	4.2	12.8	-4.3%	-24.5%	-33.7%
Oceania	0.4	2.5	5.6	0.3	2.2	4.9	-2.1%	-2.7%	-4.4%
Japan	-0.6	1.6	15.5	0.5	3.7	4.0	-1.1%	1.4%	-9.9%
Tropical/Other									
S. America	0.4	1.4	2.6	67.8	321.2	538.0	-5.1%	-22.6%	-36.1%
Central Am.	0.3	1.0	1.7	38.7	100.6	122.7	-14.9%	-35.2%	-46.0%
India	0.2	1.7	2.6	6.9	34.9	45.9	-1.5%	-11.0%	-11.5%
SE Asia	47.5	86.0	88.2	68.7	316.2	419.0	-12.2%	-39.3%	-45.8%
Central Asia	2.3	12.4	20.9	0.3	1.3	2.6	-2.7%	-12.9%	-20.6%
Africa	0.4	2.3	3.4	96.1	408.5	505.0	-9.2%	-38.9%	-46.7%
Total	100.3	398.1	588.9	285.8	1225.1	1728.4	-2.5%	-10.7%	-15.5%

Figure 1. Proportion of carbon sequestration in temperate and tropical forests resulting from forestry actions for several different carbon prices over a 20 and 100 year period. All carbon is measured in annual equivalent amounts and the proportion of total storage is shown.



Note: All carbon gains are additional to the baseline. *Set-aside %* is the proportion of carbon due to setting aside forestland from timber production; *Market %* is the proportion of carbon stored in marketed productions; *Mgmt %* is the proportion due to improved management of timber stocks; *Age %* is the proportion due to increasing the rotation age of harvests; and *Aff/Def %* is the proportion due to land use change arising either from afforestation or reductions in deforestation.

Figure 2. Biomass and Merchantable timber yield functions for upland hardwoods in the U.S. south.




Figure 3. Growth rate and average annual flow of timber from upland hardwood type in the Southern U.S.




Figure 4. Present value of upland hardwoods in U.S. South under baseline (no carbon incentive) and three alternative carbon prices (derived from equation 5), assuming management inputs remain \$56/hectare.




Figure 5. Relationship between \$/hectare management inputs, "n", and stocking density (delta, or δ).




Figure 6. Age class distribution of upland hardwood stands in the Southern U.S. in the baseline and climate mitigation scenario (high damage) from Sohngen and Mendelsohn (2006).




Figure 7. Aboveground carbon stored in natural southern pine stands in the United States in the baseline and the high damage carbon mitigation scenario.




Figure 8. Timber price index for baseline and high damage carbon policy scenario, derived from Sohngen and Mendelsohn (2007).




Figure 9. Annual timber harvests in southern upland hardwood stands in the United States in the baseline and the high damage carbon mitigation scenario. Estimates are taken from the midpoint of each decade, starting with 2010 to illustrate harvests during the 2005-2014 decade.

